Mapping of dendritic lesions in patients with herpes simplex keratitis using in vivo confocal microscopy

Hideaki Yokogawa
Akira Kobayashi
Natsuko Mori
Kazuhisa Sugiyama

Department of Ophthalmology,
Kanazawa University Graduate School
of Medical Science, Kanazawa, Japan

Purpose: To produce a two-dimensional reconstruction map of dendritic lesions in patients with herpes simplex keratitis (HSK) using in vivo confocal microscopy.

Methods: Four eyes of four patients (mean 65.8 years) with HSK presenting with a dendritic lesion were enrolled. Slit-lamp biomicroscopy and in vivo laser confocal microscopy were performed. Acquired confocal images at the level of the epithelium were arranged and mapped into subconfluent montages. Changes in the shape and degree of light reflection of abnormal cells and deposits around dendritic lesions as well as other corneal layers were qualitatively evaluated.

Results: Mapping of dendritic lesion was successful in all cases, and the subconfluent montages clearly showed the larger image of dendritic lesion. In all cases, the dendritic lesion consisted of hyperreflective irregular epithelial cells, and was surrounded by distorted and elongated epithelial cells. In three cases, hyperreflective deposits were noted at the midline of the lesion. The corneal stroma showed a hyperreflective honeycomb pattern. In two cases, inflammatory cells were observed at the level of endothelial cell layer.

Conclusion: Mapping of dendritic lesions in patients with HSK was successful in all patients using in vivo confocal microscopy. Cellular level observation of dendritic lesion at a relatively larger magnification may help understand the in vivo morphological change of HSK. Further study in more patients with HSK and nonherpetic dendritic lesion is needed to utilize confocal microscopy images in differential diagnosis and follow-up of the epithelial lesions with dendrite.

Keywords: herpetic epithelial keratitis, in vivo confocal microscopy, fluorescein slit-lamp photograph, hyperreflective irregular epithelial cells, dendritic lesion

Introduction

Herpes simplex keratitis (HSK) is still a major cause of corneal blindness in many countries.^{1,2} Clinical phenotypes of HSK include infectious epithelial keratitis, neurotrophic keratopathy, immune stromal keratitis, necrotizing stromal keratitis, and endotheliitis. Among these, the most common presentation of HSK is the dendritic lesion of infectious epithelial keratitis. Dendritic lesions may occur with a primary herpes simplex virus infection; however, they occur more commonly due to the reactivation of latent virus. Dendritic lesions are characterized by a branching, linear lesion with a terminal bulb and swollen epithelial borders, all of which are visualized well with slit-lamp biomicroscopy with fluorescein staining.

Recently, in vivo confocal microscopy has become an increasingly popular noninvasive method to facilitate the understanding of the corneal microstructure in the healthy as well as the diseased cornea.³ Prior studies have evaluated the corneal involvement of HSK with different types of confocal microscopes, each investigating

Correspondence: Akira Kobayashi
Department of Ophthalmology,
Kanazawa University Graduate School
of Medical Science, 13-1 Takara-machi,
Kanazawa, Ishikawa-ken 920-8641, Japan
Tel +81 76 265 2403
Fax +81 76 222 9660
Email kobaya@kenroku.kanazawa-u.ac.jp

the subbasal nerve plexus, the epithelium, the endothelium, or the inflammatory response.⁴⁻⁸ So far, only a single case report was published that described in vivo confocal microscopic features of the dendritic lesion.⁹ In the current study, to further elucidate the cellular level morphological change of dendritic lesion at a wider angle, we produced larger two-dimensional reconstruction confocal map of the dendritic lesion.

Materials and methods

This clinical study was approved by the Ethical Committee of Kanazawa University Graduate School of Medical Science and followed the tenets of the Helsinki Declaration. Four eyes of four patients (65.8±21.5 years of age) with HSK presenting with a dendritic lesion were evaluated in this study (Figure 1). The diagnosis of HSK was made according

to characteristic clinical features, including dendritic lesion with terminal bulb, history of recurrent HSK, and response to acyclovir treatment (Table 1). Only in Case 4 was herpes simplex virus detected by immunochromatography of an epithelial sample.

In vivo laser confocal microscopy

After the nature and possible consequences of confocal microscopic study (such as superficial punctate keratopathy) were explained to the patients, informed consent was obtained for confocal microscopic examination. For each patient, a fluorescein slit-lamp photograph demonstrating dendritic lesion was obtained. Each dendritic lesion was manually scanned at the level of the epithelium with a Heidelberg Retina Tomograph 2 Rostock Cornea Module (HRT2-RCM,

Figure 1 Fluorescein slit-lamp photographs of dendritic ulcers with terminal bulbs in Case 1 (A), Case 2 (B), Case 3 (C), and Case 4 (D).

Note: The area framed by the white rectangle corresponds to the confocal maps in Figure 2.

Table I Clinical data of four patients with herpes simplex keratitis

Case	Age (years)/sex	Eye	Systemic disease	Ocular disease history	Intraocular pressure (mmHg)	Virological test	Response to acyclovir treatment
1	61/F	R	None	Recurrent herpes simplex keratitis (from 10 years ago) Rhegmatogenesis retinal detachment surgery (3 years ago)	16	No data	Disappearance of dendritic lesion
2	88/F	R	None	Cataract surgery (3 months ago)	8	No data	Disappearance of dendritic lesion
3	38/M	L	Atopic skin	Laser in situ keratomileusis (3 years ago)	8	No data	Disappearance of dendritic lesion
4	76/F	R	HTN	Corneal endotheliitis (9 months ago)	20	Positive HSV by immunochromatography of epithelial sample	Disappearance of dendritic lesion

Abbreviations: F, female; HSV, herpes simplex virus; HTN, hypertension; M, male; R, right eye; L, left eye.

Heidelberg Engineering GmbH, Dossenheim, Germany) with a 60× water-immersion objective lens (Olympus, Tokyo, Japan) (numerical aperture, 0.9). The microscope utilizes a 670-nm red wavelength diode laser as the light source (area of observation, 400×400 μm²). The manufacturer quotes the transverse resolution and optical section thickness as 2 and 4 μm, respectively.

Two-dimensional reconstruction mapping

Acquired confocal images at the level of the epithelium were arranged and mapped into subconfluent montages with a mapping technique and computer software (Adobe Photoshop, Adobe Systems, San Jose, CA, USA) as previously described.^{10–11} Changes in the shape and degree of light reflection of abnormal cells and deposits around dendritic lesions as well as other corneal layers were qualitatively evaluated.

Results

A total of 530.5±188.0 *in vivo* confocal microscopic images (400×400 μm²) of all cell layers were obtained for each patient; of them, a mean of 8.5±2.5 images of the epithelial layer at the area of the dendritic lesion were used for subconfluent montage formation (Figure 2). The mean dimensions of the mapped areas were 1.0±0.2 mm (horizontal) and 0.8±0.2 mm (vertical).

Mapping of dendritic lesion was successful in all cases, and the subconfluent montages showed clearly the larger image of dendritic lesion. In all cases, the dendritic lesions consisted of hyperreflective irregular epithelial cells, and were surrounded by distorted and elongated epithelial cells. In three cases (75%, Cases 1, 2, and 4), hyperreflective deposits, which may represent necrotic abnormal cells, were

noted at the midline of the lesion. In Case 2, the central area of dendritic lesion seemed dark, indicating an epithelial defect. Putative dendritic inflammatory cells were sparsely noted.

At the level of deep epithelium to subbasal nerve plexus, numerous inflammatory cells including possible Langerhans cells were noted, and were not exclusively located at the dendritic lesion (Figure 3). At the level of corneal stroma, honeycomb-shaped hyperreflective keratocytes were seen in all cases, suggesting stromal edema (Figure 4A and B). At the level of endothelium, inflammatory cells were only seen in two cases (Cases 1 and 4) (Figure 4C and D).

Discussion

In the current study, we investigated the confocal microscopic characteristics of *in vivo* epithelial changes of dendritic lesion as well as other corneal layers using similar mapping technique. As a result, larger images of the dendritic lesion were successfully constructed. In all cases, the dendritic lesion consisted of hyperreflective irregular epithelial cells, and was surrounded by distorted and elongated epithelial cells. Based on previous histopathological examinations,^{12–17} these cells may represent herpes simplex virus-infected cells, as they were multinucleated giant cells and cells with intranuclear inclusions with koilocytic changes. Hyperreflective deposits were frequently observed (75%) within the dendritic lesion, especially at the branching point. We surmise that they might be infected necrotic abnormal cells. A histopathological study by Maudgal et al¹² showed that dendritic ulcers had necrotic areas with fragmented nuclei and ghost cells debris. The usefulness of a mapping technique using numerous *in vivo* confocal microscopy images was previously reported for visualizing the subbasal nerve plexus,^{18,19} Kobayashi-structures (K-structures, sub-Bowman's fibrous structures

Figure 2 Subconfluent montages of in vivo confocal microscopic images at the area of dendritic lesion in Case 1 (**A**), Case 2 (**B**), Case 3 (**C**), and Case 4 (**D**). (Bar =200 μ m). **Notes:** (**A**) The swollen epithelial borders consist of relatively hyperreflective irregular epithelial cells (white arrows). Hyperreflective deposits (black arrow heads) were noted at the midline of the lesion. Epithelial cells that surrounded the dendritic lesion were distorted. Filament-like lesion (white arrow heads) was also noted. (**B**) The central dark areas (asterisks) without cells might represent an epithelial defect area. The swollen epithelial borders consist of relatively hyperreflective irregular cells (white arrows). (**C**) Hyperreflective materials (black arrow heads) may represent necrotic abnormal cells. Relatively hyperreflective irregular cells (white arrows) were also noted. A putative dendritic inflammatory cell (black arrow) was noted between epithelial cells. (**D**) The swollen epithelial borders consists of relatively hyperreflective irregular cells (black arrows). The dendritic lesion was surrounded by distorted epithelial cells (white arrows). Putative dendritic inflammatory cells (black arrowheads) were also noted.

correlated with anterior mosaic),¹⁰ and endothelial cells in cases of cytomegalovirus corneal endotheliitis.¹¹

In addition to the epithelial findings, we found that numerous Langerhans cells were characteristically observed at the level of deep epithelium to subbasal nerve plexus. It should be noted that the Langerhans cells were not exclusively located at the dendritic ulcer lesion. At the level of corneal stroma, honeycomb-shaped hyperreflective keratocytes were seen in all cases. Also, at the level of endothelium, inflammatory cells were only seen in two cases. We surmise that these changes were nonspecific inflammatory response of the cornea caused by HSK.

To date, several in vivo confocal microscopic studies in both epithelial and nonepithelial type HSK have investigated subbasal nerve plexus abnormalities (density, branch, etc)

and epithelial/stromal cell abnormalities (density, size, etc) with different types of confocal microscopes.⁴⁻⁸ For example, Rosenberg et al⁴ reported that dendritic structures, possibly Langerhans cells, were characteristically observed, at the level of the basal epithelium in resolved HSK cases, using a tandem scanning confocal microscope. Their findings were consistent with our observation, in which there were numerous Langerhans cells at the level of deep epithelium to subbasal nerve plexus (Figure 3). In terms of corneal nerve changes due to HSK, Nagasato et al⁵ reported decreases in subbasal nerve density parameters in epithelial and stromal-type HSK using HRT2-RCM, the identical device used in this study. Additionally, Hamrah et al^{6,7} reported that HSK-induced superficial epithelial changes included increases in epithelial cell size and decreases in epithelial cell density

Figure 3 In vivo confocal microscopic images at the level of deep epithelium to subbasal nerve plexus in Case 1 (A), Case 2 (B), Case 3 (C), and Case 4 (D).

Notes: Numerous Langerhans cells as dendritic inflammatory cells (arrows) were noted, and were not exclusively located at the dendritic lesion. (Bar = 100 μ m).

with Confoscan 4, and they concluded that these epithelial cell changes correlated with corneal innervation. On the other hand, Mocan et al⁸ observed inflammatory cell infiltration and keratic precipitates in nonepithelial-type HSK with Confoscan 3.0. Interestingly, Martone et al⁹ have already published a single case report of in vivo confocal microscopic analysis of dendritic lesion in a patient with bilateral HSK. They used a same confocal device with ours (HRT2-RCM), and obtained quite similar findings with our study: a distortion of the superficial and basal epithelium, the presence of irregular hyperreflective structures, and dendritic particles near the epithelial cells.

One limitation of this study is that the sample size was quite small (n=4). Also, nonexistence of the nonherpetic controls

makes the study just an observational one. Nonetheless, we believe that further comparative studies of nonherpetic dendrites (pseudodendrite) frequently seen in early-stage *Acanthamoeba* keratitis would make the current study more clinically significant. Additionally, it might be important to study the treated healing lesions as a future research.

In conclusion, mapping of dendritic lesions in patients with HSK was successful in all patients using in vivo confocal microscopy. Cellular level observation of dendritic lesion at relatively larger image may help understand the in vivo morphological change of HSK. A further study in more patients with HSK and nonherpetic dendritic lesion is needed to utilize confocal microscopy images in the differential diagnosis and follow-up of the epithelial lesions with dendrite.

Figure 4 In vivo confocal microscopic images at the level of stroma and endothelium.

Notes: (A and B) Representative stromal image in Case 2 (A) and Case 4 (B). Honeycomb-shaped hyperreflective keratocytes were seen, suggesting stromal edema. (C and D) At the level of endothelium, inflammatory cells (arrows) were seen in Case 1 (C) and 4 (D). (Bar = 100 μ m).

Acknowledgment

This study was supported by a Grant-in-Aid for Scientific Research KAKENHI, Japan (No 23890072). The funding organizations had no role in the design or conduct of this research.

Disclosure

The authors report no conflicts of interest in this work.

References

1. Liesegang TJ. Herpes simplex virus epidemiology and ocular importance. *Cornea*. 2001;20:1–13.
2. Holland EJ, Schwartz GS, Neff KD. Herpes simplex keratitis. In: Krachmer JH, Mannis MJ, Holland EJ, editors. *Cornea*. Philadelphia, PA: Elsevier Mosby; 2011:953–984.
3. Cavanagh HD, Petroll WM, Alizadeh H, He YG, McCulley JP, Jester JV. Clinical and diagnostic use of in vivo confocal microscopy in patients with corneal disease. *Ophthalmology*. 1993;100:1444–1454.
4. Rosenberg ME, Tervo TM, Müller LJ, Moilanen JA, Vesaluoma MH. In vivo confocal microscopy after herpes keratitis. *Cornea*. 2002;21:265–269.
5. Nagasato D, Araki-Sasaki K, Kojima T, Ideta R, Dogru M. Morphological changes of corneal subepithelial nerve plexus in different types of herpetic keratitis. *Jpn J Ophthalmol*. 2011;55:444–450.
6. Hamrah P, Cruzat A, Dastjerdi MH, et al. Corneal sensation and subbasal nerve alterations in patients with herpes simplex keratitis: an in vivo confocal microscopy study. *Ophthalmology*. 2010;117:1930–1936.
7. Hamrah P, Sahin A, Dastjerdi MH, et al. Cellular changes of the corneal epithelium and stroma in herpes simplex keratitis: an in vivo confocal microscopy study. *Ophthalmology*. 2012;119:1791–1797.
8. Mocan MC, Irkec M, Mikropoulos DG, Bozkurt B, Orhan M, Konstant AG. In vivo confocal microscopic evaluation of the inflammatory response in non-epithelial herpes simplex keratitis. *Curr Eye Res*. 2012;37:1099–1106.

9. Martone G, Alegente M, Balestrazzi A, et al. In vivo confocal microscopy in bilateral herpetic keratitis: a case report. *Eur J Ophthalmol*. 2008; 18(6):994–997.
10. Yokogawa H, Kobayashi A, Sugiyama K. Mapping of normal corneal K-structures by in vivo laser confocal microscopy. *Cornea*. 2008;27(8): 879–883.
11. Yokogawa H, Kobayashi A, Sugiyama K. Mapping owl's eye cells of patients with cytomegalovirus corneal endotheliitis using in vivo laser confocal microscopy. *Jpn J Ophthalmol*. 2013;57:80–84.
12. Maudgal PC, Missotten L. Histopathology of human superficial herpes simplex keratitis. *Br J Ophthalmol*. 1978;62:46–52.
13. Nakagawa H, Uchida Y, Takamura E, et al. Diagnostic impression cytology for herpes simplex keratitis. *Jpn J Ophthalmol*. 1993;37: 505–513.
14. Kim JH, Ko MK, Shin JC. Infectivity of basal epithelial cells in herpetic dendritic epithelial keratitis. *Korean J Ophthalmol*. 1997;11:84–88.
15. Athmanathan S, Bandlapally SR, Rao GN. Collection of corneal impression cytology directly on a sterile glass slide for the detection of viral antigen: an inexpensive and simple technique for the diagnosis of HSV epithelial keratitis – a pilot study. *BMC Ophthalmol*. 2001;1:3.
16. Holbach LM, Font RL, Wilhelmus KR. Recurrent herpes simplex keratitis with concurrent epithelial and stromal involvement. Immunohistochemical and ultrastructural observations. *Arch Ophthalmol*. 1991;109: 692–695.
17. Blaum JL. Morphogenesis of the dendritic figure in herpes simplex keratitis. A negative study. *Am J Ophthalmol*. 1970;70:722–724.
18. Patel DV, McGhee CN. Mapping of the normal human corneal sub-Basal nerve plexus by in vivo laser scanning confocal microscopy. *Invest Ophthalmol Vis Sci*. 2005;46:4485–4488.
19. Patel DV, McGhee CN. Mapping the corneal sub-basal nerve plexus in keratoconus by in vivo laser scanning confocal microscopy. *Invest Ophthalmol Vis Sci*. 2006;47:1348–1351.

Clinical Ophthalmology

Publish your work in this journal

Clinical Ophthalmology is an international, peer-reviewed journal covering all subspecialties within ophthalmology. Key topics include: Optometry; Visual science; Pharmacology and drug therapy in eye diseases; Basic Sciences; Primary and Secondary eye care; Patient Safety and Quality of Care Improvements. This journal is indexed on

Submit your manuscript here: <http://www.dovepress.com/clinical-ophthalmology-journal>

Dovepress

PubMed Central and CAS, and is the official journal of The Society of Clinical Ophthalmology (SCO). The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.