Contraceptive use in women with hypertension and diabetes: cross-sectional study in northwest Ethiopia

Tensae Tadesse Mekonnen¹
Solomon Meseret
Woldeyohannes²
Tegbar Yigzaw³

¹Department of Midwifery, Tseda Health Science College, ²Department of Epidemiology and Biostatistics, Institute of Public Health, University of Gondar, Gondar, ³Jhpiego-Ethiopia, Addis Ababa, Ethiopia

Purpose: Women with diabetes and hypertension are at increased risk of pregnancy complications, including those from surgical delivery and their offspring are at risk for congenital anomalies. Thus, diabetic and hypertensive women of reproductive age are advised to use valid contraceptive methods for reducing unwanted pregnancy and its complications. However, contraceptive use among these segments of the population had not been previously assessed in Ethiopia. Hence, the aim of this study was to assess contraceptive use and associated factors among diabetic and hypertensive women of reproductive age on chronic follow-up care at University of Gondar and Felege Hiwot Hospitals.

Methods: Hospital-based cross-sectional study was conducted from April to May 2012 among diabetic and hypertensive women on follow-up at the chronic illness care center. The sample size calculated was 403. Structured and pretested questionnaire was used for data collection. Data were collected using interview supplemented by chart review. The data were entered using EPI info Version 2000 and analyzed using SPSS Version 16. Frequencies, proportion, and summary statistics were used to describe the study population in relation to relevant variables. Both bivariate and multivariate analyses were run to see the association of each independent variable with contraceptive practice.

Results: A total of 392 married women on chronic follow-up care were interviewed making the response rate of 93.3%. The contraceptive prevalence rate was found to be 53.8%. Factors such as age 25–34 years (adjusted odds ratio, AOR [95% confidence interval, CI]=3.60 [1.05–12.36]), (AOR [95% CI]=2.29 [1.15–4.53]), having middle- and high-level incomes (AOR [95% CI]=2.12 [1.19–3.77]), (AOR [95% CI]=5.03 [2.19–11.54]), receiving provider counseling (AOR [95% CI]=9.02 [4.40–18.49]), and controlled disease condition (AOR [95% CI]=4.13 [2.35–7.28]) were significantly associated with contraceptive practice.

Conclusion: The contraceptive utilization of women on diabetes and hypertension follow-up care was found to be low. Hence, strengthening counseling and education about family planning and controlling their medical conditions would help increase the contraceptive uptake of women on chronic follow-up.

Keywords: family planning, chronic disease, pregnancy

Introduction

Chronic diseases, such as heart disease, hypertension, diabetes, cancers, and chronic respiratory diseases, are by far the leading cause of mortality and morbidity in the world.¹ From the 58 million deaths that occurred in 2005, ~35 million were from chronic diseases,² and among these, 18 million were women.³ Women are uniquely affected by chronic disease. In the developing world, the effect of chronic diseases on women is even more pronounced. Women comprise 60% of the world's poor, and poverty worsens the chances of survival from chronic disease.⁴

Correspondence: Tensae Tadesse Mekonnen
Department of Midwifery, Tseda Health Science College, PO Box 102, Gondar, Ethiopia
Tel/fax +251 5 8111 6221
Email almaztade@gmail.com

Globally, there are 300 million people who have diabetes, and approximately half of these are women.⁴ A study that was conducted in Addis Ababa showed that from a total of 229 diabetic clients, 63.85% were females.⁵ Hypertension is also an important cause of mortality and morbidity in reproductive age women.⁶ Although younger, premenopausal women have lower blood pressures (BPs) than the age-matched men population, BP rises with age, and the prevalence of hypertension is higher in older women.⁷ In 2006, the prevalence of hypertension in reproductive age US women was 6.9%.⁸ A population-based study that was conducted in Addis Ababa showed that the prevalence of hypertension in adult females was found to be 29%.⁹

Women with diabetes and hypertension are at increased risk of pregnancy-related complications, which includes spontaneous abortion, preterm labor, hypertensive disorders, and delivery by cesarean section.¹⁰ Existing population-based studies demonstrated increased congenital abnormalities and prenatal mortality in diabetic mothers as compared to the non-diabetic population.^{11,12} Clinical and experimental evidence indicate that the increased risk of congenital malformation and spontaneous abortion in pregnancies complicated by diabetes is linked to disturbance in maternal metabolism around the time of conception and during the period of organogenesis.¹³ Unfortunately, unplanned pregnancies occur in two-thirds of women with diabetes, leading to persistent excess malformation in their infants.¹⁴ There are no contraceptive methods that are specifically contraindicated for women with diabetes.¹³ Thus, the selection of a method for an individual patient should be made by a provider who is familiar with contraceptive prescribing practices using the same guidelines that are relevant to a woman without diabetes. Because unexpected failure of contraception can lead to increased risk of congenital malformation and spontaneous abortion in pregnancies complicated by diabetes, the focus should be on methods with verified high degree of effectiveness.¹⁵

Although research and educational efforts addressing contraceptive use in women with medical problems are limited, pregnancy in women with medical disease in many instances exposes both the woman and her offspring to definite morbidity and possible mortality.¹⁴ Women with a medical disease should be offered the most effective methods of contraception that also safeguard their existing health status.¹⁵

The lack of contraceptive use in this high-risk group population is related to sociodemographic and other factors related to pregnancy.¹³ Approximately half of all pregnancies in USA are unintended and nearly two-thirds of pregnancies in women with diabetes are unplanned.¹⁶

Hence, diabetes and hypertension are diseases with severe complications. The problems are even more pronounced among reproductive women than in nonreproductive women. Women with diabetes and hypertension are at increased risk of pregnancy complications, including those from surgical delivery, and their offspring are at risk of congenital anomalies. Thus, diabetic and hypertensive women of reproductive age must use valid contraceptive methods for reducing unwanted pregnancy and its complications. Effective family planning used in conjunction with hypertension and glucose management may reduce the risk to both the mother and the fetus. In spite of family planning benefits to these categories of the population, practice of using contraceptive among these segments of the population in Ethiopia was not previously assessed. Therefore, this study was conducted with the purpose of assessing contraceptive practice by diabetic and hypertensive married women of reproductive age group on follow-up at the chronic illness care center, with the hope of bringing information to support and guide further development and accomplishment of specific intervention program targeted to women of reproductive age group with diabetes and hypertension. Besides, it could serve as a baseline data for planning future studies.

Materials and methods

Institution-based cross-sectional study was conducted from April to May, 2012, at the University of Gondar Referral Hospital and Felege Hiwot Referral Hospital. These two hospitals are the largest public hospitals in the Amhara National Regional State. A total of 4,096 diabetic and 2,100 hypertensive clients were having regular follow-up at the two hospitals. Among these, 2,795 (1,995 diabetic and 800 hypertensive) were women. All women who fulfilled the inclusion criteria (married women of reproductive age, ie, 15–49 years, who were not pregnant and infertile at the time of the study) and who came for follow-up visit during the data collection period were enrolled. These women came because they seek care, hence, the study has nothing to do with absence or presence of women during the study period and this could help in minimizing the selection bias.

Sample size was determined using the formula for estimating a single population proportion.¹⁷ The following assumptions were made while calculating the sample size. A 95% probability of obtaining the population proportion of women who practice contraceptive within 5% margin of error was assumed to be 50% as there were no studies done in Ethiopia about contraceptive practice and associated

factors among diabetic and hypertensive women. Following the assumptions and using the Statistical Package Epi_Info Version 2000 (Centers for Disease Control and Prevention, Atlanta, GA, USA), the required size was 385 married women in the reproductive age group. Expecting a 5% nonresponse rate, the final sample size was calculated to be 403.

Nurses collected the data via interview using structured questionnaire. To assure data quality, a number of precautions were taken. The questionnaire was prepared in English, translated to Amharic, and then translated back to English, to check for consistency. Training was given for the data collectors. Before the actual data collection, pretest was done on 21 women in the follow-up care and appropriate modifications were made on the questionnaire after analyzing the pretest result. The supervisors and the principal investigator made frequent checks on the data collection process to ensure the completeness and consistency of the gathered information.

Data were collected on women's age, place of residence, educational status, occupation of mother, monthly income, media exposure, duration of follow-up, BP or glucose level, number of comorbidities and provider counseling, and number of living children and contraceptive practice. Operational definitions were performed on the following variables.

Controlled hypertension: If a woman had maintained the average BP reading $<140/90$ mmHg at the time of data collection irrespective of measurements at other times. However, readings $\geq 140/90$ mmHg is referred as uncontrolled hypertension.

Controlled diabetes: If a woman had maintained mean fasting blood sugar <130 mg/dL for the last three consecutive visits.

Contraceptive practice: Usage of one of hormonal, barrier, or sterilization method of contraceptives. In this study, contraceptive use refers to if a woman is currently using any one of hormonal, barrier, or sterilization method. Hence, a woman was categorized as current contraceptive user if she used one of the methods, otherwise as a nonuser.

Family monthly income: Low level – if the family monthly income is $<25\%$ quartile, that is, <500 Birr; Middle level – if the family monthly income is between the 25% and 75% quartile, 500–2,000 Birr; High level – if the family monthly income is $>75\%$ quartile, that is, $>2,000$ Birr.

The data were entered and cleaned using EPI info Version 6 statistical software (Centers for Disease Control and Prevention) and exported to SPSS Version 16 statistical

packages for analysis. Frequencies, proportion and summary statistics were used to describe the study population in relation to relevant variables. Logistic regression was carried out to see the effect of each independent variable on the dependent variable. P -value <0.05 was used to declare statistical significance. Odds ratio with 95% CI was computed to assess the strength of the association.

Ethical clearance was obtained from the Ethical Review Board of Institute of Public Health. Communication with the University of Gondar Hospital and Felege Hiwot Hospital was made through formal letter. After the purpose and objective of the study have been informed, verbal consent was obtained from each study participant. Participants were also informed that participation would be on voluntary basis, and they could withdraw from the study at any time if they were not comfortable about the questionnaire. Participants were also informed that all data obtained from them would be kept confidential by using codes instead of any personal identifiers.

Results

Sociodemographic and economic characteristics of the respondents

A total of 392 women responded to the questionnaire, yielding a response rate of 97.3%. One hundred fifty-four (39.3%) respondents were from the rural areas. The mean age of the respondents was 36 ± 7.58 years. Approximately 45% of the study participants and 33.4% of their spouses did not have formal education. Most respondents (87%) belonged to the Amhara ethnic group and (78.6%) were followers of Orthodox Christianity. Furthermore, 53.1% (208) of women were housewives, 53.3% (209) had middle-level family income, and 83.8% (329) possessed radio and/or television (Table 1).

Chronic medical conditions

Approximately half (49.5%) and one-third (35.7%) of the study participants had diabetes mellitus and hypertension, respectively, while the remaining 14.8% had both. More than eight out of ten women were on follow-up for >1 year, and the disease was well controlled in 271 (69.1%) of the patients. In addition, 100 (25.5%) women had comorbidity and 68 (17.3%) reported facing pregnancy complication after they developed the disease (Table 2).

Contraceptive practice

Two hundred seventy-one (69.1%) women reported that they ever used contraceptive methods. However, only 53.8%

Table 1 Sociodemographic and economic characteristics of the study participants, May 2012

Variables	Frequency	Percentage
Residence		
Rural	154	39.3
Urban	238	60.7
Age, years		
≤24	34	8.7
25–34	95	24.2
≥35	263	67.1
Ethnicity		
Amhara	343	87.5
Tigre	27	6.9
Others	22	5.6
Religion		
Orthodox Christian	308	78.6
Muslim	70	17.9
Protestant	13	3.3
Catholic Christian	1	0.3
Educational status		
Cannot read and write	129	32.9
Read and write	49	12.5
Class 1–6	33	8.4
Class 7–10	65	16.6
Class 11–12	96	24.5
College or university	20	5.1
Women occupation		
Housewife	208	53.1
Merchant	61	15.6
Daily laborer	27	6.9
Government employee	96	24.5
Husband educational status		
Cannot read and write	45	11.5
Read and write	86	21.9
Class 1–6	34	8.7
Class 7–10	57	14.5
Class 11–12	108	27.6
College or university	62	15.8
Husband occupation		
Merchant	104	26.5
Government employee	159	40.6
Daily laborer	56	14.3
Farmer	73	18.6
Family monthly income		
Low	114	29.1
Middle	209	53.3
High	69	17.6
Having TV or radio		
Radio	97	24.7
TV	28	7.1
Radio and TV	204	52.0
None	63	16.1

(211) were using modern contraceptive methods during the time of data collection. The frequently used methods were injectable (39.8%), intrauterine contraceptive device (IUCD) (18%), pill (16.1%), and implant (14.2%). More than half (53.1%) of the women were using contraceptives for the purpose of limiting births. The most frequently mentioned

Table 2 Chronic medical conditions of the study participants, May 2012

Variables	Frequency	Percentage
Type of disease		
Diabetes	194	49.5
Hypertension	140	35.7
Both	58	14.8
Duration of follow-up		
≤1 years	68	17.3
1–4 years	178	45.4
>4 years	146	37.2
Disease controlled		
Yes	271	69.1
No	121	30.9
Comorbidity		
Yes	100	25.5
No	292	74.5
History of pregnancy complication		
Yes	68	17.3
No	324	82.7

reasons for not using contraceptive were fear of side effects (12.8%), desire for children (12.2%), religious prohibition (11.2%), husband opposition (11%), and lack of knowledge about contraceptives (8.9%; Table 3).

Table 3 Contraceptive practice of the respondents, May 2012

Variable	Frequency	Percentage
Ever use		
Yes	271	69.1
No	121	30.9
Current use		
Yes	211	53.8
No	181	46.2
Contraceptive type currently using		
Pills	34	16.1
IUCD	38	18
DMPA	84	39.8
Norplant	30	14.2
Tubal ligation	16	7.6
Emergency pills	5	2.4
Natural method	4	1.8
Purpose of using		
Spacing between children	99	46.9
Limiting the number of children	112	53.1
Husband approval		
Approve	156	39.8
Disapprove	219	55.9
Do not know	17	4.3
Reasons for nonuse		
Fear of side effect	50	12.8
Not aware of contraceptive	35	8.9
Husband opposition	43	11.0
Infrequent sex	35	8.9
Religion prohibition	44	11.2
Desire for children	48	12.2

Abbreviations: IUCD, intrauterine contraceptive device; DMPA, depot medroxy-progesterone acetate.

Factors associated with contraceptive use

The impact of selected sociodemographic and other characteristics on contraceptive practice was investigated using both the classic bivariate methods and the multivariate logistic regression technique. Variables found to be significant at *P*-value of 0.2 were included in the multivariate analysis. Finally, age, income, discussion with health worker about contraception, well-controlled disease condition, and having living children were found to be significantly and independently associated with contraceptive practice of women with chronic disease on follow-up care. Accordingly, women belonging to age groups 25–34 years (adjusted odds ratio, AOR [95% confidence interval, CI] =3.6 (1.1–12.4)) and 35–39 years (AOR [95% CI] =2.3 (1.2–4.5)) were more likely to use contraceptives as compared to those younger than 25 years. Participants in high- (AOR [95% CI] =2.1 [1.2–3.8]) and middle- (AOR [95% CI] =5.0 [2.2–11.5]) income groups were more likely to use contraceptive than those in low-level income group.

The number of living children that the women had was also a factor that had influence on contraceptive practice. Women having one to four (AOR [95% CI] =7.2 [2.4–21.9]) and five and more (AOR [95% CI] =7.6 [2.9–19.5]) living children were approximately seven times more likely to use contraceptives than women with no children. Having discussion with health professionals concerning contraceptive issues for women with chronic disease was important because it helped them to practice it. Respondents reporting having discussion concerning contraceptives were approximately nine times more likely to use contraceptive (AOR [95% CI] =9.0 [4.40–18.5]; Table 4).

Discussion

Contraception in women suffering from chronic diseases is an important question, mainly as it has been proven that pregnancy outcome both for the fetus and for the mother depends on disease condition before conception right to delivery.¹⁸ Women with chronic medical conditions are of particular concern due to significantly increased risk for pregnancy-related morbidity, mortality, and adverse pregnancy outcomes. This has made planning of future pregnancy mandatory for these patients.^{18,19} In this study, 53.8% of married women surveyed were using contraceptive methods. Although this figure was higher than contraceptive prevalence rate at national (29%) and regional state (33.9%) levels as reported in the 2011 Ethiopian Demographic and Health Survey,²⁰ the greater likelihood of risk to the mother and the fetus in this group makes the contraceptive use

unacceptably low. Studies done in Michigan, Iran, and Italy reported very high levels of contraceptive use among women with chronic medical conditions, 80%, 85%, and 89.3%,^{21–23} respectively, while a study from Algeria found comparable contraceptive prevalence.²⁴

In this study, we found an association between contraceptive use and women's age, income level, presence of children, discussion with health care providers about contraception, and status of disease control. Women ≥ 25 years of age were more likely to use contraceptive than their younger counterparts. This might be because younger married women have greater desire for more children and, hence, do not use contraceptives. This interpretation is consistent with the finding in this study that the desire for more children was the second most frequent reason for the nonuse of contraceptives. This finding was in line with other studies done in USA and Italy.^{21,22}

Women in the middle and higher income groups in this study were more likely to use contraceptives than those who have low income. This finding is consistent with results from the Ethiopian demographic and health survey that reported current use of contraceptives increases with wealth.²⁰ This may be due to situational, demographic, and behavioral factors related to economical status that increase the knowledge, motivation, and get necessary support to practice contraceptives. Having higher income has similarly been identified in other studies as having significant association with contraceptive use among women with diabetes who seek preconception care and continuous use of contraception.^{25,26}

In this study, “having living children” was strongly associated with the contraceptive practice. This finding is consistent with the pattern in the Ethiopian demographic and health survey, which reported higher contraceptive use among women having one to four living children than those without children.²⁰ This is because the main determinant to contraception continuation and type seems to be the size of the family, which is determined by the number of children. It is also considered as one of the major determinants in the developing countries. This result is similar with Algeria's study.²⁴

The study demonstrates that women who discussed with health care providers about contraceptive use were approximately nine times more likely to practice contraceptive than those who did not. This is intuitive because discussion increases women's contraceptive knowledge and addresses their concerns, thereby facilitating adoption and continuation of contraceptive use. It is known that when health care providers counsel clients well, it helps them successfully choose and use contraceptives.²⁵

Table 4 Association of selected sociodemographic and other characteristics with contraceptive practice among women on chronic follow-up care at Gondar University Hospital and Felege Hiwot Hospital, May 2012

Variable	Contraceptive use		Crude OR (95% CI)	Adjusted OR (95% CI)
	Yes	No		
Age, years				
≤24	13 (38.2%)	21 (61.8%)	1	1
25–34	56 (58.9%)	39 (41.1%)	2.3 (1.0–5.2)	3.6 (1.1–12.4)
35–49	142 (54.0%)	121 (46.0%)	1.9 (0.9–3.9)	2.3 (1.2–4.5)
Resident				
Rural	75 (48.7%)	79 (51.3%)	1	
Urban	136 (57.1%)	102 (42.9%)	1.4 (0.9–2.1)	
Educational status				
Cannot read and write	65 (50.4%)	64 (49.6%)	1	
Read and write	32 (65.3%)	17 (34.7%)	1.9 (0.9–3.7)	
Class ≤10	45 (45.9%)	53 (54.1%)	0.8 (0.5–1.4)	
Class 11+	69 (59.5%)	47 (40.5%)	1.4 (0.9–2.4)	
Occupation				
Housewife	108 (51.9%)	100 (48.1%)	1	
Merchant	35 (57.4%)	26 (42.6%)	1.3 (0.7–2.2)	
Daily Laborer	11 (40.7%)	16 (59.3%)	0.6 (0.3–1.4)	
Government employee	57 (59.4%)	39 (40.6%)	1.4 (0.8–2.2)	
Having TV/radio				
Radio	49 (50.5%)	48 (49.5%)	2.4 (1.2–4.6)	
TV	15 (53.6%)	13 (46.4%)	2.7 (1.1–6.7)	
Both	128 (62.7%)	76 (37.3%)	3.9 (2.1–7.2)	
None	19 (30.2%)	44 (69.8%)	1	
Income				
Low	40 (35.1%)	74 (64.9%)	1	1
Middle	116 (55.5%)	93 (44.5%)	2.3 (1.4, 3.7)	2.1 (1.2–3.8)
High	55 (79.7%)	14 (20.3%)	7.3 (3.6, 14.7)	5.0 (2.2–11.5)
No of living children				
None	18 (25.4%)	53 (74.6%)	1	1
1–4	158 (64.0%)	89 (36.0%)	5.2 (2.9, 9.5)	7.2 (2.4–21.9)
>5	35 (47.3%)	39 (52.7%)	2.6 (1.3, 5.3)	7.6 (2.9–19.5)
Discussion with health workers				
Yes	91 (89.2%)	11 (10.8%)	11.7 (6.0–22.9)	9.1 (4.4–18.5)
No	120 (41.4%)	170 (58.6%)	1	1
Disease controlled				
Yes	182 (67.2%)	89 (32.8%)	6.5 (3.9–10.6)	4.1 (2.4–7.4)
No	29 (24.0%)	92 (76.0%)	1	1
Duration of follow-up				
<1 years	41 (60.3%)	27 (39.7%)	1	
1–4 years	99 (55.6%)	79 (44.4%)	0.8 (0.5–1.5)	
>4 years	71 (48.6%)	75 (51.4%)	0.6 (0.4–1.1)	
Comorbidities				
Yes	55 (55.0%)	45 (45.0%)	1.1 (0.7–1.7)	
No	156 (53.4%)	136 (46.6%)	1	

Abbreviations: OR, odds ratio; CI, confidence interval.

Regarding disease condition, women whose medical condition is controlled were more likely to practice contraceptives than those who did not. The reason may be those women with better controlled disease did so because they adhere to medical advice and, by extension, are more likely to use contraceptives. It is also possible that women in good health condition are more likely to be sexually active and use contraceptives.

Limitations of the study

This study could have the following limitations. First, since the study was done only among women on follow-up care, generalization could not be made to all women with hypertension and diabetes who were not at follow-up care at the time of the study. Data regarding treatment adherence, body mass index, husband–wife communications on choice and decisions were not collected, which might have implications

on the current findings of the study. In addition, the role of mass media on contraceptive practice was not assessed. Finally, there could be a possibility for social desirability of responses.

Conclusion

Contraceptive utilization of women on diabetes and hypertension follow-up care was found to be low. Women aged 25–34 years, of high and middle level family monthly income groups, having more than five living children, providing counseling and controlling of the disease were found to have positive association with the contraceptive practice. Hence, strengthening counseling, education about family planning, and controlling their medical conditions would help increase the contraceptive uptake of women on chronic follow-up care.

Acknowledgments

The authors would like to forward our deepest gratitude to the staff of the Institute of Public Health for their material and technical support and to the women who participated in the study.

Author contributions

All authors made substantial contributions to conception and design, acquisition of data, or analysis and interpretation of data; took part in either drafting the article or revising it critically for important intellectual content; gave final approval of the version to be published; and agree to be accountable for all aspects of the work.

Disclosure

The authors report no conflicts of interest in this work.

References

1. Alwan A, MacLean DR, Riley LM, et al. Monitoring and surveillance of chronic non-communicable diseases: progress and capacity in high-burden countries. *Lancet*. 2010;376(9755):1861–1868.
2. World Health Organization. Preventing chronic diseases: a vital investment. Geneva: World Health Organization; 2005. Available from: http://www.who.int/chp/chronic_disease_report/en/. Accessed September 17, 2015.
3. World Health Organization. *Global status report on noncommunicable diseases 2010*. Geneva: World Health Organization; 2010. Available from: http://www.who.int/nmh/publications/ncd_report_full_en.pdf. Accessed September 17, 2015.
4. The NCD Alliance. *Noncommunicable diseases: a priority for women's health and development*. 2010. Available from: http://www.who.int/pmnch/topics/maternal/2011_women_ncd_report.pdf. Accessed September 17, 2015.
5. Feleke Y, Enkuselassie F. An assessment of the health care system for diabetes in Addis Ababa, Ethiopia. *Ethiop J Health Dev*. 2005;19(3): 203–210.
6. Say L, Chou D, Gemmill A, et al. Global causes of maternal death: a WHO systematic analysis. *Lancet Glob Health*. 2014;2(6):e323–e333.
7. August P, Oparil S. Acute Long term management of hypertension in non-pregnant women. *BMJ*. 2012;27(5):145–178.
8. CDC. Preventing and managing chronic disease to improve the health of women and infants; 2010. Available from: <http://www.cdc.gov/reproductivehealth>.
9. Tesfaye F, Byass P, Wall S. Population based prevalence of high blood pressure among adults in Addis Ababa: uncovering a silent epidemic. *Cardiovasc Disord*. 2009;9(39):1471–2261.
10. Casson IF, Clarke CA, Howard CV, McKendrick O, Pennycook S, Pharonah PO. Outcomes of pregnancy in insulin dependent diabetic women: result of a five year population cohort study. *BMJ*. 1997;315: 275–278.
11. Wahabi HA, Alzeidan RA, Bawazeer GA. Pre conception care for diabetic women for improving maternal and fetal outcomes systemic review. *Diabetes Care*. 2010;10(63):1471–2393.
12. American Diabetes Association. Preconception care of women with diabetes. *Diabetes Care*. 2004;27(suppl 1):S76–S78.
13. Rey E, Couturier A. The prognosis of pregnancy in women with chronic hypertension. *Am J Obstet Gynecol*. 1994;171(2):410–416.
14. Kjos SL. Contraception for women with medical problems. In: *Gynecology and Obstetrics CD-ROM*. Philadelphia: Lippincott Williams and Wilkins; 2004;1–6:Chapter 14.
15. Institute of Medicine. Weight gain during pregnancy; 2010. Available from: www.iom.edu/Reports/2009/Weight-Gain-During-PregnancyReexamining-theGuidelines.aspx. Accessed April 2, 2012.
16. Evers IM, de Valk HW, Visser GH. Risk of complications of pregnancy in women with type 1 diabetes: nationwide prospective study in the Netherlands. *BMJ*. 2004;328(7445):915.
17. Fosgate GT. Practical sample size calculations for surveillance and diagnostic investigations. *J Vet Diagn Invest*. 2009;21(1):3–14.
18. Nikolov A, Dimitrov A, Kolarov G, Todorova K, Mekhandzhiev T. Contraception in women with diabetes mellitus. *Akush Ginekolog*. 2005; 44(5):47–52.
19. Dunlop AL, Jack BW, Bottalico JN, et al. The clinical content of pre-conception care: women with chronic medical conditions. *Am J Obstet Gynecol*. 2008;34(2):S310–S327.
20. Central Statistical Agency [Ethiopia] and ICF International. *Ethiopia Demographic and Health Survey 2011*. Addis Ababa, Ethiopia and Calverton, Maryland, USA: Central Statistical Agency and ICF International; 2012.
21. Cynthia H, Chuang A. Contraceptive use by diabetic and obese women. *Womens Health Issue*. 2005;15(4):167–173.
22. Mazaheri E, Mostafazadeh F, Karimollahi M. Pattern of contraceptive use in diabetic women. *Maturitas*. 2009;63(1):128–150.
23. Janz N. Contraception in diabetic women: an Italian study. *Diabetes Res Clin Pract*. 2005;67(3):267–272.
24. Benotmane A, Faraoun K. Contraceptive practice in diabetic women in Algeria. *Diabetes Metab*. 2008;27(4):510–511.
25. Holing E. Why don't women with diabetes plan their pregnancies. *Diabetes Care*. 1998;21(6):889–895.
26. Vahratian A, Barber J, Lawrence J. Family planning practice among diabetes, over weight and obese women in 2002 national survey for family growth. *Diabetes Care*. 2009;32(6):1026–1031.

International Journal of Women's Health**Dovepress****Publish your work in this journal**

The International Journal of Women's Health is an international, peer-reviewed open-access journal publishing original research, reports, editorials, reviews and commentaries on all aspects of women's healthcare including gynecology, obstetrics, and breast cancer. The manuscript management system is completely online and includes

a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <http://www.dovepress.com/international-journal-of-womens-health-journal>