Effect of picoside II on erythrocyte deformability and lipid peroxidation in rats subjected to hind limb ischemia reperfusion injury

Faruk Metin Çomu¹Yiğit Kılıç²Abdullah Özer²Mehmet Kirişçi³Ali Doğan Dursun⁴Tolga Tatar²Mustafa Hakan Zor²Hakan Kartal²Ayşegül Küçük⁵Hakan Boyunağa⁶Mustafa Arslan⁷¹Department of Physiology, Kirikkale University Medical Faculty, Kirikkale,²Department of Cardiovascular Surgery, Gazi University Medical Faculty, Ankara, ³Department of Cardiovascular Surgery, Necip Fazıl State Hospital, Kahramanmaraş,⁴Department of Physiology, Ankara University Medical Faculty, Ankara, ⁵Department of Physiology, Dumlupınar University Medical Faculty, Kütahya, ⁶Department of Biochemistry, Kirikkale University Medical Faculty, Kirikkale,⁷Department of Anaesthesiology and Reanimation, Gazi University Medical Faculty, Ankara, Turkey

Background: Ischemia reperfusion injury (I/R) in hind limb is a frequent and important clinical phenomenon. Many structural and functional damages are observed in cells and tissues in these kinds of injuries. In this study, we aimed to evaluate the effect of picoside II on lipid peroxidation and erythrocyte deformability during I/R in rats.

Methods: Rats were randomly divided into four groups – each containing six animals (sham, I/R, sham + picoside II, and I/R + picoside II). The infrarenal section of the abdominal aorta was occluded with an atraumatic microvascular clamp in I/R groups. The clamp was removed after 120 minutes and reperfusion was provided for a further 120 minutes. Picoside II (10 mg·kg⁻¹) was administered intraperitoneally to the animals in the appropriate groups (sham + picoside II, I/R + picoside II groups). All rats were euthanized by intraperitoneal administration of ketamine (100 mg·kg⁻¹) and taking blood from the abdominal aorta. Erythrocytes were extracted from heparinized complete blood samples. Buffer (P_T) and then erythrocytes (P_E) were passed through the filtration system and the changes in pressure were measured to investigate the role of serum malondialdehyde and nitric oxide (NO) in lipid peroxidation and erythrocyte deformability index.

Results: Deformability index was significantly increased in the I/R group compared to groups sham, sham + picoside-II, and I/R + picoside-II ($P<0.0001$, $P<0.0001$, and $P=0.007$). Malondialdehyde (MDA) and NO levels were evaluated. MDA level and NO activity were also higher in the I/R group than in the other groups. Picoside II treatment before hind limb I/R prevented these changes.

Conclusion: These results support that deformability of erythrocytes is decreased in I/R injury and picoside II plays a critical role to prevent these alterations. Further experimental and clinical studies are needed to evaluate and clarify the molecular mechanisms of action and clinical importance of these findings.

Keywords: hind limb ischemia reperfusion injury, lipid peroxidation, malondialdehyde, nitric oxide, erythrocyte deformability, picoside II

Introduction

Ischemia reperfusion injury (I/R) in hind limb is a frequent and important clinical phenomenon. Reperfusion period following an ischemic insult may paradoxically cause increased rates of mortality and morbidity due to systemic complications. Local edema and muscle tissue necrosis are likely to be followed by systemic inflammatory response syndrome and multiple organ failure (kidney, respiratory, and circulatory system) as reperfusion advances.^{1–3}

Correspondence: Mustafa Arslan
Department of Anaesthesiology and
Reanimation, Gazi University Medical
Faculty, 06510, Beşevler, Ankara, Turkey
Tel +90 312 202 67 39
Email mustarslan@gmail.com

Swelling of cells, degeneration of cell skeleton structure, and loss of selective membrane permeability are characteristic features of reperfusion injury. These changes all end up with tissue edema and decreased capillary blood flow.⁴

Microcirculatory damage is the first target of I/R. Leukocyte–endothelial interactions cause transendothelial migration and tissue damage by triggering the release of reactive oxygen species and elastase.⁵ Many tissues and cells such as erythrocytes may be damaged due to the exogenous sources of reactive oxygen species.⁶ Deformability of erythrocytes is the key factor for the regulation of normal microvascular circulation. Viscoelastic properties of erythrocyte membrane are the major determinants of red blood cell deformability.⁷

Picroside II is one of the main active constituents isolated from *Picrorhiza scrophulariiflora*. The roots of this plant are of benefit and often used in traditional Chinese medicine for a number of conditions.⁸ Picroside II has been shown to possess a wide range of pharmacological effects, including effects against oxidative stress and inflammation.^{9–11} Picroside II has also been shown to protect against I/R in other organs, including the brain,¹² and kidneys,¹³ due to its antioxidative, anti-inflammatory, and antiapoptotic properties.

Previous studies have shown that picroside II prevents the I/R-induced lipid peroxidation, but as far as we know this is the first report of picroside II's effect on lipid peroxidation and erythrocyte deformability in hind limb I/R injury. The primary purpose of this study was to investigate I/R injury-induced changes of hind limb in deformability of erythrocytes and the protective role of picroside II on increased lipid peroxidation and erythrocyte deformability alterations in experimental hind limb I/R injury.

Materials and methods

Animals and experimental protocol

This study was conducted in the GUDAM (Gazi Universitesi Laboratuvar Hayvanları Yetiştirme ve Deneysel Araştırmalar Merkez) Laboratory of Gazi University with the consent of the Experimental Animals Ethics Committee at Gazi University. All procedures were performed according to the accepted standards of the Guide for the Care and Use of Laboratory Animals.

In our study, 24 male Wistar Albino rats weighing between 175 and 230 g, raised under the same environmental conditions, were used. The rats were housed between 20°C–21°C with 12-hour daylight and 12-hour darkness cycles, and had free access to food until 2 hours before the anesthesia procedure. The animals were randomly separated

into four groups, each containing six rats. Midline laparotomy was done under ketamine (100 mg·kg⁻¹) anesthesia.

Sham group: midline laparotomy was done alone without any additional surgical intervention. Blood samples were collected after 4 hours of follow-up and animals were sacrificed eventually.

Sham + picroside II group: midline laparotomy was done alone without any additional surgical intervention. Blood sample was collected after 4 hours of follow-up and animals were sacrificed eventually.

I/R group: midline laparotomy was done similarly. Infra-renal part of the abdominal aorta was clamped for 2 hours. After removing the clamp, reperfusion was established for another 2 hours. Finally, rats were sacrificed after collecting blood samples from their abdominal aorta.

I/R + picroside II group: Similar steps were followed but in addition to the procedure mentioned above, picroside II was supplied by Sigma-Aldrich Co. Ltd. (St Louis, MO, USA; CAS No: 39012-20-9, purity >98%, molecular formula: C₂₃H₂₈O₁₃). The picroside II was diluted to a 1% solution with normal saline. The animals in the treatment group were administered an intraperitoneal injection of picroside II 30 minutes before ischemia (10 mg·kg⁻¹); the animals in the control and I/R groups were injected with the same volume of 0.09% NaCl. Rats were sacrificed at the end of reperfusion period that lasted 2 hours, after collecting blood samples.

All the rats were given ketamine 100 mg·kg⁻¹ intraperitoneally and intraabdominal blood samples were taken. Erythrocyte packs were prepared using heparinized total blood samples. Measurements for deformability were performed using erythrocyte suspensions with 5% hematocrit in phosphate buffered saline (PBS).

Deformability measurements

Samples of blood were taken very carefully and the measurement process was done as fast as possible to avoid hemolysis. Collected blood was centrifuged at 1,000×g for 10 minutes. Serum and buffy coat on erythrocytes were removed. Isotonic PBS buffer was added to collapsing erythrocytes and this mixture was centrifuged at 1,000×g for 10 minutes. Liquid on the upper surface was removed. Finally pure red cell packs were obtained from the washing process, which was repeated three times. Erythrocyte packs were mixed with PBS buffer to generate a suspension with the value of 5% hematocrit. These erythrocyte suspensions were used for the measurement of deformability. Collection and deformability measurements of erythrocytes were done at 22°C.

The constant-current filtrometer system was used for measurement of erythrocyte deformability. Samples to be

measured were prepared as 10 mL of erythrocytes suspension and PBS buffer. The flow rate was held constant at 1.5 mL/min with an infusion pump. A 28 mm nucleopore polycarbonate filter with a 5 μ m pore diameter was preferred. A constant-current filterometer system was used in the measurement of the erythrocyte deformability. Pressure changes were detected by a pressure transducer while the erythrocytes passed through the filter and the data was transferred to computer with the help of MP 30 data equation systems (BIOPAC Systems Inc., Goleta, CA). The necessary calculations were performed with related computer programs by measuring the pressure changes at various times. Pressure calibration of the system was performed each time before measuring the samples. First buffer (P_T) and then erythrocytes (P_E) were passed through from the filtration system and the changes in pressure were measured. The relative refractory period value relative resistance (Rrel) was calculated by relating the pressure value of erythrocyte suspension to pressure value of buffer. The deformability index was interpreted as Rrel was increasing the ability of erythrocyte deformability was affected adversely.¹⁴

Measurements of MDA levels and nitric oxide activities

Esterbauer method was applied in order to measure lipid peroxidation. Malondialdehyde reacted with thiobarbituric acid at 90°C–95°C and resulted in pink chromogranin. After 15 minutes specimens rapidly cooled, then absorbances were read at 532 nm spectrophotometrically. Results were represented as nmol/g tissue protein.¹⁵ Concentrations of stable oxidative metabolites of nitric oxide (NO , NO_2^- , and NO_3^-) were measured in serum and NO production was determined. Measurement of nitrite concentration was performed with Griess reaction.¹⁶

Statistical analysis

The statistical analyses were performed with SPSS 17.0 software program and $P < 0.05$ was considered statistically significant. The findings were expressed as mean $\pm$ standard deviation. The data were evaluated with Kruskal–Wallis variance analysis. The variables with significance were evaluated with Bonferroni corrected Mann–Whitney U -test.

Results

Erythrocyte deformability as a significant determinant of relative resistance was increased in group I/R. Deformability index was significantly increased in the I/R group compared to groups sham, sham + picoside II, and I/R + picoside II

Figure 1 Erythrocyte deformability index values of the groups.

Notes: Each bar represents the mean $\pm$ SE. * $P < 0.05$ compared to the group ischemia/reperfusion.

Abbreviation: SE, standard error.

($P < 0.0001$, $P < 0.0001$, $P = 0.007$, respectively) (Figure 1). However, after picoside II treatment deformability index was similar with the control (groups sham, sham + picoside II, and I/R + picoside II).

Malondialdehyde (MDA) level was also higher in I/R group than the other groups ($P < 0.0001$, $P < 0.0001$, $P = 0.007$, respectively) (Figure 2). However, picoside II treatment in I/R-P group resulted in similar MDA levels to the sham, sham + picoside II groups (Figure 2).

NO activity was also higher in I/R group than the other groups ($P = 0.015$, $P = 0.020$, $P = 0.016$, respectively) (Figure 3). However, picoside II treatment in I/R + picoside II group resulted in similar NO activity to the sham, sham + picoside II groups (Figure 3).

Discussion

Picoside II (β -D-glucopyranoside, 1a, 1b, 2, 5a, 6, 6a-hexahydro-6-[(4-hydroxy-3-methoxybenzoyl)oxy]-1-a(hydroxymethyl)oxireno[4,5]cyclopenta[1,2-c]pyran-2-yl)

Figure 2 Malondialdehyde (MDA) levels of the groups.

Notes: Each bar represents the mean $\pm$ standard error. * $P < 0.05$ compared to the group ischemia/reperfusion.

Abbreviation: prot, protein.

Figure 3 Nitric oxide (NO) activity values of the groups.

Notes: Each bar represents the mean $\pm$ standard error. * $P < 0.05$ compared to the group ischemia/reperfusion.

is a major iridoid glycoside isolated from *P. scrophulariiflora* Pennell (Scrophulariaceae).¹⁷ Previous studies have shown that picoside II has a number of pharmacological effects, including neuroprotective,^{12,18} hepatoprotective,¹⁹ antiapoptotic, anticholestatic, anti-inflammatory, and immune-modulating activities.^{10,11} In addition, picoside II is also reported to possess potent antioxidant activities.^{9,18}

Unpaired electrons, free radicals, are highly reactive and readily take part in chemical reactions with virtually all cell components (lipids, proteins, complex carbohydrates and nucleic acids) in the body. These reactions occur through a chain of oxidative reactions to cause tissue injury. For most biological structures (like lipids, proteins, and nucleic acids), free radical damage is closely associated with oxidative damage, causing direct cellular injury by inducing lipid and protein peroxidation and damaging nucleic acids.²⁰ The highly unsaturated fatty acids present in cellular membranes are the most susceptible macromolecules to oxidative damage in cells. Endothelial cells, macrophages, neutrophils, and neuronal cells generate superoxide (O_2^-) and NO, which can combine to form peroxynitrite anion ($ONOO^-$). Peroxynitrite, known to oxidize sulfhydryls and to yield products indicative of hydroxyl radical ($\cdot OH$) reaction with deoxyribose and dimethyl sulfoxide, is shown herein to induce membrane lipid peroxidation.²¹ Lipid peroxidation is a free-radical initiated chain reaction resulting in sequential abstraction of hydrogen ions from polyunsaturated fatty acids. Lipid peroxides, such as MDA, are markers and intermediate products of lipid peroxidation and are used for the assessment of tissue injury made by the free radicals produced by I/R.²²

Zhang et al²³ animal experiment showed that the amount of MDA, NO, and H_2O_2 can directly reflect the changes of the free radicals content in vivo in cerebral ischemic injury. The amount of MDA, NO, and H_2O_2 increased

significantly in serum and brain tissue after establishing the animal models and reduced after injecting picoside II intraperitoneally. It was suggested that picoside II can enhance the activity of endogenous antioxidant enzymes and clear the excess oxygen radicals to protect brain tissue from oxidative injury.

Wu et al²⁴ study showed that picoside II remarkably increased the amount of NO in ischemic myocardium. Therefore, it is speculated that picoside II improves postischemic cardiac functional recovery as a result of an increase in NO production, because NO can reinforce myocardial contractility.^{25,26}

Erythrocytes must be able to extend and curve and have the capability to move in end organ capillaries to deliver the oxygen and vital molecules to the tissues and clear the metabolic wastes via final organ capillaries. This ability is termed as “deformability” and gets more important in microcirculation. Altered erythrocyte deformability is not only important for oxygen delivery capacity of the erythrocytes but it is also critical for the survival of the circulating erythrocytes.^{27–29}

Wu et al²⁴ showed conclusively that picoside II pretreatment significantly increased the values of left ventricular developed pressure, positive first-order derivative of ventricular pressure ($+dp/dt$), negative first-order derivative of ventricular pressure ($-dp/dt$), and coronary flow, and decreased the size of infarction, which suggests that picoside II could produce a protective effect on myocardial ischemia-reperfusion injury in rats.

All these data and our findings indicate that erythrocyte deformability is impaired in rats subjected to I/R and this impairment leads to disturbance of microvascular perfusion and related problems. Hence, we think that measurement of erythrocyte deformability can be useful as a parameter in cases of I/R. We also observed beneficial effect of picoside II on maintaining erythrocyte deformability during periods of I/R, but we still think these promising results should further be supported by more detailed studies with larger numbers.

Acknowledgment

The abstract of our manuscript was accepted for poster presentation at the Turkish Society of Physiological Sciences, 41th National Physiology Congress 2015, *Acta Physiologica* 2015; Volume 215, Supplement S704, PC: 117, pp: 71.

Disclosure

The authors report no conflicts of interest in this work.

References

1. Duru S, Koca U, Oztekin S, et al. Antithrombin III pretreatment reduces neutrophil recruitment into the lung and skeletal muscle tissues in the rat model of bilateral lower limb and reperfusion: a pilot study. *Acta Anaesthesiol Scand*. 2005;49:1142–1148.
2. Turchanyi B, Toth B, Racz I, Vendég Z, Fűrész J, Hamar J. Ischemia reperfusion injury of skeletal muscle after selective deafferentation. *Physiol Res*. 2005;54:25–32.
3. Lin B, Ginsberg M, Busto R, Li L. Hyperglycemia triggers massive neutrophil deposition in brain following transient ischemia in rats. *Neurosci Lett*. 2000;278:1–4.
4. Homer-Vanniasinkam S, Crinnion JN, Gough MJ. Post-ischaemic organ dysfunction: a review. *Eur J Vasc Endovasc Surg*. 1997;14:195–203.
5. Welbourn CR, Goldman G, Paterson IS. Pathophysiology of ischaemia reperfusion injury: central role of the neutrophil. *Br J Surg*. 1991;78:651–655.
6. Baskurt OK, Temiz A, Meiselman HJ. Effect of superoxide anions on red blood cell rheologic properties. *Free Radic Biol Med*. 1998;24:102–110.
7. Koter M, Franiak I, Broncel M, Chojnowska-Jezierska J. Effects of simvastatin and pravastatin on peroxidation of erythrocyte plasma membrane lipids in patients with type 2 hypercholesterolemia. *Can J Physiol Pharmacol*. 2003;81:485–492.
8. Li JX, Li P, Tezuka Y, Namba T, Kadota S. Three phenylethanoid glycosides and an iridoid glycoside from *Picrorhiza scrophulariiflora*. *Phytochemistry*. 1998;48:537–542.
9. Cao Y, Liu JW, Yu YJ, et al. Synergistic protective effect of picoside II and NGF on PC12 cells against oxidative stress induced by H₂O₂. *Pharmacol Rep*. 2007;59:573–579.
10. Smit HF, Kroes BH, van den Berg AJ, et al. Immunomodulatory and anti-inflammatory activity of *Picrorhiza scrophulariiflora*. *J Ethnopharmacol*. 2000;73:101–109.
11. He LJ, Liang M, Hou FF, Guo ZJ, Xie D, Zhang X. Ethanol extraction of *Picrorhiza scrophulariiflora* prevents renal injury in experimental diabetes via anti-inflammation action. *J Endocrinol*. 2009;200:347–355.
12. Li Q, Li Z, Xu XY, Guo YL, Du F. Neuroprotective properties of picoside II in a rat model of focal cerebral ischemia. *Int J Mol Sci*. 2010;11:4580–4590.
13. Wang L, Liu XH, Chen H, et al. Picoside II protects rat kidney against ischemia/reperfusion-induced oxidative stress and inflammation by the TLR4/NF- κ B pathway. *Exp Ther Med*. 2015;9(4):1253–1258.
14. Arslan M, Comu FM, Işık B, Unal Y, Cekmen N, Kurtipek O. Effects of a general anaesthetic agent, propofol, on erythrocyte deformability. *Bratisl Med J*. 2010;111:126–128.
15. Esterbauer H, Cheeseman KH. Determination of aldehydic lipid peroxidation products: malonaldehyde and 4-hydroxynonenal. *Methods Enzymol*. 1990;186:407–421.
16. Cortas NK, Wakid NW. Determination of inorganic nitrate in serum and urine by a kinetic cadmium-reduction method. *Clin Chem*. 1990;36:1440–1443.
17. Li P, Matsunaga K, Yamakuni T, Ohizumi Y. Potentiation of nerve growth factor-action by picosides I and II, natural iridoids, in PC12D cells. *Eur J Pharmacol*. 2000;406:203–208.
18. Li T, Liu JW, Zhang XD, Guo MC, Ji G. The neuroprotective effect of picoside II from hu-huang-lian against oxidative stress. *Am J Chin Med*. 2007;35:681–691.
19. Gao H, Zhou YW. Inhibitory effect of picoside II on hepatocyte apoptosis. *Acta Pharmacol Sin*. 2005;26:729–736.
20. Halliwell B, Gutteridge JMC. *Free Radicals in Biology and Medicine*. 3rd ed. New York: Oxford University Press; 1999:246–350.
21. Radi R, Beckman JS, Bush KM, Freeman BA. Peroxynitrite-induced membrane lipid peroxidation: the cytotoxic potential of superoxide and nitric oxide. *Arch Biochem Biophys*. 1991;288:481–487.
22. Grisham MB, Granger DN. Free radicals: reactive metabolites of oxygen as mediators of postischemic reperfusion injury. In: Martson A, Bulkeley GB, Fiddian-Green RG, Haglund U, editors. *Splanchnic Ischemia and Multiple Organ Failure*. St Louis, MO: Mosby; 1989:135–144.
23. Zhang M, Fangfang P, Zhang R, et al. The antioxidant effect of picoside II and the optimizing of therapeutic dose and time window in cerebral ischemic injury in rats. *Merit Res J Pharm Pharm Sci*. 2013;1(1):1–7.
24. Wu N, Li W, Shu W, Jia D. Protective effect of picoside II on myocardial ischemia reperfusion injury in rats. *Drug Des Devel Ther*. 2014;8:545–554.
25. Kitakaze M, Node K, Minamino T, et al. Role of nitric oxide in regulation of coronary blood flow during myocardial ischemia in dogs. *J Am Coll Cardiol*. 1996;27(7):1804–1812.
26. Beresewicz A, Karwatowska-Prokopczuk E, Lewartowski B, Cedro-Ceremążyńska K. A protective role of nitric oxide in isolated ischaemic/reperfused rat heart. *Cardiovasc Res*. 1995;30(6):1001–1008.
27. Zinchuk VV. Erythrocyte deformability: physiological aspects. *Usp Fiziol Nauk*. 2001;32(3):66–78.
28. Kuypers FA. Red cell membrane damage. *J Heart Valve Dis*. 1998;7:387–395.
29. Sivilotti ML. Oxidant stress and haemolysis of the human erythrocyte. *Toxicol Rev*. 2004;23:169–188.

Drug Design, Development and Therapy

Publish your work in this journal

Drug Design, Development and Therapy is an international, peer-reviewed open-access journal that spans the spectrum of drug design and development through to clinical applications. Clinical outcomes, patient safety, and programs for the development and effective, safe, and sustained use of medicines are a feature of the journal, which

Submit your manuscript here: <http://www.dovepress.com/drug-design-development-and-therapy-journal>

has also been accepted for indexing on PubMed Central. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Dovepress