Integrating subpathway analysis to identify candidate agents for hepatocellular carcinoma

Jiye Wang¹
Mi Li²
Yun Wang³
Xiaoping Liu⁴

¹The Criminal Science and Technology Department, Zhejiang Police College, Hangzhou, Zhejiang Province,

²Department of Nursing, Shandong College of Traditional Chinese Medicine College, Yantai, Shandong Province,

³Office Department of Gastroenterology, The First Affiliated Hospital of Xi'an Jiao Tong University, Xi'an, Shanxi Province,

⁴Key Laboratory of Systems Biology, Shanghai Institutes for Biological Sciences, Shanghai, People's Republic of China

Abstract: Hepatocellular carcinoma (HCC) is the second most common cause of cancer-associated death worldwide, characterized by a high invasiveness and resistance to normal anticancer treatments. The need to develop new therapeutic agents for HCC is urgent. Here, we developed a bioinformatics method to identify potential novel drugs for HCC by integrating HCC-related and drug-affected subpathways. By using the RNA-seq data from the TCGA (The Cancer Genome Atlas) database, we first identified 1,763 differentially expressed genes between HCC and normal samples. Next, we identified 104 significant HCC-related subpathways. We also identified the subpathways associated with small molecular drugs in the CMap database. Finally, by integrating HCC-related and drug-affected subpathways, we identified 40 novel small molecular drugs capable of targeting these HCC-involved subpathways. In addition to previously reported agents (ie, calmidazolium), our method also identified potentially novel agents for targeting HCC. We experimentally verified that one of these novel agents, prenylamine, induced HCC cell apoptosis using 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide, an acridine orange/ethidium bromide stain, and electron microscopy. In addition, we found that prenylamine not only affected several classic apoptosis-related proteins, including Bax, Bcl-2, and cytochrome c, but also increased caspase-3 activity. These candidate small molecular drugs identified by us may provide insights into novel therapeutic approaches for HCC.

Keywords: subpathways, hepatocellular carcinoma, small molecules, network

Introduction

Hepatocellular carcinoma (HCC), the sixth most common cancer worldwide, accounts for 95% of primary liver cancer.¹ HCC is the second most common cause of cancer-associated death worldwide and was estimated to be responsible for ~746,000 deaths in 2012.^{2,3} If diagnosed early, HCC can be treated curatively with surgical resection or liver transplantation; however, since the majority of HCC patients are diagnosed at an advanced stage, their median survival times are generally <1 year, leading to a poor prognosis.^{4,5} Additionally, very few patients are candidates for liver resection, and liver donors are generally uncommon.² Furthermore, other modalities such as radiofrequency ablation and transarterial chemoembolization have poor efficacy.^{2,6} Therefore, special and effective chemotherapy for treating HCC is still needed.

However, the development of novel anticancer drugs is time consuming and risky. Therefore, drug repurposing, identifying new indications for known drugs, has become an effective and innovative approach. One strategy for drug repurposing is to distinguish and investigate pathways related to HCC development and then identify potential agents that may affect these pathways. As bioinformatics databases and methods have developed, high-throughput screening of HCC has become possible, providing global cancer genome information. One example is The Cancer Genome

Correspondence: Jiye Wang
The Criminal Science and Technology Department, Zhejiang Police College, 555 Binwen Road, Binjiang District, 310000 Hangzhou, Zhejiang Province, People's Republic of China
Tel/fax +86 571 8778 7220
Email wangjiye_poli@163.com

Atlas (TCGA) database, which is a community resource project, in which data are made available rapidly after generation for community research use. Additionally, databases allowing the investigation of gene signatures after small molecule treatments, such as the CMap database, have been constructed to provide microarray data under drug-induced conditions.⁷

To explore new agents for HCC, we implemented a bioinformatics method for drug repurposing of HCC based on integrating subpathway analyses. The differentially expressed genes (DEGs) between HCC tissues and nontumor tissues were identified from the TCGA database. Then, we identified HCC-related pathways. After treatment with small molecules in CMap, affected pathways were also identified. Finally, small molecule drugs that are capable of targeting these HCC-related subpathways were considered potential novel agents for HCC treatment. We experimentally verified that prenylamine, one of the potential agents we identified, could induce apoptosis of HepG2 (HCC cell line). Finally, we explored the apoptotic mechanisms of prenylamine by measuring the expression of apoptosis-related proteins, including Bax, Bcl-2, cytochrome *c*, and caspase-3 activity. In brief, the candidate agents identified by our approach may provide important insights in improving the therapeutic status of HCC cases.

Materials and methods

Microarray data and DEG analysis

We downloaded RNA-seq data from liver HCC samples from the TCGA database (<http://cancergenome.nih.gov/>). These data are based on human HCC tissues isolated from normal and HCC patients. A total of 371 HCC samples and 49 normal samples were sequenced on the Illumina Genome Analyzer RNA Sequencing platform. We downloaded the TCGA level 3 data for further analysis. We used edgeR, a Bioconductor software package for examining differential expression of replicated count data,⁸ to identify DEGs. The genes with a fold-change value >2 or <0.5 and with a false discovery rate (FDR) <0.01 were considered DEGs.

Identifying subpathways by SubpathwayMiner

HCC-related subpathways were identified using SubpathwayMiner an R-based software package that flexibly identifies subpathways using the Kyoto Encyclopedia of Genes and Genomes (KEGG) pathway structure information database.⁹ KEGG (<http://www.genome.jp/kegg/>) is a collection of online databases containing information on

genomes, enzymatic pathways, and biological chemicals. In the SubpathwayMiner package, the pathway structure data of KEGG are converted to undirected R graph objects, in which enzymes are considered nodes; and if the two enzymes participate in the same reactions, two nodes are connected by an edge. To identify subpathways, in which all enzymes have similar functions, from entire pathways, the *k*-clique concept in social network analysis was adopted. This software can define subpathways based on distance similarity among enzymes.⁹ In social network analysis, a *k*-clique in a graph is defined as a subgraph, in which the distance between any two nodes is not greater than a parameter, specified as *k*.⁹ Here, we set $k=3$, a setting at which the distance among enzymes in one subpathway is not >3 .

We then imported the DEGs in HCC into SubpathwayMiner to identify significantly enriched subpathways. An FDR <0.1 was considered statistically significant.

Global associations between drugs and metabolic subpathways

First, we identified the global association between 1,309 bioactive small molecular agents from the CMap database and the identified subpathways from the SubpathwayMiner.⁷ We downloaded microarray profiles for each agent from the CMap database and obtained the affected genes for each small molecule. The CMap database contains 6,100 instances corresponding to 1,309 bioactive small molecules. Each instance includes treatment and control gene expression profiles, corresponding to treatment with or without a bioactive small molecule administered to the cells. We downloaded all the microarray profiles and their associated annotation files from the CMap website (<http://www.broadinstitute.org/cmap/>). For each instance, we matched perturbation and control pairs of expression profiles according to descriptions of the instances in the annotation file. Then, we used fold-change analysis to identify DEGs, for each instance with a fold change >2 or <0.5 , between the corresponding treatment and control gene expression profiles. The DEGs were merged if the corresponding instances (experiments) belonged to the same bioactive small molecule, and these genes were considered to be affected genes for this agent.

Next, we entered the affected genes for each small molecule into SubpathwayMiner to identify significantly enriched subpathways for each small molecule. An FDR <0.1 was chosen as the cutoff for statistically significant subpathways.

Identification of novel drugs for HCC

To find novel agents, which may play a role in HCC-involved subpathways, we integrated the aforementioned global associations between drugs and subpathways and HCC-involved subpathways. If drug-related subpathways were also involved in HCC, the corresponding drugs were considered potential drugs for the treatment of HCC.

HepG2 cell culture

The human hepatoma HepG2 cell line was purchased from the Cell Bank of the Chinese Academy of Sciences (Shanghai, People's Republic of China) and cultured in Dulbecco's Modified Eagle's Medium containing 10% (v/v) heat-inactivated fetal bovine serum at 37°C in an incubator containing humidified air with 5% (v/v) CO₂. No ethics statement was required from the institutional review board for the use of these cell lines.

Cell viability assay

Cells were seeded onto 96-well plates at 8×10^3 cells per well 24 hours before treatment. The cells were treated with the prenylamine (at 0, 10, 50, 100, 200, and 500 $\mu\text{mol/L}$). After 12 hours, 15 μL (5 mg/mL) 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT)¹⁰ (Sigma-Aldrich Co., St Louis, MO, USA) was added to each well, followed by incubation at 37°C for 4 hours. Then, the MTT solution was removed, and 150 μL dimethyl sulfoxide was added. The mixtures were shaken for 10 minutes to fully dissolve the crystals. A microplate reader (Tecan) was used to measure the absorbance at a wavelength of 570 nm. Cell viability was expressed as percentage changes in the absorbance values from the treatment group over the control group.

Electron microscopy

HepG2 cells were cultured in 60 mm plates, collected in phosphate-buffered saline (PBS) solution, and fixed with 2% (v/v) paraformaldehyde containing 2.5% (w/v) glutaraldehyde (Paesel & Lorei) buffered in Hank's modified salt solution at 4°C for 4 hours. The cells were further fixed in 1% (w/v) OsO₄ solution buffered by 0.1 M cacodylate (pH 7.2) at 4°C for 2 hours, after which the cells were scraped off from the plastic and dehydrated in ethanol. Dehydration was completed in propylene oxide. The specimens were embedded in Araldite (Serva). Ultrathin sections were produced on an FCR Reichert Ultracut ultra microtome (Leica Microsystems), mounted on pioloform-coated copper grids, and contrasted with lead citrate. Specimens were analyzed and documented with an EM 10A electron microscope (Carl Zeiss Meditec AG).

Acridine orange/ethidium bromide fluorescence staining

HepG2 cells in the exponential growth phase were cultivated on sterile coverslips for 24 hours and were subsequently treated with 200 $\mu\text{mol/L}$ for 12 hours. The cells were washed twice with PBS and then mixed with 1 mL of dye mixture containing 100 mg/mL acridine orange (AO) and 100 mg/mL ethidium bromide (EB) in PBS.¹¹ Cellular morphological changes were examined using fluorescence microscopy ($\times 200$). The percentage of apoptotic cells was calculated by the following formula: apoptotic rate (%) = number of apoptotic cells/number of all cells counted.^{12,13}

Cell apoptosis assays

Double staining with Annexin V-FITC and propidium iodide (BD, Franklin Lakes, NJ, USA) was performed for quantification of apoptosis of HepG2 cells. Both the attached cells and the cells in the supernatant were collected, washed twice with ice-cold PBS, and resuspended in 400 mL of binding buffer. Annexin V-FITC and propidium iodide were added to the cells and incubated for 15 minutes in the dark at 4°C. After incubation, the samples were analyzed by a flow cytometry (BD FACSCalibur; BD), and 2.0×10^4 events per sample were counted.

Western blotting

Total protein was extracted from the HepG2 cells for immunoblotting analysis. Protein samples (80 μg of protein) were separated in a 12% sodium dodecyl sulfate polyacrylamide gel electrophoresis gel and blotted onto nitrocellulose membranes. After blotting, the membranes were probed with Bax, Bcl-2, cytochrome *c*, and glyceraldehyde-3-phosphate dehydrogenase antibodies and incubated overnight at 4°C. Infrared fluorescent dye-labeled secondary antibody was incubated with the membrane for 1 hour. Western blot bands were collected using the infrared Imaging System (LI-COR Biosciences), and the band density was quantified using Odyssey 3.0 software for each group and normalized against glyceraldehyde-3-phosphate dehydrogenase.

Caspase-3 activity assay

Caspase-3 activity was analyzed using a caspase-3 activity assay kit (Beyotime Institute of Biotechnology, Shanghai, People's Republic of China) according to the manufacturer's instructions, using the substrate peptides Ac-DEVD-p-nitroanilide (pNA), Ac-IETD-pNA, and Ac-LEHD-pNA.

Briefly, the supernatant of the cell lysate was mixed with buffer containing the substrate peptides for caspase attached to pNA. The release of pNA was quantified by determining the absorbance with an enzyme-linked immunosorbent assay reader at 405 nm. Caspase activities were expressed as percentage over control.

Result

DEGs between HCC cases and healthy controls

In order to explore the mechanisms underlying HCC, we obtained the expression profiles of HCC samples and normal samples from the TCGA database. We identified a total of 1,763 genes as differentially expressed between HCC patients and controls ("Materials and methods" section).

Subpathway enrichment analysis

Using SubpathwayMiner with a *k* value set to 3, we identified 104 enriched subpathways corresponding to 42 entire pathways with an FDR <0.1 (Tables 1 and S1). These 104 metabolic subpathways and the corresponding entire pathways were considered significantly related to the

development of HCC. Furthermore, we identified global associations between 1,309 bioactive small molecular agents from the CMap database and these subpathways. Using a cutoff of FDR <0.01, a total of 840 associations between 196 small molecules and the 104 subpathways were obtained (Table S2).

Drug repurposing for HCC

To identify candidate agents that are capable of targeting HCC-related subpathways, we integrated the aforementioned global associations between drugs and metabolic subpathways ("Materials and methods" section).¹⁴ By integrating these 104 HCC-related subpathways and small molecules that affected these subpathways, we selected the overlapping subpathways that are related to both HCC and small molecules. Thus, we assumed that the small molecules, which affected the overlapping subpathway, might also play a role in perturbing the development of HCC. There were eight overlapping subpathways corresponding to a total of 40 small molecules (Table 2).

We then built a network between these repositioned drugs and the overlapping metabolic subpathways by integrating the

Table 1 The enriched subpathways and the corresponding entire pathways

Pathway ID	Pathway name	Subpathways
Path: 04920	Adipocytokine signaling pathway	Path: 04920_5
Path: 00250	Alanine, aspartate, and glutamate metabolism	Path: 00250_1
Path: 05146	Amebiasis	Path: 05146_9; path: 05146_2; path: 05146_1; path: 05146_7
Path: 00590	Arachidonic acid metabolism	Path: 00590_1; path: 00590_2; path: 00590_3; path: 00590_4
Path: 00330	Arginine and proline metabolism	Path: 00330_12; path: 00330_11; path: 00330_10
Path: 04360	Axon guidance	Path: 04360_19
Path: 00410	Beta-alanine metabolism	Path: 00410_5; path: 00410_4
Path: 00650	Butanoate metabolism	Path: 00650_8; path: 00650_7; path: 00650_4; path: 00650_13; path: 00650_3
Path: 04020	Calcium signaling pathway	Path: 04020_3
Path: 04514	CAMs	Path: 04514_81; path: 04514_84
Path: 04110	Cell cycle	Path: 04110_23; path: 04110_17; path: 04110_3; path: 04110_12
Path: 04062	Chemokine signaling pathway	Path: 04062_2; path: 04062_1; path: 04062_13
Path: 04610	Complement and coagulation cascades	Path: 04610_11; path: 04610_6; path: 04610_4
Path: 00270	Cysteine and methionine metabolism	Path: 00270_2; path: 00270_8; path: 00270_1
Path: 05414	Dilated cardiomyopathy	Path: 05414_1
Path: 00982	Drug metabolism – cytochrome P450	Path: 00982_4; path: 00982_5; path: 00982_9; path: 00982_12; path: 00982_10
Path: 04512	ECM–receptor interaction	Path: 04512_11; path: 04512_13; path: 04512_14; path: 04512_12; path: 04512_3; path: 04512_27; path: 04512_10; path: 04512_9; path: 04512_5; path: 04512_6
Path: 00071	Fatty acid metabolism	Path: 00071_11; path: 00071_2; path: 00071_7; path: 00071_8; path: 00071_4; path: 00071_9; path: 00071_10; path: 00071_3; path: 00071_5; path: 00071_6; path: 00071_1
Path: 04510	Focal adhesion	Path: 04510_1
Path: 00790	Folate biosynthesis	Path: 00790_4; path: 00790_6
Path: 00010	Glycolysis/gluconeogenesis	Path: 00010_3
Path: 00591	Linoleic acid metabolism	Path: 00591_1
Path: 04010	MAPK signaling pathway	Path: 04010_2; path: 04010_11
Path: 00980	Metabolism of xenobiotics by cytochrome P450	Path: 00980_3; path: 00980_1

(Continued)

Table 1 (Continued)

Pathway ID	Pathway name	Subpathways
Path: 00910	Nitrogen metabolism	Path: 00910_3
Path: 04114	Oocyte meiosis	Path: 04114_16; path: 04114_11; path: 04114_21
Path: 05200	Pathways in cancer	Path: 05200_52; path: 05200_14
Path: 04145	Phagosome	Path: 04145_1
Path: 03320	PPAR signaling pathway	Path: 03320_2
Path: 04914	Progesterone-mediated oocyte maturation	Path: 04914_4
Path: 00640	Propanoate metabolism	Path: 00640_5
Path: 05215	Prostate cancer	Path: 05215_7
Path: 00830	Retinol metabolism	Path: 00830_2
Path: 04970	Salivary secretion	Path: 04970_1; path: 04970_2
Path: 05222	Small cell lung cancer	Path: 05222_1
Path: 00140	Steroid hormone biosynthesis	Path: 00140_11
Path: 04350	TGF-beta signaling pathway	Path: 04350_13
Path: 04530	Tight junction	Path: 04530_10
Path: 04930	Type II diabetes mellitus	Path: 04930_5
Path: 00350	Tyrosine metabolism	Path: 00350_2; path: 00350_3; path: 00350_4
Path: 00280	Valine, leucine, and isoleucine Degradation	Path: 00280_1; path: 00280_2; path: 00280_8; path: 00280_3; path: 00280_7; path: 00280_10; path: 00280_9
Path: 04270	Vascular smooth muscle contraction	Path: 04270_9; path: 04270_8

Abbreviations: CAMs, cell adhesion molecules; ECM, extracellular matrix; MAPK, mitogen-activated protein kinases; PPAR, peroxisome proliferator-activated receptor; TGF, transforming growth factor.

Table 2 Potential novel drugs and their information

Drug name	ATC	Detailed information
5248896		
Docosahexaenoic acid ethyl ester		
Calmidazolium		
Mefloquine	P01BC02	Methanolquinolines
Dyclonine	N01BX02	Other local anesthetics
Oxyphenbutazone	M02AA04	Anti-inflammatory
5182598		
Quercetin		
C-75		
5194442		
Nifuroxazide	A07AX03	Antibacterial
Ebselen		
F0447-0125		
Rottlerin		
Myricetin		
Pizotifen	N02CX01	Serotonin inhibitor
Alclometasone	D07AB10	Corticosteroids, moderately potent (group II)
Alprostadil	C01EA01	Prostaglandins
Amphotericin B	A01AB04	Anti-infectives and antiseptics for local oral treatment
Suloctidil	C04AX19	Vasodilator
Vinburnine	C04AX17	Vasodilator, cerebral
Levomopromazine	N05AA02	Analgesic
Chloropyramine	D04AA09	Antihistamines for topical use
Mianserin	N06AX03	Other antidepressants
Triprolidine	R06AX07	Other antihistamines for systemic use
Profenamine	N04AA05	Antiparkinsonian
Prenylamine	C01DX02	Vasodilator
Anisomycin		
Metirapone	V04CD01	Tests for pituitary function
Thioguanosine		
Alexidine		Antibacterial
Parthenolide		
MG-262		
Trifluridine	S01AD02	Antivirals
Niclosamide	P02DA01	Salicylic acid derivatives
Scriptaid		
Pyrvinium	P02CX01	Anthelmintic
Ionomycin		
MG-132		
0297417-0002B		

Abbreviation: ATC, anatomical therapeutic chemical classification.

relationships mentioned earlier (Figure 1). In this network, the rectangle nodes represent drugs and the circle nodes represent subpathways. The subpathways are color coded by the entire pathway in which they participate (Figure 1).

Of 40 small molecules, prenylamine was chosen for next validation not only because of its safety and known mode of action but also due to its apoptosis potential. It was always used as antianginal drug for decades.¹⁵ This class of medications mimicked the cardiac effects of simple Ca^{2+} withdrawal in that they diminished Ca^{2+} -dependent high energy phosphate

utilization, contractile force, and oxygen requirement of the beating heart without impairing the Na-dependent action potential parameters.¹⁵ Furthermore, several studies had reported the pivotal role of Ca^{2+} homeostasis in cell apoptosis.¹⁶ Therefore, we next chose prenylamine to treat HepG2 cells.

Prenylamine suppresses the viability of HepG2 cells

The antiproliferative effect of prenylamine on HepG2 cells was examined by exposing the cells to various concentrations

Figure 1 Small molecular drug and their perturbed subpathways in HCC.

Notes: The rectangle nodes represent drugs, and the circle nodes represent subpathways. The subpathways in the same color are included in same entire pathways. The description of each corresponding entire pathway is mentioned below the network.

Abbreviation: HCC, hepatocellular carcinoma.

(0, 10, 50, 100, 200, and 500 $\mu\text{mol/L}$) of prenylamine for 12, 24, and 48 hours (Figure S1). We found that cell growth was inhibited in a dose-dependent manner (Figure 2A). In the presence of 200 $\mu\text{mol/L}$ prenylamine, HepG2 cells exhibited ~50% inhibition of proliferation after treatment for 12 hours. Hence, this concentration and a 12-hour treatment time were used in the following experiments.

Prenylamine induces apoptosis in HepG2 cells

To investigate whether prenylamine regulates apoptosis, AO/EB staining and electron microscopy were used to detect the apoptotic cells. The results from our fluorescence microscopy analysis are shown in Figure 2B. Prenylamine induced substantial cell apoptosis ($P < 0.05$). Under an electron

Figure 2 Prenylamine induced HepG2 cell death.

Notes: (A) HepG2 cells were treated with the prenylamine at the concentrations of 10, 50, 100, 200, and 500 $\mu\text{mol/L}$ for 12 hours. Relative cell viability was determined by the MTT assay. * $P < 0.05$ compared with controls; $n = 6$ independent experiments for each condition. (B) AO/EB staining to detect changes in the nucleus. A statistical bar graph of apoptotic cells by AO/EB staining is shown. Arrows indicate apoptotic cells. * $P < 0.05$ compared with controls; $n = 3$ independent experiments for each group. (C) Transmission electron microscopy to estimate micromorphological changes (magnification: $\times 8,000$).

Abbreviations: MTT, 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide; AO/EB, acridine orange/ethidium bromide; Ctrl, control; mag, magnification; HV, high voltage.

microscope, the cells treated with prenylamine exhibited robust changes in microstructure, including cell surface microvillus reduction, nuclear chromatin condensation, margination, and membrane blistering¹⁰ (Figure 2C). In addition, apoptotic body was found (the white arrow in Figure 2C).

We also determined whether prenylamine affected HCC cell apoptosis using a flow cytometry. After being treated with prenylamine for 12 hours, the late apoptosis of HepG2 cells was higher than the control group (Figure S2A and B). These data implied that prenylamine induces apoptosis of HCC cells.

Prenylamine activates proapoptotic signaling pathways

To explore the mechanisms by which prenylamine induced apoptosis in HepG2 cells, we measured the expression of downstream proteins in apoptotic pathways, including Bax, Bcl-2, and cytochrome *c*. The results demonstrated that treatment with prenylamine could upregulate Bax

and cytochrome *c* and downregulate Bcl-2 expression (Figure 3A and B). In addition, caspase-3 activity, as a marker of apoptosis, was increased (1.9-fold) by prenylamine (Figure 3C).

Discussion

Analysis of gene expression in disease can reveal the underlying gene activity changes, which contribute to the disease and can pinpoint potential targets for improving therapeutic intervention. In this study, RNA-seq data from the TCGA database were used to explore the mechanisms of HCC development. A total of 1,763 DEGs were identified between HCC samples and healthy controls. The subpathway mining results showed 104 metabolic subpathways corresponding to 42 entire pathways that were dysfunctional in HCC. Then, we identified potential therapeutic agents for HCC using a computational drug repurposing method. A total of 40 small molecules were identified that are potential drug candidates for perturbing the development of HCC.

Figure 3 Prenylamine alters Bax, Bcl-2, and cytochrome *c* expression and promotes caspase-3 activation.

Notes: Western blot was used to detect (A) Bax and Bcl-2 expression and (B) cytochrome *c* expression in HepG2 cells treated with prenylamine (200 μmol/L, 12 h). Relative expression of Bax, Bcl-2, and cytochrome *c* was normalized to GAPDH; n=3 independent experiments for each group. (C) Activation of caspase-3 by prenylamine. Data are averaged from five independent experiments for each group. Similar results were observed from another three experiments. **P*<0.05 compared with controls.

Abbreviations: h, hours; GAPDH, glyceraldehyde-3-phosphate dehydrogenase; Ctrl, control; AMC, Ac-DEVD-pNA(acetyl-Asp-Glu-Val-Asp p-nitroanilide).

The majority of current pathways enrichment approaches commonly focus on entire pathways. However, a previous study showed that subpathway analysis may be more suitable than entire pathways for the identification of drug responses.⁹ In this study, we identified 104 subpathways among 42 KEGG pathways. Some of these subpathways are well documented as involved in the development of HCC. For example, the top three significant subpathways belong to the extracellular matrix–receptor interaction pathway (path: 04512; $P=3.54\times10^{-11}$; FDR $=8.16\times10^{-8}$; Tables 1 and S1), which consists of molecules such as CD36, CD44, and CD104 that serve an important role in the direct or indirect control of cellular activities such as adhesion, migration, differentiation, proliferation, and apoptosis.¹⁷ Molecules in this pathway have been shown to drive pathological angiogenesis and tumor progression.¹⁸ The cell cycle pathway (path: 04110_17; $P=5.59\times10^{-05}$; FDR $=0.003$), referring to the reproducible sequence of events in the G1 and G2 phases of the cell cycle, was also detected. Key regulatory enzymes of this pathway are cyclin-dependent kinases, which are dysregulated and serve as potential therapeutic targets in HCC.^{19,20} Additionally, we also identified other cancer-related subpathways, such as the peroxisome proliferator-activated receptor signaling pathway (path: 03320) and pathways in cancer (path: 05200).

We then identified a group of known drugs with potential novel therapeutic efficacy for HCC by integrating the association between small molecules and subpathways. A study by Li et al also identified global associations between 403 metabolic subpathways and 488 known drugs using SubpathwayMiner with $k=3$.²¹ However, in that study, only metabolic subpathways were investigated rather than all kinds of pathways.²¹ Some signaling pathways also play an important role in drug response. Therefore, in our study, we implemented subpathway enrichment to all kinds of pathways. Of 104 drug-affected subpathways, eight subpathways were also involved in the development of HCC. These overlapping subpathways corresponded to 40 small molecules.

Some of these small molecules were found to have apoptotic effects. For example, calmidazolium induced apoptosis of human hepatoma cells,²² and niclosamide inhibited proliferation as well as inducing apoptosis and cell cycle arrest in human osteosarcoma cell lines.²³ However, the effects of some other drugs are not well characterized. Prenylamine, a calcium antagonist, has been used for the treatment of angina pectoris.²⁴ Although some early studies have investigated its effects on a colon cancer cell line,¹⁴ there have been no direct reports about the apoptotic effects of prenylamine on HCC cells. To further investigate the effects of prenylamine

on HCC, we treated HepG2 cells with different concentrations (10, 50, 100, 200, and 500 $\mu\text{mol/L}$) of prenylamine. The MTT assay indicated that prenylamine could induce HepG2 cell apoptosis in a slightly dose-dependent manner. AO/EB staining and electron microscopy also detected the apoptotic cells. Finally, we found that prenylamine not only affected some classic apoptosis-related proteins, including Bax, Bcl-2, and cytochrome *c*, but also increased caspase-3 activity, suggesting that prenylamine might regulate multiple potential targets, directly or indirectly influencing these proteins. Although our approach based on targeting subpathways successfully identified potential agents for HCC, further experiments are needed to explore the mechanisms underlying these observations and to ensure the safety of these small molecules for use in HCC clinical treatment.

Disclosure

The authors alone are responsible for the content and writing of the paper. The authors report no conflicts of interest in this work.

References

- Deng GL, Zeng S, Shen H. Chemotherapy and target therapy for hepatocellular carcinoma: new advances and challenges. *World J Hepatol*. 2015;7(5):787–798.
- Hong YP, Li ZD, Prasoon P, Zhang Q. Immunotherapy for hepatocellular carcinoma: from basic research to clinical use. *World J Hepatol*. 2015;7(7):980–992.
- Lau WY. Primary liver tumors. *Semin Surg Oncol*. 2000;19(2):135–144.
- Bruix J, Sherman M; American Association for the Study of Liver Diseases. Management of hepatocellular carcinoma: an update. *Hepatology*. 2011;53(3):1020–1022.
- Giakoustidis A, Giakoustidis D, Mudan S, Sklavos A, Williams R. Molecular signalling in hepatocellular carcinoma: role of and crosstalk among WNT/ss-catenin, Sonic Hedgehog, Notch and Dickkopf-1. *Can J Gastroenterol Hepatol*. 2015;29(4):209–217.
- Pez F, Lopez A, Kim M, Wands JR, Caron de Fromentel C, Merle P. Wnt signaling and hepatocarcinogenesis: molecular targets for the development of innovative anticancer drugs. *J Hepatol*. 2013;59(5):1107–1117.
- Lamb J, Crawford ED, Peck D, et al. The Connectivity Map: using gene-expression signatures to connect small molecules, genes, and disease. *Science*. 2006;313(5795):1929–1935.
- Robinson MD, McCarthy DJ, Smyth GK. edgeR: a Bioconductor package for differential expression analysis of digital gene expression data. *Bioinformatics*. 2010;26(1):139–140.
- Li C, Li X, Miao Y, et al. SubpathwayMiner: a software package for flexible identification of pathways. *Nucleic Acids Res*. 2009;37(19):e131.
- Zheng F, He K, Li X, et al. Transient overexpression of TGFBR3 induces apoptosis in human nasopharyngeal carcinoma CNE-2Z cells. *Biosci Rep*. 2013;33(2):e00029.
- McGahon AJ, Martin SJ, Bissonnette RP, et al. The end of the (cell) line: methods for the study of apoptosis in vitro. *Methods Cell Biol*. 1995;46:153–185.
- Chen H, Takahashi S, Imamura M, et al. Earthworm fibrinolytic enzyme: anti-tumor activity on human hepatoma cells in vitro and in vivo. *Chin Med J*. 2007;120(10):898–904.

13. Ribble D, Goldstein NB, Norris DA, Shellman YG. A simple technique for quantifying apoptosis in 96-well plates. *BMC Biotechnol.* 2005;5:12.
14. Batra S, Alenfall J. Effect of diverse categories of drugs on human colon tumour cell proliferation. *Anticancer Res.* 1991;11(3):1221–1224.
15. Fleckenstein A. History of calcium antagonists. *Circ Res.* 1983; 52(2 pt 2):I3–I16.
16. Zhang S, Chen Y, Wu X, et al. The pivotal role of Ca homeostasis in PBDE-47-induced neuronal apoptosis. *Mol Neurobiol.* Epub 2015 Dec 17.
17. Nistico P, Di Modugno F, Spada S, Bissell MJ. beta1 and beta4 integrins: from breast development to clinical practice. *Breast Cancer Res.* 2014;16(5):459.
18. Giancotti FG. Targeting integrin beta4 for cancer and anti-angiogenic therapy. *Trends Pharmacol Sci.* 2007;28(10):506–511.
19. Li W, Long J, Takikawa Y, Suzuki K, Lin S. Plasma from patients with acute liver failure dampens HepG2 cells to epidermal growth factor induced proliferation response. *Hepatogastroenterology.* 2014; 61(135):2021–2027.
20. Zhao H, Bauzon F, Bi E, et al. Substituting threonine 187 with alanine in p27Kip1 prevents pituitary tumorigenesis by two-hit loss of Rb1 and enhances humoral immunity in old age. *J Biol Chem.* 2015;290(9): 5797–5809.
21. Li C, Shang D, Wang Y, et al. Characterizing the network of drugs and their affected metabolic subpathways. *PLoS One.* 2012;7(10): e47326.
22. Liao WC, Huang CC, Cheng HH, et al. Effect of calmidazolium on [Ca²⁺]_i and viability in human hepatoma cells. *Arch Toxicol.* 2009; 83(1):61–68.
23. Li Z, Yu Y, Sun S, Wang W, Yu A. Niclosamide inhibits the proliferation of human osteosarcoma cell lines by inducing apoptosis and cell cycle arrest. *Oncol Rep.* 2015;33(4):1763–1768.
24. Beuck S, Sigmund G, Koch A, et al. Identification and characterization of urinary prenylamine metabolites by means of liquid chromatography-tandem mass spectrometry. *Drug Test Anal.* 2012;4(9):701–716.

OncoTargets and Therapy

Publish your work in this journal

OncoTargets and Therapy is an international, peer-reviewed, open access journal focusing on the pathological basis of all cancers, potential targets for therapy and treatment protocols employed to improve the management of cancer patients. The journal also focuses on the impact of management programs and new therapeutic agents and protocols on

Submit your manuscript here: <http://www.dovepress.com/oncotargets-and-therapy-journal>

patient perspectives such as quality of life, adherence and satisfaction. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Dovepress