Circulating interleukin-10 levels and human papilloma virus and Epstein–Barr virus-associated cancers: evidence from a Mendelian randomization meta-analysis based on 11,170 subjects

Kai Qu,^{1,*} Qing Pang,^{1,*} Ting Lin,¹
Li Zhang,² Mingliang Gu,³ Wenquan
Niu,⁴ Chang Liu,^{1,*} Ming Zhang⁵

¹Department of Hepatobiliary Surgery, The First Affiliated Hospital of Xi'an Jiaotong University College of Medicine, Xi'an, Shaanxi Province, People's Republic of China; ²Department of Ultrasound Diagnostics, Tangdu Hospital, Fourth Military Medical University, Xi'an, Shaanxi Province, People's Republic of China; ³Chinese Academy of Sciences Key Laboratory of Genome Sciences and Information, Beijing Institute of Genomics, Chinese Academy of Sciences, Beijing, People's Republic of China; ⁴State Key Laboratory of Medical Genomics, Ruijin Hospital, Shanghai Jiaotong University School of Medicine, Shanghai, People's Republic of China; ⁵Department of Otolaryngology, Eye, Ear, Nose and Throat Hospital, Fudan University, Shanghai, People's Republic of China

*These authors contributed equally to this work

Correspondence: Wenquan Niu
State Key Laboratory of Medical Genomics, Ruijin Hospital, Shanghai Jiaotong University School of Medicine, Ruijin Second Road 197, New Huangpu District, Shanghai 200025, People's Republic of China
Tel +86 21 6437 0045
Email niuwenquan_shcn@163.com

Ming Zhang
Department of Otolaryngology, Eye, Ear, Nose and Throat Hospital, Fudan University, 83 Fen Yang Road, Shanghai 200031, People's Republic of China
Tel +86 21 6437 7134
Email ent_zhm@126.com

Abstract: Recent studies have showed interleukin 10 (IL-10) is a critical cytokine that determines antiviral immune response and is related to virus-associated cancers. However, whether genetically elevated circulating IL-10 levels are associated with the risk of human papilloma virus and Epstein–Barr virus-associated cancers (HEACs) is still unclear. Mendelian randomization method was implemented to meta-analyze available observational studies by employing *IL-10* three variants (–592C>A, –819C>T, and –1082A>G) as instruments. A total of 24 articles encompassing 11,170 subjects were ultimately eligible for the meta-analysis. Overall, there was a significant association between *IL-10* promoter variant –1082A>G and HEACs under allelic and dominant models (both $P<0.01$). Subgroup analysis by cancer type indicated that the risk estimate of –1082A>G was significant for nasopharyngeal cancer under allelic, homozygous genotypic and dominant models (all $P<0.001$). Moreover by ethnicity, carriers of –1082G allele had a 74% increased risk for nasopharyngeal cancer in Asians under dominant model (odds ratio [OR]=1.737; 95% confidence interval [CI]: 1.280–2.358; $P<0.001$). In further Mendelian randomization analysis, the predicted OR for 10 pg/mL increment in IL-10 levels was 1.14 (95% CI: 1.01–16.99) in HEACs. Our findings provided strong evidence for a critical role of genetically elevated circulating IL-10 levels in the development of HEACs, especially in Asian population and for nasopharyngeal cancer.

Keywords: interleukin-10, human papilloma virus, Epstein–Barr virus, meta-analysis, Mendelian randomization

Introduction

There is growing recognition that viruses are capable of causing cancer in humans, and approximately 15% of all human malignancies are estimated to be attributable to viruses, creating a major global health burden.¹ Among various cancer viruses, human papilloma virus (HPV) and Epstein–Barr virus (EBV) are exhaustively investigated and are considered to account for 38% of all virus-associated cancers.² HPV infection is associated with more than 90%³ and 60%^{4,5} cases of cervical and oropharyngeal cancers, respectively, and in contrast, EBV is associated with nearly 90% of nasopharyngeal cancer,^{6,7} Burkitt's,⁸ and Hodgkin's lymphoma.^{5,7,9} Moreover, the two viruses could also exhibit synergistic or cooperative effects on carcinogenesis. For example, EBV is deemed as a “helper virus” for HPV-induced carcinogenesis.¹⁰ Coinfection of HPV and EBV was observed in 30%–50% of patients with oral cancer^{11,12} and cervical cancer.^{13,14} Although the obvious association between HPV/EBV and human papilloma virus and

Epstein–Barr virus-associated cancers (HEACs) has been universally accepted, the inherited procancer mechanisms so far remain unclear. It should be pointed out that the infection of HPV and EBV affects more than 40% of general population; however, only a small proportion of infected cases develop cancer.¹⁵ Interindividual differences of innate antiviral immunity that is affected by hereditary factors might be involved in the underlying pathological process of HEACs.¹⁶

Among the antiviral-relevant immune factors, interleukin-10 (IL-10) is a key cytokine that determines viral clearance or persistence¹⁷ and is involved in carcinogenesis.^{18,19} Recent studies have observed a marked high level of circulating IL-10 in patients with HEACs and its association with poor prognosis.^{20,21} Emerging evidence suggested that interindividual differences in circulating IL-10 levels might be due to the polymorphic defects of *IL-10*. Recently, three variants located within *IL-10* promoter region, viz, $-592C>A$ (rs1800872), $-819C>T$ (rs1800871), and $-1082A>G$ (rs1800896), have been well defined in association with the changes of IL-10 production.^{22,23} It is, therefore, reasonable to hypothesize that if IL-10 is involved in the carcinogenesis process of HEACs, the inherited genetic determinants that alter IL-10 production should affect cancer susceptibility in the direction and magnitude predicted by its circulating levels.

Mendelian randomization approach, which is based on Mendel's second law, uses measured variation in genes of known function to examine the effect of a modifiable exposure on disease in observational studies.^{24,25} This method could partially provide evidence for the causal nature of the target phenotype influenced by genetic defects. To test the hypothesis that genetically elevated levels of IL-10 due to *IL-10* genetic defects cause an increased risk of cancer, in this study, we first decided to perform a meta-analysis to evaluate the association of *IL-10* three variants with both circulating IL-10 levels and the risk for HEACs. If the variant under study is found to be predictive of both cancer and circulating IL-10 levels, Mendelian randomization approach will be further implemented to test the possible association of circulating IL-10 levels with HEACs.

Materials and methods

This meta-analysis was undertaken according to the guidelines set forth by the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) statement (Table S1).

Search strategy

A literature search for observational studies investigating the association between *IL-10* three variants ($-592C>A$

[rs1800872], $-819C>T$ [rs1800871], and $-1082A>G$ [rs1800896]) and all types of HEACs was conducted of PubMed and Google Scholar databases covering the period from the earliest possible year to August 1, 2015. Subject terms used for the search were: “interleukin 10”, “interleukin-10”, “IL 10”, “IL-10”, “oral or mouth cancer”, “nasopharyngeal cancer”, “oropharyngeal cancer”, “Hodgkin or Burkitt lymphoma”, “cervical or vaginal or vulvar cancer”, “anus or anal cancer”, combined with “polymorphism”, “genetic”, “variant”, “mutation”, “allele”, or “genotype”. The reference lists of all the retrieved articles as well as those of reviews on the same topic were also searched to supplement the additional missing articles. Searching results was limited to studies with a case–control design and articles published in the English language.

Trial selection

Two investigators (Kai Qu and Ming Zhang) independently obtained the full texts of potentially eligible articles based on the titles and abstracts. If necessary, we emailed the corresponding authors to avoid double counting of participants recruited in more than one publication. In case of more than one publication from the same study population, we abstracted data from the most recent or most complete publication.

Inclusion/exclusion criteria

For inclusion, the studies should strictly fulfill the following inclusion criteria (all points must be satisfied for inclusion): 1) clinical endpoint (dependent variable): HEACs including oropharyngeal cancer, nasopharyngeal cancer, cervical cancer, Hodgkin's lymphoma, and Burkitt's lymphoma; 2) study design: either retrospective or prospective case–control design; and 3) independent variables: the genotype and/or allele counts of at least one of *IL-10* three variants ($-592C>A$, $-819C>T$, and $-1082A>G$). Studies were excluded (one point was sufficient for exclusion) if they investigated the gene function, disease progression, severity, and the response to treatment or survival. Additionally, conference abstracts, case reports or series, editorials, narrative reviews, meta-analysis, and the non-English articles were also excluded.

Data extraction

Two investigators (Kai Qu and Ming Zhang) independently extracted data using a standardized Excel template. Disagreements were resolved by consensus or by a third investigator (Wenquan Niu). Data were collected on the

first author, year of publication, ethnicity and country of the study population, cancer type, case-control status, study design, sample size, the genotypes/alleles of *IL-10* three variants (–592C>A, –819C>T, and –1082A>G) between cases and controls, and characteristics of participants, if available, including age, sex, body mass index, smoking, drinking, family history of cancer, and virus infection status.

Statistical analysis

In this meta-analysis, three genetic models of inheritance were performed for *IL-10* variants, including allelic model (the A allele versus the C allele for –592C>A SNP [single nucleotide polymorphism]; the T allele versus the C allele for –819C>T SNP; and the G allele versus the A allele for –1082A>G SNP), homozygous (the AA genotype versus the CC genotype for –592C>A SNP; the TT genotype versus the CC genotype for –819C>T SNP; and the GG genotype versus the AA genotype for –1082A>G) and dominant model (the AA genotype plus the AC genotype versus the AA genotype for –592C>A SNP; the TT genotype plus the TC genotype versus the CC genotype for –819C>T SNP; and the GG genotype plus the GA genotype versus the AA genotype for –1082A>G).

Weighted odds ratios (ORs) and the corresponding 95% confidence intervals (CIs) were evaluated by a random-effects model using the DerSimonian and Laird method. Heterogeneity between studies was computed by the χ^2 test, and was quantified by the inconsistency index (I^2) statistic, which ranges from 0% to 100% and is defined as the percentage of the observed between-study variability that is due to heterogeneity rather than chance.

Predefined subgroup analyses were performed a priori according to the cancer types (oral cancer, nasopharyngeal cancer, cervical cancer or lymphoma [including Hodgkin and Burkitt lymphoma]), ethnicity of the study populations (Caucasian, Asian, Latinos, or African), study design (population-based or hospital-based), and the total sample size (<300 subjects or $\geq$ 300 subjects). The data were presented and summarized if there were three or more independent studies that provided the genotype or allele counts of the *IL-10* three variants between cases and controls.

Genetic association studies have been considered more closely relevant to randomized trials than other types of epidemiological study due to independent assortment of alleles that theoretically should not be confounded by environmental or behavioral factors.²⁵ Therefore, we employed Mendelian randomization model to test the hypothesis that genetically

elevated level of IL-10 because of variants in *IL-10* cause an increased risk of HEACs. In Mendelian randomization analysis, risk estimate was computed from the ratio of the coefficient of the association between a variant and a disease to that of the association between the variant and biomarker as a reflection of the potential effect of circulating IL-10 levels on cancer risk.

Publication bias was assessed by visual inspection of Begg's and Egger's funnel plots, accompanied by the corresponding Begg's and Egger's tests. The trim and fill method was implemented to estimate the number and outcomes of potentially missing trials resulting from publication bias. Data management and statistical analyses were conducted using STATA software (StataCorp, College Station, TX, USA, version 11.2 for Windows). $P<0.05$ was considered statistically significant. For Begg's and Egger's statistics, a significance level was defined as $P<0.10$.

Results

Eligible articles

A flow diagram schematizing the process of article selection with specific reasons is presented in Figure 1. In total, 103 potentially relevant articles were identified after the initial search, and 24 of them that satisfied inclusion/exclusion criteria were deemed as eligible.^{26–49} All 24 qualified articles written in English were published between 2001 and 2014.

Study characteristics

The basic characteristics of all 24 qualified articles are listed in Table 1, and the genotype distributions and allele frequencies of *IL-10* three variants (–592C>A, –819C>T, and –1082A>G) between cases and controls are listed in Table S2. In this meta-analysis, 12 articles were conducted for cervical cancer, 7 for lymphoma (including Hodgkin's lymphoma and Burkitt's lymphoma), 4 for nasopharyngeal cancer, and 1 for oral cancer. Additionally, there were 8 articles involving Asians, 9 involving Caucasians, 3 involving Latinos, 2 involving Africans, and 1 involving the mixed populations. There were 11 articles conducted on a population-based design and 13 on a hospital-based design. Of 24 qualified articles, 14 (58.33%) had the total sample size (the sum of patients and controls) of at least 300 subjects.

Overall and subgroup analysis of IL-10 variants and HEACs

Pooling all 24 qualified articles together indicated a significant association between *IL-10*–1082A>G variant and HEACs

Figure 1 Flow diagram of search strategy and study selection.

under allelic (OR=1.283; 95% CI: 1.071–1.537; $P=0.007$) and dominant models (OR=1.382; 95% CI: 1.128–1.694; $P=0.002$; Table 2). Conversely, we failed to find any significance for the other two variants (–592C>A and –819C>T) in association with HEACs (Tables S3 and S4).

To account for the potential sources of between-study heterogeneity, a set of predefined subgroup analyses were conducted (Table 2). By ethnicity, an extremely significant association between *IL-10*–1082G allele and HEACs in Asians was observed under allelic (OR=2.009; 95% CI:

Table 1 Baseline characteristics of eligible studies for association of *IL-10* three variants with HEACs

Study	Ethnicity	Cancer type	Matched	Source of controls	Sample size		Age (years)		Sex (male, %)	
					Cases	Controls	Cases	Controls	Cases	Controls
Andrie et al ²⁶	Caucasian	Lymphoma	Yes	Hospital	37	85	NA	NA	61.9	71.8
Barbisan et al ²⁷	Latinos	Cervical cancer	NA	Hospital	176	122	44.0	37.0	0.00	0.00
Chagas et al ²⁸	Latinos	Cervical cancer	NA	Hospital	171	193	34.7	34.7	0.00	0.00
Cunningham et al ²⁹	Caucasian	Lymphoma	NA	Population	49	164	NA	NA	NA	NA
da Silva et al ³⁰	Latinos	Lymphoma	NA	Hospital	65	50	31.0	7.6	64.28	50.00
Farhat et al ³¹	Caucasian	Nasopharyngeal cancer	Yes	Population	160	156	41.9	40.4	72.50	74.68
Fernandes et al ³²	Caucasian	Cervical cancer	Yes	Hospital	42	87	27.0	29.0	0.00	0.00
Govan et al ³³	Mixed	Cervical cancer	NA	Hospital	197	182	NA	NA	0.00	0.00
Ivansson et al ³⁴	Caucasian	Cervical cancer	NA	Population	1,282	288	NA	NA	0.00	0.00
Matsumoto et al ³⁵	Asian	Cervical cancer	NA	Hospital	104	173	51.7	35.6	0.00	0.00
Minnicelli et al ³⁶	Latinos	Lymphoma	Yes	Population	61	230	NA	NA	NA	NA
Munro et al ³⁷	Caucasian	Lymphoma	No	Hospital	146	111	44.0	58.3	47.26	45.94
Nieters et al ³⁸	Caucasian	Lymphoma	Yes	Population	108	660	NA	NA	NA	NA
Oduor et al ³⁹	African	Lymphoma	Yes	Hospital	117	88	5.0	7.0	65.80	55.70
Pratesi et al ⁴⁰	Caucasian	Nasopharyngeal cancer	Yes	Population	89	130	NA	NA	78.70	76.90
Roh et al ⁴¹	Asian	Cervical cancer	Yes	Hospital	144	179	NA	NA	0.00	0.00
Shekari et al ⁴²	Asian	Cervical cancer	NA	Population	200	200	48.6	48.8	0.00	0.00
Singh et al ⁴³	Asian	Cervical cancer	Yes	Hospital	150	162	48.3	47.2	0.00	0.00
Stanczuk et al ⁴⁴	African	Cervical cancer	Yes	Hospital	77	69	47.5	48.0	0.00	0.00
Tsai et al ⁴⁶	Asian	Nasopharyngeal cancer	Yes	Population	176	522	48.2	48.9	72.20	72.60
Tsai et al ⁴⁵	Asian	Oral cancer	Yes	Population	788	956	55.8	56.6	76.00	76.00
Wang et al ⁴⁷	Asian	Cervical cancer	NA	Population	186	200	54.0	42.0	0.00	0.00
Wei et al ⁴⁸	Asian	Nasopharyngeal cancer	Yes	Population	198	210	48.7	47.9	72.22	66.19
Zoodsma et al ⁴⁹	Caucasian	Cervical cancer	NA	Hospital	667	563	NA	NA	0.00	0.00

Abbreviations: IL-10, interleukin 10; HEAC, human papilloma virus and Epstein–Barr virus-associated cancers; NA, not available.

Table 2 Overall and subgroup analyses of *IL-10*-1082A>G with HEAC risk

Groups	Studies	Allelic model		Homozygous genotypic model		Dominant model	
		OR; 95% CI; P-value	I ² (%)	OR; 95% CI; P-value	I ² (%)	OR; 95% CI; P-value	I ² (%)
Overall	20	1.283; 1.071–1.537; 0.007	81.6	1.329; 0.993–1.779; 0.056	63.6	1.382; 1.128–1.694; 0.002	72.0
Ethnicity							
Caucasian	8	0.974; 0.876–1.083; 0.630	0.0	0.921; 0.742–1.142; 0.453	0.0	0.991; 0.837–1.173; 0.916	0.0
Asian	5	2.009; 1.566–2.578; 0.000	65.7	2.832; 1.831–4.379; 0.000	27.7	2.101; 1.694–2.607; 0.000	37.4
Latinos	4	1.259; 0.942–1.682; 0.119	49.0	1.575; 0.841–2.949; 0.156	49.7	1.287; 0.968–1.710; 0.082	0.0
African	2	1.652; 0.503–5.428; 0.408	87.7	0.917; 0.368–2.283; 0.853	0.0	1.843; 0.486–6.994; 0.369	87.0
Mixed	1	0.796; 0.593–1.068; 0.128	–	0.611; 0.347–1.076; 0.088	–	0.888; 0.591–1.334; 0.568	–
Sample size							
<300 subjects	11	1.315; 1.023–1.689; 0.032	68.7	1.336; 0.927–1.924; 0.120	27.0	1.417; 1.042–1.928; 0.026	58.7
≥300 subjects	9	1.252; 0.957–1.638; 0.102	88.8	1.315; 0.848–2.040; 0.222	79.0	1.353; 1.027–1.800; 0.038	81.6
Cancer type							
Cervical	8	1.275; 0.969–1.677; 0.083	79.5	1.046; 0.749–1.462; 0.792	34.5	1.389; 0.983–1.965; 0.063	75.6
Oral	1	2.003; 1.659–2.419; 0.000	–	3.268; 1.953–5.471; 0.000	–	2.054; 1.654–2.552; 0.000	–
Nasopharyngeal	4	1.530; 1.063–2.200; 0.022	76.5	1.814; 0.945–3.481; 0.074	60.3	1.737; 1.280–2.358; 0.000	44.9
Lymphoma	7	1.050; 0.828–1.311; 0.688	55.3	1.087; 0.663–1.782; 0.741	54.6	1.062; 0.827–1.365; 0.637	13.3
Case-control matched							
NA	8	1.145; 0.895–1.464; 0.282	76.2	1.014; 0.715–1.438; 0.937	44.9	1.218; 0.893–1.661; 0.213	70.0
Yes	11	1.407; 1.091–1.814; 0.009	80.8	1.612; 1.024–2.536; 0.039	66.4	1.557; 1.214–1.999; 0.000	64.5
No	1	1.107; 0.780–1.571; 0.569	–	1.200; 0.597–2.413; 0.609	–	1.076; 0.588–1.969; 0.813	–
Study design							
Population	9	1.365; 1.047–1.780; 0.022	83.7	1.598; 0.983–2.597; 0.059	74.7	1.492; 1.157–1.924; 0.002	68.6
Hospital	11	1.191; 0.958–1.481; 0.116	71.0	0.984; 0.770–1.259; 0.899	10.0	1.293; 0.972–1.722; 0.078	65.2

Note: Data in bold indicates statistical significance.

Abbreviations: IL-10, interleukin 10; HEAC, human papilloma virus and Epstein-Barr virus-associated cancers; OR, odds ratio; CI, confidence interval; NA, not available; I², inconsistency index.

1.566–2.578; $P<0.001$), homozygous genotypic (OR=2.832; 95% CI: 1.831–4.379; $P<0.001$), and dominant (OR=2.101; 95% CI: 1.694–2.607; $P<0.001$) models (Figure 2A and B). In Caucasians, Latinos, Africans, or mixed populations, there was no significant association observed in this meta-analysis. By cancer type, the magnitude of risk estimates was significant for nasopharyngeal cancer under allelic (OR=1.530; 95% CI: 1.063–2.200; $P=0.022$) and dominant models (OR=1.737; 95% CI: 1.280–2.358; $P<0.001$; Figure 2C and D), whereas no significance was reached for the other cancer types under investigation. By study design, there was a significant association between *IL-10*-1082G allele and HEACs in population-based studies under allelic (OR=1.365; 95% CI: 1.047–1.780; $P=0.022$) and dominant (OR=1.492; 95% CI: 1.157–1.924; $P=0.002$) models. Additionally, *IL-10*-1082G allele was more significantly associated with HEACs in those case-control-matched studies under allelic (OR=1.407; 95% CI: 1.091–1.814; $P=0.009$), homozygous genotypic (OR=1.612; 95% CI: 1.024–2.536; $P=0.039$), and dominant (OR=1.557; 95% CI: 1.214–1.999; $P<0.001$) models (Table 2).

There was no publication bias of three *IL-10* promoter variants either in overall or subgroup analysis as reflected by the funnel plots, Egger's tests, and Begg-Mazendard tests (Figure 3).

Association of IL-10 variants with circulating IL-10 levels

Genotype-phenotype association was based on four articles with circulating IL-10 levels measured in HEAC cancer patients (Table S5).^{37,41,50,51} We compared averaged circulating IL-10 levels under homozygous genotypic and dominant models. Circulating IL-10 level was significantly elevated in -1082G allele carriers under homozygous genotypic model (standard mean difference [SMD] =25.692; 95% CI: 1.303–50.081; $P=0.039$) and dominant model (SMD =13.313; 95% CI: 0.901–25.725; $P=0.036$; Figure 4). There were low probabilities of publication bias for both models as reflected by the Begg's funnel plots (both $P=0.296$) and the Egger's tests ($P=0.308$ and $P=0.442$, respectively). As expected, there were no significant differences in the changes of circulating IL-10 level for -592C>A and -819C>T under both models.

Predicted association of circulating IL-10 levels with HEACs from Mendelian randomization

We assumed a linear-logistic relationship between difference of circulating IL-10 level and odds of HEACs when implementing Mendelian randomization method. The predicted OR for 5 and 10 pg/mL IL-10 increment were 1.13

Figure 2 (Continued)

Figure 2 Risk estimates of IL-10-1082A>G for cancer risk by subgroup analysis.

Notes: (A) By ethnicity under allelic model; (B) by ethnicity under dominant model; (C) by cancer type under allelic model; and (D) by cancer type under dominant model. The summary OR is shown by the middle of a solid diamond whose left and right extremes represent the corresponding 95% CI. Horizontal axis represents OR values, which were calculated against healthy controls. Weights are from random effects analysis.

Abbreviations: IL-10, interleukin 10; OR, odds ratio; CI, confidence interval; I^2 , inconsistency index.

Figure 3 Funnel plots for studies investigating the effect of *IL-10* three variants on HEAC risk.

Notes: (A) *IL-10-592C>A*; (B) *IL-10-819C>T*; and (C) *IL-10-1082A>G*. Vertical axis represents the log of OR; horizontal axis represents the SE of log(OR). Funnel plots are drawn with 95% confidence limits. The graphic symbols represents the data in the plot which is sized proportional to the inverse variance.

Abbreviations: IL-10, interleukin 10; HEAC, human papilloma virus and Epstein-Barr virus-associated cancers; OR, odds ratio; SE, standard error.

(95% CI: 1.02–18.64) and 1.28 (95% CI: 1.05–347.26), respectively. Because the 95% CIs of both estimated OR did not include the null hypothesis value of 1.00, it was safe to reject the null hypothesis at a 5% significance level, and reveal a potential role of elevated circulating IL-10 level in development of HEACs.

Discussion

On the basis of a meta-analysis of the data from 24 studies involving 5,390 cases and 5,780 controls, we investigated *IL-10* three promoter variants (–592C>A, –819C>T, and –1082A>G) and circulating IL-10 levels in relation to the risk for HEACs. One principal finding of this study was the significant association of *IL-10-1082A>G* variant with HEACs, especially in Asians and for nasopharyngeal cancer. On the basis of aforementioned results, we further employed –1082A>G variant as an instrument to surrogate circulating IL-10 levels, and revealed the association of IL-10 levels with risk for HEACs. To our knowledge, this is the first meta-analysis demonstrating the association between

circulating IL-10 levels and HEACs by implementing Mendelian randomization approach.

Previous studies have revealed that HPV and EBV infection is the main etiologic risk factors for many epithelial malignancies such as oropharyngeal cancer,⁴ nasopharyngeal cancer,⁶ cervical cancer,³ and some subtypes of lymphoma such as Hodgkin's⁹ and Burkitt's lymphoma.⁸ HPV and EBV are ubiquitous, double-stranded DNA viruses, which can be found in the upper aerodigestive tract. Epidemiologic data showed HPV and EBV infection affected over 10% and 90% of the general population, respectively, but only a small percentage of those infected developed cancer, probably because of lowered immune response and virus clearance due to interindividual genetic variations that result in persistent virus infection. IL-10, a cytokine with multiple effects in immunoregulation and inflammation, has a central role in infection by limiting the immune response to pathogens. Given the essential role of IL-10 in the antiviral response in vitro⁵² and in vivo,¹⁷ it is reasonable to expect that IL-10 is implicated in the tumorigenesis of HEACs. Indeed, elevation of circulating IL-10 levels was detected in patients with cervical cancer,⁵³

Figure 4 Comparison of circulating IL-10 levels across *IL-10*–1082A>G genotypes under homozygous (A) and dominant (B) models.

Notes: The summary treatment effect (SMD) is shown by the middle of a solid diamond with the left and right extremes representing the corresponding 95% CI. Weights are from random effects analysis.

Abbreviations: IL-10, interleukin 10; SMD, standard mean difference; CI, confidence interval; WMD, weighted mean differences.

oropharyngeal cancer,⁵⁴ and Hodgkin's lymphoma,⁵⁵ and was associated with poor prognoses of these patients. Consistent with above evidence, in this study, our pooled results using Mendelian randomization approach indicated that 5 and 10 pg/mL increments in circulating IL-10 levels were 1.13 and 1.28 times more likely to develop HEACs in a significant manner, respectively. However, considering the unstable status of circulating IL-10 levels in time as previously described (plasma half-life ranged from 2.7 to 4.5 hours),⁵⁶ which may cause a weak association between IL-10 level and HEAC risk, well-designed studies with precise IL-10 measurement are required to quantify this effect size reliably.

Nasopharyngeal cancer is a quite common malignancy in Eastern Asians, especially in Chinese, as well as in migrants from those areas,⁵⁷ with its incidence rates peaking at 30 cases/100,000 in males and at 10 cases/100,000

in females.⁵⁸ Conversely, it is rare in Europe and North America, accounting for less than 1% of all cancer cases, with incidence rates generally below 2 cases/100,000 in males and 1 case/100,000 in females.⁵⁸ The obvious various incidence of nasopharyngeal cancer among different ethnic groups suggests this cancer is influenced by heredity factors. In this study, our results robustly confirmed the association between –1082A>G allele and nasopharyngeal cancer, especially in Asian population with relative low between-study heterogeneity. Our findings added a potential explanation for varying incidence of nasopharyngeal cancer worldwide. Further studies are necessary to confirm our findings, and *IL-10*–1082A>G allele, once validated, might be a specific biomarker for patients with nasopharyngeal cancer.

Despite the clear strengths including the large sample sizes and implementation of Mendelian randomization

approach, several possible limitations in the present meta-analysis should also be noted. First, to avoid the impact of low-quality studies, we only included articles written in English, which might cause publication bias, even though our funnel plots and statistical tests did not tell. Second, we only examined three promoter variants in IL-10 gene, and investigation on other variants in or flanking IL-10 gene, especially some low-penetrance genes will be encouraged. Third, the single-locus-based nature of meta-analysis precluded the possibility of gene–gene⁵⁹ and gene–environment interactions, but whether this variant integrated with other genetic or environmental risk factors will enhance prediction requires additional research. For instance, it is found that different types of HPV proteins exhibited varying abilities in inducing promoter activity of IL-10 gene.⁶⁰ Therefore, it is also necessary to perform a HPV type-stratified analysis in further study. Fourth, nearly all involved studies in this meta-analysis had circulating IL-10 measured only once and did not reflect its long-term level in the development of HEACs. Therefore, because of the above limitations, the jury must refrain from drawing a firm conclusion until a large-scale and well-designed study confirms or refutes our findings.

In summary, our findings provided evidence for a critical role of genetically elevated circulating IL-10 in the development of HEACs by employing IL-10 gene –1082A>G as an instrument, and the risk association of this variant with HEACs was more evident in Asian patients with nasopharyngeal cancer. Additional studies examining biological function of elevated circulating IL-10 level in HEACs, as well as studies seeking to provide clinical validations of our findings, are warranted.

Acknowledgments

The authors are thankful to Dr Qichao Huang in the Fourth Military Medical University for his technical assistance in statistics. This work was supported by the grants of National Natural Science Foundation of China (Grant numbers: 81201549, 81272644, and 81472247), the Project of Innovative Research Team for Key Science and Technology in Xi'an Jiaotong University (number: 2013KCJ-23), and the Scientific Research Project of Shanghai Municipal Bureau of Health (number: 20114y069).

Disclosure

The authors report no conflicts of interest in this work.

References

- McLaughlin-Drubin ME, Munger K. Viruses associated with human cancer. *Biochim Biophys Acta*. 2008;1782(3):127–150.
- Zur Hausen H. The search for infectious causes of human cancers: where and why. *Virology*. 2009;392(1):1–10.
- Bosch FX, Manos MM, Munoz N, et al. Prevalence of human papillomavirus in cervical cancer: a worldwide perspective. International biological study on cervical cancer (IBSCC) Study Group. *J Natl Cancer Inst*. 1995;87(11):796–802.
- Chaturvedi AK, Engels EA, Pfeiffer RM, et al. Human papillomavirus and rising oropharyngeal cancer incidence in the United States. *J Clin Oncol*. 2011;29(32):4294–4301.
- de Martel C, Ferlay J, Franceschi S, et al. Global burden of cancers attributable to infections in 2008: a review and synthetic analysis. *Lancet Oncol*. 2012;13(6):607–615.
- Gu AD, Zeng MS, Qian CN. The criteria to confirm the role of Epstein–Barr virus in nasopharyngeal carcinoma initiation. *Int J Mol Sci*. 2012;13(10):13737–13747.
- Parkin DM. The global health burden of infection-associated cancers in the year 2002. *Int J Cancer*. 2006;118(12):3030–3044.
- Brady G, MacArthur GJ, Farrell PJ. Epstein–Barr virus and Burkitt lymphoma. *J Clin Pathol*. 2007;60(12):1397–1402.
- Glaser SL, Lin RJ, Stewart SL, et al. Epstein–Barr virus-associated Hodgkin's disease: epidemiologic characteristics in international data. *Int J Cancer*. 1997;70(4):375–382.
- Al Moustafa AE, Chen D, Ghabreau L, Akil N. Association between human papillomavirus and Epstein–Barr virus infections in human oral carcinogenesis. *Med Hypotheses*. 2009;73(2):184–186.
- Jalouli J, Jalouli MM, Sapkota D, Ibrahim SO, Larsson PA, Sand L. Human papilloma virus, herpes simplex virus and Epstein Barr virus in oral squamous cell carcinoma from eight different countries. *Anticancer Res*. 2012;32(2):571–580.
- Tyan YS, Liu ST, Ong WR, Chen ML, Shu CH, Chang YS. Detection of Epstein–Barr virus and human papillomavirus in head and neck tumors. *J Clin Microbiol*. 1993;31(1):53–56.
- Khenchouche A, Sadouki N, Boudriche A, et al. Human papillomavirus and Epstein–Barr virus co-infection in cervical carcinoma in Algerian women. *Virol J*. 2013;10:340.
- Prayitno A. Cervical cancer with human papilloma virus and Epstein Barr virus positive. *J Carcinog*. 2006;5:13.
- Hariri S, Unger ER, Sternberg M, et al. Prevalence of genital human papillomavirus among females in the United States, the National Health And Nutrition Examination Survey, 2003–2006. *J Infect Dis*. 2011;204(4):566–573.
- Beskow AH, Gyllenstein UB. Host genetic control of HPV 16 titer in carcinoma in situ of the cervix uteri. *Int J Cancer*. 2002;101(6):526–531.
- Brooks DG, Trifilo MJ, Edelmann KH, Teyton L, McGavern DB, Oldstone MB. Interleukin-10 determines viral clearance or persistence in vivo. *Nat Med*. 2006;12(11):1301–1309.
- Dennis KL, Blatner NR, Gounari F, Khazaie K. Current status of interleukin-10 and regulatory T-cells in cancer. *Curr Opin Oncol*. 2013;25(6):637–645.
- Bermudez-Morales VH, Gutierrez LX, Alcocer-Gonzalez JM, Burguete A, Madrid-Marina V. Correlation between IL-10 gene expression and HPV infection in cervical cancer: a mechanism for immune response escape. *Cancer Invest*. 2008;26(10):1037–1043.
- El-Sherif AM, Seth R, Tighe PJ, Jenkins D. Quantitative analysis of IL-10 and IFN-gamma mRNA levels in normal cervix and human papillomavirus type 16 associated cervical precancer. *J Pathol*. 2001;195(2):179–185.
- Viviani S, Notti P, Bonfante V, Verderio P, Valagussa P, Bonadonna G. Elevated pretreatment serum levels of IL-10 are associated with a poor prognosis in Hodgkin's disease, the Milan Cancer Institute Experience. *Med Oncol*. 2000;17(1):59–63.
- Suarez A, Castro P, Alonso R, Mozo L, Gutierrez C. Interindividual variations in constitutive interleukin-10 messenger RNA and protein levels and their association with genetic polymorphisms. *Transplantation*. 2003;75(5):711–717.
- Turner DM, Williams DM, Sankaran D, Lazarus M, Sinnott PJ, Hutchinson IV. An investigation of polymorphism in the interleukin-10 gene promoter. *Eur J Immunogenet*. 1997;24(1):1–8.

24. Proitsi P, Lupton MK, Velayudhan L, et al. Genetic predisposition to increased blood cholesterol and triglyceride lipid levels and risk of Alzheimer disease: a Mendelian randomization analysis. *PLoS Med*. 2014;11(9):e1001713.
25. Niu W, Liu Y, Qi Y, Wu Z, Zhu D, Jin W. Association of interleukin-6 circulating levels with coronary artery disease: a meta-analysis implementing mendelian randomization approach. *Int J Cardiol*. 2012; 157(2):243–252.
26. Andrie E, Michos A, Kalampoki V, et al. Genetic variants in immunoregulatory genes and risk for childhood lymphomas. *Eur J Haematol*. 2009;83(4):334–342.
27. Barbisan G, Perez LO, Contreras A, Golijow CD. TNF-alpha and IL-10 promoter polymorphisms, HPV infection, and cervical cancer risk. *Tumour Biol*. 2012;33(5):1549–1556.
28. Chagas BS, Gurgel AP, da Cruz HL, et al. An interleukin-10 gene polymorphism associated with the development of cervical lesions in women infected with Human Papillomavirus and using oral contraceptives. *Infect Genet Evol*. 2013;19:32–37.
29. Cunningham LM, Chapman C, Dunstan R, Bell MC, Joske DJ. Polymorphisms in the interleukin 10 gene promoter are associated with susceptibility to aggressive non-Hodgkin's lymphoma. *Leuk Lymphoma*. 2003;44(2):251–255.
30. da Silva GN, Bacchi MM, Rainho CA, de Oliveira DE. Epstein-Barr virus infection and single nucleotide polymorphisms in the promoter region of interleukin 10 gene in patients with Hodgkin lymphoma. *Arch Pathol Lab Med*. 2007;131(11):1691–1696.
31. Farhat K, Hassen E, Gabbouj S, Bouaouina N, Chouchane L. Interleukin-10 and interferon-gamma gene polymorphisms in patients with nasopharyngeal carcinoma. *Int J Immunogenet*. 2008;35(3): 197–205.
32. Fernandes AP, Goncalves MA, Simoes RT, Mendes-Junior CT, Duarte G, Donadi EA. A pilot case-control association study of cytokine polymorphisms in Brazilian women presenting with HPV-related cervical lesions. *Eur J Obstet Gynecol Reprod Biol*. 2008;140(2):241–244.
33. Govan VA, Carrara HR, Sachs JA, Hoffman M, Stanczuk GA, Williamson AL. Ethnic differences in allelic distribution of IFN-gamma in South African women but no link with cervical cancer. *J Carcinog*. 2003;2(1):3.
34. Ivansson EL, Gustavsson IM, Magnusson JJ, et al. Variants of chemokine receptor 2 and interleukin 4 receptor, but not interleukin 10 or Fas ligand, increase risk of cervical cancer. *Int J Cancer*. 2007;121(11):2451–2457.
35. Matsumoto K, Oki A, Satoh T, et al. Interleukin-10 –1082 gene polymorphism and susceptibility to cervical cancer among Japanese women. *Jpn J Clin Oncol*. 2010;40(11):1113–1116.
36. Minnicelli C, Barros MH, Klumb CE, Romano SO, Zalcberg IR, Hassan R. Relationship of Epstein-Barr virus and interleukin 10 promoter polymorphisms with the risk and clinical outcome of childhood Burkitt lymphoma. *PLoS One*. 2012;7(9):e46005.
37. Munro LR, Johnston PW, Marshall NA, et al. Polymorphisms in the interleukin-10 and interferon gamma genes in Hodgkin lymphoma. *Leuk Lymphoma*. 2003;44(12):2083–2088.
38. Nieters A, Beckmann L, Deeg E, Becker N. Gene polymorphisms in Toll-like receptors, interleukin-10, and interleukin-10 receptor alpha and lymphoma risk. *Genes Immun*. 2006;7(8):615–624.
39. Oduor CI, Chelimo K, Ouma C, et al. Interleukin-6 and interleukin-10 gene promoter polymorphisms and risk of endemic Burkitt lymphoma. *Am J Trop Med Hyg*. 2014;91(3):649–654.
40. Pratesi C, Bortolin MT, Bidoli E, et al. Interleukin-10 and interleukin-18 promoter polymorphisms in an Italian cohort of patients with undifferentiated carcinoma of nasopharyngeal type. *Cancer Immunol Immunother*. 2006;55(1):23–30.
41. Roh JW, Kim MH, Seo SS, et al. Interleukin-10 promoter polymorphisms and cervical cancer risk in Korean women. *Cancer Lett*. 2002; 184(1):57–63.
42. Shekari M, Kordi-Tamandani DM, Malekzadeh K, Sobti RC, Karimi S, Suri V. Effect of anti-inflammatory (IL-4, IL-10) cytokine genes in relation to risk of cervical carcinoma. *Am J Clin Oncol*. 2012; 35(6):514–519.
43. Singh H, Jain M, Sachan R, Mittal B. Association of TNFA (–308G>A) and IL-10 (–819C>T) promoter polymorphisms with risk of cervical cancer. *Int J Gynecol Cancer*. 2009;19(7):1190–1194.
44. Stanczuk GA, Sibanda EN, Perrey C, et al. Cancer of the uterine cervix may be significantly associated with a gene polymorphism coding for increased IL-10 production. *Int J Cancer*. 2001;94(6):792–794.
45. Tsai CW, Chang WS, Lin KC, et al. Significant association of Interleukin-10 genotypes and oral cancer susceptibility in Taiwan. *Anticancer Res*. 2014;34(7):3731–3737.
46. Tsai CW, Tsai MH, Shih LC, Chang WS, Lin CC, Bau DT. Association of interleukin-10 (IL10) promoter genotypes with nasopharyngeal carcinoma risk in Taiwan. *Anticancer Res*. 2013;33(8):3391–3396.
47. Wang Q, Zhang C, Walayat S, Chen HW, Wang Y. Association between cytokine gene polymorphisms and cervical cancer in a Chinese population. *Eur J Obstet Gynecol Reprod Biol*. 2011;158(2):330–333.
48. Wei YS, Kuang XH, Zhu YH, et al. Interleukin-10 gene promoter polymorphisms and the risk of nasopharyngeal carcinoma. *Tissue Antigens*. 2007;70(1):12–17.
49. Zoodsma M, Nolte IM, Schipper M, et al. Interleukin-10 and Fas polymorphisms and susceptibility for (pre)neoplastic cervical disease. *Int J Gynecol Cancer*. 2005;15(Suppl 3):282–290.
50. Hohaus S, Giachelia M, Massini G, et al. Clinical significance of interleukin-10 gene polymorphisms and plasma levels in Hodgkin lymphoma. *Leuk Res*. 2009;33(10):1352–1356.
51. Jin L, Sturgis EM, Cao X, et al. Interleukin-10 promoter variants predict HPV-positive tumors and survival of squamous cell carcinoma of the oropharynx. *FASEB J*. 2013;27(6):2496–2503.
52. Santin AD, Hermonat PL, Ravaggi A, et al. Interleukin-10 increases Th1 cytokine production and cytotoxic potential in human papillomavirus-specific CD8(+) cytotoxic T lymphocytes. *J Virol*. 2000;74(10):4729–4737.
53. Chopra V, Dinh TV, Hannigan EV. Circulating serum levels of cytokines and angiogenic factors in patients with cervical cancer. *Cancer Invest*. 1998;16(3):152–159.
54. Fujieda S, Sunaga H, Tsuzuki H, Fan GK, Saito H. IL-10 expression is associated with the expression of platelet-derived endothelial cell growth factor and prognosis in oral and oropharyngeal carcinoma. *Cancer Lett*. 1999;136(1):1–9.
55. Bohlen H, Kessler M, Sextro M, Diehl V, Tesch H. Poor clinical outcome of patients with Hodgkin's disease and elevated interleukin-10 serum levels. Clinical significance of interleukin-10 serum levels for Hodgkin's disease. *Ann Hematol*. 2000;79(3):110–113.
56. Huhn RD, Radwanski E, Gallo J, et al. Pharmacodynamics of subcutaneous recombinant human interleukin-10 in healthy volunteers. *Clin Pharmacol Ther*. 1997;62(2):171–180.
57. Mousavi SM, Sundquist J, Hemminki K. Nasopharyngeal and hypopharyngeal carcinoma risk among immigrants in Sweden. *Int J Cancer*. 2010;127(12):2888–2892.
58. Polesel J, Serraino D, Negri E, et al. Consumption of fruit, vegetables, and other food groups and the risk of nasopharyngeal carcinoma. *Cancer Causes Control*. 2013;24(6):1157–1165.
59. Hayashi T, Ito R, Cologne J, et al. Effects of IL-10 haplotype and atomic bomb radiation exposure on gastric cancer risk. *Radiat Res*. 2013; 180(1):60–69.
60. Bermudez-Morales VH, Peralta-Zaragoza O, Alcocer-Gonzalez JM, Moreno J, Madrid-Marina V. IL-10 expression is regulated by HPV E2 protein in cervical cancer cells. *Mol Med Rep*. 2011;4(2):369–375.

Supplementary materials

Table S I PRISMA 2009 Checklist

Section/topic	#	Checklist item	Reported on page #
Title			
Title	1	Identify the report as a systematic review, meta-analysis, or both.	Title (Page 1)
Abstract			
Structured summary	2	Provide a structured summary including, as applicable: background; objectives; data sources; study eligibility criteria; participants, and interventions; study appraisal and synthesis methods; results; limitations; conclusions and implications of key findings; systematic review registration number.	Abstract (Page 1)
Introduction			
Rationale	3	Describe the rationale for the review in the context of what is already known.	Introduction (Page 1)
Objectives	4	Provide an explicit statement of questions being addressed with reference to participants, interventions, comparisons, outcomes, and study design (PICOS).	Introduction (Page 2)
Methods			
Protocol and registration	5	Indicate if a review protocol exists, if and where it can be accessed (eg, Web address), and, if available, provide registration information including registration number.	Methods (Page 4)
Eligibility criteria	6	Specify study characteristics (eg, PICOS, length of follow-up) and report characteristics (eg, years considered, language, publication status) used as criteria for eligibility, giving rationale.	Methods: inclusion/exclusion criteria (Page 2)
Information sources	7	Describe all information sources (eg, databases with dates of coverage, contact with study authors to identify additional studies) in the search and date last searched.	Methods: search strategy (Page 2)
Search	8	Present full electronic search strategy for at least one database, including any limits used, such that it could be repeated.	Methods: search strategy (Page 2)
Study selection	9	State the process for selecting studies (ie, screening, eligibility, included in systematic review, and, if applicable, included in the meta-analysis).	Methods: search strategy (Page 2)
Data collection process	10	Describe method of data extraction from reports (eg, piloted forms, independently, in duplicate) and any processes for obtaining and confirming data from investigators.	Methods: data extraction (Page 2)
Data items	11	List and define all variables for which data were sought (eg, PICOS, funding sources) and any assumptions and simplifications made.	Methods: data extraction (Page 2, 3)
Risk of bias in individual studies	12	Describe methods used for assessing risk of bias of individual studies (including specification of whether this was done at the study or outcome level), and how this information is to be used in any data synthesis.	Methods: statistical analysis (Page 3)
Summary measures	13	State the principal summary measures (eg, risk ratio, difference in means).	Methods: statistical analysis (Page 3)
Synthesis of results	14	Describe the methods of handling data and combining results of studies, if done, including measures of consistency (eg, I^2) for each meta-analysis.	Methods: statistical analysis (Page 3)
Risk of bias across studies	15	Specify any assessment of risk of bias that may affect the cumulative evidence (eg, publication bias, selective reporting within studies).	Methods: statistical analysis (Page 3)
Additional analyses	16	Describe methods of additional analyses (eg, sensitivity or subgroup analyses, meta-regression), if done, indicating which were pre-specified.	Methods: statistical analysis (Page 3)
Results			
Study selection	17	Give numbers of studies screened, assessed for eligibility, and included in the review, with reasons for exclusions at each stage, ideally with a flow diagram.	Figure 1

Study characteristics	18	For each study, present characteristics for which data were extracted (eg, study size, PICOS, follow-up period) and provide the citations.	Results: eligible articles Table 1, Tables S1 and S4
Risk of bias within studies	19	Present data on risk of bias of each study and, if available, any outcome level assessment (see Item 12).	Results: IL-10 gene three variants with HEACs; IL-10 variants with circulating IL-10 level; Mendelian randomization analysis (Page 3–8)
Results of individual studies	20	For all outcomes considered (benefits or harms), present, for each study: (a) simple summary data for each intervention group (b) effect estimates and confidence intervals, ideally with a forest plot.	Table 2
Synthesis of results	21	Present results of each meta-analysis done, including confidence intervals and measures of consistency.	Table 2 and Tables S2, S3
Risk of bias across studies	22	Present results of any assessment of risk of bias across studies (see Item 15).	Figures 2 and 4
Additional analysis	23	Give results of additional analyses, if done (eg, sensitivity or subgroup analyses, meta-regression [see Item 16]).	None
Discussion			
Summary of evidence	24	Summarize the main findings including the strength of evidence for each main outcome; consider their relevance to key groups (eg, health care providers, users, and policy makers).	Discussion (Page 8)
Limitations	25	Discuss limitations at study and outcome level (eg, risk of bias), and at review-level (eg, incomplete retrieval of identified research, reporting bias).	Discussion (Page 9, 10)
Conclusions	26	Provide a general interpretation of the results in the context of other evidence, and implications for future research.	Discussion (Page 10)
Funding			
Funding	27	Describe sources of funding for the systematic review and other support (eg, supply of data); role of funders for the systematic review.	Funding (Page 10)

Notes: Adapted from Moher D, Liberati A, Tetzlaff J, Altman DG; The PRISMA Group. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med*. 2009;6(6):e1000097.¹ For more information, visit: www.prisma-statement.org.

Abbreviations: IL-10, interleukin 10; HEAC, human papilloma virus and Epstein-Barr virus-associated cancers.

Table S2 The genotype distributions of three examined variants in *IL-10* between HEAC patients and controls in all qualified studies

Study	IL-10 gene -592 C>A (rs1800872)						IL-10 gene -819C>T (rs1800871)	
	Case_CC	Case_CA	Case_AA	Control_CC	Control_CA	Control_AA	Case_CC	Case_CT
Andrie et al ²	NA	NA	NA	NA	NA	NA	23	11
Barbisan et al ³	NA	NA	NA	NA	NA	NA	NA	NA
Chagas et al ⁴	NA	NA	NA	NA	NA	NA	56	90
Cunningham et al ⁵	NA	NA	NA	NA	NA	NA	NA	NA
da Silva et al ⁶	30	31	4	18	23	9	30	31
Farhat et al ⁷	NA	NA	NA	NA	NA	NA	NA	NA
Fernandes et al ⁸	NA	NA	NA	NA	NA	NA	NA	NA
Govan et al ⁹	NA	NA	NA	NA	NA	NA	NA	NA
Ivansson et al ¹⁰	736	464	82	162	112	14	NA	NA
Matsumoto et al ¹¹	NA	NA	NA	NA	NA	NA	NA	NA
Minnicelli et al ¹²	33	24	4	90	92	23	33	24
Munro et al ¹³	88	55	4	66	42	2	NA	NA
Nieters et al ¹⁴	NA	NA	NA	NA	NA	NA	NA	NA
Oduor et al ¹⁵	32	61	24	28	39	21	32	61
Pratesi et al ¹⁶	48	36	5	70	54	6	48	36
Roh et al ¹⁷	11	56	77	15	77	87	11	56
Shekari et al ¹⁸	16	96	88	17	102	81	NA	NA
Singh et al ¹⁹	NA	NA	NA	NA	NA	NA	56	94
Stanczuk et al ²⁰	NA	NA	NA	NA	NA	NA	NA	NA
Tsai et al ²²	17	66	93	56	205	261	19	69
Tsai et al ²¹	NA	NA	NA	NA	NA	NA	NA	NA
Wang et al ²³	NA	NA	NA	NA	NA	NA	NA	NA
Wei et al ²⁴	35	81	82	24	92	94	35	81
Zoodsma et al ²⁵	393	231	30	405	175	26	NA	NA

Abbreviations: IL-10, interleukin 10; HEAC, human papilloma virus and Epstein-Barr virus-associated cancers; NA, not available.

Table S3 The overall and subgroup analyses of -592C>A in *IL-10* with HEAC risk

Groups	Studies	Allelic model		Homozygous genotypic model		Dominant model	
		OR; 95% CI; P-value	I ² (%)	OR; 95% CI; P-value	I ² (%)	OR; 95% CI; P-value	I ² (%)
Overall	11	1.018; 0.913–1.135; 0.751	24.0	0.986; 0.780–1.247; 0.907	3.8	1.004; 0.853–1.183; 0.227	22.7
Ethnicity							
Caucasian	4	1.110; 0.976–1.264; 0.113	0.0	1.243; 0.857–1.804; 0.251	0.0	1.106; 0.909–1.346; 0.315	27.5
Asian	4	1.022; 0.873–1.196; 0.785	12.2	0.954; 0.668–1.365; 0.798	10.9	0.905; 0.655–1.250; 0.544	1.5
Latinos	2	0.665; 0.469–0.943; 0.022	0.0	0.371; 0.157–0.876; 0.024	0.0	0.661; 0.419–1.044; 0.076	0.0
African	1	1.023; 0.691–1.514; 0.911	–	1.000; 0.461–2.170; 1.000	–	1.240; 0.677–2.271; 0.487	–
Sample size							
<300	5	0.891; 0.728–1.090; 0.262	3.3	0.750; 0.429–1.312; 0.314	15.5	0.911; 0.703–1.181; 0.482	0.0
≥300	6	1.070; 0.952–1.204; 0.256	20.9	1.058; 0.818–1.369; 0.665	0.0	1.042; 0.828–1.312; 0.724	40.7
Cancer type							
Cervical	4	1.126; 0.998–1.270; 0.054	0.0	1.214; 0.877–1.681; 0.242	0.0	1.136; 0.929–1.388; 0.215	19.6
Nasopharyngeal	3	0.965; 0.792–1.177; 0.728	16.3	0.884; 0.540–1.445; 0.622	27.8	0.876; 0.603–1.272; 0.487	25.5
Lymphoma	4	0.854; 0.667–1.093; 0.210	17.8	0.667; 0.341–1.305; 0.237	26.6	0.887; 0.660–1.192; 0.427	0.0
Case-control matched							
NA	4	1.049; 0.859–1.279; 0.640	53.0	1.040; 0.659–1.640; 0.866	38.0	1.066; 0.816–1.394; 0.638	47.6
Yes	6	0.970; 0.839–1.122; 0.681	6.7	0.892; 0.652–1.220; 0.474	0.0	0.901; 0.704–1.152; 0.404	2.2
No	1	1.032; 0.672–1.583; 0.887	–	1.500; 0.267–8.437; 0.645	–	1.006; 0.853–1.665; 0.982	–
Study design							
Population	6	0.977; 0.866–1.104; 0.712	2.8	0.959; 0.700–1.247; 0.797	11.6	0.901; 0.750–1.082; 0.265	0.0
Hospital	5	1.070; 0.884–1.296; 0.486	33.5	1.010; 0.667–1.529; 0.964	14.1	1.211; 1.005–1.459; 0.044	0.0

Note: Data in bold indicates statistical significance.

Abbreviations: IL-10, interleukin 10; HEAC, human papilloma virus and Epstein-Barr virus-associated cancers; OR, odds ratio; CI, confidence interval; NA, not available; I², inconsistency index.

				IL-10 gene –1082A>G (rs1800896)					
Case_TT	Control_CC	Control_CT	Control_TT	Case_AA	Case_AG	Case_GG	Control_AA	Control_AG	Control_GG
3	45	35	5	5	16	16	12	32	40
NA	NA	NA	NA	50	61	11	79	83	14
25	76	87	30	56	78	37	26	36	20
NA	NA	NA	NA	16	24	9	41	82	41
4	18	23	9	26	30	9	27	19	4
NA	NA	NA	NA	58	80	22	70	60	26
NA	NA	NA	NA	18	19	5	41	38	8
NA	NA	NA	NA	88	80	29	76	65	41
NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
NA	NA	NA	NA	73	26	5	156	16	1
4	90	92	23	21	26	14	102	92	22
NA	NA	NA	NA	30	69	48	24	55	32
NA	NA	NA	NA	38	53	17	208	302	150
24	28	39	21	53	53	11	39	39	10
5	70	54	6	29	41	19	46	58	26
77	15	77	87	NA	NA	NA	NA	NA	NA
NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
0	77	85	0	NA	NA	NA	NA	NA	NA
NA	NA	NA	NA	45	31	1	58	11	0
88	52	185	285	117	49	10	419	92	11
NA	NA	NA	NA	522	217	49	766	168	22
NA	NA	NA	NA	77	85	24	103	76	21
82	24	92	94	123	61	14	167	38	5
NA	NA	NA	NA	154	326	187	130	307	169

Table S4 The overall and subgroup analyses of –819C>T in *IL-10* with HEAC risk

Groups	Studies	Allelic model		Homozygous genotypic model		Dominant model	
		OR; 95% CI; P-value	I ² (%)	OR; 95% CI; P-value	I ² (%)	OR; 95% CI; P-value	I ² (%)
Overall	10	0.955; 0.837–1.088; 0.487	20.7	0.834; 0.637–1.091; 0.185	0.0	0.970; 0.778–1.208; 0.783	30.0
Ethnicity							
Caucasian	2	0.958; 0.668–1.373; 0.814	0.0	1.198; 0.458–3.314; 0.712	0.0	0.884; 0.566–1.381; 0.588	0.0
Asian	4	0.990; 0.804–1.218; 0.921	45.0	0.793; 0.547–1.151; 0.222	0.0	0.991; 0.646–1.519; 0.966	54.6
Latinos	3	0.825; 0.556–1.225; 0.341	61.5	0.606; 0.251–1.459; 0.264	56.8	0.881; 0.527–1.472; 0.628	58.3
African	1	1.023; 0.691–1.514; 0.911	–	1.000; 0.461–2.170; 1.000	–	1.240; 0.677–2.271; 0.487	–
Sample size							
<300	5	0.854; 0.691–1.055; 0.143	0.0	0.746; 0.439–1.268; 0.279	10.3	0.852; 0.642–1.130; 0.265	0.0
≥300	5	1.013; 0.855–1.201; 0.879	35.5	0.869; 0.630–1.197; 0.390	0.0	1.071; 0.767–1.494; 0.687	47.6
Cancer type							
Cervical	3	1.177; 0.975–1.422; 0.090	0.0	1.158; 0.699–1.918; 0.569	0.0	1.375; 1.028–1.839; 0.032	0.0
Nasopharyngeal	3	0.873; 0.731–1.043; 0.134	0.0	0.754; 0.509–1.119; 0.161	0.0	0.822; 0.598–1.131; 0.229	0.0
Lymphoma	4	0.808; 0.635–1.028; 0.083	0.0	0.665; 0.355–1.245; 0.202	20.8	0.803; 0.577–1.118; 0.194	0.0
Case-control matched							
NA	2	0.872; 0.487–1.564; 0.647	72.2	0.619; 0.153–2.503; 0.501	73.5	1.004; 0.508–1.985; 0.990	60.9
Yes	8	0.953; 0.833–1.090; 0.478	9.2	0.826; 0.609–1.120; 0.219	0.0	0.945; 0.738–1.209; 0.651	28.5
No							
Study design	4	0.848; 0.719–1.000; 0.050	0.0	0.718; 0.495–1.041; 0.081	0.0	0.782; 0.592–1.033; 0.084	0.0
Population	6	1.058; 0.889–1.259; 0.524	15.0	0.975; 0.649–1.464; 0.903	6.1	1.164; 0.899–1.506; 0.249	12.4

Note: Data in bold indicates statistical significance.

Abbreviations: IL-10, interleukin 10; HEAC, human papilloma virus and Epstein–Barr virus-associated cancers; OR, odds ratio; CI, confidence interval; NA, not available; I², inconsistency index.

Table S5 Changes of circulating IL-10 level across genotypes of three examined variants in *IL-10*

Study	Ethnicity	Status	Sample size	Number	IL-10 level (pg/mL)	Number	IL-10 level (pg/mL)	Number	IL-10 level (pg/mL)
IL-10 gene -592C>A				CC genotype		CA genotype		AA genotype	
Hohaus et al ²⁶	Caucasian	Cases	95	NA	NA	85	29.20	10	53.10
Jin et al ²⁷	East Asian	Cases	180	96	10.1	84	13.20	NA	NA
Munro et al ¹³	Caucasian	Cases	25	15	106	10	35.90	NA	NA
Roh et al ¹⁷	East Asian	Cases	144	11	2.55	56	4.22	77	3.17
IL-10 gene -819C>T				CC genotype		CT genotype		TT genotype	
Jin et al ²⁷	East Asian	Cases	180	99	9.60	81	12.80	NA	NA
Roh et al ¹⁷	East Asian	Cases	144	11	2.55	56	4.22	77	3.17
IL-10 gene -1082A>G				AA genotype		AG genotype		GG genotype	
Hohaus et al ²⁶	Caucasian	Cases	95	NA	NA	87	29.20	8	56.20
Jin et al ²⁷	East Asian	Cases	180	68	10.70	112	12.60	NA	NA
Munro et al ¹³	Caucasian	Cases	26	3	43.00	11	67.80	12	91.70

Abbreviations: IL-10, interleukin 10; NA, not available.

References

- Moher D, Liberati A, Tetzlaff J, Altman DG; The PRISMA Group. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med*. 2009;6(6):e1000097.
- Andrie E, Michos A, Kalampoki V, et al. Genetic variants in immunoregulatory genes and risk for childhood lymphomas. *Eur J Haematol*. 2009;83(4):334–342.
- Barbisan G, Perez LO, Contreras A, Golijow CD. TNF-alpha and IL-10 promoter polymorphisms, HPV infection, and cervical cancer risk. *Tumour Biol*. 2012;33(5):1549–1556.
- Chagas BS, Gurgel AP, da Cruz HL, et al. An interleukin-10 gene polymorphism associated with the development of cervical lesions in women infected with Human Papillomavirus and using oral contraceptives. *Infect Genet Evol*. 2013;19:32–37.
- Cunningham LM, Chapman C, Dunstan R, Bell MC, Joske DJ. Polymorphisms in the interleukin 10 gene promoter are associated with susceptibility to aggressive non-Hodgkin's lymphoma. *Leuk Lymphoma*. 2003;44(2):251–255.
- da Silva GN, Bacchi MM, Rainho CA, de Oliveira DE. Epstein-Barr virus infection and single nucleotide polymorphisms in the promoter region of interleukin 10 gene in patients with Hodgkin lymphoma. *Arch Pathol Lab Med*. 2007;131(11):1691–1696.
- Farhat K, Hassen E, Gabbouj S, Bouaouina N, Chouchane L. Interleukin-10 and interferon-gamma gene polymorphisms in patients with nasopharyngeal carcinoma. *Int J Immunogenet*. 2008;35(3):197–205.
- Fernandes AP, Goncalves MA, Simoes RT, Mendes-Junior CT, Duarte G, Donadi EA. A pilot case-control association study of cytokine polymorphisms in Brazilian women presenting with HPV-related cervical lesions. *Eur J Obstet Gynecol Reprod Biol*. 2008;140(2):241–244.
- Govan VA, Carrara HR, Sachs JA, Hoffman M, Stanczuk GA, Williamson AL. Ethnic differences in allelic distribution of IFN-gamma in South African women but no link with cervical cancer. *J Carcinog*. 2003;2(1):3.
- Ivansson EL, Gustavsson IM, Magnusson JJ, et al. Variants of chemokine receptor 2 and interleukin 4 receptor, but not interleukin 10 or Fas ligand, increase risk of cervical cancer. *Int J Cancer*. 2007;121(11):2451–2457.
- Matsumoto K, Oki A, Satoh T, et al. Interleukin-10 -1082 gene polymorphism and susceptibility to cervical cancer among Japanese women. *Jpn J Clin Oncol*. 2010;40(11):1113–1116.
- Minnicelli C, Barros MH, Klumb CE, Romano SO, Zalberg IR, Hassan R. Relationship of Epstein-Barr virus and interleukin 10 promoter polymorphisms with the risk and clinical outcome of childhood Burkitt lymphoma. *PLoS One*. 2012;7(9):e46005.
- Munro LR, Johnston PW, Marshall NA, et al. Polymorphisms in the interleukin-10 and interferon gamma genes in Hodgkin lymphoma. *Leuk Lymphoma*. 2003;44(12):2083–2088.
- Nieters A, Beckmann L, Deeg E, Becker N. Gene polymorphisms in Toll-like receptors, interleukin-10, and interleukin-10 receptor alpha and lymphoma risk. *Genes Immun*. 2006;7(8):615–624.
- Oduor CI, Chelimo K, Ouma C, et al. Interleukin-6 and interleukin-10 gene promoter polymorphisms and risk of endemic Burkitt lymphoma. *Am J Trop Med Hyg*. 2014;91(3):649–654.
- Pratesi C, Bortolin MT, Bidoli E, et al. Interleukin-10 and interleukin-18 promoter polymorphisms in an Italian cohort of patients with undifferentiated carcinoma of nasopharyngeal type. *Cancer Immunol Immunother*. 2006;55(1):23–30.
- Roh JW, Kim MH, Seo SS, et al. Interleukin-10 promoter polymorphisms and cervical cancer risk in Korean women. *Cancer Lett*. 2002;184(1):57–63.
- Shekari M, Kordi-Tamandani DM, MalekZadeh K, Sobti RC, Karimi S, Suri V. Effect of anti-inflammatory (IL-4, IL-10) cytokine genes in relation to risk of cervical carcinoma. *Am J Clin Oncol*. 2012;35(6):514–519.
- Singh H, Jain M, Sachan R, Mittal B. Association of TNFA (-308G>A) and IL-10 (-819C>T) promoter polymorphisms with risk of cervical cancer. *Int J Gynecol Cancer*. 2009;19(7):1190–1194.
- Stanczuk GA, Sibanda EN, Perrey C, et al. Cancer of the uterine cervix may be significantly associated with a gene polymorphism coding for increased IL-10 production. *Int J Cancer*. 2001;94(6):792–794.
- Tsai CW, Chang WS, Lin KC, et al. Significant association of Interleukin-10 genotypes and oral cancer susceptibility in Taiwan. *Anticancer Res*. 2014;34(7):3731–3737.
- Tsai CW, Tsai MH, Shih LC, Chang WS, Lin CC, Bau DT. Association of interleukin-10 (IL10) promoter genotypes with nasopharyngeal carcinoma risk in Taiwan. *Anticancer Res*. 2013;33(8):3391–3396.
- Wang Q, Zhang C, Walayat S, Chen HW, Wang Y. Association between cytokine gene polymorphisms and cervical cancer in a Chinese population. *Eur J Obstet Gynecol Reprod Biol*. 2011;158(2):330–333.
- Wei YS, Kuang XH, Zhu YH, et al. Interleukin-10 gene promoter polymorphisms and the risk of nasopharyngeal carcinoma. *Tissue Antigens*. 2007;70(1):12–17.
- Zoodsma M, Nolte IM, Schipper M, et al. Interleukin-10 and Fas polymorphisms and susceptibility for (pre)neoplastic cervical disease. *Int J Gynecol Cancer*. 2005;15(Suppl 3):282–290.
- Hohaus S, Giachelia M, Massini G, et al. Clinical significance of interleukin-10 gene polymorphisms and plasma levels in Hodgkin lymphoma. *Leuk Res*. 2009;33(10):1352–1356.
- Jin L, Sturgis EM, Cao X, et al. Interleukin-10 promoter variants predict HPV-positive tumors and survival of squamous cell carcinoma of the oropharynx. *FASEB J*. 2013;27(6):2496–2503.

OncoTargets and Therapy**Publish your work in this journal**

OncoTargets and Therapy is an international, peer-reviewed, open access journal focusing on the pathological basis of all cancers, potential targets for therapy and treatment protocols employed to improve the management of cancer patients. The journal also focuses on the impact of management programs and new therapeutic agents and protocols on

Submit your manuscript here: <http://www.dovepress.com/oncotargets-and-therapy-journal>

patient perspectives such as quality of life, adherence and satisfaction. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Dovepress