Six-minute stepper test: a valid clinical exercise tolerance test for COPD patients

JM Grosbois^{1,2}C Riquier³B Chehere⁴J Coquart⁵H Béhal⁶F Bart²B Wallaert^{2,3}C Chenivresse³

¹FormAction Santé, Pérenchies, France; ²Department of Respiratory Medicine, Centre Hospitalier Germon et Gauthier, Béthune, France; ³Department of Respiratory Medicine Immunology and Allergy, Centre Hospitalier Universitaire de Lille, Competence Center for rare lung diseases, University Lille 2, Lille, France; ⁴EA 7369, URePSSS, Multidisciplinary Research Unit in Sport Health Society, University Lille 2, Lille, France; ⁵Faculty of Sport Sciences, Sports and Physical Activity, Center for Ecology and Transformation, University of Rouen, Mont Saint Aignan, France; ⁶Department of Statistical Methods and Biostatistics, Centre Hospitalier Universitaire de Lille, University of Lille Nord, Lille, France

Introduction: Exercise tolerance testing is an integral part of the pulmonary rehabilitation (PR) management of patients with chronic obstructive pulmonary disease (COPD). The 6-minute stepper test (6MST) is a new, well-tolerated, reproducible exercise test, which can be performed without any spatial constraints.

Objective: The aim of this study was to compare the results of the 6MST to those obtained during a 6-minute walk test (6MWT) and cardiopulmonary exercise testing (CPET) in a cohort of COPD patients.

Methods: Ninety-one COPD patients managed by outpatient PR and assessed by 6MST, 6MWT, and CPET were retrospectively included in this study. Correlations between the number of steps on the 6MST, the distance covered on the 6MWT, oxygen consumption, and power at the ventilatory threshold and at maximum effort during CPET were analyzed before starting PR, and the improvement on the 6MST and 6MWT was compared after PR.

Results: The number of steps on the 6MST was significantly correlated with the distance covered on the 6MWT ($r=0.56$; $P<0.0001$), the power at maximum effort ($r=0.46$; $P<0.0001$), and oxygen consumption at maximum effort ($r=0.39$; $P<0.005$). Performances on the 6MST and 6MWT were significantly improved after PR (570 vs 488 steps, $P=0.001$ and 448 vs 406 m, respectively; $P<0.0001$). Improvements of the 6MST and 6MWT after PR were significantly correlated ($r=0.34$; $P=0.03$).

Conclusion: The results of this study show that the 6MST is a valid test to evaluate exercise tolerance in COPD patients. The use of this test in clinical practice appears to be particularly relevant for the assessment of patients managed by home PR.

Keywords: 6-minute stepper test, 6-minute walk test, exercise tolerance, pulmonary rehabilitation, cardiopulmonary exercise testing, validity

Introduction

Chronic obstructive pulmonary disease (COPD) is a chronic lung disease marked by breathlessness on exertion, with systemic consequences and an increased risk of disability and death, especially as COPD is frequently associated with comorbidities (cardiovascular disease, obesity, and Type 2 diabetes).^{1,2} Pulmonary rehabilitation (PR), comprising exercise retraining, resumption of physical activity, therapeutic education, and psychosocial support, is an integral part of the management of COPD patients. This global management allows improvement of dyspnea, exercise tolerance, and quality of life.³⁻⁶

Various exercise tests are commonly used to assess the exercise tolerance of patients with lung disease in the context of PR. Cardiopulmonary exercise testing (CPET) is the standard test allowing investigation of the pathophysiological mechanisms responsible for dyspnea and evaluation of aerobic capacity.⁷ However, CPET requires specialized and expensive equipment and qualified personnel, which limit its availability. The 6-minute

Correspondence: JM Grosbois
Service de Pneumologie, Centre
Hospitalier Germon et Gauthier,
Béthune 62408, France
Email jmgrosbois@formactionsante.com

walk test (6MWT) is a validated test for most chronic lung diseases; it is easy to perform and does not require any specific equipment. The 6MWT is sensitive and reproducible and is commonly used to evaluate the course of exercise tolerance in the context of out/inpatient PR with a well-defined minimal clinically important difference.^{8–11} According to the American Thoracic Society (ATS) guidelines,¹² the 6MWT must be performed in an unobstructed 30 m hallway, but such a space is rarely available in the patient's home¹³ or in the physician's office, or even in some respiratory medicine departments. A recent study has however proposed performing the 6MWT over a distance of 10 m.¹⁴

To overcome these technical and spatial limitations, a 6-minute stepper test (6MST) has been proposed to evaluate exercise tolerance. This test requires only a limited amount of space and equipment and is feasible, easy to perform, sensitive, well tolerated, and reproducible in COPD patients.^{15,16} The 6MST has been used to assess exercise tolerance in the context of PR.^{17,18} However, the correlation between the 6MST and validated indices used to measure exercise capacity (6MWT and CPET) has not yet been established.

The primary objective of this study was to analyze the correlations between exercise tolerance data provided by the 6MST, 6MWT, and CPET in a population of COPD patients managed by outpatient PR. The secondary objective was to analyze and compare the course of exercise tolerance measured by 6MST and 6MWT before and after PR.

Material and methods

This monocentric retrospective study included COPD patients managed by outpatient PR. Patients included in this study were those who had completed the entire PR program and those for whom pre-rehabilitation 6MWT, 6MST, and CPET assessment data were available.

The severity of COPD was evaluated by Global Initiative for Chronic Obstructive Lung Disease (GOLD) stage.¹⁹ Patients were divided into two groups according to GOLD stage: mild-to-moderate COPD (Stages 1 and 2) and severe-to-very-severe COPD (Stages 3 and 4).

All data were collected prospectively and were entered into our rehabilitation computerized medical records. The study protocol was approved by the Observational Research Protocol Evaluation Committee of the French Language Society of Pulmonology, France. As the study data was collected as part of a pulmonary rehabilitation program prior to this retrospective study commencing the Observational Research Protocol Evaluation Committee advised written

informed consent did not need to be obtained from the patients.

Pulmonary function tests

Spirometry was performed before PR in clinically stable patients, according to European guidelines.^{20,21}

Cardiopulmonary exercise testing

CPET was performed according to an incremental protocol on a cycle ergometer (Ergometrics 800; ergoline, Bitz, Germany), with blood pressure and electrocardiographic monitoring (Medcard; Medisoftware®, Dinant, Belgium), according to the usual guidelines.^{22,23} The protocol included a 3-minute rest period, a 3-minute unloaded pedaling period, an incremental ramp exercise (10–15 W/min) pursued to a symptom-limited end point, and a 3-minute recovery time. Inspired and expired gases were collected by a mouthpiece connected to a gas analyzer (Ergocard, Medisoftware®). Maximal oxygen consumption ($\dot{V}O_{2\max}$) and oxygen consumption at the ventilatory threshold (VT) were expressed in mL/min/kg. VT was defined by a nonlinear increase in ventilation ($\dot{V}_E$) during exercise and was determined by the level of effort at which carbon dioxide production ($\dot{V}CO_2$) increased more rapidly than $\dot{V}O_2$,²⁴ or by an increase in the respiratory equivalent for oxygen ($\dot{V}_E/\dot{V}O_2$) with no increase in the respiratory equivalent for carbon dioxide ($\dot{V}_E/\dot{V}CO_2$).²⁵ All CPET were limited by dyspnea, fatigue, or both and provided at least one objective criterion of maximum among the following: breathing reserve <15%, peak heart rate >90% of predicted, peak lactate >7 mEq/L, peak exercise partial pressure of oxygen (PaO_2) in arterial blood <55 mmHg, or peak $\dot{V}_E/\dot{V}O_2$ >35.²²

Six-minute walk test

The 6MWT was performed over a “rectangular” distance of 70 m in the respiratory medicine department. With the exception of this point, the ATS guidelines for 6MWT were followed.¹² After resting for 5 minutes, continuous monitoring of transcutaneous oxygen saturation (SpO_2) and heart rate by a pulse oximeter (Oxymontre 3100; Nonin®, Plymouth, MN, USA) were performed.

Six-minute stepper test

The 6MST aims at measuring the number of steps performed on a stepper in 6 minutes using a protocol that is equivalent to the 6MWT protocol. A step was defined as a single complete movement of raising one foot and putting it down. The stepper (Go Sport®, Grenoble, France)¹⁶ was placed on the ground facing a wall to allow patients to maintain their balance by

placing their fingers on the wall. The step height was set at 20 cm. Before starting the test, patients got accustomed to the stepper for 2 minutes. The protocol included a 3-minute rest period and a 6-minute stepping period. Patients received standardized instructions adapted from the 6MWT ATS instructions,¹⁵ advising them to make the most number of steps they could in 6 minutes. The number of steps performed in 6 minutes was recorded. Monitoring of heart rate and SpO₂ by a pulse oximeter (Oxymontre 3100; Nonin) was performed each minute. An investigator, a member of the paramedical team, remained behind the patient throughout the test.

The 6MST and 6MWT were conducted by the same team to avoid between-investigator variability.

Pulmonary rehabilitation

Patients completed a 6-week course of outpatient PR at a rate of four sessions per week, 3 hours per session, comprising a combination of exercise retraining, therapeutic education, and psychosocial support.^{6,17}

Statistical analysis

Statistical analysis was performed with PRISM software (GraphPad®, La Jolla, CA, USA) and SAS version 9.3 software (SAS Institute, Cary, NC, USA). Quantitative parameters were expressed in terms of mean and standard deviation. Qualitative parameters were expressed in terms of frequencies and percentages. The distribution of quantitative parameters was verified graphically and by a Shapiro–Wilk test.

Pearson's or Spearman's correlation coefficient was used to evaluate the correlation between the results of the various tests performed by the patients (CPET, 6MWT, and 6MST). A linear regression was performed to analyze the independent correlation between the 6MST (dependent variable) and the 6MWT distance according to age, body mass index, forced expiratory volume in one second/forced vital capacity (independent variables).

Student's *t*-test for paired data or Wilcoxon signed rank test was used to compare the course of the 6MST and 6MWT results before and after PR, according to the distribution of the variables. All tests were initially performed on the overall population, and then according to the severity of COPD exclusively for the 6MST and 6MWT results. The significance level was set at 5%.

Results

One hundred seventy-seven COPD patients were managed by outpatient PR between January 2010 and December 2014. Ninety-one (51%) of these patients completed a pre-rehabilitation assessment comprising all three tests

Table 1 Patient characteristics (n=91)

Male ^a	76 (83.5)
Age (years) ^b	60.3±9.3
Current smoker	8 (9)
Ex-smoker	83 (91)
COPD stage ^a	
GOLD 1	7 (8)
GOLD 2	53 (58)
GOLD 3	20 (22)
GOLD 4	11 (12)
BODE Index ^b	3.7±1.82
LTOT ^a	4 (4)
NIV ^a	1 (1)
FVC (L) ^b	2.89±0.85
FVC (% predicted) ^b	73±24
FEV ₁ (L) ^b	1.69±0.59
FEV ₁ (% predicted) ^b	55±19
FEV ₁ /FVC (% ^b)	58.2±11.7
BMI (kg/m ²) ^b	27.3±6.3
>30 ^a	30 (33)
25–29.9 ^a	25 (27.5)
21–24.9 ^a	20 (22)
<21 ^a	16 (17.5)
Cardiovascular comorbidities ^a	60 (66)
Hypertension ^a	26 (29)
Peripheral arterial disease ^a	8 (8.7)
Myocardial infarction ^a	7 (8)
Diabetes ^a	11 (12)

Notes: ^aResults expressed as number (%), ^bmean ± standard deviation.

Abbreviations: BMI, body mass index; GOLD, Global Initiative for Chronic Obstructive Lung Disease; LTOT, long term oxygen therapy; NIV, non invasive ventilation; FVC, forced vital capacity; FEV₁, forced expiratory volume in one second.

(6MST, 6MWT, and CPET) and were included in this study. Another 86 patients (49%) were excluded due to missing CPET (n=39) or 6MWT data (n=21), or because a stress test rather than CPET was performed (n=26). The characteristics of the 91 patients are presented in Table 1.

A statistically significant correlation was observed between the number of steps on the 6MST and the distance covered on the 6MWT ($r=0.56$; $P<0.0001$; Figure 1),

Figure 1 Correlation between the number of steps on the 6MST and the distance covered on the 6MWT before PR.

Abbreviations: 6MST, 6-minute stepper test; 6MWT, 6-minute walk test; PR, pulmonary rehabilitation.

Table 2 Test results before PR and correlations

Test parameters	Correlation with 6MST			Correlation with 6MWT	
	All (n=91), mean $\pm$ SD	r	P-value	r	P-value
6MST (steps)	488 $\pm$ 149	NA	NA	0.56	<0.0001
6MWT (m)	406 $\pm$ 86	0.56	<0.0001	NA	NA
Power-VT (W)	52 $\pm$ 23	0.36	<0.005	0.32	0.009
VO ₂ -VT (mL/min/kg)	11.4 $\pm$ 3.3	0.26	NS	0.35	0.001
Power-max (W)	86 $\pm$ 33	0.46	<0.0001	0.39	0.0002
VO ₂ max (mL/min/kg)	15.2 $\pm$ 4.8	0.39	<0.005	0.38	0.03

Abbreviations: PR, pulmonary rehabilitation; 6MST, 6-minute stepper test; 6MWT, 6-minute walk test; VO₂-VT, oxygen consumption at ventilatory threshold; VO₂max, oxygen consumption at maximum effort; Power-VT, Power at ventilatory threshold; Power-max, Power at maximum effort; NA, not applicable; NS, not significant.

independent of the severity of COPD (Table 2). This correlation was observed in patients with mild-to-moderate COPD and in patients with severe-to-very-severe COPD (Table 3). The number of steps on the 6MST was correlated with power (W) at the VT and at maximum effort, and VO₂max (Tables 2 and 4). Similarly, the distance covered on the 6MWT was correlated with power and VO₂ at the VT and at VO₂max (Table 2). A linear regression showed that the 6MST was related to the 6MWT distance and peak exercise power, independent of age, lung function, and body mass index (Table 4).

For the overall population, a significant improvement of the 6MWT distance was observed, increasing from 406 $\pm$ 86 m before PR to 448 $\pm$ 95 m after PR ($P<0.0001$), together with a significant increase in the number of steps on the 6MST, increasing from 488 $\pm$ 149 steps before PR to 570 $\pm$ 161 steps after PR ($P<0.0001$). A statistically significant correlation was observed for all patients between the number of steps on the 6MST and the distance covered on the 6MWT after PR ($r=0.75$; $P<0.0001$; Figure 2). Subgroup analysis revealed similar results for patients with mild-to-moderate COPD ($r=0.77$; $P<0.0001$) and patients with severe-to-very-severe COPD ($r=0.76$; $P<0.0001$). As shown in Figure 3, improvement of the 6MST after PR was correlated with improvement of the 6MWT in all patients (absolute values: $r=0.34$; $P=0.03$; Figure 3; percentage change: $r=0.32$; $P=0.001$).

Discussion

This study shows that the number of steps on the 6MST was significantly correlated with the distance covered on the 6MWT, VO₂max, and power at VT and at maximum effort, independent of the severity of COPD. These results also show a significant correlation between improvements of performances on the 6MWT and 6MST after outpatient PR.

Assessment of exercise tolerance is an integral part of the evaluation of COPD patients during PR.^{4,6} Despite the demonstrated value of the 6MWT, its technical constraints do not always allow use and regular repetition of this test during patient follow-up, especially in the physician's or physiotherapist's office, or at the patient's home.^{12,13} Consequently, various teams have developed alternatives to these tests, including the 6MST.

In line with previous studies,^{15,26} our results demonstrated a significant correlation between the number of steps on the 6MST and the distance covered on the 6MWT. These correlations were independent of the severity of COPD. Delourme et al²⁶ also showed that these correlations remained significant independent of the type of interstitial lung disease (ILD). All of these results indicate that the 6MST can be used in a wide range of patients with varying degrees of severity of chronic lung disease (COPD and ILD).

The 6MWT is considered to be a submaximal test, as the subject is free to walk at his or her own rate, but it has been shown that the most severe COPD patients achieve an

Table 3 Test results before PR according to the severity of COPD and correlations with the 6MST

Test parameters	Stage 1 and 2 COPD (n=60), mean $\pm$ SD			Stage 3 and 4 COPD (n=31), mean $\pm$ SD		
		r	P-value		r	P-value
6MST (steps)	502.6 $\pm$ 145.9	NA	NA	461.1 $\pm$ 154.5	NA	NA
6MWT (m)	401.1 $\pm$ 89.3	0.53	$P<0.0001$	414.6 $\pm$ 81.7	0.55	$P<0.005$
Power-VT (W)	56.2 $\pm$ 24.08	0.43	$P<0.005$	44.5 $\pm$ 20	0.22	NS
VO ₂ -VT (mL/min/kg)	11.5 $\pm$ 3.8	0.15	NS	11.1 $\pm$ 2.2	0.52	$P<0.05$
Power-max (W)	92.3 $\pm$ 32.3	0.45	$P<0.0005$	73.9 $\pm$ 30	0.44	$P<0.05$
VO ₂ max (mL/min/kg)	15.7 $\pm$ 5.3	0.29	NS	14.4 $\pm$ 3.7	0.62	$P<0.005$

Abbreviations: PR, pulmonary rehabilitation; 6MST, 6-minute stepper test; 6MWT, 6-minute walk test; VO₂-VT, oxygen consumption at ventilatory threshold; VO₂max, oxygen consumption at maximum effort; Power-VT, Power at ventilatory threshold; Power-max, Power at maximum effort; NA, not applicable; NS, not significant.

Table 4 Linear regression using the 6MST as a dependent variable

Test parameters	Estimate	Standard error	P-value	Squared partial correlation
Power-max (W)	1.27	0.53	0.0188	0.06
6MWT (m)	0.72	0.17	0.0001	0.16
Age (years)	0.15	0.58	0.79	0.0008
BMI (kg/m ²)	-1.60	2.52	0.52	0.005
FEV ₁ /FVC (%)	152	125	0.22	0.02

Abbreviations: 6MST, 6-minute stepper test; 6MWT, 6-minute walk test; Power-max, power at maximum effort; BMI, body mass index; FVC, forced vital capacity; FEV₁, forced expiratory volume in one second.

identical VO₂ to that observed at the end of a cycle ergometer test, by the third minute of walking.^{27–30} McGavin et al³¹ have previously validated the 12MWT in COPD patients based on the significant correlation ($r=0.52$) between the distance covered and VO_{2max}, as also confirmed by Carter et al³² and, more recently, by Luxton et al³³ ($r=0.54$ and $r=0.63$, respectively). Similar results have been reported in patients with heart failure ($r=0.42$) and respiratory insufficiency ($r=0.57$).^{34,35} This study demonstrated identical correlations between maximum values obtained during CPET and the distance covered on the 6MWT, as well as the number of steps on the 6MST.

Other clinical tests have been proposed in the literature, such as the stair climbing test,^{36,37} the sit-to-stand test,³⁸ and the step test (walking up and down a 20 cm platform for a determined duration).³⁹ In a population of COPD patients, the performance on the step test is correlated with the performance on CPET, and the distance covered on the 12MWT⁴⁰ and the 6MWT.⁴¹ We decided to perform this test on a stepper device rather than on a 20 cm platform, because the stepper allows the patient to maintain balance by lightly placing the fingers on the wall, which is more reassuring for elderly patients and/or patients with balance disorders.

Figure 2 Correlation between the number of steps on the 6MST and the distance covered on the 6MWT after PR.

Abbreviations: 6MST, 6-minute stepper test; 6MWT, 6-minute walk test; PR, pulmonary rehabilitation.

Figure 3 Correlation between improvement of the number of steps on the 6MST and improvement of the distance covered on the 6MWT after PR.

Abbreviations: 6MST, 6-minute stepper test; 6MWT, 6-minute walk test; PR, pulmonary rehabilitation.

Many studies have demonstrated the good sensitivity of the 6MWT at assessing the efficacy of outpatient or home PR in COPD patients.^{4,6} Lacasse et al^{3,42} reported gains of 56 and 48 m for outpatient PR and home PR, respectively. No consensus has been reached concerning the minimal clinically important difference (MCID) of the 6MWT: 54 m in the study by Redelmeier et al⁸ and 35 m⁹ and 26 m¹¹ in more recent studies. A 14% relative improvement has also been proposed, but appears to be less sensitive than an absolute value of 25 m.¹⁰ The gain of 42 m observed in this study is consistent with the data of the literature.^{8–11} Furthermore, a significant increase in the number of steps on the 6MST (from 488 to 570) was observed not only after outpatient PR (in agreement with previously published results in 30 COPD patients),¹⁶ but also after home PR in a population of patients with idiopathic pulmonary fibrosis,⁴³ unselected patients with more severe chronic lung disease (80% on long term oxygen therapy and/or noninvasive ventilation),¹⁷ and in COPD patients.¹⁸ The mean improvement of 79 steps on the 6MST in this study is superior to the 40-step MCID proposed by Pichon et al⁴⁴ based on the analysis of the results of 62 COPD patients managed by inpatient PR. The performances on the two clinical exercise tolerance tests were significantly improved after PR: 79 steps on the 6MST and 42 m on the 6MWT, and these improvements were significantly correlated with each other. Coquart et al¹⁶ suggested that familiarization with the 6MWT would have an impact on improved performance on the 6MST after PR due to a possible learning effect. However, it should be stressed that the patients in this study did not use the stepper as a retraining tool. The improvements observed on the 6MST were consequently the result of improvement of the subject's aerobic and muscular capacities. These results suggest that,

like the 6MWT,^{4,6} the 6MST is a sensitive clinical exercise test to assess the efficacy of a PR program on the exercise tolerance of COPD patients.

This study was mainly limited by its retrospective nature. However, there were only a few data missing, and most of the variables were objective measurements collected from standardized tests. Almost half the patients were excluded from the study because all the required examinations were not performed on them. Although this could introduce a selection bias, there was no statistical difference between excluded and included patients concerning demographic, clinical, comorbidities, and pulmonary function data (data not shown).

Conclusion

This study demonstrates not only a correlation between the number of steps on the 6MST and the distance covered on the 6MWT, but also a correlation with power and $\text{VO}_{2\text{max}}$ of the CPET for COPD patients regardless of their stage of severity. The improvements on the 6MST and 6MWT observed after PR were also correlated, confirming the value of this new, sensitive, and reproducible test, which is easy to perform without any spatial constraints, for assessment of exercise tolerance in COPD patients, in routine respiratory medicine practice, and especially, before and after short- and long-term PR management.

Disclosure

JM Grosbois received financial support from Adair, France Oxygène, Homeperf, LVL, Orkyn, Santélys, SOS Oxygène, Sysmed, and VitalAire for home-based PR program. The authors report no other conflicts of interest in this work.

References

1. Vanfleteren LE, Spruit MA, Groenen M, et al. Clusters of comorbidities based on validated objective measurements and systemic inflammation in patients with chronic obstructive pulmonary disease. *Am J Respir Crit Care Med*. 2013;187(7):728–735.
2. Van Remoortel H, Hornikx M, Langer D, et al. Risk factors and comorbidities in the preclinical stages of chronic obstructive pulmonary disease. *Am J Respir Crit Care Med*. 2013;189(1):30–38.
3. Lacasse Y, Goldstein R, Lasserson TJ, Martin S. Pulmonary rehabilitation for chronic obstructive pulmonary disease. *Cochrane Database Syst Rev*. 2006;(4):CD003793.
4. Société de Pneumologie de Langue Française. Recommandations pour la pratique clinique. Prise en charge de la BPCO [Recommendation for the clinical practice management of COPD]. *Rev Mal Respir*. 2010;27(Suppl 1):S1–S76, I–XXXI. French.
5. van Ranst D, Otten H, Meijer JW, van't Hul AJ. Outcome of pulmonary rehabilitation in COPD patients with severely impaired health status. *Int J Chron Obstruct Pulmon Dis*. 2011;6:647–657.
6. Spruit MA, Singh SJ, Garvey C, et al. An official American Thoracic Society/European Respiratory Society statement: key concepts and advances in pulmonary rehabilitation. *Am J Respir Crit Care Med*. 2013;188(8):e13–e64.
7. Aguilaniu B, Wallaert B. *EFX: De L'interprétation à La Décision Médicale*. Paris, France: Margaux Orange; 2015.
8. Redelmeier DA, Bayoumi AM, Goldstein RS, Guyatt GH. Interpreting small differences in functional status: the six minute walk test in chronic lung disease patients. *Am J Respir Crit Care Med*. 1997;155(4):1278–1282.
9. Puhan MA, Mador MJ, Held U, Goldstein R, Guyatt GH, Schünemann HJ. Interpretation of treatment changes in 6-minute walk distance in patients with COPD. *Eur Respir J*. 2008;32(3):637–643.
10. Holland AE, Hill CJ, Rasekaba T, Lee A, Naughton MT, McDonald CF. Updating the minimal important difference for six-minute walk distance in patients with chronic obstructive pulmonary disease. *Arch Phys Med Rehabil*. 2010;91(2):221–225.
11. Puhan MA, Chandra D, Mosenifar Z, et al. The minimal important difference of exercise tests in severe COPD. *Eur Respir J*. 2011;37(4):784–790.
12. ATS Committee on Proficiency Standards for Clinical Pulmonary Function Laboratories. ATS statement: guidelines for the six-minute walk test. *Am J Respir Crit Care Med*. 2002;166(1):111–117.
13. Holland AE, Rasekaba T, Fiore JF, Burge AT, Lee AL. The 6-minute walk distance cannot be accurately assessed at home in people with COPD. *Disabil Rehabil*. 2015;37(12):1102–1106.
14. Beekman E, Mesters I, Gosselink R, et al. The first reference equations for the 6-minute walk distance over a 10 m course. *Thorax*. 2014;69(9):867–868.
15. Borel B, Fabre C, Saison S, Bart F, Grosbois J-M. An original field evaluation test for chronic obstructive pulmonary disease population: the six-minute stepper test. *Clin Rehabil*. 2010;24(1):82–93.
16. Coquart JB, Lemaître F, Castres I, Saison S, Bart F, Grosbois J-M. Reproducibility and sensitivity of the 6-minute stepper test in patients with COPD. *COPD*. 2015;12(5):533–538.
17. Grosbois J-M, Le Rouzic O, Monge E, Bart F, Wallaert B. [Comparison of home-based and outpatient, hospital-based, pulmonary rehabilitation in patients with chronic respiratory diseases.] *Rev Pneumol Clin*. 2013;69(1):10–17. French.
18. Grosbois JM, Gicquello A, Langlois C, et al. Long term evaluation of home-based pulmonary rehabilitation in patients with chronic obstructive pulmonary disease. *Int J Chron Obstruct Pulmon Dis*. 2015;10:2037–2044.
19. Rabe KF, Hurd S, Anzueto A, et al. Global strategy for the diagnosis, management, and prevention of chronic obstructive pulmonary disease: GOLD executive summary. *Am J Respir Crit Care Med*. 2007;176(6):532–555.
20. Quanjer PH, Tammeling GJ, Cotes JE, Pedersen OF, Peslin R, Yernault JC. Lung volumes and forced ventilatory flows. Report Working Party Standardization of Lung Function Tests, European Community for Steel and Coal. Official Statement of the European Respiratory Society. *Eur Respir J Suppl*. 1993;16:5–40.
21. Miller MR, Hankinson J, Brusasco V, et al. Standardisation of spirometry. *Eur Respir J*. 2005;26(2):319–338.
22. Aguilaniu B, Richard R, Costes F, et al. Méthodologie et Pratique de l'Exploration Fonctionnelle à l'eXercice (EFX) [Cardiopulmonary exercise testing]. *Rev Mal Respir*. 2007;24(3 Pt 2):2S111–2S160. French.
23. American Thoracic Society; American College of Chest Physicians. ATS/ACCP statement on cardiopulmonary exercise testing. *Am J Respir Crit Care Med*. 2003;167(2):211–277.
24. Beaver WL, Wasserman K, Whipp BJ. A new method for detecting anaerobic threshold by gas exchange. *J Appl Physiol*. 1986;60(6):2020–2027.
25. Wasserman K, Whipp BJ, Koyl SN, Beaver WL. Anaerobic threshold and respiratory gas exchange during exercise. *J Appl Physiol*. 1973;35(2):236–243.
26. Delourme J, Stervinou-Wemeau L, Salleron J, Grosbois JM, Wallaert B. Six-minute stepper test to assess effort intolerance in interstitial lung diseases. *Sarcoidosis Vasc Diffuse Lung Dis*. 2012;29(2):107–112.
27. Troosters T, Vilaro J, Rabinovich R, et al. Physiological responses to the 6-min walk test in patients with chronic obstructive pulmonary disease. *Eur Respir J*. 2002;20(3):564–569.

28. Turner SE, Eastwood PR, Cecins NM, Hillman DR, Jenkins SC. Physiologic responses to incremental and self-paced exercise in COPD: a comparison of three tests. *Chest*. 2004;126(3):766–773.
29. Casas A, Vilaro J, Rabinovich R, et al. Encouraged 6-min walking test indicates maximum sustainable exercise in COPD patients. *Chest*. 2005;128(1):55–61.
30. Cote CG, Pinto-Plata VM, Marin JM, Nekach H, Dordelly LJ, Celli BR. The modified BODE index: validation with mortality in COPD. *Eur Respir J*. 2008;32(5):1269–1274.
31. McGavin CR, Gupta SP, McHardy GJ. Twelve-minute walking test for assessing disability in chronic bronchitis. *Br Med J*. 1976;1(6013):822–823.
32. Carter R, Holiday DB, Stocks J, Grothues C, Tiep B. Predicting oxygen uptake for men and women with moderate to severe chronic obstructive pulmonary disease. *Arch Phys Med Rehabil*. 2003;84(8):1158–1164.
33. Luxton N, Alison JA, Wu J, Mackey MG. Relationship between field walking tests and incremental cycle ergometry in COPD. *Respirology*. 2008;13(6):856–862.
34. Guyatt GH, Sullivan MJ, Thompson PJ, et al. The 6-minute walk: a new measure of exercise capacity in patients with chronic heart failure. *Can Med Assoc J*. 1985;132(8):919–923.
35. Guyatt GH, Thompson PJ, Berman LB, et al. How should we measure function in patients with chronic heart and lung disease? *J Chronic Dis*. 1985;38(6):517–524.
36. Pollock M, Roa J, Benditt J, Celli B. Estimation of ventilatory reserve by stair climbing. A study in patients with chronic airflow obstruction. *Chest*. 1993;104(5):1378–1383.
37. Villiot-Danger E. [A stairclimbing test in COPD patients assessment]. *Rev Mal Respir*. 2009;26(5):530–536. French.
38. Jones SE, Kon SS, Canavan JL, et al. The five-repetition sit-to-stand test as a functional outcome measure in COPD. *Thorax*. 2013;68(11):1015–1020.
39. Dal Corso S, Duarte SR, Neder JA, et al. A step test to assess exercise-related oxygen desaturation in interstitial lung disease. *Eur Respir J*. 2007;29(2):330–336.
40. Swinburn CR, Wakefield JM, Jones PW. Performance, ventilation, and oxygen consumption in three different types of exercise test in patients with chronic obstructive lung disease. *Thorax*. 1985;40(8):581–586.
41. Pessoa BV, Arcuri JF, Labadessa IG, Costa JN, Santanin AC, Di Lorenzo VA. Validity of the six-minute step test of free cadence in patients with chronic obstructive pulmonary disease. *Braz J Phys Ther*. 2014;18(3):228–236.
42. Lacasse Y, Wong E, Guyatt GH, King D, Cook DJ, Goldstein RS. Meta-analysis of respiratory rehabilitation in chronic obstructive pulmonary disease. *Lancet*. 1996;348(9035):1115–1119.
43. Rammaert B, Leroy S, Cavestri B, Wallaert B, Grosbois J-M. Home-based pulmonary rehabilitation in idiopathic pulmonary fibrosis. *Rev Mal Respir*. 2011;28(7):e52–e57.
44. Pichon R, Beaumont M, Le Ber-Moy C, Péran L, Couturaud F. Détermination d'une différence minimale importante pour le stepper-test de six minutes chez les patients atteints de BPCO [Determination of a significant minimal difference for the 6 minute stepper-test in patients with COPD]. *Rev Mal Respir*. 2015;32:A75. French.

International Journal of COPD

Publish your work in this journal

The International Journal of COPD is an international, peer-reviewed journal of therapeutics and pharmacology focusing on concise rapid reporting of clinical studies and reviews in COPD. Special focus is given to the pathophysiological processes underlying the disease, intervention programs, patient focused education, and self management protocols.

Submit your manuscript here: <http://www.dovepress.com/international-journal-of-chronic-obstructive-pulmonary-disease-journal>

Dovepress

This journal is indexed on PubMed Central, MedLine and CAS. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.