Clinically relevant genetic variants of drug-metabolizing enzyme and transporter genes detected in Thai children and adolescents with autism spectrum disorder

Sadeep Medhasi¹⁻³
 Ekawat Pasomsub⁴
 Natchaya Vanwong^{1,2}
 Nattawat Ngamsamut⁵
 Apichaya Puangpetch^{1,2}
 Montri Chamnanphon^{1,2}
 Yaowaluck Hongkaew^{1,2}
 Penkhae Limsila⁵
 Darawan Pinthong³
 Chonlaphat Sukasem^{1,2}

¹Division of Pharmacogenomics and Personalized Medicine, Department of Pathology, Faculty of Medicine Ramathibodi Hospital, Mahidol University, Bangkok, Thailand; ²Laboratory for Pharmacogenomics, Somdech Phra Debaratana Medical Center, Faculty of Medicine Ramathibodi Hospital, Mahidol University, Bangkok, Thailand; ³Department of Pharmacology, Faculty of Science, Mahidol University, Bangkok, Thailand; ⁴Division of Virology, Department of Pathology, Faculty of Medicine Ramathibodi Hospital, Mahidol University, Bangkok, Thailand; ⁵Yuwaprasart Waithayopatham Child and Adolescent Psychiatric Hospital, Department of Mental Health Services, Ministry of Public Health, Samut Prakarn, Thailand

Correspondence: Chonlaphat Sukasem
 Division of Pharmacogenomics and Personalized Medicine, Department of Pathology, Faculty of Medicine Ramathibodi Hospital, Mahidol University, 270, Rama 6 Road, Thung Phaya Thai, Ratchathewi, Bangkok, 10400, Thailand
 Tel +66 2 200 4331
 Fax +66 2 200 4332
 Email chonlaphat.suk@mahidol.ac.th

Abstract: Single-nucleotide polymorphisms (SNPs) among drug-metabolizing enzymes and transporters (DMETs) influence the pharmacokinetic profile of drugs and exhibit intra- and interethnic variations in drug response in terms of efficacy and safety profile. The main objective of this study was to assess the frequency of allelic variants of drug absorption, distribution, metabolism, and elimination-related genes in Thai children and adolescents with autism spectrum disorder. Blood samples were drawn from 119 patients, and DNA was extracted. Genotyping was performed using the DMET Plus microarray platform. The allele frequencies of the DMET markers were generated using the DMET Console software. Thereafter, the genetic variations of significant DMET genes were assessed. The frequencies of SNPs across the genes coding for DMETs were determined. After filtering the SNPs, 489 of the 1,931 SNPs passed quality control. Many clinically relevant SNPs, including *CYP2C19**2, *CYP2D6**10, *CYP3A5**3, and *SLCO1B1**5, were found to have frequencies similar to those in the Chinese population. These data are important for further research to investigate the interpatient variability in pharmacokinetics and pharmacodynamics of drugs in clinical practice.

Keywords: Thai population, ADME, pharmacokinetics, autism spectrum disorder, microarray, pharmacogenetics

Introduction

Autism spectrum disorder (ASD) is a global health concern where the patient demonstrates three core features: social withdrawal, verbal and nonverbal communication impairment, and stereotype behavior with restricted and repetitive activities and/or interests.¹ Results have shown that males are more likely to have autism than females, with a prevalence ratio of 4:1.^{2,3} Atypical antipsychotics, including risperidone and aripiprazole, both US Food and Drug Administration approved, have been used as a pharmacologic therapy for the treatment of behavioral symptoms (irritability and aggression) in addition to the psychoeducational therapy among children and adolescents.^{4,5} However, children and adolescents treated with antipsychotic drugs may be predisposed to side effects, such as weight gain, metabolic effects, endocrinal dysfunction, and tardive dyskinesia.^{6,7} Genetic variations in drug-metabolizing enzymes (DMEs), drug transporters, and drug targets are important determinants of efficacy and toxicity in pharmacotherapy among patients.⁸ Individual variability in drug response poses a major challenge in drug therapy, drug development, and drug regulation.⁹ A better account of genetic variability and implementation of personalized

medicine are essential in individual patients to improve the drug therapy by enabling clinicians to prescribe the right medication in the right dose to achieve maximum therapeutic efficacy with minimum adverse drug reactions.¹⁰ Pharmacogenetics is the study of contribution of genetic differences to drug response between individuals. Advances in the field of molecular genotyping, such as DNA chips and microarrays, have facilitated the analysis of large genetic variations.¹¹ Single-nucleotide polymorphisms (SNPs) are an important marker for pharmacogenetic studies, which can be used for phenotypic association with a functional variant and for optimizing personalized drug therapy.^{12,13} Investigations on animal models of autism have revealed the association between polymorphism in serotonin transporter (5-HTT) and autism.¹⁴ In humans, genetic studies in patients with ASD have revealed the polymorphisms in *CYP1A2*1F* and its association with susceptibility to ASD,¹⁵ and another polymorphism contributing to ASD is *COMT* Val58Met.¹⁶

Polymorphisms in DMEs and transporters (DMETs) across different populations have been reported with varying frequencies of allelic variation, suggesting inter-ethnic differences in response to drug therapy. Patients are classified as poor metabolizer (PM), intermediate metabolizer, extensive metabolizer, and ultrarapid metabolizer, leading to no activity, reduced activity, normal activity, and increased activity, respectively, based on the DME genetic polymorphisms.¹⁷ The most common cytochrome P450 polymorphisms are found in *CYP2C9*, *CYP2C19*, and *CYP2D6* as identified to date (<http://www.cypalleles.ki.se>). The important Phase II enzymes exhibiting genetic polymorphisms, which determine the fate of the drugs, include uridine diphosphate glucuronosyltransferase (UGT), glutathione *S*-transferase (GST), sulfotransferase, *N*-acetyltransferase (NAT), and thiopurine methyltransferase (TPMT). Some of the best characterized drug transporter genes implicated in drug metabolism and response include the adenosine triphosphate (ATP)-binding cassette (ABC) family, such as *ABCB1*, *ABCC1*, *ABCC2*, and *ABCG2*, and solute carrier (SLC) family proteins, such as *OCT1*, *OATP1B1*, and *OAT*.^{18,19} Several pharmacogenetic biomarkers of DMETs are documented in PharmGKB (<https://www.pharmgkb.org/>), with evidence from different populations. There is a considerable difference in the frequency of genetic polymorphisms associated with drug response, across and within populations.²⁰ The DMET genes have greater interindividual variability in expression as compared to non-DMET genes.²¹ A study investigating the frequencies of DMET SNPs among Thais as well as

compared to other ethnic populations revealed significant differences in the distribution of several clinically important genetic polymorphisms among absorption, distribution, metabolism, and elimination genes.²² Integrating pharmacogenetics into clinical practice has proven to be a cost-saving and improved therapeutic outcome for patients, with the avoidance of medication adverse effects.²³ There is a critical need to understand and establish the frequency of genetic variants and implement the knowledge of genetic variation in DMET genes in the Thai population to construct dosing guidelines, so as to improve the clinical efficacy and minimize the occurrence of adverse effects.

In this study, we used the DMET™ Plus GeneChip microarray platform (Affymetrix Inc., Santa Clara, CA, USA), which is used to genotype markers across genes involved in drug pharmacokinetics. The main aim of this study was to investigate the frequency of DMET genetic variations in the Thai population. The genetic database derived from this study will help in determining the pattern of SNPs in DMET genes among Thai population and correlate the genetic information with the drug response and adverse side effects to produce the significant clinical associations.

Materials and methods

Participants

Blood samples were collected from the 119 Thai children and adolescents with ASD (according to the *Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition* criteria) at Yuwaprasart Waithayopatham Child and Adolescent Psychiatric Hospital. Participants were included who were prescribed with risperidone once or twice a day and the dosing was flexible as well. Patients with severe physical disabilities and currently being treated with other antipsychotic drugs were excluded from the study. However, stimulants were allowed in the case of co-occurring attention deficit/hyperactivity disorder. The purpose and experimental procedures were explained to the parents of the participants before obtaining written informed consent from all of the participants' parents. The study was conducted in accordance with the principles of the Declaration of Helsinki and locally approved by the Ramathibodi Ethics Committee (Bangkok, Thailand).

Sample collection and genomic DNA (gDNA) extraction

Blood samples were collected into EDTA tubes with the assistance of a trained laboratory technician and were then brought to the laboratory of Pharmacogenomics and Personalized Medicine, Ramathibodi Hospital, in a well-insulated

ice box. The samples were then stored at $<-70^{\circ}\text{C}$ for the subsequent extraction of gDNA.

gDNA was isolated from EDTA-whole blood using a MagNA Pure Compact System (Hoffman-La Roche Ltd., Basel, Switzerland), according to the manufacturer's instructions, and quantified by a NanoDrop™ 1000 Spectrophotometer at 260 nm (Thermo Fisher Scientific, Waltham, MA, USA).

Genotyping and marker selection

Samples were genotyped using the DMET platform (DMET Plus) according to the standard protocol described by the manufacturer. The DMET™ Plus GeneChip enables the genotyping of 1,936 genetic variants (1,931 SNPs and five copy number variations) of 231 genes having functional significance in Phase I and Phase II DME genes and drug transporter coding genes.²⁰ The DMET Plus platform examines the genetic variations, including biallelic and triallelic SNPs, copy number variations, and insertions/deletions.²⁴ Genotype analysis using the DMET Plus platform is based on an efficient and comprehensive molecular inversion probe technology.^{25,26} DNA samples on the DMET Plus array were scanned with the GeneChip Scanner 3000 (Affymetrix Inc.). The genotype profile of 1,931 SNPs was generated by DMET™ Console Software® (Version 1.3; Affymetrix Inc.).

Quality control

Quality control of the samples and SNPs was performed as shown in Figure 1. Individual samples were considered passed or in bounds if they had genotyping calls $>90\%$. The markers with genotyping call rate $<95\%$, deviation from Hardy-Weinberg equilibrium (HWE) at $P>0.001$, minor allele frequency (MAF) <0.05 , and variants on chromosome X were discarded.

Statistical analysis

Data are presented as median and ranges, or as otherwise specified. The Haploview 4.2 software was used for checking HWE, along with MAF, and filtering markers according to the criteria set for quality control.²⁷

Results

Patient characteristics and selection of markers

Most of the participants were males (87.1%) and had a median age of 8.8 years (range 3.3–18.3 years). A total of 1,681 SNPs passed the first quality control filter with call rate of $>95\%$. Only 489 SNPs were included for the assessment

Figure 1 Flow diagram of genetic marker selection.

Abbreviations: DMET, drug-metabolizing enzymes and transporters; HWE, Hardy-Weinberg equilibrium; MAF, minor allele frequency; SNPs, single-nucleotide polymorphisms.

of genetic prevalence after considering the quality control filter removing the SNPs that were not in HWE, $\text{MAF} < 0.05$, and SNPs on the X chromosome (46 SNPs) (Figure 1). The average call rate of the sample was 99.23%.

Allele frequency distribution of common functional polymorphisms of Phase I enzymes

The observed minor allele frequencies of common polymorphisms in CYP enzymes with their clinical or functional relevance are presented in Table 1. We identified 32 CYP genes with different markers prevalent in the Thai population. The prevalence rates of some pharmacologically and clinically relevant CYP polymorphisms observed were as follows: *CYP1A1*2C* (30.3%), *CYP1A2*1C* (30.3%), *CYP1A2*1F* (23.9%), *CYP2A6*9* (15.3%), *CYP2B6*4* (32.8%), *CYP2B6*6* (28.4%), *CYP2C19*2* (33.2%), *CYP2C19*3* (5%), *CYP2D6 100C>T* (41.2%), *CYP2D6 2850C>T* (10.1%), *CYP3A5*3* (37%), and *CYP3A7*2* (35.8%).

Allele frequency distribution of common functional polymorphisms of Phase II DMEs

We detected different allele frequencies of common Phase II DMEs, with a wide frequency range from 5% to 49% in

Table 1 Minor allele frequencies of common functional polymorphisms in major human *CYP* genes in Thai children and adolescents with autism spectrum disorder (sample size =119)

Gene	dbSNP RS ID	Common name	Minor allele frequency	Clinical relevance
<i>CYP1A1</i>	rs1048943	<i>CYP1A1</i> *2C_2454A>G(I462V)	0.303	<ul style="list-style-type: none"> In *2C variant, the AA genotype has decreased progression-free survival time compared to AG and GG genotypes in women with breast cancer treated with capecitabine and docetaxel⁴¹
	rs4646422	<i>CYP1A1</i> _134G>A(G45D)	0.08	
<i>CYP1A2</i>	rs2069514	<i>CYP1A2</i> *1C_-3860G>A(Promoter)	0.303	<ul style="list-style-type: none"> Genotypes AA and AG are associated with decreased metabolism of theophylline as compared to genotype GG in patients with asthma for *1C⁴² Allele A is associated with tardive dyskinesia in patients taking antipsychotic medications compared to allele G among smokers for *1C polymorphism⁴³ Patients with *1F variants are associated with reduced serum concentration of olanzapine and subsequent decreased response⁴⁴
	rs762551	<i>CYP1A2</i> *1F_-163C>A(Promoter)	0.239	
	rs2470890	<i>CYP1A2</i> _5347T>C(N516N)	0.206	
<i>CYP2A6</i>	rs28399433	<i>CYP2A6</i> *9_-48T>G(Promoter)	0.153	<ul style="list-style-type: none"> <i>CYP2A6</i>*9 is associated with increased plasma concentration of efavirenz⁴⁵
	rs1137115	<i>CYP2A6</i> _51G>A(V17V)	0.261	
	rs4079369	<i>CYP2A6</i> _3570C>G	0.076	
	rs8192729	<i>CYP2A6</i> _5336G>A	0.076	
<i>CYP2B6</i>	rs8192709	<i>CYP2B6</i> *2_64C>T(R22C)	0.055	<ul style="list-style-type: none"> <i>CYP2B6</i>*4 is associated with increased plasma concentration of efavirenz⁴⁶ <i>CYP2B6</i>*6 is associated with increased efavirenz plasma concentrations⁴⁷
	rs4803418	<i>CYP2B6</i> _14593C>G	0.382	
	rs2279343	<i>CYP2B6</i> *4_18053A>G(K262R)	0.328	
	rs3745274	<i>CYP2B6</i> *6_15631G>T(Q172H)	0.284	
	rs2279344	<i>CYP2B6</i> _18273G>A	0.286	
	rs8192719	<i>CYP2B6</i> _21563C>T	0.282	
<i>CYP2C19</i>	rs4244285	<i>CYP2C19</i> *2_19154G>A(P227P)	0.332	<ul style="list-style-type: none"> <i>CYP2C19</i>*2 and *3 is known to affect the metabolism and/or responses of several drugs, like amitriptyline, clopidogrel, sertraline, citalopram, and escitalopram⁴⁸⁻⁵²
	rs4986893	<i>CYP2C19</i> *3_17948G>A(W212X)	0.05	
<i>CYP2D6</i>	rs1135840	<i>CYP2D6</i> _4180G>C(S486T)	0.298	<ul style="list-style-type: none"> Amitriptyline, nortriptyline, paroxetine, codeine, and tramadol are some of the drugs whose pharmacokinetics, response, and toxicity are known to be influenced by <i>CYP2D6</i> polymorphisms⁵³⁻⁵⁷
	rs16947	<i>CYP2D6</i> _2850C>T(R296C)	0.101	
	rs1058164	<i>CYP2D6</i> _1661G>C(V136V)	0.286	
	rs1065852	<i>CYP2D6</i> _100C>T(P34S)	0.412	
	rs1080985	<i>CYP2D6</i> _1584C>G	0.071	
	rs28360521	<i>CYP2D6</i> _2178G>A	0.397	
<i>CYP3A5</i>	rs776746	<i>CYP3A5</i> *3_6986A>G(SpliceDefect)	0.37	<ul style="list-style-type: none"> Patients with *3/*3 genotype show higher concentration of tacrolimus as compared to */*1 and */*3 genotypes⁵⁸
<i>CYP3A7</i>	rs2257401	<i>CYP3A7</i> *2_26041C>G(T409R)	0.358	<ul style="list-style-type: none"> Presence of <i>CYP3A7</i>*2 is associated with higher concentration/dose ratio of tacrolimus⁵⁹

Note: dbSNP RS ID is an identification tag assigned by the National Center for Biotechnology Information to a group of single nucleotide polymorphisms (SNPs) that map to an identical location in the SNP database.

this Thai population, as shown in Table 2. Among *UGT1A1* polymorphisms, *UGT1A1**60 was the most prevalent (36.1%) in this study group. The prevalence rates of other notable clinically significant variants among several Phase II enzymes were as follows: *UGT2B7**2 (31.5%), *UGT1A7**12 (15.1%), *GSTA5* c.-31+2057C>T (35.7%), *GSTP1**B (27.5%), *GSTZ1* c.94G>A (39.5%), *NAT2**13 (45.4%), and *NAT2**6 (27.7%).

Allele frequency distribution of common functional polymorphisms of drug transporters

Table 3 shows the allele frequencies of drug transporter genes in Thai population. Among the important efflux

ABCB1 transporters, the MAF of *ABCB1* 1236C>T and 3435C>T exhibited prevalence rates of 34.5% and 43.7%, respectively. Other ABC SNPs with clinical relevance exhibited prevalence rates of rs212091 (15.7%), rs717620 (22.3%), rs2273697 (6.7%), rs3740066 (24.3%), and 30.7% for *ABCG2* 421C>A.

The uptake SLC transporters (OCT1, OCT2, OATP1A2, and OATP1B2) were present with at least one variant allele in this study group (data for all variants not shown). Among the OCT1 family, the two variants rs628031 (Val408Met, 1222G>A) and rs1867351 (Ser52Ser, 156T>C) had higher frequency of 33.9% and 31.9%, respectively. The pharmacologically important variants *SLCO1B1**1B, *SLCO1B1**5,

Table 2 Minor allele frequencies of common functional polymorphisms in major human Phase II drug-metabolizing enzyme genes in Thai children and adolescents with autism spectrum disorder (sample size = 119)

Gene	dbSNP RS ID	Common name	Minor allele frequency	Clinical relevance
UGT1A1	rs4148323	UGT1A1*6_c.211G>A(G71R)	0.067	<ul style="list-style-type: none"> UGT1A1*6 is associated with increased risk of neutropenia when treated with irinotecan³⁴ UGT1A1*93 is associated with increased risk of hematologic toxicity in irinotecan-treated patients³³
	rs4124874	UGT1A1*60_c.-3279T>G(Promoter)	0.361	
	rs10929303	UGT1A1*76_c.*211C>T(3'UTR)	0.088	
	rs8330	UGT1A1*79_c.*440C>G(3'UTR)	0.082	
	rs887829	UGT1A1*80_c.-364C>T	0.13	
	rs10929302	UGT1A1*93_c.-3156G>A(Promoter)	0.126	
UGT1A7	rs7586110	UGT1A7*12_c.-57T>G(5'UTR)	0.151	<ul style="list-style-type: none"> UGT1A7*12 is associated with increased risk of hyperbilirubinemia when treated with atazanavir³⁵
UGT2B7	rs7439366	UGT2B7*2_c.802C>T(H268Y)	0.315	<ul style="list-style-type: none"> Genotype TT shows better response to lorazepam and valproic acid as compared to genotype CC in UGT2B7*2 polymorphism⁶⁰ UGT2B7*3 is associated with decreased clearance of carvedilol⁶¹
	rs12233719	UGT2B7*3_c.211G>T(A71S)	0.084	
GSTA5	rs4715354	GSTA5_c.-31+2057C>T	0.357	<ul style="list-style-type: none"> rs4715354 and rs7746993, both combined, are associated with decreased busulfan clearance⁶²
	rs7746993	GSTA5_-8526G>T	0.147	
GSTP1	rs1695	GSTP1*B_c.313A>G(I105V)	0.275	<ul style="list-style-type: none"> GSTP1*B is associated with increased response and decreased severity of toxicity among patients with breast cancer treated with cyclophosphamide and epirubicin⁶³ Efficacy and/or toxicity of oxaliplatin, methotrexate, and fluorouracil are associated with GSTP1*B variant⁶⁴
GSTZ1	rs1046428	GSTZ1_c.245C>T(T82M)	0.055	<ul style="list-style-type: none"> rs7975 and rs1046428 are associated with clearance of dichloroacetic acid⁶⁵
	rs7975	GSTZ1_c.94G>A(E32K)	0.395	
NAT2	rs1799929	NAT2_c.481C>T(L161L)	0.078	<ul style="list-style-type: none"> rs1799929 variant is associated with hepatotoxicity when treated with antituberculosis drugs⁶⁶ NAT2*5, *6, *7 and *13 influence the metabolism of isoniazid⁶⁷
	rs1801280	NAT2*5_c.341T>C(I114T)	0.092	
	rs1799930	NAT2*6_c.590G>A(R197Q)	0.277	
	rs1799931	NAT2*7_c.857G>A(G286E)	0.176	
	rs1041983	NAT2*13_c.282C>T(Y94Y)	0.454	

Note: dbSNP RS ID is an identification tag assigned by the National Center for Biotechnology Information to a group of single nucleotide polymorphisms (SNPs) that map to an identical location in the SNP database.

and *SLCO1B1*17* had frequencies of 26.9%, 12.2%, and 8.4%, respectively.

Discussion

In this study population of 119 Thai children and adolescents with ASD, we have reported allele frequency of clinically relevant DMET SNPs. Among the 1,931 markers genotyped by the DMET assay, we detected ~26% of SNPs distributed along the pharmacokinetic genes. Although, 87% of SNPs passed the 95% call rate criteria, many of the SNPs were discarded because they were monomorphic and had a prevalence of <5%. Our study sample was comprised of mostly males (87%), and hence 46 SNPs on the X chromosome were excluded, as well. The SNPs reported in this study could serve as informative data in assessing the potential risk of therapeutic failure or adverse events in the Thai

population, where there is little information available about the prevalence of variations across DMET genes. The DMET microarray technique proved to be a powerful tool enabling us to characterize the clinically important genetic variants among DMET genes.

The frequency of some alleles in this study showed similar prevalence with previous studies conducted in the Thai population. The allele frequencies of *CYP2C19*3* (PM), *CYP2D6*2*, and *CYP2D6*10* (decreased enzyme activity) were 5%, 10%, and 41%, respectively, in this study. Similar findings were reported by the group studying *CYP2C19* and *CYP2D6* polymorphisms and tamoxifen efficacy in Thai patients with breast cancer.²⁸ *CYP2C19*2*, a PM, which has been shown to cause severe acute adverse drug reactions among Thai children treated with phenobarbital,²⁹ showed a higher frequency of 33.2% in

Table 3 Minor allele frequencies of common functional polymorphisms of drug transporter genes (ABC and SLC) in Thai children and adolescents with autism spectrum disorder (sample size =119)

Gene	dbSNP RS ID	Common name	Minor allele frequency	Clinical relevance
ABCB1 (MDR1)	rs1128503	ABCB1_c.1236C>T(G412G)	0.345	<ul style="list-style-type: none"> ABCB1 1236C>T is associated with overall increased survival period among oxaliplatin-treated patients with colorectal neoplasms⁶⁸ ABCB1 3435C>T is associated with increased serum concentration of digoxin⁶⁹ and nevirapine-induced hepatotoxicity⁷⁰ rs10276036 is associated with increased risk of death in patients with osteosarcoma after chemotherapy⁷¹
	rs1045642	ABCB1_c.3435C>T(11145I)	0.437	
	rs10276036	ABCB1_c.IVS9-44A>G	0.339	
ABCC1 (MRP1)	rs212091	ABCC1_c.*1512T>C	0.157	<ul style="list-style-type: none"> rs212091 is associated with virological failure in antiretroviral drugs therapy⁷²
ABCC2 (MRP2)	rs717620	ABCC2_c.-24C>T(5'UTR)	0.223	<ul style="list-style-type: none"> rs717620 and rs3740066 affect the response to antiepileptic drugs⁷³ rs717620 influences the metabolism of erythromycin⁷⁴ and is associated with toxicity among patients treated with fluorouracil, leucovorin, and oxaliplatin⁷⁵ rs2273697 influences the pharmacokinetics of talinolol and irinotecan^{76,77}
	rs2273697	ABCC2_c.1249G>A(V417I)	0.067	
	rs3740066	ABCC2_c.3972C>T(11324I)	0.243	
ABCG2 (BCRP)	rs2231142	ABCG2_c.421C>A(Q141K)	0.307	<ul style="list-style-type: none"> rs2231142 is associated with the plasma concentration of rosuvastatin^{78,79}
SLC22A1 (OCT1)	rs1867351	SLC22A1_c.156T>C(S52S)	0.319	<ul style="list-style-type: none"> c.480C>G polymorphism is associated with pharmacokinetics of imatinib⁸⁰ rs628031 is associated with gastrointestinal side effects when treated with metformin⁸¹
	rs683369	SLC22A1_c.480C>G(F160L)	0.148	
	rs2282143	SLC22A1_c.1022C>T(P341L)	0.13	
	rs628031	SLC22A1_c.1222G>A(V408M)	0.339	
SLCO1B1 (OATP1B1)	rs4149015	SLCO1B1*17_c.-11187G>A(Promoter)	0.084	<ul style="list-style-type: none"> SLCO1B1*17 is associated with the efficacy and pharmacokinetics of pravastatin^{82,83} SLCO1B1*5 is associated with high risk of muscular diseases when treated with simvastatin^{84,85} Other drugs associated with *5 variant include cerivastatin, pravastatin, and rosuvastatin^{78,86,87}
	rs2306283	SLCO1B1*1B_c.388A>G(N130D)	0.269	
	rs4149056	SLCO1B1*5_c.521T>C(V174A)	0.122	
	rs4149057	SLCO1B1_c.571T>C(L191L)	0.282	

Note: dbSNP RS ID is an identification tag assigned by the National Center for Biotechnology Information to a group of single nucleotide polymorphisms (SNPs) that map to an identical location in the SNP database.

Abbreviations: ABC, adenosine triphosphate-binding cassette; SLC, solute carrier.

this study, which is in line with the previously reported frequency of 27%.³⁰ *CYP2B6* variants, c.516G>T and c.785A>G, which were strongly correlated with plasma efavirenz levels in the Thai population, are not reported in the findings of this study.³¹ Another variant of a major *CYP2B6* enzyme, c.64C>T, which is also associated with efavirenz plasma concentration, was present at a frequency of 0.055, which is also reported by the earlier study to occur at similar prevalence of 0.06.

Polymorphisms in Phase II metabolic enzymes and transporters are of clinical significance, because of altered function affecting the drug safety and drug efficacy.³² UGT enzyme polymorphisms have been associated with several drug substrates and widely studied among several

ethnic groups.³³⁻³⁵ The prevalence of *UGT1A1**6, which is more commonly observed in Asian populations and absent among Caucasians, resembled earlier studies that reported a frequency of 9%.³⁶ A previous study of *UGT2B7**2 among Thai patients with breast cancer reported an allele frequency of 28%,³⁷ which is similar to the 31.5% reported in this study. *CYP1A2**1F was associated with an increased risk of cholangiocarcinoma (CCA), while *NAT2**6 and *7 were associated with a lower risk of CCA in a study by Prawan et al,³⁸ who evaluated the relationship between *CYP1A2* and *NAT2* alleles and CCA among individuals living in the northeastern region of Thailand. The allelic frequencies of *CYP1A2**1F, *NAT2**6, and *NAT2**7 in this study group were found to be 23.9%, 27.7%, and 17.6%, respectively. Genetic variants of

a widely documented polymorphic enzyme TPMT did not pass quality control in this study group with the exception of one variant, *TPMT_c.474C>T*, with a prevalence of 22.7% (data not shown).

Among the genetic polymorphisms of *ABC* and *SLC* transporters, we found a modest replication of allele frequency, as was reported in a previous comprehensive study of the frequency of several functional variants known to affect plasma concentration of anti-human immunodeficiency virus drugs among the Thai population, including *ABCB1* (1236C>T, 2677G>T, and 3435C>T), *ABCG2* 421C>A, and *SLCO1B1* 521T>C.³⁹

Limitations in this study include lack of validation data for the markers extracted from the DMET assay. Second, we did not evaluate the clinical role of DMET SNPs in determining the influence on the ASD symptoms. A study by Jiao et al⁴⁰ has accumulated the evidence of the role of SNPs in the classification of symptom severities among ASD children. It is necessary to analyze the effects of genetic polymorphisms in DMET genes on clinical efficacy and adverse events to amplify the strength of this study. Finally, the small size and inability to compare with the diverse ethnic Thai populations may limit the scope of application of our findings. Further inclusion of control population would have provided more evidence for translating the findings from this study into clinical practice, which was lacking in this study.

Conclusion

In this study, we have analyzed the DMET SNPs that are increasingly significant in identifying the predicted drug metabolizer phenotypes among individuals. Knowledge of these variations and implementation of this information to correlate these variants with drug response in the Thai population will help us to understand the interindividual variations in response to pharmacological agents. Identification of several functional SNPs in the clinical setting, rather than just the few widely studied SNPs, will provide more concise and predictive value for pharmacotherapy.

Acknowledgments

This study was supported by grants from the following organizations: 1) Pharmacogenomics for Autistic Child Project, Khoon Poom Foundation, the Project of Her Royal Highness Princess Ubolratana Rajakanya Siriwatana Bhanawadee; 2) Office of the National Research Council of Thailand; 3) Faculty of Medicine Ramathibodi Hospital; and 4) Mahidol University. The authors thank all staff

in Yuwaprasart Waithayopatum Child and Adolescent Psychiatric Hospital and all the children and adolescents with ASD who participated in this study.

Disclosure

The authors report no conflicts of interest in this work.

References

- Li J, You Y, Yue W, et al. Genetic evidence for possible involvement of the calcium channel gene CACNA1A in autism pathogenesis in Chinese Han population. *PLoS One*. 2015;10(11):e0142887.
- Rosemeyer R. A review of the serotonin transporter and prenatal cortisol in the development of autism spectrum disorders. *Mol Autism*. 2013;4(1):37.
- Mezzelani A, Raggi ME, Marabotti A, Milanese L. Ochratoxin A as possible factor triggering autism and its male prevalence via epigenetic mechanism. *Nutr Neurosci*. 2016;19(1):43–46.
- Ghanizadeh A, Tordjman S, Jaafari N. Aripiprazole for treating irritability in children & adolescents with autism: a systematic review. *Indian J Med Res*. 2015;142(3):269–275.
- McQuire C, Hassiotis A, Harrison B, Pilling S. Pharmacological interventions for challenging behaviour in children with intellectual disabilities: a systematic review and meta-analysis. *BMC Psychiatry*. 2015;15(1):303.
- Hongkaew Y, Ngamsamut N, Puangpetch A, et al. Hyperprolactinemia in Thai children and adolescents with autism spectrum disorder treated with risperidone. *Neuropsychiatr Dis Treat*. 2015;11:191–196.
- Benvenuto A, Battan B, Porfirio MC, Curatolo P. Pharmacotherapy of autism spectrum disorders. *Brain Dev*. 2013;35(2):119–127.
- Kapedanovska Nestorovska A, Jakovski K, Naumovska Z, et al. Distribution of the most common genetic variants associated with a variable drug response in the population of the Republic of Macedonia. *Balkan J Med Genet*. 2014;17(2):5–14.
- Ma Q, Lu AY. Pharmacogenetics, pharmacogenomics, and individualized medicine. *Pharmacol Rev*. 2011;63(2):437–459.
- Gupta PD. Pharmacogenetics, pharmacogenomics and ayurgenomics for personalized medicine: a paradigm shift. *Indian J Pharm Sci*. 2015;77(2):135–141.
- Mango R, Vecchione L, Raso B, et al. Pharmacogenomics in cardiovascular disease: the role of single nucleotide polymorphisms in improving drug therapy. *Expert Opin Pharmacother*. 2005;6(15):2565–2576.
- Gray IC, Campbell DA, Spurr NK. Single nucleotide polymorphisms as tools in human genetics. *Hum Mol Genet*. 2000;9(16):2403–2408.
- Chaudhary R, Singh B, Kumar M, et al. Role of single nucleotide polymorphisms in pharmacogenomics and their association with human diseases. *Drug Metab Rev*. 2015;47(3):281–290.
- Holmes A, Murphy DL, Crawley JN. Abnormal behavioral phenotypes of serotonin transporter knockout mice: parallels with human anxiety and depression. *Biol Psychiatry*. 2003;54(10):953–959.
- Braam W, Keijzer H, Struik Boudier H, Didden R, Smits M, Curfs L. CYP1A2 polymorphisms in slow melatonin metabolisers: a possible relationship with autism spectrum disorder? *J Intellect Disabil Res*. 2013;57(11):993–1000.
- Karam RA, Rezk NA, Abdelrahman HM, et al. Catechol-O-methyltransferase Val158Met polymorphism and hyperactivity symptoms in Egyptian children with autism spectrum disorder. *Res Dev Disabil*. 2013;34(7):2092–2097.
- Umamaheswaran G, Kumar DK, Adithan C. Distribution of genetic polymorphisms of genes encoding drug metabolizing enzymes & drug transporters – a review with Indian perspective. *Indian J Med Res*. 2014;139(1):27–65.
- Li J, Bluth MH. Pharmacogenomics of drug metabolizing enzymes and transporters: implications for cancer therapy. *Pharmacogenomics Pers Med*. 2011;4:11–33.

19. Sim SC, Kacevska M, Ingelman-Sundberg M. Pharmacogenomics of drug-metabolizing enzymes: a recent update on clinical implications and endogenous effects. *Pharmacogenomics J*. 2013;13(1):1–11.
20. Bonifaz-Pena V, Contreras AV, Struchiner CJ, et al. Exploring the distribution of genetic markers of pharmacogenomics relevance in Brazilian and Mexican populations. *PLoS One*. 2014;9(11):e112640.
21. Yang L, Price ET, Chang CW, et al. Gene expression variability in human hepatic drug metabolizing enzymes and transporters. *PLoS One*. 2013;8(4):e60368.
22. Jittikoon J, Mahasirimongkol S, Charoenyingwattana A, et al. Comparison of genetic variation in drug ADME-related genes in Thais with Caucasian, African and Asian HapMap populations. *J Hum Genet*. Epub 2015 Oct 1.
23. Saldivar JS, Taylor D, Sugarman EA, et al. Initial assessment of the benefits of implementing pharmacogenetics into the medical management of patients in a long-term care facility. *Pharmacogenomics Pers Med*. 2016;9:1–6.
24. Fernandez CA, Smith C, Yang W, et al. Concordance of DMET plus genotyping results with those of orthogonal genotyping methods. *Clin Pharmacol Ther*. 2012;92(3):360–365.
25. Hardenbol P, Baner J, Jain M, et al. Multiplexed genotyping with sequence-tagged molecular inversion probes. *Nat Biotechnol*. 2003;21(6):673–678.
26. Syvanen AC. Toward genome-wide SNP genotyping. *Nat Genet*. 2005;37(suppl):S5–S10.
27. Barrett JC, Fry B, Maller J, Daly MJ. Haploview: analysis and visualization of LD and haplotype maps. *Bioinformatics*. 2005;21(2):263–265.
28. Chamnanphon M, Pechatanan K, Sirachainan E, et al. Association of CYP2D6 and CYP2C19 polymorphisms and disease-free survival of Thai post-menopausal breast cancer patients who received adjuvant tamoxifen. *Pharmacogenomics Pers Med*. 2013;6:37–48.
29. Manuyakorn W, Siripool K, Kamchaisatien W, et al. Phenobarbital-induced severe cutaneous adverse drug reactions are associated with CYP2C19*2 in Thai children. *Pediatr Allergy Immunol*. 2013;24(3):299–303.
30. Tassaneeyakul W, Tawalee A, Kukongviriyapan V, Blaisdell J, Goldstein JA, Gaysornsir D. Analysis of the CYP2C19 polymorphism in a North-eastern Thai population. *Pharmacogenetics*. 2002;12(3):221–225.
31. Sukasem C, Chamnanphon M, Koomdee N, et al. High plasma efavirenz concentration and CYP2B6 polymorphisms in Thai HIV-1 infections. *Drug Metab Pharmacokinet*. 2013;28(5):391–397.
32. Yiannakopoulou E. Pharmacogenomics of phase II metabolizing enzymes and drug transporters: clinical implications. *Pharmacogenomics J*. 2013;13(2):105–109.
33. Cote JF, Kirzin S, Kramar A, et al. UGT1A1 polymorphism can predict hematologic toxicity in patients treated with irinotecan. *Clin Cancer Res*. 2007;13(11):3269–3275.
34. Han FF, Guo CL, Yu D, et al. Associations between UGT1A1*6 or UGT1A1*6/*28 polymorphisms and irinotecan-induced neutropenia in Asian cancer patients. *Cancer Chemother Pharmacol*. 2014;73(4):779–788.
35. Lankisch TO, Moebius U, Wehmeier M, et al. Gilbert's disease and atazanavir: from phenotype to UDP-glucuronosyltransferase haplotype. *Hepatology*. 2006;44(5):1324–1332.
36. Sukasem C, Atasilp C, Chansriwong P, Chamnanphon M, Puangpetch A, Sirachainan E. Development of pyrosequencing method for detection of UGT1A1 polymorphisms in Thai colorectal cancers. *J Clin Lab Anal*. 2016;30(1):84–89.
37. Areepium N, Panomvana D, Rungwanonchai P, Sathaporn S, Voravud N. Effects of CYP2D6 and UGT2B7 polymorphisms on pharmacokinetics of tamoxifen in Thai breast cancer patients. *Breast Cancer (Dove Med Press)*. 2013;5:73–78.
38. Prawan A, Kukongviriyapan V, Tassaneeyakul W, Pairojkul C, Bhudhisawasdi V. Association between genetic polymorphisms of CYP1A2, arylamine N-acetyltransferase 1 and 2 and susceptibility to cholangiocarcinoma. *Eur J Cancer Prev*. 2005;14(3):245–250.
39. Chaikan A, Chierakul N, Saguenwong N, et al. Frequency of functional drug disposition gene polymorphism in Thai population: relevance to antiretroviral drugs. In: Ninth International Congress on Drug Therapy in HIV Infection; 2008; Glasgow.
40. Jiao Y, Chen R, Ke X, et al. Single nucleotide polymorphisms predict symptom severity of autism spectrum disorder. *J Autism Dev Disord*. 2012;42(6):971–983.
41. Dong N, Yu J, Wang C, et al. Pharmacogenetic assessment of clinical outcome in patients with metastatic breast cancer treated with docetaxel plus capecitabine. *J Cancer Res Clin Oncol*. 2012;138(7):1197–1203.
42. Yoon Y, Park HD, Park KU, Kim JQ, Chang YS, Song J. Associations between CYP2E1 promoter polymorphisms and plasma 1,3-dimethyluric acid/theophylline ratios. *Eur J Clin Pharmacol*. 2006;62(8):627–631.
43. Tiwari AK, Deshpande SN, Rao AR, et al. Genetic susceptibility to tardive dyskinesia in chronic schizophrenia subjects: I. Association of CYP1A2 gene polymorphism. *Pharmacogenomics J*. 2005;5(1):60–69.
44. Laika B, Leucht S, Heres S, Schneider H, Steimer W. Pharmacogenetics and olanzapine treatment: CYP1A2*1F and serotonergic polymorphisms influence therapeutic outcome. *Pharmacogenomics J*. 2010;10(1):20–29.
45. Haas DW, Kwara A, Richardson DM, et al. Secondary metabolism pathway polymorphisms and plasma efavirenz concentrations in HIV-infected adults with CYP2B6 slow metabolizer genotypes. *J Antimicrob Chemother*. 2014;69(8):2175–2182.
46. Manosuthi W, Sukasem C, Lueangniyomkul A, et al. Impact of pharmacogenetic markers of CYP2B6, clinical factors, and drug-drug interaction on efavirenz concentrations in HIV/tuberculosis-coinfected patients. *Antimicrob Agents Chemother*. 2013;57(2):1019–1024.
47. Mukonzo JK, Nanzigu S, Waako P, Ogwal-Okeng J, Gustafson LL, Aklilu E. CYP2B6 genotype, but not rifampicin-based anti-TB cotreatments, explains variability in long-term efavirenz plasma exposure. *Pharmacogenomics*. 2014;15(11):1423–1435.
48. van der Weide J, van Baalen-Benedek EH, Kootstra-Ros JE. Metabolic ratios of psychotropics as indication of cytochrome P450 2D6/2C19 genotype. *Ther Drug Monit*. 2005;27(4):478–483.
49. Cui H, Lin S, Chen X, et al. Correlation between SNPs in candidate genes and VerifyNow-detected platelet responsiveness to aspirin and clopidogrel treatment. *Cardiovasc Drugs Ther*. 2015;29(2):137–146.
50. Horenstein RB, Madabushi R, Zineh I, et al. Effectiveness of clopidogrel dose escalation to normalize active metabolite exposure and antiplatelet effects in CYP2C19 poor metabolizers. *J Clin Pharmacol*. 2014;54(8):865–873.
51. Rudberg I, Hermann M, Refsum H, Molden E. Serum concentrations of sertraline and N-desmethyl sertraline in relation to CYP2C19 genotype in psychiatric patients. *Eur J Clin Pharmacol*. 2008;64(12):1181–1188.
52. Chang M, Tybring G, Dahl ML, Lindh JD. Impact of cytochrome P450 2C19 polymorphisms on citalopram/escitalopram exposure: a systematic review and meta-analysis. *Clin Pharmacokinet*. 2014;53(9):801–811.
53. Penas-Lledo EM, Trejo HD, Dorado P, et al. CYP2D6 ultrarapid metabolism and early dropout from fluoxetine or amitriptyline monotherapy treatment in major depressive patients. *Mol Psychiatry*. 2013;18(1):8–9.
54. Wu X, Yuan L, Zuo J, Lv J, Guo T. The impact of CYP2D6 polymorphisms on the pharmacokinetics of codeine and its metabolites in Mongolian Chinese subjects. *Eur J Clin Pharmacol*. 2014;70(1):57–63.
55. Hodgson K, Tansey K, Dernovsek MZ, et al. Genetic differences in cytochrome P450 enzymes and antidepressant treatment response. *J Psychopharmacol*. 2014;28(2):133–141.
56. Jurica J, Zourkova A. Dynamics and persistence of CYP2D6 inhibition by paroxetine. *J Clin Pharm Ther*. 2013;38(4):294–300.
57. Kirchheiner J, Keulen JT, Bauer S, Roots I, Brockmoller J. Effects of the CYP2D6 gene duplication on the pharmacokinetics and pharmacodynamics of tramadol. *J Clin Psychopharmacol*. 2008;28(1):78–83.

58. Nair SS, Sarasamma S, Gracious N, George J, Anish TS, Radhakrishnan R. Polymorphism of the CYP3A5 gene and its effect on tacrolimus blood level. *Exp Clin Transplant*. 2015;13(suppl 1):197–200.
59. Birdwell KA, Grady B, Choi L, et al. The use of a DNA biobank linked to electronic medical records to characterize pharmacogenomic predictors of tacrolimus dose requirement in kidney transplant recipients. *Pharmacogenet Genomics*. 2012;22(1):32–42.
60. Chung JY, Cho JY, Yu KS, et al. Pharmacokinetic and pharmacodynamic interaction of lorazepam and valproic acid in relation to UGT2B7 genetic polymorphism in healthy subjects. *Clin Pharmacol Ther*. 2008; 83(4):595–600.
61. Takekuma Y, Takenaka T, Kiyokawa M, et al. Evaluation of effects of polymorphism for metabolic enzymes on pharmacokinetics of carvedilol by population pharmacokinetic analysis. *Biol Pharm Bull*. 2007; 30(3):537–542.
62. Ten Brink MH, Swen JJ, Bohringer S, et al. Exploratory analysis of 1936 SNPs in ADME genes for association with busulfan clearance in adult hematopoietic stem cell recipients. *Pharmacogenet Genomics*. 2013;23(12):675–683.
63. Zhang BL, Sun T, Zhang BN, et al. Polymorphisms of GSTP1 is associated with differences of chemotherapy response and toxicity in breast cancer. *Chin Med J (Engl)*. 2011;124(2):199–204.
64. Lamas MJ, Duran G, Balboa E, et al. Use of a comprehensive panel of biomarkers to predict response to a fluorouracil-oxaliplatin regimen in patients with metastatic colorectal cancer. *Pharmacogenomics*. 2011; 12(3):433–442.
65. Shroads AL, Langae T, Coats BS, et al. Human polymorphisms in the glutathione transferase zeta 1/maleylacetoacetate isomerase gene influence the toxicokinetics of dichloroacetate. *J Clin Pharmacol*. 2012; 52(6):837–849.
66. Gupta VH, Amarapurkar DN, Singh M, et al. Association of N-acetyltransferase 2 and cytochrome P450 2E1 gene polymorphisms with antituberculosis drug-induced hepatotoxicity in Western India. *J Gastroenterol Hepatol*. 2013;28(8):1368–1374.
67. Verhagen LM, Coenen MJ, Lopez D, et al. Full-gene sequencing analysis of NAT2 and its relationship with isoniazid pharmacokinetics in Venezuelan children with tuberculosis. *Pharmacogenomics*. 2014; 15(3):285–296.
68. Wu H, Kang H, Liu Y, et al. Association of ABCB1 genetic polymorphisms with susceptibility to colorectal cancer and therapeutic prognosis. *Pharmacogenomics*. 2013;14(8):897–911.
69. Aarnoudse AJ, Dieleman JP, Visser LE, et al. Common ATP-binding cassette B1 variants are associated with increased digoxin serum concentration. *Pharmacogenet Genomics*. 2008;18(4):299–305.
70. Ciccacci C, Borgiani P, Ceffa S, et al. Nevirapine-induced hepatotoxicity and pharmacogenetics: a retrospective study in a population from Mozambique. *Pharmacogenomics*. 2010;11(1):23–31.
71. Caronia D, Patino-Garcia A, Perez-Martinez A, et al. Effect of ABCB1 and ABCC3 polymorphisms on osteosarcoma survival after chemotherapy: a pharmacogenetic study. *PLoS One*. 2011;6(10):e26091.
72. Coelho AV, Silva SP, de Alencar LC, et al. ABCB1 and ABCC1 variants associated with virological failure of first-line protease inhibitors antiretroviral regimens in Northeast Brazil patients. *J Clin Pharmacol*. 2013;53(12):1286–1293.
73. Qu J, Zhou BT, Yin JY, et al. ABCC2 polymorphisms and haplotype are associated with drug resistance in Chinese epileptic patients. *CNS Neurosci Ther*. 2012;18(8):647–651.
74. Franke RM, Lancaster CS, Peer CJ, et al. Effect of ABCC2 (MRP2) transport function on erythromycin metabolism. *Clin Pharmacol Ther*. 2011;89(5):693–701.
75. Lee KH, Chang HJ, Han SW, et al. Pharmacogenetic analysis of adjuvant FOLFOX for Korean patients with colon cancer. *Cancer Chemother Pharmacol*. 2013;71(4):843–851.
76. Haenisch S, May K, Wegner D, Caliebe A, Cascorbi I, Siegmund W. Influence of genetic polymorphisms on intestinal expression and rifampicin-type induction of ABCC2 and on bioavailability of talinolol. *Pharmacogenet Genomics*. 2008;18(4):357–365.
77. Fujita K, Nagashima F, Yamamoto W, et al. Association of ATP-binding cassette, sub-family C, number 2 (ABCC2) genotype with pharmacokinetics of irinotecan in Japanese patients with metastatic colorectal cancer treated with irinotecan plus infusional 5-fluorouracil/leucovorin (FOLFIRI). *Biol Pharm Bull*. 2008;31(11):2137–2142.
78. DeGorter MK, Tirona RG, Schwarz UI, et al. Clinical and pharmacogenetic predictors of circulating atorvastatin and rosuvastatin concentrations in routine clinical care. *Circ Cardiovasc Genet*. 2013; 6(4):400–408.
79. Lee HK, Hu M, Lui S, Ho CS, Wong CK, Tomlinson B. Effects of polymorphisms in ABCG2, SLCO1B1, SLC10A1 and CYP2C9/19 on plasma concentrations of rosuvastatin and lipid response in Chinese patients. *Pharmacogenomics*. 2013;14(11):1283–1294.
80. Di Paolo A, Polillo M, Capecchi M, et al. The c.480C>G polymorphism of hOCT1 influences imatinib clearance in patients affected by chronic myeloid leukemia. *Pharmacogenomics J*. 2014;14(4):328–335.
81. Tarasova L, Kalnina I, Geldner K, et al. Association of genetic variation in the organic cation transporters OCT1, OCT2 and multidrug and toxin extrusion 1 transporter protein genes with the gastrointestinal side effects and lower BMI in metformin-treated type 2 diabetes patients. *Pharmacogenet Genomics*. 2012;22(9):659–666.
82. Niemi M, Neuvonen PJ, Hofmann U, et al. Acute effects of pravastatin on cholesterol synthesis are associated with SLCO1B1 (encoding OATP1B1) haplotype *17. *Pharmacogenet Genomics*. 2005; 15(5):303–309.
83. Niemi M, Schaeffeler E, Lang T, et al. High plasma pravastatin concentrations are associated with single nucleotide polymorphisms and haplotypes of organic anion transporting polypeptide-C (OATP-C, SLCO1B1). *Pharmacogenetics*. 2004;14(7):429–440.
84. Brunham LR, Lansberg PJ, Zhang L, et al. Differential effect of the rs4149056 variant in SLCO1B1 on myopathy associated with simvastatin and atorvastatin. *Pharmacogenomics J*. 2012;12(3):233–237.
85. SEARCH Collaborative Group, Link E, Parish S, et al. SLCO1B1 variants and statin-induced myopathy – a genomewide study. *N Engl J Med*. 2008;359(8):789–799.
86. Marcianti KD, Durda JP, Heckbert SR, et al. Cerivastatin, genetic variants, and the risk of rhabdomyolysis. *Pharmacogenet Genomics*. 2011; 21(5):280–288.
87. Ho RH, Choi L, Lee W, et al. Effect of drug transporter genotypes on pravastatin disposition in European- and African-American participants. *Pharmacogenet Genomics*. 2007;17(8):647–656.

Neuropsychiatric Disease and Treatment

Publish your work in this journal

Neuropsychiatric Disease and Treatment is an international, peer-reviewed journal of clinical therapeutics and pharmacology focusing on concise rapid reporting of clinical or pre-clinical studies on a range of neuropsychiatric and neurological disorders. This journal is indexed on PubMed Central, the 'PsycINFO' database and CAS,

Submit your manuscript here: <http://www.dovepress.com/neuropsychiatric-disease-and-treatment-journal>

Dovepress

and is the official journal of The International Neuropsychiatric Association (INA). The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.