Fungistatic activity of all-*trans* retinoic acid against *Aspergillus fumigatus* and *Candida albicans*

Elena Campione¹

Roberta Gaziano²

Daniele Marino²

Augusto Orlandi³

¹Department of Dermatology,

²Department of Microbiology,

³Department of Anatomic Pathology,
University of Rome Tor Vergata, Rome,
Italy

→ Video abstract

Point your Smartphone at the code above. If you have a
QR code reader the video abstract will appear. Or use:

<http://youtu.be/njCcs00YtFA>

Purpose: Fungal infections are a major complication in hematologic and neoplastic patients causing severe morbidity and mortality. *Aspergillus fumigatus* and *Candida albicans* are among the most invasive opportunistic pathogens in immunocompromised patients, and classic antifungal drugs are frequently unsuccessful in these patients. Recent reports hypothesize that the antifungal efficacy of all-*trans* retinoic acid (ATRA) is mainly related to its strong capacity to stimulate monocyte-mediated immunity, but no consideration was given to its potential direct fungistatic activity. Moreover, ATRA offers the opportunity for systemic therapy.

Methods and results: We investigated the efficacy of ATRA at different concentrations for its antifungal activity against opportunistic *A. fumigatus* and *C. albicans* obtained from clinical samples according to standard protocols. A fungistatic activity of ATRA on *A. fumigatus* and *C. albicans* at 0.5–1 mM concentration was documented up to 7 days.

Conclusion: This is the first evidence of a direct and strong fungistatic activity of ATRA against *A. fumigatus* and *C. albicans*. The potential adjuvant therapeutic application of ATRA might be useful in the treatment and/or prevention of systemic mycoses in immunocompromised patients. The discovery of a direct fungistatic activity, in association with its reported immunomodulatory properties, makes ATRA an excellent candidate for new combined antifungal strategies for systemic mycoses in immunocompromised and cancer patients.

Keywords: all-*trans* retinoic acid, fungistatic activity, fungal infections

Introduction

Fungal infections are a major complication in hematologic and neoplastic patients undergoing prolonged chemotherapy and often cause severe morbidity and mortality. In humans, *Aspergillus fumigatus* and *Candida albicans* are important fungal pathogens frequently causing severe infections with invasive growth in immunocompromised patients.¹ Unfortunately, the use of classic antifungal drugs is unsuccessful in many of these patients. Retinoic acid – the biologically active metabolite of vitamin A – controls the normal immune system development as a modulator of both innate and adaptive immune responses.² Vitamin A is a nutrient obtained through the diet either as provitamin-A (carotenoids) or as preformed vitamin A (retinol and retinyl esters). After its transport through the cytoplasmic binding with specific receptors, liver dehydrogenases convert retinol into retinoic acid, its biologically active metabolite.^{3–7} Over the last decades, an increasing effort has been devoted to better define the involvement in the regulation of immune response, since vitamin A deficiency has been associated with an increased susceptibility to severe infectious diseases.^{8,9} The adding of all-*trans* retinoic acid (ATRA) to the therapy of acute promyelocytic leukemia resulted in a lower incidence of total episodes of fungemias in the patients.¹⁰ Recent studies support the hypothesis that efficacy of vitamin A or ATRA alone or in combination with other

Correspondence: Elena Campione
Department of Dermatology, University
of Rome Tor Vergata, Viale Oxford 81,
00133 Rome, Italy
Email campioneelena@hotmail.com

drugs is related to its capacity to stimulate both innate and the adaptive immunity, in particular the monocyte-mediated immune response.^{4,10–12} We recently reported that tazarotene, a retinoic acid derivative, inhibits *Tricophyton* and *C. albicans* growth in vitro and a topical treatment with 0.1% tazarotene gel is effective in lateral onychomycosis.¹³ Since the contribution of inflammatory response is quite limited in the affected nail, we hypothesized that the beneficial effects of retinoids derive from a direct fungistatic activity. In comparison with tazarotene, ATRA offers the advantage of systemic therapy and is widely used in many patients safely. We aimed to investigate the efficacy of ATRA against opportunistic fungal pathogens such as *A. fumigatus* and *C. albicans*.

Methods

Fungal culture conditions

A. fumigatus and *C. albicans* strains used in this study were obtained from a clinical sample. The fungal strains were grown on Sabouraud dextrose agar (Difco Laboratories, Detroit, MI, USA) supplemented with chloramphenicol. Resting conidia of *Aspergillus* and *Candida* yeasts were harvested by washing the slant cultures with sterile saline. Swollen conidia were obtained by shaking at 300 rpm at 37°C for 8 hours in Roswell Park Memorial Institute medium (RPMI) 1640 with 10% of fetal calf serum (FCS), until the spores swelled to almost twice their resting diameter. For generation of hyphae, swollen conidia were allowed to germinate (98% germination) by further incubation in RPMI 1640 with 10% FCS (~6 hours). For the generation of hyphae, *C. albicans* yeasts were allowed to germinate (98% germination) by incubation at 37°C for 4 hours in RPMI 1640, pH 6.8–7.2, with 10% FCS. To assess the effect of ATRA on the germination of *A. fumigatus* or *C. albicans*, resting conidia or yeast cells were plated on a 24-well microplate at a density of 1×10^5 cells in 1 mL of RPMI 1640 with 10% FCS and supplemented with penicillin (100 U/mL) and streptomycin (100 µg/mL). ATRA (Sigma-Aldrich, St Louis, MO, USA) was added to the fungal suspension culture at time 0 at different concentrations (1, 0.5, 0.25, 0.12, and 0.06 mM). Fungal cultures were maintained at 37°C under continuous shaking at 300 rpm and monitored every day until 1 week following treatment. The vitality of *C. albicans* yeasts and *A. fumigatus* conidia was evaluated by light microscopy using the trypan blue dye exclusion method. To confirm vitality after exposure to higher doses of ATRA for 7 days, conidia and yeast cells were harvested, washed, and again cultured with fresh medium containing 10% FCS, in the

absence of ATRA. Under this condition, conidia and yeasts were able to germinate into hyphae within 24 hours.

Evaluation of fungistatic activity

The development of *A. fumigatus* and *C. albicans* was followed using an optical microscope (Carl Zeiss Meditec AG, Jena, Germany) with a 40× magnification objective lens. Microscopic images were recorded after 8 and 4 hours of incubation with ATRA at 37°C for *A. fumigatus* conidial swelling and *C. albicans* yeast cell germination, respectively.

Results

Light optical microscopy revealed that, at the earliest time point (8 hours) in the presence of ATRA (1 mM), *A. fumigatus* conidia did not swell and remained similar in shape and size to the resting ones (Figure 1B). On the contrary, as shown in Figure 1A, most of the control untreated conidia (~80%) were swollen and some developed short germ tube similar to those observed in the ATRA-treated cultures of non-synchronized population of *A. fumigatus* at doses <1 mM (Figure 1C). After more prolonged incubation (24 hours), all control and ATRA-treated conidia at doses <1 mM were swollen and produced a germ tube (data not shown). ATRA at 1 and 0.5 mM also inhibited germination and hyphal outgrowth of *C. albicans* (Figure 1E and F). Instead, control untreated yeast cells developed germ tube and subsequently hyphal growth (Figure 1D). Doses of ATRA <0.5 mM did not affect the ability of *C. albicans* to germinate (Figure 1G). The inhibitory effect of ATRA on the germination of *A. fumigatus* and *C. albicans* was maintained and examined at a later time point (day 7). These data suggested that the fungistatic activity of ATRA at 0.5–1 mM concentration was obtained without any adverse effects on *A. fumigatus* conidia and *C. albicans* yeast, since their viability was >90% as assessed by trypan blue. Similar results were observed at other time points (data not shown). To confirm the vitality of *A. fumigatus* and *C. albicans* after exposure to higher (0.5–1 mM) doses, conidia and yeast cells were harvested, washed after 7 days, and again cultured in the absence of ATRA. *A. fumigatus* conidia and *C. albicans* yeasts were able to germinate into hyphae within 24 hours (data not shown), again supporting the fungistatic activity of ATRA in vitro.

Discussion

Our results documented the fungistatic activity of ATRA at 0.5–1 mM concentration on *A. fumigatus* and *C. albicans*.

Figure 1 Inhibitory effect of ATRA on *Aspergillus* and *Candida* germination.

Notes: Resting *Aspergillus fumigatus* conidia or *Candida albicans* yeast cells were cultured in RPMI 1640 with 10% FCS at 37°C in the presence or absence of ATRA. (A) Conidia untreated, (B) conidia treated with ATRA 1 mM, (C) conidia treated with ATRA 0.5 mM, (D) *Candida* yeasts untreated, (E) *Candida* yeasts treated with ATRA 1 mM, (F) *Candida* yeasts treated with ATRA 0.5 mM, and (G) *Candida* yeasts treated with ATRA 0.25 mM. The germination of both fungi was followed using an optical microscope. Microscopic images were recorded after 8 and 4 hours of incubation for *A. fumigatus* conidia and *C. albicans* yeasts, respectively. Magnification: $\times 40$.

Abbreviations: FCS, fetal calf serum; ATRA, all-*trans* retinoic acid.

ATRA offers the opportunity for systemic therapy against opportunistic fungal pathogens such as *A. fumigatus* and *C. albicans*. Despite an increasing interest in the immunomodulatory role of ATRA, its specific role in the immune response to fungal infections was never explored. Our study is the first evidence of a direct and high fungistatic activity of ATRA against *A. fumigatus* and *C. albicans*. Till now, several in vitro studies highlighted the function of vitamin A as an important factor for immune system development and a modulator of the innate and the adaptive immune response.^{6–9} As a matter of fact, vitamin A regulates the development of B lymphocytes and immunoglobulin production.¹⁴ Retinoic acid also shows specific immunomodulatory activities. In T lymphocytes, retinoic acid attenuated the Th1-associated gene expression and skewed the immune response toward a Th2 profile.¹⁵ Retinoic acid also modulated the production of LPS-induced cytokines and/or chemokines in monocytes, macrophages, and dendritic cells.^{16,17} Moreover, ATRA induced proliferative arrest in nonmyeloid mesenchymal cells.^{18,19} ATRA has been shown to stimulate the differentiation of myeloid-derived suppressor cells to dendritic cells and macrophages. At therapeutic concentrations, ATRA could substantially drop the number of myeloid-derived suppressor cells in tumor-bearing mice and in patients with cancer, thus improving their antigen-specific response.^{20,21} Lei et al¹¹ hypothesized that ATRA treatment, alone and in combination with an 8-aminoquinoline primaquine, can push lung myeloid-derived suppressor cells to differentiate in functional macrophages helping to clear pneumonia from the *Pneumocystis carinii* infection. Klassert et al¹² recently reported the effects of ATRA on the immune response of human monocytes during *C. albicans* infection. The authors documented an immunomodulatory effect of ATRA leading to a highly significant suppression of the fungi-induced expression of TNF α , IL6, and IL12 at both the transcriptional and posttranslational levels.¹²

The effectiveness of ATRA and its possible toxicity can be easily monitored in the blood. Our results in vitro are already widely proven in the clinical practice since patients affected by promyelocytic leukemia were treated with ATRA, they were less subjected to systemic infections, with a more preserved immunological profile.¹⁰ The adjuvant application of ATRA might also be useful in the treatment and/or prevention of systemic mycoses in immunocompromised patients. Therefore, the discovery of a direct fungistatic activity associated with its reported immunomodulatory properties makes ATRA an excellent candidate for new and efficient antifungal strategies. ATRA may be administered alone or in

combination with lower doses of conventional antimycotic agents, thus reducing their side effects. Although the therapeutic systemic use of ATRA as an adjuvant for antifungal therapies requires further validation in preclinical models, the considerable health burden represented by invasive mycosis and the emerging multidrug resistance justifies every effort directed to the development of new therapeutic solutions.

Acknowledgment

The authors wish to thank Denis Mariano for editing assistance. In memory of Professor Chimenti who recently passed away.

Disclosure

The authors report no conflicts of interest in this work.

References

1. Pfaller MA, Diekema DJ. Epidemiology of invasive candidiasis: a persistent public health problem. *Clin Microbiol Rev*. 2007;20(1):133–163.
2. Majer O, Bourgeois C, Zwolanek F, et al. Type I interferons promote fatal immunopathology by regulating inflammatory monocytes and neutrophils during *Candida* infections. *PLoS Pathog*. 2012;8(7):e1002811.
3. D'Ambrosio DN, Clugston RD, Blaner WS. Vitamin A metabolism: an update. *Nutrients*. 2011;3(1):63–103.
4. Wolf G. A history of vitamin A and retinoids. *FASEB J*. 1996;10(9):1102–1107.
5. Doldo E, Costanza G, Agostinelli S, et al. Vitamin A, cancer treatment and prevention: the new role of cellular retinol binding proteins. *Biomed Res Int*. 2015;2015:624627.
6. Hall JA, Grainger JR, Spencer SP, Belkaid Y. The role of retinoic acid in tolerance and immunity. *Immunity*. 2011;35(1):13–22.
7. Theodosiou M, Laudet V, Schubert M. From carrot to clinic: an overview of the retinoic acid signaling pathway. *Cell Mol Life Sci*. 2010;67(9):1423–1445.
8. Clagett-Dame M, Knutson D. Vitamin A in reproduction and development. *Nutrients*. 2011;3(4):385–428.
9. Mark M, Ghyselinck NB, Chambon P. Function of retinoic acid receptors during embryonic development. *Nucl Recept Signal*. 2009;7:e002.
10. Girmenia C, Lo Coco F, Breccia M, et al. Infectious complications in patients with acute promyelocytic leukaemia treated with the AIDA regimen. *Leukemia*. 2003;17(5):925–930.
11. Lei GS, Zhang C, Shao S, Jung HW, Durant PJ, Lee CH. All-transretinoic acid in combination with primaquine clears pneumocystis infection. *PLoS One*. 2013;8(1):e53479.
12. Klassert TE, Hanisch A, Bräuer J, et al. Modulatory role of vitamin A on the *Candida albicans*-induced immune response in human monocytes. *Med Microbiol Immunol*. 2014;203(6):415–424.
13. Campione E, Paternò EJ, Costanza G, et al. Tazarotene as alternative topical treatment for onychomycosis. *Drug Des Devel Ther*. 2015;9:879–886.
14. Sommer A, Vyas KS. A global clinical view on vitamin A and carotenoids. *Am J Clin Nutr*. 2012;96(5):1204S–1206S.
15. Dawson HD, Collins G, Pyle R, et al. Direct and indirect effects of retinoic acid on human Th2 cytokine and chemokine expression by human T lymphocytes. *BMC Immunol*. 2006;7:27.
16. Tsai YC, Chang HW, Chang TT, et al. Effects of all-trans retinoic acid on Th1- and Th2-related chemokines production in monocytes. *Inflammation*. 2008;31(6):428–433.
17. Wojtal KA, Wolfram L, Frey-Wagner I, et al. The effects of vitamin A on cells of innate immunity in vitro. *Toxicol In Vitro*. 2013;27(5):1525–1532.

18. Orlandi A, Francesconi A, Cocchia D, et al. Phenotypic heterogeneity influences apoptotic susceptibility to retinoic acid and cis-platinum of rat arterial smooth muscle cells in vitro: Implications for the evolution of experimental intimal thickening. *Arterioscler Thromb Vasc Biol.* 2001;21(7):1118–1123.
19. Orlandi A, Pucci S, Ciucci A, Pichiorri F, et al. Modulation of clusterin isoforms is associated with all-trans retinoic acid-induced proliferative arrest and apoptosis of intimal smooth muscle cells. *Arterioscler Thromb Vasc Biol.* 2005;25(2):348–353.
20. Nefedova Y, Fishman M, Sherman S, et al. Mechanism of all-trans retinoic acid effect on tumor-associated myeloid-derived suppressor cells. *Cancer Res.* 2007;67(22):11021–11028.
21. Lee JM, Seo JH, Kim YJ, et al. The restoration of myeloid-derived suppressor cells as functional antigen-presenting cells by NKT cell help and all-trans-retinoic acid treatment. *Int J Cancer.* 2012;131(3):741–751.

Drug Design, Development and Therapy

Publish your work in this journal

Drug Design, Development and Therapy is an international, peer-reviewed open-access journal that spans the spectrum of drug design and development through to clinical applications. Clinical outcomes, patient safety, and programs for the development and effective, safe, and sustained use of medicines are a feature of the journal, which

Submit your manuscript here: <http://www.dovepress.com/drug-design-development-and-therapy-journal>

has also been accepted for indexing on PubMed Central. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Dovepress