Immediate results and long-term cardiovascular outcomes of endovascular therapy in octogenarians and nonoctogenarians with peripheral arterial diseases

Hsuan-Li Huang¹
 Jyh-Ming Jimmy Juang²
 Hsin-Hua Chou^{1,3}
 Chien-An Hsieh¹
 Shih-Jung Jang¹
 Shih-Tsung Cheng^{1,3}
 Yu-Lin Ko^{1,3}

¹Division of Cardiology, Department of Internal Medicine, Taipei Tzu Chi Hospital, Buddhist Tzu Chi Medical Foundation, New Taipei City,

²Cardiovascular Center and Division of Cardiology, Department of Internal Medicine, National Taiwan University Hospital, National Taiwan University College of Medicine, Taipei, ³School of Medicine, Tzu Chi University, Hualien, Taiwan

Purpose: To investigate the clinical outcomes of endovascular therapy (EVT) in octogenarians and nonoctogenarians with peripheral arterial disease.

Methods: A retrospective analysis of 511 patients (654 affected legs) who underwent EVT between July 2005 and December 2013 was conducted in a prospectively maintained database. Immediate results and long-term vascular outcomes were analyzed and compared between octogenarians and nonoctogenarians.

Results: Octogenarians were more likely to be female and have atrial fibrillation (AF), whereas nonoctogenarians had higher rates of obesity, claudication, and medical comorbidities. There were no differences in the rates of EVT success, 30-day major adverse vascular events, and 6-month functional improvement between groups. Over the 10-year follow-up period, the rates of 3-year limb salvage, sustained clinical success, freedom from major cerebrovascular and cardiovascular events, and composite vascular events were similar between groups, but the survival rate was better in nonoctogenarians than in octogenarians (73% vs 63%, respectively, $P=0.004$). In Cox regression analysis, dependence on dialysis and AF were significant predictors of death (odds ratio [OR] 4.44 in dialyzed and 2.83 in AF patients), major cerebrovascular and cardiovascular events (OR 3.49 and 2.45), and composite vascular events (OR 3.14 and 2.25).

Conclusion: EVT in octogenarians was feasible, without an increased risk of periprocedural complications. The rates of limb salvage, sustained clinical success, and long-term vascular events were comparable between groups. Dialysis dependence and AF are independent predictors for poor prognosis in patients with peripheral arterial disease. However, these observations require further confirmation in larger scale studies.

Keywords: peripheral artery disease, octogenarians, endovascular therapy, cardiovascular outcomes, dialysis, atrial fibrillation

Introduction

Patients with peripheral arterial disease (PAD) have significantly increased rates of myocardial infarction (MI), cardiovascular mortality, and stroke.¹ Cross-sectional studies have shown that approximately half of all patients with PAD have some clinical evidence of coronary artery disease or cerebrovascular disease.^{2,3} The risk of mortality and major cardiovascular events (eg, heart attack and stroke) is approximately threefold higher in PAD patients than in those without PAD.⁴⁻⁶

With an aging population and improved quality of medical care, physicians face an ever-increasing number of elderly patients presenting with progressed forms of PAD.^{7,8}

Correspondence: Hsuan-Li Huang; Yu-Lin Ko
 Division of Cardiology, Department of Internal Medicine, Taipei Tzu Chi Hospital, Buddhist Tzu Chi Medical Foundation, 289 Jiang Kuo Road, Xindian District, New Taipei City 231, Taiwan
 Tel +886 2 6628 9779 (ext 5709)
 Fax +886 2 6628 9009
 Email huang304@gmail.com;
 yulinkotw@yahoo.com.tw

Advanced age is associated with increased perioperative and postoperative mortality after vascular operations because of multiple comorbidities.^{9–12} Untreated chronic critical limb ischemia (CLI), an advanced stage of PAD, is associated with a dismal prognosis.^{1,13} On the contrary, CLI patients with successful revascularization have a better quality of life and longer survival when compared to those treated conservatively or with primary amputation.^{5,14} The wide use of endovascular therapy (EVT) and evidence of better short-term survival compared with bypass surgery^{9,10,15} render angioplasty a tempting first-choice treatment in very elderly vascular patients. There are no specific guidelines for the treatment of elderly patients with PAD, and information regarding long-term vascular events following EVT in elderly versus younger patients is scarce in Asian countries.

In this study, we performed a retrospective review using a prospective registry of all patients undergoing EVT at our institution over a 10-year period. We compared the baseline demographics and procedural outcomes between octogenarians and nonoctogenarians. The immediate results, functional improvement, and long-term cardiovascular outcomes were analyzed to determine the safety and effectiveness of EVT between groups.

Methods

Study population

Subjects for this study were derived from the Tzuchi Registry of ENDovascular Intervention for Peripheral Artery Disease (TRENDPAD), which is an ongoing, prospective, physician-initiated, single-center observational registry of patients who underwent EVT for lower limb ischemia starting from July 2005. This database was interrogated to identify adult patients with symptomatic PAD treated between July 2005 and December 2013.

To be eligible for this analysis, patients were required to have an age ≥ 18 years, provide written consent to participate in this study, and a candidate for EVT originating from atherosclerotic PAD. Patients who had acute limb ischemia, nonatherosclerotic PAD, a life-threatening infection, follow-up duration < 3 months in surviving patients, or refused to participate were not eligible. The protocol was approved by the local ethics committee and the institutional review board of Taipei Tzu Chi Hospital (IRB 03-X27-098), and all the study procedures were performed according to good clinical practices and the applicable laws of various governing bodies.

All patients received complete baseline medical examinations before EVT, which included a clinical examination, hemodynamic evaluation (ankle or toe pressure, pulse volume

recording, and duplex ultrasound), and anatomic assessment (computed tomographic angiography, magnetic resonance angiography, or diagnostic angiography). Toe pressure, pulse volume recording, and Doppler waveform pattern measurements were performed to assess the hemodynamic changes in patients with a falsely elevated ankle brachial index.

Interventions

The interventional procedure was usually conducted using an antegrade or crossover approach, but multiple access sites (brachial or pedal puncture) were used for complex cases. Detailed procedures for endovascular intervention have been described previously.¹⁶ Plain balloon angioplasty and bare metal stents were the mainstream treatment in these study participants. With advances in and introduction of new devices, additional procedures including cutting balloons (Boston Scientific, Natick, MA, USA), excimer laser angioplasty (Spectranetics, Colorado Springs, CO, USA), and TurboHawk atherectomy (eV3, Irvine, CA, USA) were performed at the discretion of the operator to obtain better immediate success. Drug-eluting devices, introduced to improve vessel patency in femoropopliteal disease, have been used since November 2011.

The major variables of demographics included in statistical analyses were age at the time of procedure (≥ 80 years); sex (male); presence of comorbidities including hypertension (systolic blood pressure > 140 mmHg or diastolic pressure > 90 mmHg), diabetes mellitus (use of oral hypoglycemic agent or insulin, fasting plasma sugar ≥ 126 mg/dL, or glycated hemoglobin level $\geq 6.5\%$), atrial fibrillation (AF), congestive heart failure (CHF) (ejection fraction $< 40\%$ by echocardiography or radionuclide ventriculography), coronary artery disease (diameter stenosis $> 50\%$ in at least one of the three main coronary arteries), stroke, and dialysis dependence; and history of smoking (defined as smoking within the previous year). Serum C-reactive protein (CRP) values, determined by high-sensitivity assay, were obtained before EVT or at admission. Normolipidemia was defined as total cholesterol < 200 mg/dL or low-density lipoprotein (LDL) cholesterol < 100 mg/dL without the use of statin treatment, and baseline serum LDL cholesterol levels were determined using the most recent value within 3 months preprocedure.

Definition and outcome end points

A single-level intervention was defined as EVT for isolated aortoiliac, femoropopliteal, or below-the-knee lesions. A multilevel intervention was defined as EVT in more than one area.

Functional status was classified as ambulatory independently, ambulatory with assistive device, wheelchair-bound, and bed-ridden status. The changes in functional status at presentation and 6 months after EVT were compared between groups.

Nonfatal cardiovascular events included nonfatal MI, stroke, or transient ischemic attack; hospitalization with decompensated heart failure; percutaneous coronary intervention or coronary artery bypass surgery; and hospitalization with angina pectoris or ischemic hands.

Vascular death included fatal stroke, or sudden death due to MI, CHF, AF, nonrheumatic heart valve disease, and ruptured aortic aneurysm.

The safety end point was 30-day major adverse vascular events (MAVEs) (death, MI, stroke, emergent surgery, contrast-induced acute kidney injury requiring dialysis, subacute arterial occlusion, groin complications requiring manual compression or surgical repair, and unplanned reintervention), while efficacy was based on procedure success (successful access and deployment of the device and $\leq 30\%$ residual stenosis by quantitative angiography with evidence of at least one patent tibial artery to the foot).

Outcomes in follow-up included major cerebrovascular and cardiovascular events (MACCEs) (any death and nonfatal cardiovascular events) and composite vascular events (CVEs) (vascular deaths, nonfatal cardiovascular events, and major amputation). Clinical improvement was defined as a ≥ 2 Rutherford category improvement, an increase in ankle brachial index (ABI) of at least 0.15, and wound healing in ≤ 4 months after the index intervention for patients with tissue loss. Sustained clinical success (SCS) was defined as clinical improvement without target vessel revascularization or major amputation (limb loss above the ankle).

Follow-up

At 1 week, 1 month, and thereafter every 3 months after EVT, each patient was assessed by symptoms, ankle or toe brachial pressure index, and duplex ultrasounds. Repeat interventions were performed if recurrent symptoms, significant vessel stenosis ($\geq 70\%$) with dampened Doppler waveform patterns by duplex ultrasound, and an ABI decrease of ≥ 0.15 were observed. The main events (death, amputation, failure of SCS, and late vascular events) were documented at discharge and at follow-up visits. If office follow-up visits were not feasible, alternate data sources included telephone interviews, medical records, the local electronic medical database, and the referring physician.

Statistical analysis

All continuous data were expressed as mean $\pm$ standard deviation and were analyzed using independent *t*-tests. A frequency comparison was performed using the chi-square test or Fisher's exact tests. Variables with $P < 0.2$ in the univariate analyses were backward selected into the multivariate analysis. Independent sample *t*-test was used to compare the functional changes after EVT between groups. Survival curves, MACCE, and CVE were plotted using the Kaplan–Meier method and analyzed using the log-rank test (SAS software version 9.2; SAS Institute Inc., Cary, NC, USA). A two-tailed *P*-value of < 0.05 was considered significant. Odds ratios (ORs) were given with 95% confidence intervals (CIs).

Results

Study participants

During the study period, 573 patients with 728 affected legs were treated with EVT. Of them, ten patients were excluded (seven with acute limb ischemia and three with vasculitis-related PAD). The remaining 563 patients were divided into two groups based on age (≥ 80 years or < 80 years). Fifty-two patients (64 legs) were not entered into the analysis of long-term outcomes: 34 patients (40 legs) with failed procedures and 18 patients (24 legs) with a follow-up time < 30 days or in-hospital mortality. The final cohort for long-term outcome analysis included 136 patients (170 legs) in octogenarians and 375 patients (484 legs) in nonoctogenarians. The flowchart of study participants is shown in Figure 1.

Baseline patient and lesion characteristics

Table 1 summarizes the baseline demographics and clinical characteristics between groups. Octogenarians were more likely to be female and have higher incidence of AF, whereas nonoctogenarians were more likely to have diabetes mellitus, dialysis dependence, history of smoking, hyperlipidemia, and a high body mass index (BMI). The hematocrit, neutrophil-to-lymphocyte ratio, platelet-to-lymphocyte ratio, and levels of serum albumin and LDL cholesterol were similar between groups, but nonoctogenarians had higher levels of glycated hemoglobin ($6.5\% \pm 1.4\%$ vs $7.6\% \pm 1.9\%$, $P < 0.001$) and CRP (3.19 ± 4.94 mg/dL vs 5.13 ± 6.96 mg/dL, $P < 0.001$). Compared to nonoctogenarians, less usage of statin was noted in octogenarians during the follow-up period (21% vs 37%, $P = 0.001$).

Table 2 summarizes the lesion characteristics in both groups. The ABI index was lower in octogenarians than

Figure 1 Flowchart of study participants.

Abbreviations: ALI, acute limb ischemia; PAD, peripheral artery disease.

in nonoctogenarians (0.46 ± 0.17 vs 0.51 ± 0.20 , $P=0.002$). Octogenarians had more patients presenting with resting pain (26% vs 18%, $P=0.022$), in contrast with more disabling claudication in nonoctogenarians (12% vs 20%, $P=0.008$). There was no difference in the presentation of tissue loss between groups. The rates of procedure success, single or multilevel intervention, and stent implantation were similar between groups.

Thirty-day outcomes and changes in functional status after EVT

Three patients in octogenarians (two with sepsis and one with rupture of an abdominal aortic aneurysm) and seven in nonoctogenarians (four with sepsis, two with cardiogenic shock, and one with hemorrhagic stroke) died in hospital. Detailed data regarding perioperative complications are shown in Table 3. No differences were observed with regard

to the number of intensive care unit (ICU) transfers after EVT (8.8% vs 9.5%, $P=0.792$) and 30-day MAVE rate (12.4% vs 10.3%, $P=0.466$) between groups. The length of hospital stay was longer in nonoctogenarians than in octogenarians (8 ± 8 vs 11 ± 12 days, $P=0.002$). Octogenarians included a higher rate of wheelchair-bound patients (68% vs 48%, $P<0.001$) at presentation; however, percentage changes in functional status after EVT were not different between the groups ($P=0.24$).

Long-term outcomes

Over a mean follow-up period of 34 ± 25 months (range 3–112), 157 patients died (50 in octogenarians and 107 in nonoctogenarians), and the rates of overall survival in both groups were significantly different at 3 (63% vs 73%) and 5 years (45% vs 67%, $P=0.004$) (Figure 2). There were no significant differences between groups in the rates of limb

Table 1 Baseline demographics

	Octogenarians	Nonoctogenarians	P-value
No of patients	136	375	
Age (years)	85±4	66±9	<0.001*
Male sex	65 (48)	218 (58)	0.038*
Underlying disease			
Diabetes mellitus	77 (57)	317 (85)	<0.001*
Hypertension	112 (82)	319 (85)	0.456
CAD	54 (40)	186 (50)	0.047*
CVD	35 (26)	79 (21)	0.252
ESRD	30 (22)	169 (45)	<0.001*
CHF	23 (17)	65 (17)	0.614
Smoking	38 (28)	149 (40)	0.014*
Hyperlipidemia	48 (35)	194 (52)	0.001*
Atrial fibrillation	35 (26)	43 (11)	<0.001*
Body height (m)	1.57±0.09	1.60±0.08	<0.001*
Body weight (kg)	57±11	62±10	<0.001*
BMI (kg/m ²)	23.1±3.5	24.2±3.3	<0.001*
Hematocrit (mg/dL)	33.8±4.9	34.0±5.9	0.645
CRP (mg/dL)	3.19±4.94	5.13±6.96	<0.001*
NLR	4.32±4.48	5.46±8.58	0.120
PLR	183±235	190±137	0.618
HbA1C (%)	6.5±1.4	7.6±1.9	<0.001*
LDL-C (mg/dL)	93±30	98±35	0.063
Albumin (mg/dL)	3.19±0.61	3.15±0.67	0.183
Medications			
Aspirin	70 (51)	213 (57)	0.284
Thienopyridine	109 (80)	305 (81)	0.762
Cilostazole	103 (76)	281 (75)	0.383
Statin	29 (21)	137 (37)	0.001*
Beta-blocker	47 (35)	148 (39)	0.313
ACEI or ARB	68 (50)	186 (50)	0.936

Notes: Values are mean ± SD or n (%); **P*<0.05 indicates a significant difference between groups.

Abbreviations: ACEI, angiotensin-converting enzyme inhibitor; ARB, angiotensin receptor blocker; BMI, body mass index; CAD, coronary artery disease; CHF, congestive heart failure; CRP, C-reactive protein; CVD, cerebrovascular disease; ESRD, end-stage renal disease; HbA1C, glycated hemoglobin; LDL-C, low-density lipoprotein cholesterol; NLR, neutrophil-to-lymphocyte ratio; PLR, platelet-to-lymphocyte ratio; SD, standard deviation.

salvage (91% vs 87%, *P*=0.07) and SCS at 3 years (51% vs 47%, *P*=0.297). The rates of freedom from cardiovascular death (84% vs 89%, *P*=0.122), nonfatal cardiovascular events (78% vs 73%, *P*=0.121), and stroke (86% vs 87%, *P*=0.504) at 3 years were similar between groups. More octogenarians died owing to noncardiovascular problems, mainly from sepsis followed by malignancy. There were no between-group differences at 3 years in the rates of freedom from MACCEs (50% vs 55%, *P*=0.47) or CVEs (63% vs 61%, *P*=0.38) (Figures 3 and 4). However, non-octogenarians had higher rates of nonfatal MI during the follow-up period compared with octogenarians (19% vs 8%, *P*=0.049). Multivariate analysis showed that the octogenarian factor was not significantly associated with

Table 2 Characteristics of the affected limb

	Octogenarians	Nonoctogenarians	P-value
No of affected limbs	170	484	
Target-limb ABI	0.46±0.17	0.51±0.20	0.002*
Stiff artery	24 (14%)	71 (15%)	0.861
Clinical presentation			
Intermittent claudication	20 (12%)	97 (20%)	0.008*
Rest pain	44 (26%)	86 (18%)	0.022*
Nonhealing ulcer	79 (46%)	219 (45%)	0.783
Gangrene	27 (16%)	82 (17%)	0.458
Procedure success	175 (94%)	503 (95%)	0.557
Single level	83 (49%)	272 (56%)	0.096
Iliac	12	55	0.111
FP	36	95	0.664
BTK	35	122	0.225
Multilevel	87 (51%)	212 (44%)	0.096
Iliac + FP	7	13	0.351
FP + BTK	79	192	0.121
Ilio-FP-BTK	1	7	0.381
Iliac and BTK	0	2	0.401
Stenting rate	99 (58%)	272 (56%)	0.645

Notes: Values are mean ± SD or n (%); **P*<0.05 indicates a significant difference between groups.

Abbreviations: ABI, ankle brachial index; FP, femoropopliteal; BTK, below-the-knee; SD, standard deviation.

MACCEs or CVEs. Dialysis dependence and AF were strong independent predictors of overall survival (OR 4.44; 95% CI 1.79–11.1; *P*=0.001 and OR 2.83; 95% CI 1.49–5.35; *P*=0.001, respectively), MACCE (OR 3.49; 95% CI 1.48–8.19; *P*=0.004 and OR 2.45; 95% CI 1.31–4.45; *P*=0.005, respectively), and CVE (OR 3.14; 95% CI 1.32–7.48; *P*=0.009 and OR 2.25; 95% CI 1.21–4.17; *P*=0.010, respectively) (Table 4).

Discussion

Our results show that the safety and efficacy of EVT in terms of procedure success, 30-day MAVEs, functional improvement, limb salvage, and SCS are comparable between octogenarians and nonoctogenarians. Although younger patients had more traditional risk factors, age did not increase the risk of long-term vascular events over the 10-year study period. Nonoctogenarians have a survival advantage as compared to octogenarians. Regular dialysis and AF were independent predictors for long-term survival, MACCEs, and CVEs.

PAD is a progressive disease of atherosclerosis, and PAD-related death and disability has increased globally and regionally in the past 20 years.¹³ There are few reports comparing cardiovascular risk factor profiles between octogenarians and nonoctogenarians, especially in the Asian countries. Our findings that nonoctogenarians with PAD had a higher BMI

Table 3 Thirty-day outcomes and changes in functional status after EVT

	Octogenarians	Nonoctogenarians	P-value
No of affected limbs	170	484	
Death	3	7	0.771
MI	2	6	0.949
CVA	2	3	0.609
Emergent surgery	0	2	0.401
AKI requiring dialysis	0	3	0.304
Groin complications	10	24	0.641
Subacute arterial occlusion	2	1	0.107
Unplanned reintervention	0	3	0.304
UGI bleeding with shock	1	0	0.553
Compartment syndrome	1	1	0.438
Length of stay (days)	8±8	11±12	0.002*
Number of ICU transfers	15 (8.8%)	46 (9.5%)	0.792
MAVE	21 (12.4%)	50 (10.3%)	0.466
Changes in functional status after EVT	Octogenarians N=115	Nonoctogenarians N=333	P-value 0.24
	Admission	Admission	6 months
AI	2 (2%)	42 (13%)	193 (58%)
AA	23 (20%)	161 (36%)	75 (23%)
WH	79 (68%)	121 (48%)	60 (18%)
BED	11 (10%)	9 (3%)	5 (2%)

Notes: Values are mean ± SD or n (%); * $P < 0.05$ indicates a significant difference between groups.

Abbreviations: EVT, endovascular therapy; AI, ambulatory independently; AA, ambulatory with assisting device; BED, bed-ridden status; CVA, cerebrovascular accident; MI, myocardial infarction; AKI, acute kidney injury; UGI, upper gastrointestinal; ICU, intensive care unit; MAVE, major adverse vascular event; WH, wheelchair-bound; SD, standard deviation.

and more atherosclerotic risk factors are in agreement with an earlier study of octogenarians and septuagenarians, which suggested that younger PAD patients have more comorbid disease.¹⁷ On the contrary, more patients with AF and female sex were noted with PAD in octogenarians.

Patients with symptomatic PAD were considered candidates for revascularization, either endovascular or surgical therapy. Untreated chronic CLI, an advanced stage of PAD, is associated with a dismal prognosis.^{1,13} Instead, patients with CLI who undergo successful revascularization survive longer and experience greater improvements in quality of life when

compared to patients treated conservatively or with primary amputation.^{5,12} The benefits of revascularization may be attenuated when taking into account the aging of the population, which significantly increases the number of elderly patients at high risk of perioperative mortality due to limb-saving procedures, adverse cardiovascular events, and other medical comorbidities.^{9–12} There are no specific guidelines regarding revascularization procedures in elderly patients with symptomatic PAD. Our study highlights the comparable safety of EVT in both treatment groups. The rates of procedure success, number of ICU transfers, and 30-day MAVEs were similar

Figure 2 Kaplan–Meier curves for overall survival.

Notes: The survival rates for octogenarians and nonoctogenarians were 63% and 73% at 3 years and 45% and 67% at 5 years, respectively ($P=0.004$). Yellow line and blue line represent octogenarians and nonoctogenarians, respectively.

Figure 3 Kaplan–Meier curves for freedom from major cerebrovascular and cardiovascular events (MACCEs).

Notes: The rates of 3-year freedom from MACCEs were similar between groups (50% in octogenarians vs 55% in nonoctogenarians, $P=0.564$). Yellow line and blue line represent octogenarians and nonoctogenarians, respectively.

Figure 4 Kaplan–Meier curves for freedom from composite vascular events (CVEs).

Notes: The rates of 3-year freedom from CVEs were similar between groups (63% in octogenarians vs 61% in nonoctogenarians, $P=0.308$). Yellow line and blue line represent octogenarians and nonoctogenarians, respectively.

between groups. Nonoctogenarians had a longer hospital stay when compared to octogenarians, which was related to more tissue inflammation and debridement surgery in these patients. Our results are consistent with previous reports^{9–11,14} showing that minimally invasive EVT is safe in octogenarians and does not increase the risk of periprocedural complications as compared to nonoctogenarians. Recent advances in endovascular techniques and devices, coupled with improved quality of medical care, render EVT a tempting first-choice treatment in very elderly vascular patients.

Although more octogenarians, associated with age-related neurologic or musculoskeletal problems, were wheelchair-bound at presentation than nonoctogenarians, no significant difference was found in both groups regarding the changes of functional status 6 months after EVT. These findings suggested that EVT has beneficial effect in improving ambulatory function in elderly patients.

Table 4 Crude and significant multivariate risk factors for long-term events

Outcome predictors	OR (95% CI)	P-value
Overall survival		
Atrial fibrillation	2.83 (1.49–5.35)	0.001
Dialysis dependence	4.44 (1.79–11.1)	0.001
Critical limb ischemia	3.19 (1.31–7.74)	0.001
Major cerebrovascular and cardiovascular events		
Atrial fibrillation	2.45 (1.31–4.45)	0.005
Dialysis dependence	3.49 (1.48–8.19)	0.004
Coronary artery disease	1.74 (1.13–2.66)	0.011
C-reactive protein	1.05 (1.01–1.09)	0.019
Composite vascular events		
Atrial fibrillation	2.25 (1.21–4.17)	0.010
Dialysis dependence	3.14 (1.32–7.48)	0.009
Coronary artery disease	1.84 (1.18–2.86)	0.007

Abbreviations: OR, odds ratio; CI, confidence interval.

The long-term survival rate of octogenarians in our study remained lower than that of nonoctogenarians (63% and 73% at 3 years and 45% and 67% at 5 years, respectively). Although cardiovascular death remained a major cause of death in this study cohort (35%, 55/157), no differences were observed in subsequent cardiovascular deaths between groups. The leading cause of death in octogenarians was infectious disease (52% due to sepsis and pneumonia), which reflects the impaired immune and functional status in the aging population.

Previous reports have shown that elderly patients are at higher risk of adverse events following PAD treatment because of multiple comorbidities and relatively reduced physiologic reserve.² However, reported outcomes of EVT for octogenarians mainly focused on mortality, amputation, and reintervention in prior studies.^{9–15,17,18} Long-term data regarding late CVEs in symptomatic PAD patients are scarce.¹⁹ We included major amputation in our CVE outcomes, which led to more PAD-related predictors. In this study, only nonfatal MI reached the marginal significance between groups ($P=0.049$), which may be associated with the longer life expectancy and more atherosclerosis risk factors in nonoctogenarians. After successful EVT, octogenarians showed no increase in the rates of MACCEs and CVEs as compared to nonoctogenarians. In addition, late vascular events can be prevented by optimal medical care, such as increased use of beta-blockers periprocedure, coupled with antiplatelet agents, statin and angiotensin-converting enzyme inhibitors, or angiotensin receptor blockers during long-term management.^{20–24}

The Cox regression model showed that dialysis dependence and AF are independent risk factors predicting subsequent cardiovascular events despite age stratification. AF is an independent risk factor for survival in patients with CLI.²⁵ In this study, AF remained an independent predictor of survival, MACCE, and CVE outcomes. The prevalence of AF is higher in patients with systemic atherothrombosis.^{26,27} Severe PAD and a high CHADS₂ (congestive heart failure, hypertension, age ≥ 75 years, diabetes mellitus, stroke) score²⁸ were observed in many of our patients, which might explain why AF played an important role in late cardiovascular events. Dialyzed patients with PAD have a higher rate of mortality and cardiovascular events.^{29,30} Despite improved limb salvage from EVT in dialyzed patients with PAD, the overall survival in this group remained poor.^{31,32} Our results are consistent with prior reports,^{33,34} wherein dialysis is an important predictor of mortality and subsequent cardiovascular events in PAD patients when other risk factors are adjusted. Traditional risk factors, coupled with vascular inflammation and malnutrition, were associated with general atherosclerosis in dialyzed patients with PAD.

Study limitations

This study has some limitations. First, it was a single-center, observational study using a prospective database. Treatment allocation was made at the discretion of the operator and the patient's policy of reimbursement. Second, the fact that all patients were treated at a single center in Taiwan opens up the possibility of referral/selection bias based on the current practice of this group. As with all observational studies, the reported association may not represent the underlying causality. The unavoidable risk of selective bias and differences in long-term survival not due to exposure to the treatment being studied threatens the observations made in a cohort study. Third, underutilization of statins may have negatively impacted our ability to prevent late ischemic events, which may relate to reimbursed policy and less hyperlipidemia in octogenarians and dialyzed patients. Finally, we did not investigate the mechanisms underlying the effect of AF on PAD.

Conclusion

In conclusion, EVT in octogenarians is feasible, without an increased risk of periprocedural complications. Nonoctogenarians have a survival advantage as compared to octogenarians. However, the rates of limb salvage, SCS, long-term CVEs, and MACCEs were comparable between groups. Dialysis dependence and AF are independent predictors of long-term survival, MACCEs, and CVEs.

Acknowledgments

The authors thank the cardiac catheterization laboratory medical staff and clinical research coordinators of Taipei Tzu Chi hospital who had participated in this study.

Disclosure

The authors report no conflicts of interest in this work.

References

- Hirsch AT, Haskal ZJ, Hertzner NR, et al. ACC/AHA 2005 Practice Guidelines for the management of patients with peripheral arterial disease (lower extremity, renal, mesenteric, and abdominal aortic): a collaborative report from the American Association for Vascular Surgery/Society for Vascular Surgery, Society for Cardiovascular Angiography and Interventions, Society for Vascular Medicine and Biology, Society of Interventional Radiology, and the ACC/AHA Task Force on Practice Guidelines (Writing Committee to Develop Guidelines for the Management of Patients With Peripheral Arterial Disease); endorsed by the American Association of Cardiovascular and Pulmonary Rehabilitation; National Heart, Lung, and Blood Institute; Society for Vascular Nursing; TransAtlantic Inter-Society Consensus; and Vascular Disease Foundation. *Circulation*. 2006;113(11):e463–e654.
- Hooi JD, Stoffers HE, Knottnerus JA, et al. The prognosis of non-critical limb ischaemia: a systematic review of population-based evidence. *Br J Gen Pract*. 1999;49(438):49–55.
- Hooi JD, Stoffers HE, Kester AD, et al. Risk factors and cardiovascular diseases associated with asymptomatic peripheral arterial occlusive disease. The Limburg PAOD Study. *Peripheral Arterial Occlusive Disease. Scand J Prim Health Care*. 1998;16(3):177–182.
- Pande RL, Perlstein TS, Beckman JA, et al. Secondary prevention and mortality in peripheral artery disease: National Health and Nutrition Examination Study, 1999 to 2004. *Circulation*. 2011;124(1):17–23.
- Smith GD, Shipley MJ, Rose G. Intermittent claudication, heart disease risk factors, and mortality. The Whitehall Study. *Circulation*. 1990;82(6):1925–1931.
- Ankle Brachial Index Collaboration. Ankle brachial index combined with Framingham Risk Score to predict cardiovascular events and mortality: a meta-analysis. *JAMA*. 2008;300(2):197–208.
- Fowkes FG, Rudan D, Rudan I, et al. Comparison of global estimates of prevalence and risk factors for peripheral artery disease in 2000 and 2010: a systematic review and analysis. *Lancet*. 2013;382(9901):1329–1340.
- Diehm C, Schuster A, Allenberg JR, et al. High prevalence of peripheral arterial disease and co-morbidity in 6880 primary care patients: cross-sectional study. *Atherosclerosis*. 2004;172(9):95–105.
- Brosi P, Dick F, Do DD, et al. Revascularization for chronic critical lower limb ischemia in octogenarians is worthwhile. *J Vasc Surg*. 2007;46(6):1198–1207.
- Dosluoglu HH, Lall P, Cherr GS, et al. Superior limb salvage with endovascular therapy in octogenarians with critical limb ischemia. *J Vasc Surg*. 2009;50(2):305–315.
- Arvela E, Venermo M, Söderström M, et al. Infrainguinal percutaneous transluminal angioplasty or bypass surgery in patients aged 80 years and older with critical leg ischaemia. *Br J Surg*. 2011;98(4):518–526.
- Dick P, Barth B, Mlekusch W, et al. Complications after peripheral vascular interventions in octogenarians. *J Endovasc Ther*. 2008;15(4):383–389.
- Norgren L, Hiatt WR, Dormandy JA; for TASC II Working Group. Inter-Society Consensus for the Management of Peripheral Arterial Disease (TASC II). *J Vasc Surg*. 2007;45(Suppl S):S5–S67.
- Klevsgård R, Risberg BO, Thomsen MB, et al. A 1-year follow-up quality of life study after hemodynamically successful or unsuccessful surgical revascularization of lower limb ischemia. *J Vasc Surg*. 2001;33(1):114–122.
- Adam DJ, Beard JD, Cleveland T; for BASIL trial participants. Bypass versus angioplasty in severe ischaemia of the leg (BASIL): multicentre, randomized controlled trial. *Lancet*. 2005;366(9501):1925–1934.
- Huang HL, Chou HH, Wu TY, et al. Endovascular intervention in Taiwanese patients with critical limb ischemia: patient outcomes in 333 consecutive limb procedures with a 3-year follow-up. *J Formos Med Assoc*. 2014;113(10):688–695.
- Chang JB, Stein TA. Infrainguinal revascularizations in octogenarians and septuagenarians. *J Vasc Surg*. 2001;34(1):133–138.
- Jones DW, Siracuse JJ, Graham A, et al. Safety and effectiveness of endovascular therapy for claudication in octogenarians. *Ann Vasc Surg*. 2015;29(1):34–41.
- Van Hattum ES, Tangelder MJ, Lawson JA, et al. Long-term risk of vascular events after peripheral bypass surgery. A cohort study. *Thromb Haemost*. 2012;108(3):543–553.
- Westin GG, Armstrong EJ, Bang H, et al. Association between statin medications and mortality, major adverse cardiovascular event, and amputation-free survival in patients with critical limb ischemia. *J Am Coll Cardiol*. 2014;63(7):682–690.
- Chung J, Timaran DA, Modrall JG, et al. Optimal medical therapy predicts amputation-free survival in chronic critical limb ischemia. *J Vasc Surg*. 2013;58(4):972–980.
- Lindenauer PK, Pekow P, Wang K, et al. Perioperative beta-blocker therapy and mortality after major non-cardiac surgery. *N Engl J Med*. 2005;353(4):349–361.
- Poldermans D, Bax JJ, Kertai MD, et al. Statins are associated with a reduced incidence of perioperative mortality in patients undergoing major noncardiac vascular surgery. *Circulation*. 2003;107(14):1848–1851.
- Yusuf S, Sleight P, Pogue J, et al. Effects of an angiotensin-converting-enzyme inhibitor, ramipril, on cardiovascular events in high-risk patients. The Heart Outcomes Prevention Evaluation Study Investigators. *N Engl J Med*. 2000;342(3):145–153.

25. Chang SH, Tsai YJ, Chou HH, et al. Clinical predictors of long-term outcomes in patients with critical limb ischemia who have undergone endovascular therapy. *Angiology*. 2014;65(4):315–322.
26. Winkel TA, Hoeks SE, Schouten O, et al. Prognosis of atrial fibrillation in patients with symptomatic peripheral arterial disease: data from the REduction of Atherothrombosis for Continued Health (REACH) Registry. *Eur J Vasc Endovasc Surg*. 2010;40(1):9–16.
27. Goto S, Bhatt DL, Röther J; for REACH Registry Investigators. Prevalence, clinical profile, and cardiovascular outcomes of atrial fibrillation patients with atherothrombosis. *Am Heart J*. 2008;156(5):855–863.
28. Gage BF, Waterman AD, Shannon W, et al. Validation of clinical classification schemes for predicting stroke: results from the National Registry of Atrial Fibrillation. *JAMA*. 2001;285(22):2864–2870.
29. Rajagopalan S, DelleGrottaglie S, Furniss AL, et al. Peripheral arterial disease in patients with end-stage renal disease: observations from the Dialysis Outcomes and Practice Patterns Study (DOPPS). *Circulation*. 2006;114(18):1914–1922.
30. O'Hare A, Johansen K. Lower-extremity peripheral arterial disease among patients with end-stage renal disease. *J Am Soc Nephrol*. 2001;12(12):2838–2847.
31. Nakano M, Hirano K, Iida O, et al. Prognosis of critical limb ischemia in hemodialysis patients after isolated infrapopliteal balloon angioplasty: results from the Japan below-the-knee artery treatment (J-BEAT) registry. *J Endovasc Ther*. 2013;20(1):113–124.
32. Kawarada O, Yokoi Y, Higashimori A, et al. Impact of end-stage renal disease in patients with critical limb ischaemia undergoing infrapopliteal intervention. *Eurointervention*. 2014;10(6):753–760.
33. Schanzer A, Mega J, Meadows J, et al. Risk stratification in critical limb ischemia: derivation and validation of a model to predict amputation-free survival using multicenter surgical outcomes data. *J Vasc Surg*. 2008;48(6):1464–1471.
34. Owens CD, Ho KJ, Kim S, et al. Refinement of survival prediction in patients undergoing lower extremity bypass surgery: stratification by chronic kidney disease classification. *J Vasc Surg*. 2007;45(5):944–952.

Clinical Interventions in Aging

Publish your work in this journal

Clinical Interventions in Aging is an international, peer-reviewed journal focusing on evidence-based reports on the value or lack thereof of treatments intended to prevent or delay the onset of maladaptive correlates of aging in human beings. This journal is indexed on PubMed Central, MedLine,

CAS, Scopus and the Elsevier Bibliographic databases. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <http://www.dovepress.com/clinical-interventions-in-aging-journal>

Dovepress