Reliability of clinical monitoring for the diagnosis of babesiosis in dogs in Nigeria

Olufunke Omowunmi Adebayo,¹
 Rasheed Adetola Ajadi,² Temidayo
 Olutayo Omobowale,³ Samuel
 Olatunbosun Omotainse,⁴
 Morenike Atinuke Dipeolu,⁵ Helen
 Oyebukola Nottidge,³ Ebenezer
 Babatunde Otesile²

¹Veterinary Teaching Hospital,

²Department of Veterinary Medicine and
 Surgery, College of Veterinary Medicine,
 Federal University of Agriculture,
 Abeokuta, ³Faculty of Veterinary
 Medicine, University of Ibadan, Ibadan,
⁴Department of Veterinary Pathology,
⁵Department of Veterinary Public
 Health and Reproduction, College of
 Veterinary Medicine, Federal University
 of Agriculture, Abeokuta, Nigeria

→ Video abstract

Point your SmartPhone at the code above. If you have a
 QR code reader the video abstract will appear. Or use:
<http://youtu.be/Dk-GfJgEtY>

Abstract: Babesiosis accounts for a high percentage of hospital cases in canines in Africa, with about 40% mortality in the cases presented. In Nigeria, records show an estimated 30% annual morbidity when diagnosis is largely based on clinical and laboratory findings. This study monitored clinical indices associated with canine babesiosis. One hundred and three babesiosis-suspected dogs were selected on the basis of clinical signs of anorexia, fever, presence of ticks, and enlarged lymph nodes or spleen when clinical parameters were recorded at the time of presentation. Parasite detection was done using thin blood smears; that is, the presence of *Babesia* merozoites was compared between capillary and cephalic blood. Blood was also assayed for hematology and blood chemistry using automated blood analyzers. The babesiosis-infected dogs' outcome was monitored. Data obtained were analyzed using chi-square test, analysis of variance, and Pearson's correlation. Results based on thin blood smears showed that 61.1% of the dogs were positive for *Babesia* species. Breed disposition, sex, and age did not significantly influence the incidence of *Babesia canis*, while mean rectal temperatures did not differ significantly between the cases ($P>0.05$). Heart rate and pulse rates of *Babesia*-positive dogs were significantly ($P<0.05$) higher than those that were negative. The packed cell volume between the cases was not significantly different, with the values in the positive and negative case obtained being $26.4\% \pm 11.26\%$ and $31.6\% \pm 11.9\%$, respectively, with a range of 6% to 50% and 10% to 47% observed, respectively. Normal leukogram was also observed in 62% of the *Babesia*-positive cases while 22.2% and 15.8% had leukocytosis and leukopenia, respectively. Most of the positive cases whose results were based on thin blood smear were treated with 5% oxytetracycline for 5 days and fully recovered. Pearson's correlation was used to give relationship in the observed data. This study concluded that clinical indices are not reliable markers in the diagnosis of canine babesiosis.

Keywords: canine babesiosis, *Babesia* species, dogs, clinical parameters, markers.

Introduction

Canine babesiosis, an important tick-borne protozoan disease caused mainly by both large and small *Babesia* species, is very endemic to different parts of the world, and presents varying clinical, hematological, and pathological manifestation depending on the species and subspecies involved.^{1,2}

Diagnosis of canine babesiosis is done mainly using clinical signs, history, and identification of the parasite in thin blood smears.³ Recently, the use of molecular means of identification through DNA extraction, amplification, and sequencing of the *Babesia* spp. has become a common practice in different parts of the world.^{4,5} However, in Nigeria, where *Babesia canis rossi* accounts for the majority of the *Babesia* spp. involved in canine babesiosis,⁶ diagnosis of the disease is routinely done using nonspecific clinical signs and identification of the parasites on thin

Correspondence: Olufunke Omowunmi
 Adebayo
 Veterinary Teaching Hospital, College of
 Veterinary Medicine, Federal University of
 Agriculture, Abeokuta, PMB 2240 Alabata,
 Abeokuta Ogun, Nigeria
 Tel +234 80 3566 1285
 Email adebayo.fk@gmail.com

submit your manuscript | www.dovepress.com

Dovepress

<http://dx.doi.org/10.2147/VMR.104072>

Veterinary Medicine: Research and Reports 2016:7 85–90

© 2016 Adebayo et al. This work is published and licensed by Dove Medical Press Limited. The full terms of this license are available at <http://www.dovepress.com/terms.php> and incorporate the Creative Commons Attribution – Non Commercial (unported, v3.0) License (<http://creativecommons.org/licenses/by-nc/3.0/>). By accessing the work you hereby accept the Terms. Non-commercial uses of the work are permitted without any further permission from Dove Medical Press Limited, provided the work is properly attributed. For permission for commercial use of this work, please see paragraphs 4.2 and 5 of our Terms (<http://www.dovepress.com/terms.php>).

blood smears, which is mainly subjective as identification depends on the expertise of the laboratory personnel.⁷ With the prevalence of canine babesiosis reportedly recorded between 2% and 43.6% in all parts of Nigeria and the high cost of molecular means of identification of the *Babesia* spp., there is a dire need for prompt, easy, and cost-effective means of detecting this disease.

This study evaluated the sensitivity and specificity of clinical parameters associated with the diagnosis of naturally occurring babesiosis in Nigeria.

Materials and methods

One hundred and three dogs, of which 48 were males and 55 were females, of different ages and breeds, and having clinical signs suggestive of canine babesiosis, were included in this study following informed owners' consent.

The dogs were located in the southwestern part of Nigeria in three different locations (Veterinary Teaching Hospital of the Federal University of Agriculture, Abeokuta; Ministry of Agriculture, Veterinary Hospital Complex, Ita-eko; Mokola Veterinary Clinic, Ibadan, Nigeria). The inclusion criteria consisted of presence or history of tick infestation, anorexia, and any two of the following clinical signs: persistent fever, splenomegaly, anemia, hemoglobinuria, and jaundice. Excluded were dogs with a history of treatment in the preceding 3 weeks, presence of concurrent disease, or detection of *Ehrlichia canis* morula on capillary blood smears.

Pertinent signalment such as breed, age, and sex was noted; also recorded were other clinical parameters such as rectal temperature, heart rate, pulse rate, capillary refill time, and respiratory rates.

Blood was collected in a 5 mL sodium ethylenediamine-tetraacetic acid bottle for the use of parasite detection from a thin blood smear and to determine the packed cell volume (PCV). Giemsa staining technique was employed for the thin blood smear while an automated blood analyzer was used for the PCV.

Statistical analysis

Data were analyzed using Student's *t*-test and were presented as mean $\pm$ standard deviation for clinical parameters and levels of plasma bilirubin, plasma proteins, and serum enzymes. Level of significance was set at 0.05. One-way analysis of variance was used for data from the *Babesia* positive and negative cases. Pearson's correlation coefficient was used to determine correlation between clinical parameters in the *Babesia*-positive case. Age and sex distribution between cases was analyzed using the chi-square test.

Results

On the basis of the thin blood smears, the overall prevalence of canine babesiosis in this study population was 61.1%. Parasite detection was observed more on cephalic (57.4%) than on capillary (34.4%) thin blood smears, while 8.2% were observed on both sites. Most observed clinical findings associated with the disease were tick infestation (86.6%), fever (57.2%), and anorexia (49%) in positive cases. In the negative cases, however, tick infestation, fever, and anorexia had percentages of 90.1%, 46.4%, and 36%, respectively. Table 1 shows the distribution of the study population using clinical parameters.

The breed distribution (Table 2) revealed a higher number of positive cases in three breeds, namely, Alsatian (62.2%), Rottweiler (55.0%), and Boerboel (64.3%), out of the 14 breeds present.

Among sexes, 66.5% prevalence was observed in males (31/48) and in females prevalence was observed in 56.2% (30/55). Adult dogs (≥ 9 months old) showed a prevalence

Table 1 Distribution of study population according to clinical signs and parameters used for their selection

Clinical signs, n (%)	Number of dogs with clinical manifestation	
	<i>Babesia</i> -positive dogs (n=63)	<i>Babesia</i> -negative dogs (n=40)
Tick infestation	53 (86.6)	35 (90.1)
Anorexia	30 (49.0)	14 (36.0)
Fever	35 (57.2)	18 (46.4)
Splenomegaly	12 (19.6)	8 (20.6)
Anemia (pale mucosa)	17 (27.7)	9 (23.1)
Hemoglobinuria	3 (4.9)	2 (5.2)
Jaundice	6 (9.8)	2 (5.2)

Table 2 Breed distribution of dogs with or without *Babesia* spp.

SN	Breed	NP	%P	NN	%N	Total
1	Alsatian	32	62.2	21	40.8	53
2	Belgian Malamute	0	0	1	100	1
3	Boerboel	5	64.3	3	38.6	8
4	Bullmastiff	1	50.0	1	50.0	2
5	Caucasian	5	100	0	0	5
6	Crossbreed	3	60.0	2	40.0	5
7	Dalmatian	1	50.0	1	50.0	2
8	Doberman	1	–	0	0	1
9	Labrador Retriever	0	0	2	100	2
10	Maltese	1	–	0	0	1
11	Pit Bull Terrier	1	–	0	0	1
12	Rottweiler	11	55.0	9	45.0	20
13	Samoyed	1	–	0	0	1
14	Terrier Cross	1	–	0	0	1
	Total	63	–	40	–	103

Abbreviations: SN, serial number; NP, number of positive cases; %P, percentage of positive cases; NN, number of negative cases; %N, percentage of negative cases.

of 59.1% (39/68) while puppies (<9 months old) had 64.7% (22/35), with equally no significant difference between the cases and the outcome ($P=0.67$; odds ratio 0.79).

Mean rectal temperatures were not significantly different between cases ($P>0.05$) with positive cases having a mean $\pm$ standard deviation of $39.8^{\circ}\text{C}\pm 0.95^{\circ}\text{C}$ and negative cases having $39.6^{\circ}\text{C}\pm 0.89^{\circ}\text{C}$. Higher temperatures ($>39.4^{\circ}\text{C}$) were seen in positive cases (40/63) compared to the negative case (23/40). A total of 61% were pyrexemic while 35% and 4% had normal and subnormal temperatures, respectively. Forty-seven-point-five percent (19/40) of pyrexemic cases were males, while 52.5% (21/40) were females. Regarding the age case, 67.5% (27/40) of pyrexemic cases were adults while 32.5% (13/40) were puppies. An odds ratio of 0.90 was arrived at for the relationship of pyrexia observed between adults and puppies within cases.

The study revealed significantly higher values of mean heart rate in positive case ($P<0.05$; 128 ± 19.7 bpm) compared to the negative case (118 ± 17.7 bpm) while rates >130 bpm of both heart and pulse rates were seen more often (33/63) in cases that were positive with canine babesiosis. Normal rates (90–120 bpm) were observed in 28/63 and 23/40 of *Babesia*-positive and *Babesia*-negative cases, respectively.

Conversely, the mean respiratory rates were not significantly different between those that were positive and those negative for *Babesia* spp. However, increased respiratory rates (>34 cycles per minute) were observed in 21/63 of positive and 14/40 of the negative cases, while decreased respiratory rates (<18 cycles per minute) were seen in 3% of the total number of cases. Approximately 62% of the total number of canines had normal respiratory rates (18–34 cycles per minute).

The PCV and capillary refill time did not differ significantly between dogs who were positive or negative for *Babesia* spp. The severity of anemia varied between the canines that were positive for *Babesia* spp.; based on the result of the variations in packed cell volume it was observed that out of the 63 dogs, 21 (34.3%), 14 (22%), eight (12.7%), and 20 (31.7%) had no anemia ($>30\%$), mild anemia (25% to 30%), moderate anemia (18% to 24%), and severe anemia ($<18\%$), respectively. Table 3 compares the severity of anemia observed in the positive and negative cases.

The mean PCV in the positive and negative cases were 26.4% (26.4 ± 11.26) and 28.8% (28.8 ± 10.5), respectively, with ranges of 6% to 50% and 10% to 47%, respectively. In the study, three out of the 63 positive cases had hypoxia and respiratory distress, while two within this case had severe

Table 3 Severity of anemia in *Babesia*-positive and *Babesia*-negative cases

SN	Severity of anemia (PCV, %)	NP	%P	NN	%N
1	Severe (<18)	20	31.7	7	17.9
2	Moderate (18–24)	8	12.7	6	15.4
3	Mild (25–30)	14	22.2	12	30.8
4	Not anemic (>30)	21	33.3	14	35.9
	Total	63	100	39	100

Abbreviations: SN, serial number; PCV, packed cell volume; NP, number of positive cases; %P, percentage of positive cases; NN, number of negative cases; %N, percentage of negative cases.

Table 4 Severity of anemia within breeds positive with *Babesia* spp.

Breed	No anemia	Mild anemia	Moderate anemia	Severe anemia	Total
Alsatian	12	2	3	10	27
Dalmatian	1	–	–	1	2
Boerboel	–	3	1	2	6
Pit Bull	–	–	1	–	1
Maltese	–	1	–	–	1
Rottweiler Cross	1	–	–	–	1
Caucasian	1	–	2	1	4
Samoyed	–	1	–	–	1
Terrier	1	–	–	–	1
Neapolitan Mastiff	1	–	–	–	1
Rottweiler	2	3	–	5	10
Bullmastiff	–	2	–	–	2
Mongrel	2	1	–	1	4
Alsatian Cross	–	1	1	–	2
Total	21	14	8	20	63

Table 5 Leukocytic changes in *Babesia*-positive dogs

Leukocyte change	Number of positive dogs	% Positive
Leukopenia ($<5,000$)	10	15.8
Leukocytosis ($>17,000$)	14	22.2
Normal (5,000–17,000)	39	62
Total	63	100

anemia. Only three of the positive case had congestion of the mucous membranes. Table 4 shows variation in PCV values within the breed of dogs.

In this report, the white blood cell (WBC) count, the mean neutrophil count, and the mean lymphocyte count did not significantly differ between dogs that were positive and negative to *Babesia* spp. In addition, normal leukogram was also observed in 62% of the *Babesia*-positive cases. Approximately 22% and 15.8% had leukocytosis and leukopenia, respectively, within the positive case, as shown in Table 5. In this report, variable leukogram was observed as *Babesia*-positive cases showed as normal leukogram (62%), leukocytosis (22%) and leukopenia (15.8%) in cases observed, thus, supporting the report by Di Cicco and Birkenheuer.¹ Table 6 shows breed-wise leukocytic changes in dogs.

Most of the positive cases were treated with 5% oxytetracycline (5 mg/kg) for 5 days and a total of 50/63 of the positive cases recovered, while eleven dogs died. Of the positive cases, three were treated with a combination of berenil (3.5 mg/kg) and oxytetracycline (5 mg/kg) or doxycycline (5 mg/kg); 44/63 were treated with only oxytetracycline of 5% or 5.5% concentration. Out of the 44/63 cases treated with oxytetracycline (5% or 5.5%) alone, a mortality of seven was recorded.

One-way analysis of variance showed a value of 0.34, revealing no significant findings between the PCV values in both cases. However, the PCV in the positive case on the basis of the Pearson's correlation table (Table 7) showed a significant

positive correlation (0.005) with temperatures of affected dogs and a negative nonsignificance (−0.462) with heart rate.

Table 7 shows that the rectal temperatures of the dogs infected with *Babesia* spp. were positively correlated to the PCV ($P=0.005$, $r=0.358$) and negatively correlated to the WBC counts ($P=0.003$, $r=-0.390$). However, the PCV was negatively correlated to WBC counts ($P=0.009$, $r=-0.342$). Total plasma protein level was significantly and positively correlated to plasma albumin ($P=0.000$, $r=0.763$) and globulin ($P=0.000$, $r=0.775$) levels but negatively correlated to total bilirubin level ($P=0.000$, $r=-0.550$). Also, the plasma globulin level was significantly and negatively correlated to total bilirubin level ($P=0.000$, $r=-0.600$).

Discussion

From this study, clinical findings such as tick infestation, fever, and anorexia were most commonly observed in both *Babesia*-positive and negative dogs (tick infestation: 90.1% [35/40]; fever: 46.4% [18/40]; and anorexia 36% [14/40]). This result revealed that the aforementioned clinical findings of canine babesiosis are nonspecific, as previously observed among Egyptian dogs.¹¹ However, an observable difference was seen with splenomegaly and jaundice between the cases (splenomegaly: positive case 19.6% [12/63], negative case 20.6% [8/40]; jaundice: positive case 9.8% [6/63], negative case 5.2% [2/40]). Among the 14 breeds of dogs present, the most affected were the foreign and smaller breeds, while moderate susceptibility was observed among the Alsatian breeds. Ogo et al reported that although breed susceptibility has not been established for canine *Babesia* infection,⁵ a 50% to 55% seropositivity of

Table 6 Leukocytic changes within breeds positive to *Babesia* spp.

Breed	Normal leukocyte count	Leukocytosis	Leukopenia	Total
Alsatian	18	8	6	32
Dalmatian	–	1	–	1
Boerboel	2	2	1	5
Pit Bull Terrier	1	–	–	1
Maltese	1	–	–	1
Caucasian	4	–	1	5
Samoyed	1	–	–	1
Terrier Cross	1	–	–	1
Rottweiler	7	3	1	11
Bullmastiff	–	–	1	1
Crossbreed	3	–	–	3
Doberman	1	–	–	1
Total	39	14	10	63

Table 7 Pearson's correlation results of measured parameters

Measured parameters	r or P	Temp	HR	PCV	WBC	TB	TPP	Alb	Glob
Temp	r	1	−0.246	0.358**	−0.387**	0.120	0.046	0.156	−0.112
	P	0	0.059	0.005	0.003	0.384	0.741	0.257	0.419
HR	r	−0.246	1	−0.098	−0.050	0.089	−0.115	−0.074	−0.084
	P	0.059	0	0.462	0.712	0.521	0.410	0.594	0.548
PCV	r	0.358**	−0.098	1	−0.343**	0.097	.120	−0.011	0.161
	P	0.005	0.462	0	0.009	0.486	0.393	0.937	0.251
WBC	r	−0.387**	−0.050	−0.343**	1	−0.038	0.028	−0.035	0.088
	P	0.003	0.712	0.009	0	0.789	0.847	0.805	0.539
TB	r	0.120	0.089	0.097	−0.038	1	−0.552**	−0.254	−0.601**
	P	0.384	0.521	0.486	0.789	0	0.000	0.061	0.000
TPP	r	0.046	−0.115	0.120	0.028	−0.552**	1	0.765**	0.774**
	P	0.741	0.410	0.393	0.847	0.000	0	0.000	0.000
Alb	r	0.156	−0.074	−0.011	−0.035	−0.254	0.765**	1	0.210
	P	0.257	0.594	0.937	0.805	0.061	0.000	0	0.127
Glob	r	−0.112	−0.084	0.161	0.088	−0.601**	0.774**	0.210	1
	P	0.419	0.548	0.251	0.539	0.000	0.000	0.127	0

Note: **Correlation is significant at the 0.01 level (two-tailed).

Abbreviations: Temp, temperature; HR, heart rate; PCV, packed cell volume; WBC, white blood cell; TB, total bilirubin; TPP, total plasma protein; Alb, albumin; Glob, globulin.

Babesia canis vogeli infection in Greyhounds was observed,^{12–14} while American Staffordshire Terrier and American Pit Bull Terrier breeds had 15% to 93% seropositivity.^{14–16} Furthermore, a study conducted in 2009 showed that prevalence was highest in crossbreeds and lowest in Doberman.¹⁷ In this study, Alsatian and Rottweiler breeds were overpresented as 58.8% and 55%, respectively, while only one indigenous breed was presented and positive for *Babesia* infection.

For the sexes, outcome observed for positive males and females was 65% (31/48) and 55% (30/55), respectively, with an 1.20 odds ratio of male dogs having the disease more than females. Bashir et al stated that the odds of canine babesiosis were 2.63 times higher in male than female dogs,¹⁷ which is similar to this study. Furthermore, it has been suggested that male dogs were more prone to the disease than the females due to bite wounds and blood transmission during fighting among male dogs.^{11,17} Inasmuch as the fact that blood transmission of canine babesiosis has not been established in Nigeria, there is a probability that this could occur. In addition, *Babesia*-positive cases were observed in 65% males and 55% females.

For susceptibility among ages, 57.3% (39/68) and 62.8% (22/35) were positive among adults and puppies, respectively, while the odds ratio was 0.79; $P > 0.05$. This result agrees with an earlier report which stated that age does not have any influence on the animals' susceptibility to the disease.⁸ In addition, nonspecific or innate factors (genetics or age) possessed by the host act as natural protective elements;^{9,10} that is, the immunity of an individual could be primed to withstand or be susceptible to a disease challenge at a particular point in time.

Other clinical parameters, such as pulse and heart rates, in the positive case gave significantly higher values compared to the negative case, and this could be due to the cardiac pathology reported in cases of canine babesiosis; also the heart rate of dogs that were positive to *Babesia* parasites was significantly higher than ($P < 0.05$) those that were negative for the parasite. Increased heart rate was reported as a common finding in uncomplicated canine babesiosis.^{18–22}

Based on the study, in the positive cases, a higher number of dogs were observed to have severe anemia (31.7%) and normal packed cell volume (33.3%) when compared to those with moderate or mild anemia. While the presence of anemia in this study supports the reports by Lobetti² and Zygnier et al,⁴ variable hematology was observed in both positive and negative cases and the presenting state of the dog could account for this result. Also, factors such as nutrition or the presence of unidentified causes during investigation using thin blood smears could account for this result.

The Pearson's correlation shown in Table 7 reveals that the temperature of dogs positive for *Babesia* spp. was

significantly correlated with the PCV ($P < 0.05$). This agrees with reports stating significant correlation between body temperature and heart rate, red blood cell count, PCV, and platelet count.²² In addition to this study, the body temperature of dogs positive to the *Babesia* parasite showed negative correlation with WBC count from positive dogs ($P < 0.05$).

Conclusion

Canine babesiosis with a prevalence of 61% appears to be a common cause of illness among the dog population in Nigeria. Commonly determined physiological parameters such as temperature and respiratory rates did not change significantly between *Babesia*-positive and *Babesia*-negative dogs, which suggests that these parameters are not sensitive indicators for monitoring canine babesiosis, however, pulse and heart rates appear to be positive and sensitive indicators. Hematological tests that determine PCV and WBC counts are often performed in the diagnosis of canine babesiosis. In this study, both PCV and WBC counts are not sensitive indicators for monitoring the progression and prognosis of babesiosis in dogs.

Disclosure

The authors report no conflicts of interest in this work.

References

- Di Cicco MF, Birkenheuer AJ. Canine babesiosis. Infectious disease/parasitology. *Consultants on Call/NAVCC Clinician's Brief*; 2012.
- Lobetti R. Hematological changes associated with tick-borne diseases (infectious/parasitic diseases). In: World Small Animal Veterinary Association World Congress Proceedings. October 6–9, 2004, for 29th WSAVA Congress Proceedings.
- Jacobson LS, Lobetti R. Glucose, lactate and pyruvate concentrations in dogs with babesiosis. *Am J Vet Res*. 2005;66(2):244–250.
- Zygnier W, Gojska-Zygnier O, Długosz E, Wedrychowicz H. Liver enzyme activity in dogs infected with *Babesia canis*. *Bull Vet Inst Pulawy*. 2011;55(3):423–427.
- Ogo NI, Lawal AI, Okubanjo OO, Kamani J, Ajayi OO. Current status of canine babesiosis and the situation in Nigeria: a review. *Niger Vet J*. 2011;32(2):69–78.
- Sasaki M, Omobowale O, Tozuka M, et al. Molecular survey of *Babesia canis* in dogs in Nigeria. *J Vet Med Sci*. 2007;69(11):1191–1193.
- Vannier E, Gewurz BE, Krause PJ. Human babesiosis. *Infect Dis Clin North Am*. 2008;22(3):469–488, viii–ix.
- Martinod S, Laurent N, Moreau Y. Resistance and immunity of dogs against *Babesia canis* in an endemic area. *J Vet Parasitol*. 1986;19(3–4):245–254.
- Johnston LAY, Leach G, Jones PN. The duration of latent infection and functional immunity in Droughmaster and Hereford cattle following natural infection with *Babesia argentina* and *Babesia bigemina*. *Aust Vet J*. 1978;56(1):14–18.
- Levy MG, Clabaugh G, Ristic M. Age resistance in bovine babesiosis: role of blood factors in resistance to *Babesia bovis*. *Infect Immun*. 1982;37(3):1127–1131.
- Salem NY, Farag HS. Clinical, hematologic and molecular findings in naturally occurring *Babesia canis vogeli* in Egyptian dogs. *Vet Med Int*. 2014;2014(2014).
- Breitschwerdt EB, Malone JB, MacWilliams P, Levy MG, Quails CW, Prudich MJ. Babesiosis in the Greyhound. *J Am Vet Med Assoc*. 1983;182(9):978–982.

13. Yamane I, Gardener I, Ryan C, Levy M, Urrico J, Conrad P. Survey of *Babesia canis*, *Babesia gibsoni* and *Ehrlichia canis* in pound dogs in California, USA. *Prev Vet Med*. 1994;18(4):293–304.
14. Birkenheuer AJ, Correa MT, Levy MG, Breitschwerdt EB. Geographical distribution of babesiosis among dogs in the United States and association with dog bites: 150 cases (2000–2003). *J Am Vet Med Assoc*. 2005;227(6):942–947.
15. Macintire DK, Bourdreaux MX, West GD, Bournec C, Wright JC, Conrad PA. *Babesia gibsoni* infection among dogs in the Southeastern United States. *J Am Vet Med Assoc*. 2002;220(3):325–329.
16. Birkenheuer AJ, Levy MG, Stebbins M, Poore M, Breitschwerdt EB. Serosurvey of anti-*Babesia* antibodies in stray dogs and American pit bull terriers and American Staffordshire terriers from North Carolina. *J Am Anim Hosp Assoc*. 2003;39(6):551–557.
17. Bashir IN, Chaudhry ZI, Ahmed S, Saeed MA. Epidemiological and vector identification studies on canine babesiosis. *Pakistan Vet J*. 2009; 29(2):51–54.
18. Lobetti GR. Canine babesiosis. In: Day MJ, Littelwood JD, Mackin A, editors. *Manual of Canine and Feline Haematology and Transfusion Medicine*. Cheltenham: British Small Animal Veterinary Association; 2000:85–91.
19. Vaughan-Scott T. *Serum Concentration of Tumor Necrosis Factor in Dogs Naturally Infected with Babesia canis and Its Relation to Severity of Disease*. [master's thesis]. Cape Town: University of Pretoria; 2001.
20. Böhm M. *The Comparative Assessment of Capillary and Venous Babesia rossi Parasitaemias on Thin Blood Smears and Their Association with Disease Manifestation* [master's thesis]. Pretoria: University of Pretoria; 2006.
21. Ayoob AL, Hackner SG, Prittie J. Clinical management of canine babesiosis. *J Vet Emerg Crit Care*. 2010;20(1):77–89.
22. Torbica G, Bedrica L, Samardzija M, et al. Canine babesiosis treatment with three different medicines. *Acta Vet (Beograd)*. 2013;63(2–3): 279–290.

Veterinary Medicine: Research and Reports

Publish your work in this journal

Veterinary Medicine: Research and Reports is an international, peer-reviewed, open access journal publishing original research, case reports, editorials, reviews and commentaries on all areas of veterinary medicine. The manuscript management system is completely online and includes a very quick and fair peer-review system.

Submit your manuscript here: <http://www.dovepress.com/veterinary-medicine-research-and-reports-journal>

Dovepress

Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.