The association between vitamin D receptor gene polymorphisms (*TaqI* and *FokI*), Type 2 diabetes, and micro-/macrovascular complications in postmenopausal women

Juliana Maia¹
 Andreia Soares da Silva²
 Rodrigo Feliciano do Carmo³
 Taciana Furtado de Mendonça²
 Luiz Henrique Maciel Griz¹
 Patricia Moura²
 Francisco Bandeira¹

¹Division of Endocrinology and Diabetes, Agamenon Magalhães Hospital, ²Institute of Biological Sciences, University of Pernambuco Medical School, Recife, ³Federal University of São Francisco Valley, Petrolina, Pernambuco, Brazil

Introduction: Since there is evidence of the action of vitamin D as a modulator of insulin release and atherosclerosis, it may well be that the vitamin D receptor polymorphisms are associated with diabetes and its chronic complications.

Aims: To examine the associations between vitamin D receptor polymorphisms (*FokI* and *TaqI*) and Type 2 diabetes (T2DM) and its associated chronic complications in postmenopausal women.

Methods: This cross-sectional study analyzed 100 postmenopausal women with T2DM (mean age 65.7±7.18 years) and 100 postmenopausal women without diabetes in the control group (mean age 65.1±9.18 years; $P=0.1608$). We evaluated clinical and metabolic parameters and analyzed *TaqI* and *FokI* polymorphisms.

Results: There were no significant differences in genotype and allele frequencies between patients and controls in either of the polymorphisms studied. In the group of patients with diabetes, there were no significant differences in either polymorphism in relation to stroke, retinopathy, nephropathy, or neuropathy. However, in patients with T2DM and coronary artery disease, *f* genotype ($P=0.0361$) and the combination of *Ff* + *ff* genotypes were observed less frequently ($P=0.0462$).

Conclusion: This study suggests the potential protective factor of *FokI* polymorphism for coronary artery disease in postmenopausal women with T2DM in the recessive model.

Keywords: Type 2 diabetes, VDR, polymorphism, vitamin D

Introduction

Although vitamin D is recognized mainly for its effects on bone metabolism, it is also known that it regulates multiple aspects of cellular differentiation and replication in many target tissues, including endocrine pancreas.^{1,2} Some studies have suggested that vitamin D plays an important role in the mechanism of insulin release and in the maintenance of glucose tolerance and have also shown the association between vitamin D deficiency and Type 2 diabetes (T2DM).^{3,4}

Cardiovascular disease (CVD) is the leading cause of mortality and morbidity in patients with diabetes and the largest contributor to the cost of diabetes.⁵ Various studies have also suggested the association between reduced 25-hydroxyvitamin D (25OHD) concentrations and an increased risk of atherosclerosis, endothelial dysfunction, and coronary artery disease (CAD).⁶⁻⁹

Correspondence: Juliana Maia
 Division of Endocrinology and Diabetes,
 Agamenon Magalhães Hospital, University
 of Pernambuco Medical School, Rua
 Marechal Rondon, 120, ap 1602, Recife,
 Pernambuco 52061-050, Brazil
 Email maia.juliana@gmail.com

Recent developments have led public health authorities to take an increasing interest in vitamin D research, since vitamin D deficiency is highly prevalent in the general population,¹⁰ even in those living in areas with abundant sunlight.^{11,12}

It is well known that metabolic actions related to vitamin D, on target cells, are mediated by the binding of 1,25-dihydroxyvitamin D to its specific cytosolic/nuclear receptor, which has been identified in many tissues.¹³ The vitamin D receptor (VDR) is a member of the steroid/thyroid hormone receptor family and functions as a transcriptional activator of many genes.¹⁴ The VDR gene is located on chromosome 12q13.11 in humans and consists of eleven exons.¹⁵ Several polymorphisms of the VDR gene have been identified, but four single-nucleotide polymorphisms (SNPs) of this gene are the ones that have been most studied, namely, *BsmI* (rs1544410), *ApaI* (rs7975232), *TaqI* (T>C; rs731236), and *FokI* (C>T; rs2228570, formerly known as rs10735810).

The *FokI* polymorphism occurs in the start codon of VDR and its polymorphic form (f) leads to the translation of a longer VDR protein variant (427 amino acid), and some in vitro studies suggest that it functions less effectively than the shorter VDR protein (424 amino acid) encoded by the F allele.^{15–17} The functional activity of this SNP is yet to be fully understood, since other in vitro experiments have failed to show this difference.¹⁸ *TaqI* polymorphism results from a mutation in a transcriber region located in exon 9 of the VDR gene. However, there are no studies demonstrating its functionality.¹⁵

Several studies in recent years have attempted to elucidate the possible relationship between polymorphisms of the VDR gene and diabetes and CVD, but the results are still controversial and substantial differences have been observed between SNPs and the ethnic groups analyzed.^{19–26} This further emphasizes the need to conduct studies to analyze VDR polymorphisms in different ethnic populations.

The aims of this study were, therefore, to access the genotypic distribution of *FokI* and *TaqI* polymorphisms of VDR in a population of postmenopausal women with T2DM and in a control group, and to evaluate whether these two SNPs of VDR (*FokI* and *TaqI*) are associated with micro- and macrovascular complications.

Methods

Study design

This cross-sectional study was carried out between September 2013 and August 2014 in the city of Recife (Brazil) and comprised 100 female patients with physiological menopause

and diagnosis of T2DM, according to the American Diabetes Association criteria.⁵ A total of 100 postmenopausal women without diabetes were enrolled as the control group. The subjects of the study were recruited from the community. The study was approved by the Ethics in Research Committee of Agamenon Magalhães Hospital (CEP-HAM, Recife, Brazil). All patients gave their written informed consent to take part in this study.

The exclusion criteria were as follows: patients with Type 1 diabetes; end-stage renal disease; use of glucocorticoids; use of vitamin D replacement therapy; use of hormone replacement therapy; primary hyperparathyroidism; Paget's disease of bone; malabsorption syndrome; and/or active neoplasia.

The patients were interviewed by a physician, answered a standard questionnaire, and underwent a complete physical examination, including measurement of height, weight, waist circumference, and blood pressure. Skin phototypes were classified in six types, according to Fitzpatrick's classification (Type 1 as the lightest and Type 6 as the darkest).²⁷ Blood samples were obtained from all participants, after overnight fasting. Low-density lipoprotein cholesterol, high-density lipoprotein cholesterol, triglyceride levels, fasting blood glucose, and glycated hemoglobin (HbA1c) were evaluated. Biochemical tests were processed in a dry chemistry autoanalyzer (Vitro System, Johnson & Johnson, New Brunswick, NJ, USA) and HbA1c concentrations were determined by immunoturbidimetric assay (Cobas 501, Roche Diagnostic, Rotkreuz, Switzerland).

Serum 25OHD levels were measured by electrochemiluminescent competitive immunoassay (Diasorin-Liaison, Saluggia, Italy), with the interassay and intra-assay coefficients of variation of 8%–15% and 8%–13%, respectively, with a minimum limit of detection of 2 ng/mL.²⁸

DNA extraction and genotyping

The DNA was extracted from whole blood samples (ethylenediaminetetraacetic acid) using the QIAamp Mini Spin Columns kit (QIAGEN, Venlo, the Netherlands) in accordance with the manufacturer's instructions. For detection of polymorphic variable, the real-time polymerase chain reaction method in the TAQMAN® (Thermo Fisher Scientific, Waltham, MA, USA) system was used.

We genotyped the following SNP of the VDR gene: *FokI* (rs10735810/ID: C__12060045_20) and *TaqI* (rs731236/ID: C__2404008_10) according to the manufacturer's instructions (<http://www.thermofisher.com/order/genome-database>). *FokI* polymorphism results in different translation

initiation sites due to the replacement of cytosine (C) by thymine (T) and *TaqI* polymorphism is constituted by the silent replacement of thymine (T) by cytosine (C).

Statistical evaluation

Data for categorical variables are presented as numbers (percentages), and data for continuous variables are presented as mean values (standard deviations). The differences in genotype and allele frequencies were analyzed by Pearson's chi-square test and Fisher's exact test. Differences were considered statistically significant for *P*-values of less than 0.05. Odds ratio (OR) with a 95% confidence interval (CI) was determined to estimate the association between variables. The GraphPad Prism program (version 6.0) for Windows (GraphPad Software, San Diego, CA, USA) was used for the analysis. To determine the Hardy–Weinberg equilibrium in the population, the ARLEQUIN program was used (version 3.11, Bern, Swiss).

Results

The clinical characteristics of the patients and control group are summarized in Table 1. Age distribution between T2DM patients (65.7±7.18 years) and controls (65.1±9.18 years) was not significantly different (*P*=0.1608). The frequency of hypertension and body mass index >25 kg/m² were higher in the T2DM patients than in the controls (*P*=0.0001 and *P*=0.0355, respectively). Mean waist circumference, serum triglyceride levels, and HbA1c were elevated in patients compared with controls. There were no significant differences in skin phototype between patients and controls. Serum 25OHD levels below 30 ng/mL were found to be significantly more prevalent in T2DM patients than in the control group (74% and 48%, respectively; *P*=0.0001). Using cutoffs for serum 25OHD levels below 20 ng/mL, the prevalence of vitamin D deficiency was 12% in patients with diabetes and 7% in the control group (*P*=0.3350).

There were no significant differences in genotype and allele frequency distribution between T2DM patients and controls in either the *TaqI* or *FokI* analysis (Tables 2 and 3). The prevalence of *TaqI* genotypes was 45% TT, 46% Tt, 9% tt in T2DM patients, and 49% TT, 39% Tt, and 12% tt in the control group (Table 2). The frequency of the T allele in T2DM patients was 68% and that of the t allele 32%, while in the control group, it was 68.5% and 31.5%, respectively (*P*=1.0000; Table 2). The prevalence of *FokI* was 46% FF, 42% Ff, and 12% ff in T2DM patients and 38% FF, 47% Ff, and 15% ff in the control group. The frequency of the F allele in T2DM patients was 67% and that of the f allele 33%, while

Table 1 Characteristics of study participants

Characteristics	Cases (n=100)	Control (n=100)	P-value
Age			
Mean (±SD)	65.7 (±7.18)	65.1 (±9.18)	<i>P</i> =0.16
Hypertension (n)			
Yes	86	60	
No	14	40	<i>P</i> ^b =0.0001*
BMI (kg/m ²) (n)			
Normal (<25)	19	33	
Overweight/obese (>25)	81	67	<i>P</i> ^b =0.0355*
WC (cm)			
Mean (±SD)	99 (±12)	89 (±14)	<i>P</i> ^a <0.0001*
HbA1c (%)			
Mean (±SD)	7.8 (±1.7)	5.5 (±0.3)	<i>P</i> ^a <0.0001*
Triglycerides (mg/dL)			
Mean (±SD)	160 (±68)	135 (±62)	<i>P</i> ^a =0.0200*
Skin phototype(n)			
I + II	6	3	
III + IV	70	58	<i>P</i> ^b =0.73
V + VI	24	39	<i>P</i> ^b =0.15
25OHD (ng/mL) (n)			
Deficiency/insufficiency (<30 ng/mL)	74	48	
Normal (>30 ng/mL)	26	52	<i>P</i> ^b =0.0001*

Notes: *Statistically significant difference (*P*<0.05); ^aStudent's *t*-test; ^bFisher's exact test. **Abbreviations:** SD, standard deviation; BMI, body mass index; WC, waist circumference; HbA1c, glycated hemoglobin; 25OHD, 25-hydroxyvitamin D.

Table 2 Distribution of VDR (*TaqI*) polymorphism genotypes and allele frequencies between T2DM patients and control groups

VDR	T2DM patients n=100	Control n=100	P-value ^a	OR	95% CI
Genotype					
AA (TT)	45 (45%)	49 (49%)	—		
AG (Tt)	46 (46%)	39 (39%)	0.45	1.28	0.68–2.41
GG (tt)	9 (9%)	12 (12%)	0.81	0.82	0.28–2.33
AG + GG (Tt + tt)	55 (55%)	52 (52%)	0.67	1.17	0.65–2.13
Alleles					
A (T)	136 (68%)	137 (68.5%)	—		
G (t)	64 (32%)	63 (31.5%)	1.00	1.02	0.66–1.59

Notes: ^aChi-square test. "—" indicates no data.

Abbreviations: VDR, vitamin D receptor; T2DM, Type 2 diabetes; OR, odds ratio; CI, confidence interval.

in the control group, it was 61.5% and 38.5%, respectively (*P*=0.1256). The study subjects were in Hardy–Weinberg equilibrium for both *TaqI* and *FokI*.

In addition, we also compared the VDR (*TaqI* and *FokI*) genotypes and allele frequencies with the presence of macro- and microvascular complications of diabetes. There were no significant differences in relation to stroke, retinopathy, nephropathy, or neuropathy. No statistically significant differences were found in the *TaqI* genotype or allele frequency in the groups of patients with diabetes

Table 3 Distribution of VDR (*FokI*) polymorphism genotypes and allele frequencies between T2DM patients and control groups

VDR	T2DM patients n=100	Control n=100	P-value ^a	OR	95% CI
Genotype					
GG (FF)	46 (46%)	38 (38%)	–		
AG (Ff)	42 (42%)	47 (47%)	0.13	1.4	0.77–2.56
AA (ff)	12 (12%)	15 (15%)	0.17	1.5	0.63–3.62
AG + GG (Ff + ff)	54 (54%)	62 (62%)	0.13	1.39	0.79–2.44
Alleles					
G (F)	134 (67%)	123 (61.5%)	–		
A (f)	66 (33%)	77 (38.5%)	0.13	1.2	0.84–1.91

Notes: ^aChi-square test. “–” indicates no data.

Abbreviations: VDR, vitamin D receptor; T2DM, Type 2 diabetes; OR, odds ratio; CI, confidence interval.

and without a history of CAD (Table 4), but in the *FokI* analysis (Table 5), compared with genotype FF, genotype Ff was likely to be a protective factor for CAD (FF vs Ff, $P=0.0391$, OR 0.29, 95% CI 0.09–1.01). We also found a significant association between *FokI* polymorphism and CAD in the recessive model (FF vs Ff + ff, $P=0.0462$, OR 0.35, 95% CI 0.12–1.02).

Discussion

In our study, no significant differences were found in genotype and allele frequency distribution of VDR (*TaqI* and *FokI*) genes between postmenopausal Brazilian women with T2DM and those of the control group. However, in T2DM patients with a history of CAD, we observed a lower frequency of the Ff genotype ($P=0.0361$) and the combination of genotypes Ff + ff ($P=0.0462$).

The prevalence of VDR *TaqI* and *FokI* polymorphism genotypes was similar to data reported previously in another study that also analyzed Brazilian postmenopausal women.²⁹

Reports on the association between VDR gene polymorphisms and T2DM have been inconsistent. Our analysis did not find any evidence of an association between *TaqI* and *FokI* polymorphism and T2DM among Brazilian postmenopausal women. Two meta-analyses showed data similar to ours regarding the lack of evidence of any association of the *TaqI* polymorphism with T2DM.^{30,31} However, with regard to the *FokI* polymorphism, these meta-analyses found some statistical evidence that the f allele may be a susceptibility allele for T2DM in an Asian population.^{30,31} It is important to emphasize that this result may vary according to the study population because of substantial differences in VDR polymorphisms between races, which have already been demonstrated.¹⁵ In fact, a study in Polish Caucasians found a result similar to ours, with no evidence for an association of *FokI* polymorphism with T2DM.²²

Table 4 Distribution of VDR (*TaqI*) polymorphism genotypes and allele frequencies in T2DM patients according to the presence of CAD

VDR	Yes n=18	No n=82	P-value ^a
Genotype			
AA (TT)	8 (44%)	37 (45%)	
AG (Tt)	7 (39%)	39 (48%)	0.78
GG (tt)	3 (17%)	6 (7%)	0.37
AG + GG (Tt + tt)	10 (56%)	45 (55%)	1.00
Alleles n (%)			
A (T)	23 (64%)	113 (69%)	
G (t)	13 (36%)	51 (31%)	0.56

Note: ^aChi-square test.

Abbreviations: VDR, vitamin D receptor; T2DM, Type 2 diabetes; CAD, coronary artery disease.

Table 5 Distribution of VDR (*FokI*) polymorphism genotypes and allele frequencies in T2DM patients according to the presence of CAD

VDR	Yes n=18	No n=82	P-value ^a	OR	95% CI
Genotype					
GG (FF)	12 (67%)	34 (42%)	–		
AG (Ff)	4 (22%)	38 (46%)	0.0361*	0.29	0.09–1.01
AA (ff)	2 (11%)	10 (12%)	0.39	0.57	0.11–2.96
AG + AA (Ff + ff)	6 (33%)	48 (58%)	0.0462*	0.35	0.12–1.02
Alleles n (%)					
G (F)	28 (78%)	106 (65%)	–		
A (f)	8 (22%)	58 (35%)	0.09	0.52	0.22–1.22

Notes: *Statistically significant difference ($P<0.05$); ^aChi-square test. “–” indicates no data.

Abbreviations: VDR, vitamin D receptor; T2DM, Type 2 diabetes; OR, odds ratio; CI, confidence interval; CAD, coronary artery disease.

We likewise found no association between *TaqI* polymorphism and CAD among postmenopausal Brazilian women with T2DM. A prospective study found a significant association between the t allele of *TaqI* VDR and baseline cases of CAD in a French population.²⁴ On the other hand, in a study of the Egyptian population, the *TaqI* showed no significant association with CAD.³² As it has been suggested that the *TaqI* polymorphism is a silent mutation, it results in no amino acid sequence change.¹⁵ This evidence may partly explain the result of this study.

The analysis of *FokI* SNPs of VDR, in our study, showed that the Ff genotype is probably a protective factor for CAD in postmenopausal women with T2DM; moreover, when we compared a recessive model for analysis (Ff + ff), there was a significant association with CAD.

Accumulating evidence suggests that low serum levels of vitamin D are associated with an increased risk of

atherosclerosis, endothelial dysfunction, and CVD.^{6,7,33,34} The epigenetic effects of vitamin D include modulation of smooth muscle cell proliferation, vascular calcification, inflammation, and thrombosis.^{35–37} In addition, vitamin D is a negative regulator of renin–angiotensin.^{38–40} VDR polymorphism is thus a potential candidate for coronary atherosclerosis.

Ortlepp et al²⁰ reported no association between *BsmI* polymorphism of VDR and coronary angiographic phenotypes in a large study population. A cohort from a multinational prospective study found that the haplotype comprising the minor allele of *BsmI*, the major allele of *ApaI*, and the minor allele of *TaqI* of VDR was associated with an increased risk of CAD in T2DM patients.²⁴ Our study did not evaluate the *ApaI* and *BsmI* polymorphisms; it did, however, analyze the *FokI* SNP, which is considered to be an independent marker of the VDR gene because it is not in linkage disequilibrium with other VDR SNPs, and there is speculation on its possible relationship with the functionality of the VDR protein.¹⁵

With regard to the association of genotype and allele frequencies of the *FokI* polymorphism with CAD, a Chinese study observed no significant evidence.⁴¹ Another study in a Caucasian population found no significant differences in anthropometric and biochemical features of CVD between genotypes of *FokI* in female and male groups,⁴² but it is difficult to compare those findings with ours because of differences in the study population. Both studies analyzed men and women in general, while our population is specific to postmenopausal women with T2DM.

Our findings cannot be related to the VDR functionality of *FokI* SNP, since the studies on functional differences in the VDR protein encoded by the polymorphic form are conflicting, suggesting the need for further studies on this topic.

Vaidya et al⁴³ observed that both higher serum 25OHD levels and the f allele at *FokI* were independently associated with lower plasma renin activity in a large population of hypertensives.⁴³ As plasma renin activity is associated with CVD, this finding is consistent with our finding of an association between the recessive model of *FokI* (Ff + ff) and CAD.

Our data therefore suggest that the association observed between the recessive model of *FokI* polymorphism and CAD in postmenopausal women with T2DM as a protective factor may be mediated by modulation of the renin–angiotensin system. However, additional studies with a larger sample size across various ethnic groups are needed to fully determine such an association and clarify the mechanism involved.

Some potential limitations should be considered in this study, such as the small size of the study population, the

cross-sectional design of the study, and the fact that patients did not undergo diagnostic angiography, and so there is a possibility that patients with asymptomatic CAD were overlooked.

Conclusion

Our study suggests that the *FokI* polymorphism in the VDR gene is a protective factor for CAD in postmenopausal women with T2DM in the recessive model. Our data evidenced no association between the polymorphisms analyzed (*FokI* and *TaqI*) and T2DM and other chronic complications of diabetes.

Disclosure

The authors report no conflicts of interests in this work.

References

- Walters MR. Newly identified actions of the vitamin D endocrine system. *Endocr Rev.* 1992;13(4):719–764.
- Lee S, Clark SA, Gill RK, Christakos S. 1,25-Dihydroxyvitamin D3 and pancreatic beta-cell function: vitamin D receptors, gene expression, and insulin secretion. *Endocrinology.* 1994;134(4):1602–1610.
- Mattila C, Knekt P, Männistö S, et al. Serum 25-hydroxyvitamin D concentration and subsequent risk of type 2 diabetes. *Diabetes Care.* 2007;30(10):2569–2570.
- Song Y, Wang L, Pittas AG, et al. Blood 25-hydroxy vitamin D levels and incident type 2 diabetes: a meta-analysis of prospective studies. *Diabetes Care.* 2013;36(5):1422–1428.
- American Diabetes Association. Standards of medical care in diabetes – 2014. *Diabetes Care.* 2014;37 (Suppl 1):S14–S80.
- Brewer LC, Michos ED, Reis JP. Vitamin D in atherosclerosis, vascular disease, and endothelial function. *Curr Drug Targets.* 2011;12(1):54–60.
- Grandi NC, Breitling LP, Brenner H. Vitamin D and cardiovascular disease: systematic review and meta-analysis of prospective studies. *Prev Med.* 2010;51(3–4):228–233.
- Kim DH, Sabour S, Sagar UN, Adams S, Whellan DJ. Prevalence of hypovitaminosis D in cardiovascular diseases (from the National Health and Nutrition Examination Survey 2001 to 2004). *Am J Cardiol.* 2008;102(11):1540–1544.
- Wang TJ, Pencina MJ, Booth SL, et al. Vitamin D deficiency and risk of cardiovascular disease. *Circulation.* 2008;117(4):503–511.
- Mithal A, Wahl DA, Bonjour JP, et al. Global vitamin D status and determinants of hypovitaminosis D. *Osteoporos Int.* 2009;20(11):1807–1820.
- Bandeira F, Griz L, Freese E, et al. Vitamin D deficiency and its relationship with bone mineral density among postmenopausal women living in the tropics. *Arq Bras Endocrinol Metabol.* 2010;54(2):227–232.
- Cabral MA, Borges CN, Maia JM, Aires CA, Bandeira F. Prevalence of vitamin D deficiency during the summer and its relationship with sun exposure and skin phototype in elderly men living in the tropics. *Clin Interv Aging.* 2013;8:1347–1351.
- Bouillon R, Carmeliet G, Verlinden L, et al. Vitamin D and human health: lessons from vitamin D receptor null mice. *Endocr Rev.* 2008;29(6):726–776.
- Mangelsdorf DJ, Thummel C, Beato M, et al. The nuclear receptor superfamily: the second decade. *Cell.* 1995;83(6):835–839.
- Uitterlinden AG, Fang Y, Van Meurs JB, Pols HA, Van Leeuwen JP. Genetics and biology of vitamin D receptor polymorphisms. *Gene.* 2004;338(2):143–156.
- Arai H, Miyamoto K, Taketani Y, et al. A vitamin D receptor gene polymorphism in the translation initiation codon: effect on protein activity and relation to bone mineral density in Japanese women. *J Bone Miner Res.* 1997;12(6):915–921.

17. Nejentsev S, Godfrey L, Snook H, et al. Comparative high-resolution analysis of linkage disequilibrium and tag single nucleotide polymorphisms between populations in the vitamin D receptor gene. *Hum Mol Genet.* 2004;13(15):1633–1639.
18. Gross C, Krishnan AV, Malloy PJ, Eccleshall TR, Zhao XY, Feldman D. The vitamin D receptor gene start codon polymorphism: a functional analysis of FokI variants. *J Bone Miner Res.* 1998;13(11):1691–1699.
19. Bid HK, Konwar R, Aggarwal CG, et al. Vitamin D receptor (FokI, BsmI and TaqI) gene polymorphisms and type 2 diabetes mellitus: a North Indian study. *Indian J Med Sci.* 2009;63(5):187–194.
20. Ortlepp JR, Lauscher J, Hoffmann R, Hanrath P, Joost HG. The VDR gene variant is associated with the prevalence of Type 2 diabetes mellitus and coronary artery disease. *Diabet Med.* 2001;18:842–845.
21. Dilmeç F, Uzer E, Akkafa F, Kose E, van Kuilenburg AB. Detection of VDR gene ApaI and TaqI polymorphisms in patients with type 2 diabetes mellitus using PCR-RFLP method in a Turkish population. *J Diabetes Complications.* 2010;24(3):186–191.
22. Malecki MT, Frey J, Moczulski D, Klupa T, Kozek E, Sieradzki J. Vitamin D receptor gene polymorphisms and association with type 2 diabetes mellitus in a Polish population. *Exp Clin Endocrinol Diabetes.* 2003;111(8):505–509.
23. Ortlepp JR, von Korff A, Hanrath P, Zerres K, Hoffmann R. Vitamin D receptor gene polymorphism BsmI is not associated with the prevalence and severity of CAD in a large-scale angiographic cohort of 3441 patients. *Eur J Clin Invest.* 2003;33(2):106–109.
24. Ferrarezi DA, Bellili-Muñoz N, Dubois-Laforgue D, et al. Allelic variations of the vitamin D receptor (VDR) gene are associated with increased risk of coronary artery disease in type 2 diabetics: the DIABHYCAR prospective study. *Diabetes Metab.* 2013;39(3):263–270.
25. Jia J, Ding H, Yang K, et al. Vitamin D receptor genetic polymorphism is significantly associated with risk of type 2 diabetes mellitus in Chinese Han population. *Arch Med Res.* 2015;46(7):572–579.
26. Khodaeian M, Enayati S, Tabatabaei-Malazy O, Amoli MM. Association between genetic variants and diabetes mellitus in Iranian populations: a systematic review of observational studies. *J Diabetes Res.* 2015;2015:585917.
27. Fitzpatrick TB. The validity and practicality of sun-reactive skin types I through VI. *Arch Dermatol.* 1988;124(6):869–871.
28. Ersfeld DL, Rao DS, Body JJ, et al. Analytical and clinical validation of the 25 OH vitamin D assay for the LIASON automated analyzer. *Clin Biochem.* 2004;37(10):867–874.
29. Gentil P, de Lima Lins TC, Lima RM, et al. Vitamin-d-receptor genotypes and bone-mineral density in postmenopausal women: interaction with physical activity. *J Aging Phys Act.* 2009;17(1):31–45.
30. Wang Q, Xi B, Reilly KH, Liu M, Fu M. Quantitative assessment of the associations between four polymorphisms (FokI, ApaI, BsmI, TaqI) of vitamin D receptor gene and risk of diabetes mellitus. *Mol Biol Rep.* 2012;39(10):9405–9414.
31. Li L, Wu B, Liu JY, Yang LB. Vitamin D receptor gene polymorphisms and type 2 diabetes: a meta-analysis. *Arch Med Res.* 2013;44(3):235–241.
32. Abu el Maaty MA, Hassanein SI, Sleem HM, Gad MZ. Vitamin D receptor gene polymorphisms (TaqI and ApaI) in relation to 25-hydroxyvitamin D levels and coronary artery disease incidence. *J Recept Signal Transduct Res.* 2014;1–5.
33. Pilz S, Tomaschitz A, März W, et al. Vitamin D, cardiovascular disease and mortality. *Clin Endocrinol (Oxf).* 2011;75(5):575–584.
34. Anderson JL, May HT, Horne BD, et al. Relation of vitamin D deficiency to cardiovascular risk factors, disease status, and incident events in a general healthcare population. *Am J Cardiol.* 2010;106(7):963–968.
35. Zehnder D, Bland R, Chana RS, et al. Synthesis of 1,25 dihydroxyvitamin D(3) by human endothelial cells is regulated by inflammatory cytokines: a novel autocrine determinant of vascular cell adhesion. *J Am Soc Nephrol.* 2002;13:621–629.
36. Aihara K, Azuma H, Akaike M, et al. Disruption of nuclear vitamin D receptor gene causes enhanced thrombogenicity in mice. *J Biol Chem.* 2004;279:35798–802.
37. Zittermann A, Schleithoff SS, Koerfer R. Putting cardiovascular disease and vitamin D insufficiency into perspective. *Br J Nutr.* 2005;94(4):483–492.
38. Li YC, Kong J, Wei M, Chen ZF, Liu SQ, Cao LP. 1,25-Dihydroxyvitamin D(3) is a negative endocrine regulator of the renin-angiotensin system. *J Clin Invest.* 2002;110(2):229–238.
39. Tomaschitz A, Pilz S, Ritz E, et al. Independent association between 1,25-dihydroxyvitamin D, 25-hydroxyvitamin D and the renin-angiotensin system: the Ludwigshafen Risk and Cardiovascular Health (LURIC) Study. *Clin Chim Acta.* 2010;411(17–18):13541360.
40. Vaidya A, Forman JP, Hopkins PN, Seely EW, Williams JS. 25-hydroxyvitamin D is associated with plasma renin activity and the pressor response to dietary sodium intake in Caucasians. *J Renin Angiotensin Aldosterone Syst.* 2011;12(3):311–319.
41. Pan XM, Li DR, Yang L, et al. No association between vitamin D receptor polymorphisms and coronary artery disease in a Chinese population. *DNA Cell Biol.* 2009;28(10):521–525.
42. Laczmański L, Milewicz A, Lwow F, et al. Vitamin D receptor gene polymorphism and cardiovascular risk variables in elderly Polish subjects. *Gynecol Endocrinol.* 2013;29(3):268–272.
43. Vaidya A, Sun B, Forman JP, et al. The FokI vitamin D receptor gene polymorphism is associated with plasma renin activity in Caucasians. *Clin Endocrinol (Oxf).* 2011;74(6):783–790.

The Application of Clinical Genetics

Publish your work in this journal

The Application of Clinical Genetics is an international, peer-reviewed open access journal that welcomes laboratory and clinical findings in the field of human genetics. Specific topics include: Population genetics; Functional genetics; Natural history of genetic disease; Management of genetic disease; Mechanisms of genetic disease; Counselling and ethical

issues; Animal models; Pharmacogenetics; Prenatal diagnosis; Dysmorphology. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/the-application-of-clinical-genetics-journal>

Dovepress