Evaluating vancomycin susceptibility in *Staphylococcus aureus*

Marcelo J Mimica
Alessandra Navarini

Department of Pathology, Division of
Microbiology, Santa Casa de São Paulo
School of Medicine, São Paulo, Brazil

Dear editor

We read the report by Phillips et al¹ with great interest and would like to discuss it in comparison with our previous published data on the subject.^{2,3}

We have also studied a number of *Staphylococcus aureus* clinical isolates (n=125), comparing different vancomycin susceptibility tests, including microdilution, Etest® (bio-Mérieux, Marcy-l'Étoile, France), and brain heart infusion vancomycin screening plates.

We found only one isolate with reduced susceptibility with a minimum inhibitory concentration (MIC) =4 mg/L when tested with Etest and 2 mg/L when tested with microdilution.^{2,3} Our results showed a tendency of higher lethality when higher MICs were present, even within the susceptible range,³ as some previous studies have shown.^{4,5}

Concordant to Phillips et al¹ and other authors,^{6,7} we also reported a poor correlation between different tests. Comparing Etest and microdilution (approximating an Etest MIC value between two twofold dilutions up to the highest value), 58% of the isolates had similar MICs, whereas 38% had an MIC by Etest one dilution higher than microdilution. One isolate had an Etest MIC twofold higher and four isolates an Etest MIC onefold lower than microdilution.²

However, in our study, a brain heart infusion screening plate with 2.0 mg/L of vancomycin showed a sensitivity of 100% to detect isolates with an MIC ≥2.0 by Etest and 91% to detect an MIC ≥2.0 by microdilution, making this test an interesting option for initial screening of *S. aureus* isolates for reduced vancomycin susceptibility. Specificities were 63% and 38%, respectively, which would still make necessary the further testing with an MIC method, but in a much smaller number of isolates.² This approach would be suitable for a large number of laboratories throughout the world where the routine MIC testing of all *S. aureus* isolates is not feasible.

Disclosure

The authors report no conflicts of interest in this communication.

References

1. Phillips CJ, Wells NA, Martinello M, Smith S, Woodman RJ, Gordon DL. Optimizing the detection of methicillin-resistant *Staphylococcus aureus* with elevated vancomycin minimum inhibitory concentrations within the susceptible range. *Infect Drug Resist*. 2016;9:87–92.
2. Navarini A, Martino MD, Sasagawa SM, Massaia IF, Mimica MJ. Accuracy of a vancomycin brain heart infusion screening plate for the screening of *Staphylococcus aureus* isolates with increased vancomycin minimum inhibitory concentrations. *New Microbiol*. 2015;38(3):423–426.
3. Sulla F, Bussius DT, Acquesta F, Navarini A, Sasagawa SM, Mimica MJ. Vancomycin minimum inhibitory concentrations and lethality in *Staphylococcus aureus* bacteremia. *Germes*. 2015;5(2):39–43.

Correspondence: Marcelo J Mimica
Department of Pathology, Division of
Microbiology, Santa Casa de São Paulo
School of Medicine, Rua Doutor Cesário
Motta Júnior, 112, São Paulo
CEP 01221-020, SP, Brazil
Email marcelo.mimica@fmsantacasasp.edu.br

4. Aguado JM, San-Juan R, Lalueza A, Sanz F, Rodríguez-Otero J, Gómez-Gonzalez C, Chaves F. High vancomycin MIC and complicated methicillin-susceptible *Staphylococcus aureus* bacteremia. *Emerg Infect Dis*. 2011;17(6):1099–1102.
5. van Hal SJ, Lodise TP, Paterson DL. The clinical significance of vancomycin minimum inhibitory concentration in *Staphylococcus aureus* infections: a systematic review and meta-analysis. *Clin Infect Dis*. 2012;54(6):755–771.
6. Vaudaux P, Huggler E, Bernard L, Ferry T, Renzoni A, Lew DP. Underestimation of vancomycin and teicoplanin MICs by broth microdilution leads to underdetection of glycopeptide-intermediate isolates of *Staphylococcus aureus*. *Antimicrob Agents Chemother*. 2010;54(9):3861–3870.
7. Mason EO, Lamberth LB, Hammerman WA, Hulten KG, Versalovic J, Kaplan SL. Vancomycin MICs for *Staphylococcus aureus* vary by detection method and have subtly increased in a pediatric population since 2005. *J Clin Microbiol*. 2009;47(6):1628–1630.

Authors' reply

Cameron J Phillips¹⁻³

Nicholas A Wells⁴

Marianne Martinello⁴

Simon Smith⁴

Richard J Woodman⁵

David L Gordon^{2,4}

¹SA Pharmacy, Flinders Medical Centre, Bedford Park,

²Department of Microbiology and Infectious Diseases, School of Medicine, Flinders University, ³School of Pharmacy and Medical Sciences, University of South Australia, Adelaide, ⁴SA Pathology, Microbiology and Infectious Diseases, Flinders Medical Centre, Bedford Park, ⁵Flinders Centre for Epidemiology and Biostatistics, School of Medicine, Flinders University, Adelaide, SA, Australia

Correspondence: Cameron J Phillips

SA Pharmacy, Flinders Medical Centre, 1 Flinders Drive, Bedford Park, SA, 5042, Australia

Email Cameron.Phillips@sa.gov.au

Dear editor

We thank Mimica and Navarini¹ for their comments on our article.² We note with interest their findings consistent with our study regarding the poor correlation between methods for determining vancomycin minimum inhibitory concentration (MIC) also reported elsewhere.³ The need to obtain good susceptibility methods that provide both high sensitivity and high specificity is indeed challenging. We have assessed diagnostic

accuracy for two commonly used susceptibility methods (Etest[®] and Vitek[®]2) measured against the gold standard broth microdilution to give microbiologists and clinicians further insight into the sensitivity and specificity at incremental MICs. Employing two susceptibility methods is likely to become a more common practice when testing vancomycin MIC ≥ 1 and < 2 $\mu\text{g/mL}$ in an effort to more appropriately dose and monitor vancomycin in patients with these infections. Investigators of future laboratory and clinical studies that report MICs using two methods may also consider the reporting of diagnostic accuracy using a combined test approach, which might improve the interpretation of overall sensitivity and specificity.

Disclosure

The authors report no conflicts of interest in this communication.

References

1. Mimica MJ, Navarini A. Evaluating vancomycin susceptibility in *Staphylococcus aureus*. *Infect Drug Resist*. 2016;4:27–32.
2. Phillips CJ, Wells NA, Martinello M, Smith S, Woodman RJ, Gordon DL. Optimizing the detection of methicillin-resistant *Staphylococcus aureus* with elevated vancomycin minimum inhibitory concentrations within the susceptible range. *Infect Drug Resist*. 2016;9:87–92.
3. Keel RA, Sutherland CA, Aslanzadeh J, Nicolau DP, Kuti JL. Correlation between vancomycin and daptomycin MIC values for methicillin-susceptible and methicillin-resistant *Staphylococcus aureus* by 3 testing methodologies. *Diag Microbiol Infect Dis*. 2010;68(3):326–329.

Infection and Drug Resistance

Publish your work in this journal

Infection and Drug Resistance is an international, peer-reviewed open-access journal that focuses on the optimal treatment of infection (bacterial, fungal and viral) and the development and institution of preventive strategies to minimize the development and spread of resistance. The journal is specifically concerned with the epidemiology of antibiotic

Submit your manuscript here: <https://www.dovepress.com/infection-and-drug-resistance-journal>

resistance and the mechanisms of resistance development and diffusion in both hospitals and the community. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Dovepress