The potential role for use of mitochondrial DNA copy number as predictive biomarker in presbycusis

Masoumeh Falah^{1,2}
 Massoud Houshmand³
 Mohammad Najafi²
 Maryam Balali¹
 Saeid Mahmoudian¹
 Alimohamad Asghari⁴
 Hessamaldin Emamdjomeh¹
 Mohammad Farhadi¹

¹ENT and Head & Neck Research Center and Department, Iran University of Medical Sciences, Tehran, Iran; ²Cellular and Molecular Research Center, Biochemistry Department, Iran University of Medical Sciences, Tehran, Iran; ³Department of Medical Genetics, National Institute for Genetic Engineering and Biotechnology, Tehran, Iran; ⁴Skull base research center, Iran University of Medical Sciences, Tehran, Iran

Objectives: Age-related hearing impairment, or presbycusis, is the most common communication disorder and neurodegenerative disease in the elderly. Its prevalence is expected to increase, due to the trend of growth of the elderly population. The current diagnostic test for detection of presbycusis is implemented after there has been a change in hearing sensitivity. Identification of a pre-diagnostic biomarker would raise the possibility of preserving hearing sensitivity before damage occurs. Mitochondrial dysfunction, including the production of reactive oxygen species and induction of expression of apoptotic genes, participates in the progression of presbycusis. Mitochondrial DNA sequence variation has a critical role in presbycusis. However, the nature of the relationship between mitochondrial DNA copy number, an important biomarker in many other diseases, and presbycusis is undetermined.

Methods: Fifty-four subjects with presbycusis and 29 healthy controls were selected after ear, nose, throat examination and pure-tone audiometry. DNA was extracted from peripheral blood samples. The copy number of mitochondrial DNA relative to the nuclear genome was measured by quantitative real-time polymerase chain reaction.

Results: Subjects with presbycusis had a lower median mitochondrial DNA copy number than healthy subjects and the difference was statistically significant ($P=0.007$). Mitochondrial DNA copy number was also significantly associated with degree of hearing impairment ($P=0.025$) and audiogram configuration ($P=0.022$).

Conclusion: The findings of this study suggest that lower mitochondrial DNA copy number is responsible for presbycusis through alteration of mitochondrial function. Moreover, the significant association of mitochondrial DNA copy number in peripheral blood samples with the degree of hearing impairment and audiogram configuration has potential for use as a standard test for presbycusis, providing the possibility of the development of an easy-to-use biomarker for the early detection of this condition.

Keywords: age-related hearing impairment, presbycusis, biomarker, mtDNA

Introduction

Presbycusis, or age-related hearing impairment, manifests as symmetric, bilateral, sensorineural hearing impairment, occurring slowly over a lifetime.¹ It is a major reason for hearing impairment in adults. Recently, Homans et al conducted a large cohort study that showed that 33% of men and 29% of women aged >65 years suffer from presbycusis.² Presbycusis affects communication skills, gradually leading to isolation, depression, and frustration. Thus, the condition greatly impacts the quality of human life and general health.^{1,3} Panza et al demonstrated that presbycusis is a risk factor for dementia and cognitive decline.⁴ There is, as yet, no treatment for

Correspondence: Mohammad Farhadi
 ENT and Head & Neck Research Center
 and Department, Iran University of
 Medical Sciences, Hazrat Rasool Akram
 Hospital, Niayesh Street, Sattarkhan
 Street, Tehran, Iran
 Tel +98 21 6655 2828
 Fax +98 21 6652 5329
 Email mfa.ent@gmail.com

presbycusis, which develops over many years and people are usually identified after their hearing sensitivity change.⁵ Various research groups are seeking a biomarker for this common impairment;^{6–9} development of a sensitive and specific biomarker for detection and prediction of presbycusis could reduce the impact of the condition on the health of society in the future.

Mitochondria are intracellular bodies, and their impairment contributes to presbycusis.¹⁰ These are the only cellular organelles that contain their own genomes, known as mtDNA. According to the mitochondrial theory of aging proposed by Harman,¹¹ accumulation of reactive oxygen species (ROS) produced by mitochondria causes damage to intracellular macromolecules, particularly mtDNA. A lack of introns and protective histones make mtDNA more susceptible to ROS and other types of damage, which could lead to sequence mutations or copy number alterations.¹² Variation of both quality (mutations)^{13,14} and quantity (copy number)¹⁵ of mtDNA has been associated with many human diseases, including neurodegenerative diseases,¹⁶ metabolic diseases, and various types of cancer,¹⁷ as well as in the process of aging.¹⁸ Therefore, the application of mtDNA alteration as a biomarker in aging, age-related diseases, cancers, metabolic diseases, and even forensic science is currently being investigated.^{19,20}

In recent years, studies have identified some chromosomal loci and genes associated with presbycusis.^{3,21,22} A comparison of the mtDNA from temporal bones of presbycusis patients with that of healthy controls revealed a significant relationship between large-scale deletions in mtDNA and presbycusis.^{23–25} In addition to large deletions, there are reports of single nucleotide changes to mitochondrial genomes associated with presbycusis.²⁶ To determine the role of mtDNA copy number in the development of presbycusis and investigate the possibility of using mtDNA copy number as a biomarker for this impairment, we compared the mtDNA copy number in subjects with presbycusis and healthy controls by quantitative real-time polymerase chain reaction (PCR).

Materials and methods

All 50- to 90-year-old volunteers participating in the molecular investigation of presbycusis at the ENT and Head and Neck Research Center and Department, Iran University of Medical Sciences completed a questionnaire, based on that developed by the European presbycusis consortium, which included questions to determine a full medical history and details of exposure to environmental factors that affect

hearing.^{21,27} Participants with confounding factors (any known reason for hearing loss, exposure to ototoxic drugs, and other pathologies correlated with hearing loss, such as cardiovascular disorder, liver cirrhosis, renal failure, and head and neck tumor), neurodegenerative diseases, and cancers were excluded. Otoscopic examination and pure-tone audiometry were performed on all volunteers. Air conduction was measured at octave frequencies of 250–8,000 Hz and bone conduction at 250–4,000 Hz using an Amplaid 319 audiometer (Milan, Italy). Calibration was performed according to the American National Standards Institute (ANSI, 2004a). Subjects with any kind of middle ear pathology affecting hearing sensitivity, conductive hearing loss (mean air-bone differences at 500, 1,000, and 2,000 Hz of ≥ 15 dB), and extensive noise exposure (self-reported) were excluded.

Presbycusis patients ($n=54$) with symmetric, bilateral hearing impairment with the pure-tone average of mean frequencies 1, 2, 4, and 8 kHz of ≥ 30 dB HL were selected. Healthy subjects ($n=29$) with normal ear examination and the pure-tone average of mean frequencies ≤ 25 dB HL were selected. The mean age of patients was 66.87 ± 8.9 years and that of healthy subjects was 63 ± 5.3 years. The gender ratio was 54.2% male and 45.8% female, and there were no statistically significant differences in age and gender between presbycusis and healthy subjects ($P > 0.05$). This study was approved by the Ethics Committee of Iran University of Medical Sciences, and all subjects provided signed informed consent.

Genomic DNA was extracted from peripheral blood samples using the QIAamp DNA Mini Kit (Cat No 51304; Qiagen, Hilden, Germany) according to the manufacturer's protocol.

Relative mtDNA copy number was determined through simultaneous assessment of the nuclear gene, *TMIE* (NC_000003.12), and the mitochondrial gene, *MT-RNR1* (NC_012920.1), using quantitative PCR (qPCR). Primers were as follows: *MT-RNR1*-F, 5'-AGTAG AGTGCTTAGTTGAGC-3'; *MT-RNR1*-R, 5'-AG TACACTTCCATGTTACG-3'; *TMIE*-F, 5'-CCATTC CTTGGGTCTCTGAA-3'; and *TMIE*-R, 5'-AGCAGA GGAACAGGGTGAC-3'. qPCR was performed using a Rotor-gene 6000 instrument (Corbett Life Science, Sydney, Australia) and QuantiFast SYBR Green PCR master mix (Cat No 204054, Qiagen). All reactions were performed in duplicate with total reaction volumes of 10 μ L, containing 5 μ L of 1X Quanti Fast SYBR Green PCR master mix, 0.12 μ L of each primer (10 pmol), 0.7 μ L of DNA, and 4 μ L of RNase-free water. After initial heating for 1 min at 95°C as an activation step, 40 cycles of denaturation at 95°C for

10 s, followed by annealing and extension at 60°C for 45 s, were performed.

The threshold cycle (Ct) of each sample was determined for *MT-RNR1* and *TMIE*. Relative mtDNA copy number was determined using the following equation: $2^{-\Delta Ct}$ ($\Delta Ct = Ct_{MT-RNR1} - Ct_{TMIE}$).

Standard curves were prepared for both nuclear and mitochondrial PCR. R^2 correlation for each standard curve was ≥ 0.99 . Melting curve analysis of qPCR and running of PCR products on agarose gels (1.5%) were performed to confirm amplification of a specific product without primer dimmer or unwanted PCR product.

Statistical analyses were performed using SPSS 22 for Windows (IBM, Chicago, IL, USA). All numerical variables were checked for normal distribution using the one-sample Kolmogorov–Smirnov test. The association between the mtDNA copy number in presbycusis and healthy subjects was evaluated using the Mann–Whitney *U*-test. The associations between mtDNA copy number and degree of hearing impairment and audiogram configuration were evaluated using Kruskal–Wallis H-test. Statistical significance was defined as $P < 0.05$.

Results

The degree of hearing impairment of subjects with presbycusis was classified into five categories according to the American Speech–Language–Hearing Association: mild, moderate, moderately severe, severe, and profound.²⁸ Patients in this study suffered from mild, moderate, and moderately severe hearing impairment (Figure 1A).

Based on audiogram configuration described previously, patients were divided into five groups: flat, high-frequency

gently sloping (HFGS), high-frequency steeply sloping (HFSS), low-frequency ascending, and mid-frequency U-shape.²⁹ Presbycusis subjects in this study exhibited flat, HFGS, and HFSS shapes (Figure 1B).

The mean audiograms of presbycusis patients and healthy subjects are presented in Figure 2.

The mitochondrial copy number of study participants was evaluated by measuring the proportion of mtDNA relative to nuclear DNA using the level of *MT-RNR1* as an evaluation index of the mitochondrial genome and that of *TMIE* for nuclear DNA. Median mtDNA copy number for subjects with presbycusis was 0.76, whereas that of healthy subjects was 1.45, indicating that the mitochondrial copy number of subjects with presbycusis was almost halved compared with the healthy controls (Figure 3). The difference in mtDNA copy number between the two groups was statistically significant ($P=0.007$).

A Kruskal–Wallis H-test showed that there was a statistically significant difference in mtDNA copy number between the different hearing degrees, $\chi^2(3)=9.33$, $P=0.025$, with a mean rank mtDNA copy number of 50.79 for normal, 29.72 for mild, 38.20 for moderate, and 44.64 for moderately severe.

A Kruskal–Wallis H-test showed that there was a statistically significant difference in mtDNA copy number between the different audiogram configurations, $\chi^2(2)=7.62$, $P=0.022$, with a mean rank mtDNA copy number of 51.23 for flat shape, 32.96 for HFGS shape, and 37.68 for HFSS shape.

Discussion

Presbycusis is categorized as a complex or multifactorial disorder.^{30,31} Factors affecting its development include genetic background, environmental factors, ototoxic drugs,

Figure 1 (A) Classification of presbycusis patients by degree of hearing impairment (n=54). (B) Classification of patients by shape of audiogram. **Abbreviations:** HFGS, high-frequency gently sloping; HFSS, high-frequency steeply sloping.

Figure 2 Mean audiograms of patients with presbycusis and healthy subjects. Error bars represent SDs.

Abbreviation: SD, standard deviation.

medical conditions, and hormonal factors.^{32–34} Presbycusis occurs as a result of degeneration of both the inner ear structure and the central auditory system of the brain.^{35–37} Mitochondria play a key role in cellular damage and death in the aging process of hearing.^{38,39} Mitochondrial dysfunction is related to the development of certain neurodegenerative, age-related diseases, and cancer.^{14,40} Mutations in the *MT-RNR1* and *MT-TF* genes can cause hearing loss.^{26,41} Investigation

Figure 3 Distribution of peripheral blood mtDNA copy number in Iranian patients with presbycusis and healthy controls. The relative mtDNA copy number was evaluated by measuring the proportion of mtDNA relative to nuclear DNA. The difference between mtDNA copy number in the presbycusis and healthy subject groups was statistically significant (Mann–Whitney *U*-test, $P=0.007$). The box plots show the median (solid line across the box), inter-quartile range (shaded boxes), and range (whiskers).

of the temporal bone of presbycusis patients showed an increase in deletions and point mutations in mtDNA. Bai et al detected a 4,977 bp deletion, known as the common deletion, in the mtDNA of human cochlear tissue of presbycusis patients.^{23,24} Markaryan demonstrated that the expression of the *MT-CO3* gene is decreased in spiral ganglion cells from individuals with presbycusis, suggesting that deficits in the electron transport chain of spiral ganglion cells contribute to presbycusis.²⁵

The results of this study indicate a statistically significant decrease in mtDNA copy number in presbycusis subjects. Our study is the first to investigate the relationship between mtDNA copy number and presbycusis. Previously, Santos et al proposed a decrease in mtDNA copy number as an important marker for oocyte quality, having shown that such a decrease is due to insufficient mitochondrial biogenesis or cytoplasmic maturation, causing fertilization failure.⁴² Studies of the relationship between neurodegenerative disease and mtDNA content have produced similar results to those of our study. Podlesniy et al suggested low content of mtDNA in cerebrospinal fluid as a novel biomarker for the early detection of preclinical Alzheimer's disease.⁴³ Petersen et al demonstrated reduced mtDNA copy number in blood leukocytes from patients with Huntington's disease.¹⁶

Our results are consistent with previous studies demonstrating that presbycusis is commonly a high-frequency hearing impairment, as HFSS was the most common configuration in patients in this study (70.9%), followed by HFGS (23.64%) (Figure 1B).^{1,44}

According to previous studies, a decrease in the number of mitochondrial genomes can cause impaired or reduced mitochondrial activity. A study using HeLa cells indicated that depletion of the mitochondrial genome by decreasing catalase expression enhances oxidative damage to nuclear DNA and influences nuclear genome stability.⁴⁵ Oxidative stress is associated with excessive ROS production in the mitochondrial matrix; ROS increases death of cellular structures of the cochlear, leading to an increase in mutations in the mitochondrial genome.⁴⁰ In transgenic mice, accumulation of mtDNA mutations by mitochondrial dysfunction influences energy metabolism. The consequent reduction in energy production induces apoptosis, leading to a loss of critical cochlear cells and resulting in age-related hearing impairment.⁴⁶

The results of this study, together with those of previous reports demonstrating the relationship between mtDNA sequence variation and presbycusis,^{23,26,47} highlight the role of mtDNA in presbycusis progression. A significant reduction

in the number of mitochondrial genomes in presbycusis subjects, due to mitochondrial dysfunction, may be one of the molecular mechanisms involved in the development of this impairment. In addition, quantification of mtDNA in peripheral blood samples may constitute an easily assessed biomarker of presbycusis, since the high-sensitivity qPCR-based assay can detect even minute variations in mtDNA copy number. This blood biomarker could easily be incorporated into regular physical examinations as it employs a simple and cheap screening method. Moreover, the results of this study demonstrate that mtDNA copy number is statistically significantly associated with the degree of hearing ($P=0.025$) and audiogram configuration ($P=0.022$). Clearly, these results require further confirmation in a larger study population to determine whether the method is appropriate for clinical application. In addition, more research is required to understand the relationship between the decrease in mtDNA copy number and the onset of presbycusis.

Age-related hearing impairment is the most common sensory disease in the elderly population.¹⁰ With the growth of the aging population, the prevalence of age-related disorders such as presbycusis is increasing.¹ The World Health Organization estimates that in 2025, more than 500 million individuals will suffer from presbycusis.⁴⁸ Presbycusis begins slowly, without clinical manifestation. Further human studies are required to elucidate methods for early diagnosis and prevention.

Acknowledgments

We appreciate the participation of all the volunteers in this research and would also like to thank those in Audiometry and Clinical Department, Rasoul Akram Hospital, Iran University of Medical Sciences for their kind cooperation during this research.

Disclosure

The authors report no conflicts of interest in this work.

References

- Huang Q, Tang J. Age-related hearing loss or presbycusis. *Eur Arch Otorhinolaryngol*. 2010;267(8):1179–1191.
- Homans NC, Metselaar RM, Dingemanse JG, et al. Prevalence of age-related hearing loss, including sex differences, in older adults in a large cohort study. *Laryngoscope*. Epub 2016 Jul 5.
- Ciorba A, Hatzopoulos S, Bianchini C, Aimoni C, Skarzynski H, Skarzynski PH. Genetics of presbycusis and presbystasis. *Int J Immunopathol Pharmacol*. 2015;28(1):29–35.
- Panza F, Solfrizzi V, Logroscino G. Age-related hearing impairment-a risk factor and frailty marker for dementia and AD. *Nat Rev Neurol*. 2015; 11(3):166–175.
- Carson AJ. “What brings you here today?” The role of self-assessment in help-seeking for age-related hearing loss. *J Aging Stud*. 2005;19(2): 185–200.
- Michikawa T, Nakamura T, Imamura H, et al. Markers of overall nutritional status and incident hearing impairment in community-dwelling older Japanese: the Kurabuchi Study. *J Am Geriatr Soc*. 2016;64(7): 1480–1485.
- Williamson TT, Zhu X, Walton JP, Frisina RD. Auditory brainstem gap responses start to decline in mice in middle age: a novel physiological biomarker for age-related hearing loss. *Cell Tissue Res*. 2015;361(1): 359–369.
- Pang J, Xiong H, Yang H, et al. Circulating miR-34a levels correlate with age-related hearing loss in mice and humans. *Exp Gerontol*. 2016;76: 58–67.
- Kuchinsky SE, Ahlstrom JB, Cute SL, Humes LE, Dubno JR, Eckert MA. Speech-perception training for older adults with hearing loss impacts word recognition and effort. *Psychophysiology*. 2014;51(10): 1046–1057.
- Fujimoto C, Yamasoba T. Oxidative stresses and mitochondrial dysfunction in age-related hearing loss. *Oxid Med Cell Longev*. 2014; 2014:582849.
- Harman D. The biologic clock: the mitochondria? *J Am Geriatr Soc*. 1972;20(4):145–147.
- Lee HC, Wei YH. Mitochondrial biogenesis and mitochondrial DNA maintenance of mammalian cells under oxidative stress. *Int J Biochem Cell Biol*. 2005;37(4):822–834.
- Dowlati MA, Derakhshandeh-Peykar P, Houshmand M, et al. Novel nucleotide changes in mutational analysis of mitochondrial 12S rRNA gene in patients with nonsyndromic and aminoglycoside-induced hearing loss. *Mol Biol Rep*. 2013;40(3):2689–2695.
- Greaves LC, Reeve AK, Taylor RW, Turnbull DM. Mitochondrial DNA and disease. *J Pathol*. 2012;226(2):274–286.
- Clay Montier LL, Deng JJ, Bai Y. Number matters: control of mammalian mitochondrial DNA copy number. *J Genet Genomics*. 2009;36(3): 125–131.
- Petersen MH, Budtz-Jørgensen E, Sørensen SA, et al. Reduction in mitochondrial DNA copy number in peripheral leukocytes after onset of Huntington’s disease. *Mitochondrion*. 2014;17:14–21.
- Zhou W, Zhu M, Gui M, et al. Peripheral blood mitochondrial DNA copy number is associated with prostate cancer risk and tumor burden. *PLoS One*. 2014;9(10):e109470.
- Zabihi Diba L, Mohaddes Ardebili SM, Ghahesouran J, Houshmand M. Age-related decrease in mtDNA content as a consequence of mtDNA 4977 bp deletion. *Mitochondrial DNA A DNA Mapp Seq Anal*. 2016; 27(4):3008–3012.
- Wallace DC. A mitochondrial paradigm of metabolic and degenerative diseases, aging, and cancer: a dawn for evolutionary medicine. *Annu Rev Genet*. 2005;39:359–407.
- Meissner C, Bruse P, Mohamed SA, et al. The 4977 bp deletion of mitochondrial DNA in human skeletal muscle, heart and different areas of the brain: a useful biomarker or more? *Exp Gerontol*. 2008; 43(7):645–652.
- Van Laer L, Huyghe JR, Hannula S, et al. A genome-wide association study for age-related hearing impairment in the Saami. *Eur J Hum Genet*. 2010;18(6):685–693.
- Newman DL, Fisher LM, Ohmen J, et al. GRM7 variants associated with age-related hearing loss based on auditory perception. *Hear Res*. 2012;294(1–2):125–132.
- Markaryan A, Nelson EG, Hinojosa R. Quantification of the mitochondrial DNA common deletion in presbycusis. *Laryngoscope*. 2009; 119(6):1184–1189.
- Bai U, Seidman MD, Hinojosa R, Quirk WS. Mitochondrial DNA deletions associated with aging and possibly presbycusis: a human archival temporal bone study. *Am J Otol*. 1997;18(4):449–453.
- Markaryan A, Nelson EG, Hinojosa R. Major arc mitochondrial DNA deletions in cytochrome c oxidase-deficient human cochlear spiral ganglion cells. *Acta Otolaryngol*. 2010;130(7):780–787.
- Zhu Y, Zhao J, Feng B, et al. Mutations in the mitochondrial 12S rRNA gene in elderly Chinese people. *Acta Otolaryngol*. 2015;135(1):26–34.

27. Van Eyken E, Van Laer L, Fransen E, et al. KCNQ4: a gene for age-related hearing impairment? *Hum Mutat*. 2006;27(10):1007–1016.
28. Clark JG. Uses and abuses of hearing loss classification. *ASHA*. 1981;23(7):493–500.
29. Mostafa H, Saad M, El-Attar A, et al. Mitochondrial DNA (mtDNA) haplotypes and dysfunctions in presbycusis. *Acta Otorhinolaryngol Ital*. 2014;34(1):54–61.
30. Ohlemiller KK. Age-related hearing loss: the status of Schuknecht's typology. *Curr Opin Otolaryngol Head Neck Surg*. 2004;12(5):439–443.
31. Falah M, Houshmand M, Farhadi M. Presbycusis: From current knowledge to future treatment prospects. *J Isfahan Med School*. 2016;34(382):526–535. Persian.
32. Lin FR, Thorpe R, Gordon-Salant S, Ferrucci L. Hearing loss prevalence and risk factors among older adults in the United States. *J Gerontol A Biol Sci Med Sci*. 2011;66(5):582–590.
33. Charitidi K, Frisina RD, Vasilyeva ON, Zhu X, Canlon B. Expression patterns of estrogen receptors in the central auditory system change in prepubertal and aged mice. *Neuroscience*. 2010;170(4):1270–1281.
34. Tadros SF, D'Souza M, Zettel ML, Zhu X, Lynch-Erhardt M, Frisina RD. Serotonin 2B receptor: upregulated with age and hearing loss in mouse auditory system. *Neurobiol Aging*. 2007;28(7):1112–1123.
35. Ohlemiller KK, Frisina RD. Age-related hearing loss and its cellular and molecular bases. In: Schacht J, Popper AN, Fay RR, editors. *Auditory Trauma, Protection, and Repair*. Boston, MA: Springer US; 2008: 145–194.
36. Frisina DR, Frisina RD. Speech recognition in noise and presbycusis: relations to possible neural mechanisms. *Hear Res*. 1997;106(1–2): 95–104.
37. Zettel ML, Zhu X, O'Neill WE, Frisina RD. Age-related decline in Kv3.1b expression in the mouse auditory brainstem correlates with functional deficits in the medial olivocochlear efferent system. *J Assoc Res Otolaryngol*. 2007;8(2):280–293.
38. Wong AC, Ryan AF. Mechanisms of sensorineural cell damage, death and survival in the cochlea. *Front Aging Neurosci*. 2015;7:58.
39. Falah M, Najafi M, Houshmand M, Farhadi M. Expression levels of the BAK1 and BCL2 genes highlight the role of apoptosis in age-related hearing impairment. *Clin Interv Aging*. 2016;11:1003–1008.
40. Kamogashira T, Fujimoto C, Yamasoba T. Reactive oxygen species, apoptosis, and mitochondrial dysfunction in hearing loss. *Biomed Res Int*. 2015;2015:617207.
41. Balali M, Kamalidehghan B, Farhadi M, et al. Association of nuclear and mitochondrial genes with audiological examinations in Iranian patients with nonaminoglycoside antibiotics-induced hearing loss. *Ther Clin Risk Manag*. 2016;12:117–128.
42. Santos TA, El Shourbagy S, St John JC. Mitochondrial content reflects oocyte variability and fertilization outcome. *Fertil Steril*. 2006;85(3): 584–591.
43. Podlesniy P, Figueiro-Silva J, Llado A, et al. Low cerebrospinal fluid concentration of mitochondrial DNA in preclinical Alzheimer disease. *Ann Neurol*. 2013;74(5):655–668.
44. Uchida Y, Sugiura S, Sone M, Ueda H, Nakashima T. Progress and prospects in human genetic research into age-related hearing impairment. *Biomed Res Int*. 2014;2014:390601.
45. Delsite RL, Rasmussen LJ, Rasmussen AK, Kalen A, Goswami PC, Singh KK. Mitochondrial impairment is accompanied by impaired oxidative DNA repair in the nucleus. *Mutagenesis*. 2003;18(6):497–503.
46. Someya S, Yamasoba T, Kujoth GC, et al. The role of mtDNA mutations in the pathogenesis of age-related hearing loss in mice carrying a mutator DNA polymerase gamma. *Neurobiol Aging*. 2008;29(7):1080–1092.
47. Manwaring N, Jones MM, Wang JJ, et al. Mitochondrial DNA haplogroups and age-related hearing loss. *Arch Otolaryngol Head Neck Surg*. 2007;133(9):929–933.
48. Sprinzl GM, Riechelmann H. Current trends in treating hearing loss in elderly people: a review of the technology and treatment options – a mini-review. *Gerontology*. 2010;56(3):351–358.

Therapeutics and Clinical Risk Management

Publish your work in this journal

Therapeutics and Clinical Risk Management is an international, peer-reviewed journal of clinical therapeutics and risk management, focusing on concise rapid reporting of clinical studies in all therapeutic areas, outcomes, safety, and programs for the effective, safe, and sustained use of medicines. This journal is indexed on PubMed Central, CAS,

Submit your manuscript here: <http://www.dovepress.com/therapeutics-and-clinical-risk-management-journal>

Dovepress

EMBASE, Scopus and the Elsevier Bibliographic databases. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.