The Danish nationwide clinical register for patients with rheumatoid arthritis: DANBIO

Else Helene Ibfelt¹
Dorte Vendelbo Jensen^{2,3}
Merete Lund Hetland^{2,4}

¹Registry Support Centre (East), Epidemiology and Biostatistics, Research Centre for Prevention and Health, Rigshospitalet, Glostrup University Hospital, ²DANBIO Registry and Copenhagen Center for Arthritis Research, Center for Rheumatology and Spine Diseases, Centre of Head and Orthopedics, Rigshospitalet, Glostrup, ³Department of Rheumatology, Herlev and Gentofte University Hospital, Hellerup, ⁴Department of Clinical Medicine, Faculty of Health and Medical Sciences, University of Copenhagen, Copenhagen, Denmark

Introduction: DANBIO is a research register and a data source for rheumatologic diseases (rheumatoid arthritis [RA], axial spondyloarthritis, and psoriatic arthritis) for monitoring clinical quality at the national, regional, and hospital levels.

Study population: The register includes patients with rheumatologic diseases who are treated at a hospital or a private rheumatologic clinic. Registration is mandatory for all patients with RA regardless of treatment and also for patients with other diagnoses if treated with biological disease-modifying antirheumatic drugs. Since 2006, the registration has been done electronically, including patient-reported outcome measures registered electronically by the patients with the use of touch screens.

Main variables: Core variables such as diagnosis, year of diagnosis, age, and sex are registered at the beginning. Data entered at later visits included the following: patient-reported outcomes for disease activity, pain, fatigue, functional status, and physician-reported objective measures of disease activity, treatment, C-reactive protein, and, when indicated, imaging. For subgroups of patients, the variables such as quality of life, sociodemographic factors, lifestyle, and comorbidity are also registered.

Descriptive data: The DANBIO cohort comprised ~26,000 patients with RA, 3,200 patients with axial spondyloarthritis, and 6,200 patients with psoriatic arthritis in 2015. DANBIO has high nationwide coverage and completeness on key data variables. More than 60 original papers as well as annual reports of clinical quality (since 2005) have been published.

Conclusion: DANBIO is a powerful register for research in rheumatologic diseases and furthermore serves as a Clinical Quality Register with the aim of monitoring treatment quality in patients with RA in Denmark.

Keywords: rheumatoid arthritis, axial spondyloarthritis, psoriatic arthritis, DANBIO register clinical quality, epidemiology

Introduction

The database, DANBIO, is a nationwide, Danish register for research use in rheumatologic diseases such as rheumatoid arthritis (RA), axial spondyloarthritis (Ax SpA), and psoriatic arthritis (PsA). It also serves as a clinical database that monitors clinical quality of treatment by the use of selected quality indicators for patients with RA in Denmark.

Study population and data collection

Up to the year 2000, no routine-based nationwide reporting of patients with RA existed in Denmark, and patient files rarely comprised quantitative patient-reported outcome

Correspondence: Dorte Vendelbo Jensen
DANBIO Registry and Copenhagen
Center for Arthritis Research, Center for
Rheumatology and Spine Diseases, Centre
of Head and Orthopedics, Ndr. Ringvej 57,
DK-2600 Glostrup, Denmark
Tel +45 3863 3103
Fax +45 3863 4192
Email dvj@dadlnet.dk

data. The introduction of new biological disease-modifying antirheumatic drugs (bDMARDs) triggered the formation of a nationwide voluntary register, which aimed to survey indications for treatment, efficacy, and adverse events in rheumatologic patients who received biological therapies in routine care.

In the beginning, data were collected on paper forms, which were subsequently scanned into the register at the DANBIO office.¹ In 2006, an online version of DANBIO was introduced, and the aims were extended to include patients regardless of treatment and also to collect patient-reported outcomes regarding, eg disability, pain, lifestyle, and quality of life by the use of dedicated touch screens in the waiting room (Figure 1). This is routinely done before consultation with the doctor, and the information from the touch screen is available during the consultation. DANBIO has been approved by the National Board of Health as a Clinical Quality Register, and since 2006, reporting to the register has been mandatory. For clinical quality registers, the usual requirement for obtaining an informed patient consent before registration is not needed. Only rarely do patients object to be included in the register, and in these cases, it is up to the physician to decide whether to register the patient or not.

RA, Ax SpA, and PsA are chronic diseases that require lifelong monitoring and treatment. A patient is therefore reported to the database at the time of diagnosis, referral to specialized treatment, including biological therapy at the hospital, or in a private rheumatologic clinic. At the first registration, the diagnosis, date of diagnosis, age, sex, and previous medical treatment are registered. At later visits, and at least once yearly, information regarding patient's disease activity (including pain and functional status) is collected

via touch screens, and objective measures (eg, swollen and tender joint counts, C-reactive protein) are entered by the physician.

The patient groups with Ax SpA and PsA are not included at the moment in the National Clinical Quality Program, and therefore reporting is only mandatory for patients with RA and for the other patient groups only when treated with bDMARDs. In the beginning of 2015, the DANBIO cohort comprised ~26,000 patients with RA, 3,200 patients with Ax SpA, and 6,200 patients with PsA treated with conventional synthetic DMARDs and/or bDMARDs.

Main variables

Table 1 gives the key variables of the register. Some variables are collected via touch screens (patient-reported outcome measures [PROMs]), and others are entered by the physician.

The use of touch screens for data collection has been validated to perform equally well as paper forms (Figure 1).² The variables are collected in a standardized format, often using tick boxes, which have been shown to have a high validity.¹ Figure 2 shows the scoreboard for entering and reviewing individual patient data. In connection with ie, queries from DANBIO and research projects, the departments fill in missing data by checking against the hospital patient records.

The completeness of data is generally high (ie, 96%, 93%, 83%, 90%, and 93% for information on the name of the treating doctor, diagnosis, year of diagnosis, DAS28, and HAQ, respectively). National coverage of DANBIO is assessed annually in two ways: for patients in biological treatment by comparing to those registered in the patient record systems of each rheumatologic department (94% in 2013) and by

Figure 1 A patient using the touch screen.
Note: Reprinted with permission from DANBIO.

ADD NEW		Last 10 visits											
		Click here for: All visits or 1st visit + last 9 visits											
		A	B	C	D	E	F	G	H	I	J		
		07feb	15feb	29feb	01apr	29apr	27may	24jun	30jul	21aug	24sep		
		2008	2008	2008	2008	2008	2008	2008	2008	2008	2008		
1. Biologic drug		etan	infix	infix	infix	infix	infix	infix	infix	infix	infix		
2. Dose of biol. drug		50	280	280	280	280	280	280	290	300	300		
3. Freq. of biol. drug		weekly	026+8	026+8	e4w	e5w	e4w	e4w	e4w	e5w	e5w		
4. Prednisolone		-	-	-	-	-	-	5	5	5	5		
5. Methotrexate oral		7.5	7.5	7.5	20	20	20	20	20	20	20		
6. Vas pain		65	61	75	70	5	2	6	10	3	5		
7. Vas fatigue					69	16	15	16	24	31	11		
8. Vas global		70	68	80	81	16	11	8	21	16	16		
9. HAQ		2.0	2.0	2.0	2.0	1.375	1.375	1.625	1.625	1.5	1.25		
10. CRP		33	20	43	43	4	5	4	4	4	5		
11. X-ray											+/0		
12. TJC28		13	19	20	22	0	0	0	0	0	1		
13. SJC28		8	14	14	7	0	0	0	0	0	0		
14. Doctor's global		80	65	60	95	8	5	8	0	10	41		
15. DAS28crp		6.4	6.8	7.2	6.4								
16. I.a./I.v./i.m.													
17. Other inj.		1											
18. SAE													

Figure 2 Overview of patient data.
Note: Reprinted with permission from DANBIO.

Table 1 Key variables in the DANBIO register

Type of variable	Name of variable	Disease group
Baseline variables		
Sociodemographic factors		
	Age	RA, Ax SpA, PsA
	Sex	RA, Ax SpA, PsA
PROMs	Marital status	RA, Ax SpA, PsA
PROMs	Educational level	RA, Ax SpA, PsA
Disease-related variables		
	Diagnosis	RA, Ax SpA, PsA
	Year of diagnosis	RA, Ax SpA, PsA
	Previous DMARD treatment	
	Anti-CCP and IgM-RF status ($\pm$)	RA
	MRI	Ax SpA
Health behavioral factors (once a year)		
PROMs	Tobacco use	RA, Ax SpA, PsA
PROMs	Alcohol consumption	RA, Ax SpA, PsA
PROMs	Physical activity	RA, Ax SpA, PsA
PROMs	EQ-5D	RA, Ax SpA, PsA
	X-ray status (when indicated)	RA, PsA
Variables at each visit		
Disease activity measures		
	DAS28 ^a	RA, PsA
	Tender joints count	RA, PsA
	Swollen joints count	RA, PsA
PROMs	VAS for pain	RA, Ax SpA, PsA
PROMs	VAS for fatigue	RA, Ax SpA, PsA
PROMs	VAS for patient's global	RA, Ax SpA, PsA
	VAS for physician's global	RA, Ax SpA, PsA
PROMs	Health Assessment Questionnaire (HAQ) ^b	RA, PsA
PROMs	BASDAI ^c	Ax SpA
PROMs	BASFI ^d	Ax SpA
	BASMI ^e	Ax SpA
	Serum CRP	RA, Ax SpA, PsA
Current treatment (start date, dose, interval)		
	NSAID	Ax SpA
	csDMARDs	RA, Ax SpA, PsA
	bDMARDs	RA, Ax SpA, PsA
	Glucocorticoids	RA, Ax SpA, PsA
Variables at treatment termination		
	Relevant disease activity measures	RA, Ax SpA, PsA
	Date and reason for withdrawal	RA, Ax SpA, PsA
Adverse effect variables		
	Serious/not serious	RA, Ax SpA, PsA

Notes: ^aDAS based on a 28 joint count (tender and swollen joint counts), serum CRP, and patient global health. DAS28 <3.2 indicates low disease activity; DAS28 between 3.3 and 5.1, moderate disease activity; and DAS28 >5.2 high disease activity (further explanation of DAS28 is found on www.das-score.nl). ^bHAQ: 20 questions in eight categories of functioning that represent a comprehensive set of functional activities. ^cBath Ankylosing Spondylitis Disease Activity Index: a combined index comprising six questions about five major symptoms of Ax SpA. ^dBath Ankylosing Spondylitis Function Index: a combined index comprising ten questions about daily activities and ability to cope with everyday life. ^eBath Ankylosing Spondylitis Metrology Index: a combined index comprising five assessments of spinal mobility.

Abbreviations: Ax SpA, axial spondyloarthritis; DMARD, disease-modifying antirheumatic drugs; bDMARDs, biological disease-modifying antirheumatic drugs; CRP, C-reactive protein; csDMARDs, conventional synthetic disease-modifying antirheumatic drugs; DAS, disease activity score; HAQ, Health Assessment Questionnaire; NSAIDs, nonsteroid anti-inflammatory drugs; PROMs, patient-reported outcome measures; PsA, psoriatic arthritis; RA, rheumatoid arthritis; VAS, visual analog scale; Anti-CCP, anti-cyclic citrullinated peptide; IgM-RF, IgM rheumatoid factor; MRI, magnetic resonance imaging; EQ-5D, EuroQol-5D.

calculation of all incident patients with RA in DANBIO by comparison to the Danish National Patient Registry (NPR; 85% in 2013). Since NPR is an administrative register³ in which RA diagnosis may be registered for patient contacts before the final diagnosis of the patient has been established, it cannot serve as a golden standard; however, it is the most valid nationwide register available for such comparison. Further, local medical record audits have shown that the diagnosis in DANBIO is valid. Therefore, it is assumed that the true coverage of DANBIO is higher than this estimate.⁴

Follow-up

At least once-a-year follow-up information is collected in DANBIO, including PROMs such as the level of function and level of pain. Disease activity, type of treatment, treatment

Table 2 Overview of the DANBIO quality indicators monitored in the Annual Clinical Quality Report

Indicator number	Indicator title	Standard (goal set for the indicator)
1	Close follow-up of newly diagnosed patients with RA	At least 80% of patients registered twice during the first year with disease activity level (DAS28 ^a), functional status (HAQ ^b), level of pain (VAS pain ^c), and treatment
2	Longitudinal follow-up of all patients with RA	At least 90% of patients registered once a year with disease activity level (DAS28 ^a), functional status (HAQ ^b), level of pain (VAS pain ^c), and treatment
3	Effective treatment of all patients with RA evaluated by disease activity level	Low disease activity level (DAS28 ^a <3.2) in at least 66% of the registered patients
4	High functional status in all patients with RA	High functional status (HAQ ^b <1) in at least 60% of the registered patients
5	Effective treatment of pain in all patients with RA	Low pain score (VAS pain ^c <33 mm) in at least 60% of the registered patients
6	High-life quality in all patients with RA	High life quality score (VAS global ^d <33 mm) in at least 60% of the registered patients
7	Imaging by X-ray performed for all patients with RA (of hands, wrists, and feet)	X-ray status performed in at least 80% of the registered patients at the time of diagnosis and at shift between types of medical treatment

Notes: ^aDAS based on a 28 joint count (tender and swollen joint counts), serum CRP, and patient global health. DAS28 <3.2 indicates low disease activity; DAS28 between 3.3 and 5.1, moderate disease activity; and DAS28 >5.2, high disease activity. ^bHAQ: 20 questions in eight categories of functioning that represent a comprehensive set of functional activities. ^cVAS for pain. ^dVAS for patient's global. **Abbreviations:** CRP, C-reactive protein; DAS, disease activity score; HAQ, Health Assessment Questionnaire; RA, rheumatoid arthritis; VAS, visual analog scale.

effectiveness, and side effects of treatment (if any) are registered by the physician.⁵

Systematic monitoring of patients with RA with real-time feedback to the physician is feasible. It is documented that regular registration of disease activity for patients with RA results in improvement in disease activity, although the goal of treat to target is not achieved in a substantial proportion of patients in routine care.⁶

Quality indicators

The current quality indicators for RA are shown in Table 2. The indicators include goals for whether patients are “treated to target” (ie, remission or low disease activity) by follow-up of patients with measurement and registration of disease activity level, functional status, pain, quality of life, and X-ray status. A standard for each indicator is set by the DANBIO steering committee and by consensus among the members of the Danish Society of Rheumatology setting goals for how large a fraction of patients who should meet the indicator criteria. The indicator results are published at country, regional, and hospital department levels in the Annual Clinical Quality Report.⁴ An example of supplementary results from the quality report is presented in Figure 3 that gives the number of adult rheumatologic patients initiating bDMARDs in Denmark over time and the differences in prescription pattern between geographical regions.⁴

Examples of research

DANBIO is used in routine care for systematic monitoring of the patients, but it is also an extensive research database.

DANBIO data have been used for research with focus on both short- and long-term evaluation of treatment responses, remission rates, and drug adherence.^{7–10} Furthermore, research has been performed on cancer incidence and precancerous development following treatments with biological treatment regimes,^{11,12} as well as on the influence of tobacco smoking on treatment response.¹³

More than 60 peer-reviewed articles have been published to date with the use of DANBIO data and >170 abstracts have been presented at international conferences and congresses.

DANBIO is easily linked to other data sources by the use of the unique personal identification number, assigned to all Danish citizens. With regulatory approval for specific research purposes, DANBIO data can be merged with ie, the NPR, the Danish Cancer Registry, or administrative registers holding socioeconomic information in Statistic Denmark.

A nationwide biobank (ie, blood and synovial fluid) associated with DANBIO has been established since 2015 through public funding.

Administrative issues and funding

DANBIO has an independent steering committee with representatives from the Danish Society of Rheumatology, DRFO

Figure 3 Number of adult rheumatologic patients initiating bDMARDs in Denmark over time across the geographical regions.

Note: Reprinted with permission from DANBIO.

Abbreviation: bDMARDs, biological disease-modifying antirheumatic drugs.

(Danske Reumatologer og Fysiurgers Organisation), Junior Rheumatologists (Yngre Reumatologer), and the hospital owners. The daily administration is handled in the DANBIO general office at Rigshospitalet, Glostrup, and is staffed by the Head of Secretariat as well as a secretary.

DANBIO has been approved as a clinical quality register by the Danish authorities – the National Board of Health and the Danish Data Protection Agency – and is sponsored by the public hospital owners (Danish Regions) under the organization of the Danish Clinical Registries – a national improvement program. The pharmaceutical companies that provide biological treatments in rheumatology contribute to the development of the information technology platform through unrestricted grants, which have been approved by the public hospital owners. The sponsors have no influence on the register setup, data collection, data analysis, or publication of results. These issues are all administered by the steering committee.

Conclusion

DANBIO serves as a powerful register for research in rheumatologic diseases and additionally as a clinical quality register with the aim of monitoring quality by the use of selected clinical indicators for patients with RA in Denmark.

The register has existed since 2000, and patients diagnosed at the hospital or at private rheumatologic clinic are registered. Since 2006, the registration, including PROMs, has been electronically registered directly into the database with the use of touch screens.

DANBIO has high nationwide coverage of patients with rheumatologic diseases and high completeness on key data variables. Data have been used for a large number of research projects and each year a National Clinical Quality Report is published.

Acknowledgments

This paper was funded by the Program for Clinical Research Infrastructure established by the Lundbeck Foundation and the Novo Nordisk Foundation and administered by the Danish Regions. DANBIO thanks the Danish Regions for supporting the register. All the Departments of Rheumatology in Denmark and the private rheumatologic clinics are acknowledged for using the register. Niels Steen Krogh, Zitellab ApS, is acknowledged for the continuous technical development and support of the register.

Disclosure

The authors report no conflicts of interest in this work.

References

1. Hetland ML, Unkerskov J, Ravn T, et al. Routine database registration of biological therapy increases the reporting of adverse events twentyfold in clinical practice. First results from the Danish Database (DANBIO). *Scand J Rheumatol*. 2005;34:40–44.
2. Scheffe DB, Hetland ML. An open-source, self-explanatory touch screen in routine care. Validity of filling in the bath measures on ankylosing spondylitis disease activity index, function index, the health assessment questionnaire and visual analogue scales in comparison with paper versions. *Rheumatology*. 2010;49:99–104.
3. Lyng E, Sandegaard JL, Rebolj M. The Danish national patient register. *Scand J Public Health Suppl*. 2011;39(7 suppl):30–33.
4. DANBIO [homepage on the Internet]. Danbio's National Clinical Quality Report 2014 (Report in Danish). Available from: www.danbio-online.dk. Accessed July 01, 2015.
5. Hetland ML. DANBIO – powerful research database and electronic patient record. *Rheumatology*. 2011;50:69–77.
6. Hetland MH, Jensen DV, Krogh NS. Monitoring patients with rheumatoid arthritis in routine care: experiences from a treat-to-target strategy using the DANBIO registry. *Clin Exp Rheumatol*. 2014;32(suppl 85):141–146.
7. Hetland ML, Christensen IJ, Tarp U, et al. All Departments of Rheumatology in Denmark. Direct comparison of treatment responses, remission rates, and drug adherence in patients with rheumatoid arthritis treated with adalimumab, etanercept, or infliximab: results from eight years of surveillance of clinical practice in the nationwide Danish DANBIO registry. *Arthritis Rheum*. 2010;62(1):22–32.
8. Ørnberg LM, Østergaard M, Boyesen P, et al. Impact of tumour necrosis factor inhibitor treatment on radiographic progression in rheumatoid arthritis patients in clinical practice: results from the nationwide Danish DANBIO registry. *Ann Rheum Dis*. 2013;72(1):57–63.
9. Grintborg B, Østergaard M, Dreyer L, et al. Treatment response, drug survival, and predictors thereof in 764 patients with psoriatic arthritis treated with anti-tumour necrosis factor α therapy: results from the nationwide Danish DANBIO registry. *Arthritis Rheum*. 2011;63(2):382–390.
10. Grintborg B, Østergaard M, Krogh NS, Dreyer L, Kristensen HL, Hetland ML. Predictors of treatment response and drug continuation in 842 patients with ankylosing spondylitis treated with anti-tumour necrosis factor: results from 8 years' surveillance in the Danish nationwide DANBIO registry. *Ann Rheum Dis*. 2010;69(11):2002–2008.
11. Dreyer L, Mellemkjær L, Andersen AR, et al. Incidences of overall and site specific cancers in TNF α inhibitor treated patients with rheumatoid arthritis and other arthritides – a follow-up study from the DANBIO registry. *Ann Rheum Dis*. 2013;72(1):79–82.
12. Cordtz R, Mellemkjær L, Grintborg B, Hetland ML, Dreyer L. Malignant progression of precancerous lesions of the uterine cervix following biological DMARD therapy in patients with arthritis. *Ann Rheum Dis*. 2015;74(7):1479–1480.
13. Højgaard P, Grintborg B, Hetland ML, et al. Association between tobacco smoking and response to tumour necrosis factor α inhibitor treatment in psoriatic arthritis: results from the DANBIO registry. *Ann Rheum Dis*. 2014;74(12):2130–2136.

Clinical Epidemiology**Dovepress****Publish your work in this journal**

Clinical Epidemiology is an international, peer-reviewed, open access, online journal focusing on disease and drug epidemiology, identification of risk factors and screening procedures to develop optimal preventative initiatives and programs. Specific topics include: diagnosis, prognosis, treatment, screening, prevention, risk factor modification,

Submit your manuscript here: <http://www.dovepress.com/clinical-epidemiology-journal>

systematic reviews, risk & safety of medical interventions, epidemiology & biostatistical methods, and evaluation of guidelines, translational medicine, health policies & economic evaluations. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use.