miR182 activates the Ras–MEK–ERK pathway in human oral cavity squamous cell carcinoma by suppressing *RASA1* and *SPRED1*

Jinhui Wang^{1,*}Wei Wang^{2,*}Jichen Li²Liji Wu²Mei Song¹Qinggang Meng³

¹Department of Clinical Laboratory, Harbin First Hospital, ²Department of Oral and Maxillofacial Surgery, The First Affiliated Hospital of Harbin Medical University, ³Department of Osteological Surgery, Harbin First Hospital, Harbin, People's Republic of China

*These authors contributed equally to this work

Purpose: The constitutive activation of the Ras–MEK–ERK signaling pathway in oral cavity squamous cell carcinoma (OSCC) has been found to be tightly controlled at multiple levels under physiological conditions. *RASA1* and *SPRED1* are two important negative regulators of this pathway, but the exact regulating mechanism remains unclear. In this study, we aimed to explore the potential regulating mechanisms involved in the Ras–MEK–ERK signaling pathway in OSCC.

Materials and methods: MicroRNA (miRNA) expression was detected by quantitative reverse-transcription polymerase chain reaction. The protein levels of *RASA1*, *SPRED1*, and signaling proteins were detected by Western blot. Cell growth was determined using CCK-8 reagent, colony formation was stained by crystal violet, and cell invasion was tested using transwell chambers. Cell apoptosis and the cell cycle were then analyzed by flow cytometry. The binding of miR182 with *RASA1* or *SPRED1* was evaluated by luciferase reporter assays on a dual-luciferase reporter system.

Results: The expression of miR182 was found to be upregulated significantly in malignant oral carcinoma tissues compared with the adjacent nonmalignant tissues, and was inversely correlated with protein levels of *RASA1* and *SPRED1*. Overexpression of miR182 in OSCC cell lines sustained Ras–MEK–ERK signaling-pathway activation, and promoted cell proliferation, cell-cycle progression, colony formation, and invasion capacity, whereas miR182 downregulation alleviated these properties significantly in vitro. Furthermore, we demonstrated that miR182 exerted its oncogenic role in OSCC by directly targeting and suppressing *RASA1* and *SPRED1*.

Conclusion: Our results bring new insights into the important role of miR182 in the activation of the Ras–MEK–ERK signaling pathway, and suggest that miR182 may be used as a potential target for treatment of OSCC, prompting further investigation into miRNA antisense oligonucleotides for cancer therapy.

Keywords: miR182, Ras–MEK–ERK, *RASA1*, *SPRED1*, OSCC

Introduction

Oral cavity squamous cell carcinoma (OSCC), the most common type of head and neck cancer, is a leading cause of cancer-related mortality worldwide. OSCC constitutes approximately 90% of all oral malignancies, which mainly occur in the tongue and the floor of the mouth.^{1,2} Despite advances in diagnostic techniques and therapeutic regimens, the prognosis of OSCC remains poor, and is marked by an average 5-year survival rate of 50%, locoregional recurrence of 25%, and distant metastasis of 25%.³ Carcinogenesis, including OSCC,⁴ is thought to be prompted by the progressive

Correspondence: Qinggang Meng
Department of Osteological Surgery,
Harbin First Hospital, 151 Diduan Street,
Daoli, Harbin, Heilongjiang 150010,
People's Republic of China
Tel/fax +86 451 8488 3136
Email qinggang_meng@sina.com

accumulation of multiple genetic abnormalities, which include the deregulation of oncogenes or tumor-suppressing genes that play crucial roles in the pathogenesis of cancer. Therefore, there is an urgent need to gain a better understanding of the molecular mechanisms underlying the development of OSCC for improving early diagnosis and identifying new therapeutic targets.

MicroRNAs (miRNAs) are small, highly conserved, noncoding RNAs that are considered to play crucial roles in various cellular processes, such as proliferation, development, differentiation, and apoptosis. These single-strand RNAs can induce direct degradation of target messenger RNAs (mRNAs) or translational inhibition by negatively regulating gene expression at the posttranscriptional level.^{5,6} A single miRNA can regulate hundreds of protein-coding genes, while more than a third of all protein-coding genes are thought to be under translational control of miRNAs.⁷

In recent years, miRNAs have been studied most intensively in the regulation of tumorigenesis, and growing evidence shows that the deregulation of certain miRNAs regulating oncogenes or tumor-suppressing genes is involved in the pathogenesis of cancers.⁸ Emerging evidence has suggested that miRNAs can regulate the cellular signaling pathways involved in tumorigenesis through targeting the 3'-untranslated region (3'UTR) of corresponding mRNAs.⁹⁻¹² Accordingly, the deregulation of certain miRNAs in OSCC has been reported to influence cell growth and development. For instance, transfection of an miR21 inhibitor to tongue SCC cell lines inhibits anchorage-independent growth, induces apoptosis, and suppresses tumor formation.¹³ miR-31 has been found to be upregulated significantly in OSCC and head and neck squamous cell carcinoma,^{14,15} and overexpression of miR-31 represses the expression of factor-inhibiting HIF, thus activating the HIF pathway.¹⁵ In addition, miR137 and miR193a inhibit cell growth by inducing apoptosis and G₁-S arrest of OSCC cells by targeting *CDK6* and *E2F6*.¹⁶

The MAPK pathway is commonly activated in various cellular processes, of which the MEK-ERK pathway is one of the most deregulated in human cancers.¹⁷ Upon stimulation, receptor tyrosine kinases translate the ligand-binding event into a signal and Ras is activated to a GTP-bound state. GTP-bound RAS (RAS-GTP) can activate the MAPKKK (Raf) family, leading to the sequential phosphorylation and activation of MAPKK (MEK) and ERK1/2. Notably, Ras is constitutively activated in various human cancer types and plays important roles in cell growth and development.¹⁸ RASA1 and SPRED1, two negative regulators of the Ras-MEK-ERK signaling pathway, have been proved to be

involved in the regulation of Ras activation. RASA1 can enhance the weak intrinsic GTPase activity of Ras, resulting in an increase in the inactive GDP-bound form of Ras, thereby leading to an aberrant intracellular signaling through the Raf-MEK-ERK pathway.¹⁹ Whereas SPRED1 interacts with neurofibromin (*NF1* gene products) and recruits it to the plasma membrane, where NF1 performs its function as an Ras GTPase-activating protein, hydrolyzing active Ras-GTP to inactive Ras-GDP.²⁰ Although mutations of *RAS* genes occur in approximately 15%–30% of all human cancers,¹⁹ it appears that activated *RAS* mutations are rarely involved in head and neck tumors.^{21,22} Therefore, further exploring the regulatory mechanisms of key components of the Ras-MEK-ERK cascade, such as RASA1 and SPRED1, would increase our knowledge of the biological basis of activation of Ras in cancer and provide novel insights for tumor therapy.

Various miRNAs have been demonstrated to target members of the Ras-MEK-ERK pathway. Therefore, deregulation of such miRNAs in cancer cells most likely contributes to tumorigenesis by leading to an aberrant activation of the Ras-MEK-ERK pathway. In the present study, miR182 was revealed as a potential regulator of *RASA1* and *SPRED1* by in silico analysis. The expression of miR182, *RASA1*, and *SPRED1* on malignant tissues and adjacent normal tissues from OSCC patients were examined by quantitative reverse-transcription polymerase chain reaction (qRT-PCR) and Western blot analysis. Direct inhibition of RASA1 and SPRED1 translation by miR182 and a potential role of MIR182 as an oncogene in OSCC tumorigenesis were experimentally validated in vitro. Notably, *RASA1* and *SPRED1* were determined to be direct targets of miR182 in regulating the Ras-MEK-ERK signaling pathway. Our data demonstrated that MIR182 functioned as an oncogene through regulating RASA1 and SPRED1, and uncovered a novel mechanism for constitutive Ras activation in OSCC.

Materials and methods

Human tongue SCC Tca8113 cells were obtained from the Cell Resource Center of Peking Union Medical College (Beijing, People's Republic of China [PRC]). Fetal bovine serum was purchased from Biological Industries (Cromwell, CT, USA). An miRNA-isolation kit, miRNA RT kit, and miRNA qRT-PCR kit were all purchased from HaiGene Inc (Harbin, PRC). A cell-cycle assay kit and annexin V-fluorescein isothiocyanate (FITC)/propidium iodide (PI) kit for apoptosis analysis were also purchased from HaiGene. The primary antibodies against RASA1, SPRED1, β -actin, and secondary antibodies were all purchased from Santa

Cruz Biotechnology Inc (Dallas, TX, USA), and primary antibodies against ERK1/2 and phospho-ERK1/2 were purchased from Cell Signaling Technology (Danvers, MA, USA). The transfection reagent Lipofectamine 2000 was obtained from Thermo Fisher Scientific (Waltham, MA, USA), and miR182 mimics, anti-miR182 oligonucleotides, and corresponding controls were all obtained from Ruibo Inc (Guangzhou, PRC). CCK-8 reagent was purchased from Dojindo (Kumamoto, Japan). Matrigel basement membrane matrix was purchased from BD (Franklin Lakes, NJ, USA). An active Ras pull-down assay kit was purchased from EMD Millipore (Billerica, MA, USA). A luciferase-activity assay kit was obtained from Promega Corporation (Fitchburg, WI, USA).

Patient samples and cell culture

Fresh malignant tissues and adjacent noncancerous tissues were collected from ten OSCC patients (tongue, gum, and floor of the mouth) who underwent surgery at Harbin First Hospital (Harbin, PRC). Written informed consent of tissue donation for research purposes and for this study were obtained from each patient, and this study was approved by the institutional review board of Harbin First Hospital. None of the patients had received chemotherapy, radiotherapy, or immunotherapy before surgery. The clinical and pathological profiles of patients are shown in Table 1. Tumor tissues from the tongue, gum, and floor of the mouth were histopathologically diagnosed and verified by experienced pathologists. Adjacent noncancerous tissues were examined by routine histopathological analysis and confirmed to be tumor-free. Specimens were snap-frozen in liquid nitrogen immediately after surgery and stored at -80°C . Tca8113 cells were cultured in Roswell Park Memorial Institute 1640 medium supplemented with 10% fetal bovine serum, penicillin, and streptomycin (100 IU/mL) at 37°C under a 5% CO_2 atmosphere in a humidified incubator.

Table 1 Clinical and pathological data of patients in this study

Patient	Site	Sex	Age (years)	Pathologic stage
1	Tongue	Male	65	$\text{T}_2\text{N}_{2b}\text{M}_0$
2	Tongue	Male	76	$\text{T}_3\text{N}_{2c}\text{M}_0$
3	Floor of mouth	Female	48	$\text{T}_2\text{N}_1\text{M}_0$
4	Tongue	Male	67	$\text{T}_1\text{N}_{2b}\text{M}_0$
5	Gum	Female	57	$\text{T}_2\text{N}_1\text{M}_0$
6	Gum	Female	59	$\text{T}_2\text{N}_{2b}\text{M}_0$
7	Floor of mouth	Male	55	$\text{T}_1\text{N}_1\text{M}_0$
8	Tongue	Male	64	$\text{T}_3\text{N}_{2c}\text{M}_0$
9	Tongue	Male	38	$\text{T}_3\text{N}_{2b}\text{M}_0$
10	Tongue	Female	72	$\text{T}_3\text{N}_0\text{M}_0$

miRNA qRT-PCR analysis

miRNAs were isolated from the paired tissues according to the protocol provided by the manufacturer and reverse-transcribed to complementary DNA with TaqMan miRNA RT primer. Expression differentiation of miRNAs in paired tissues was analyzed by quantitative PCR analysis on a Bio-Rad Opticon 2 real-time PCR system. The specific forward and reverse primers were included in the kit. Relative expressions of miRNAs were calculated by the method of $2^{\Delta\text{Ct}_{\text{U6b}} - \text{Ct}_{\text{target miRNA}}}$, and small nuclear RNA U6b was used for normalization.

Western blot analysis

Tissue samples or cells were lysed in radioimmunoprecipitation-assay buffer supplemented with 1 mM phenylmethylsulfonyl fluoride. Protein concentrations were determined using a bicinchoninic acid protein-assay kit. Total proteins were separated with sodium dodecyl sulfate-polyacrylamide gel electrophoresis and transferred onto nitrocellulose membrane in 20 mM Tris-HCl (pH 8) containing 150 mM glycine and 20% (v/v) methanol. Membranes were blocked with 5% non-fat dry milk in Tris-buffered saline containing 0.05% Tween 20 at room temperature for 1 hour. Then, the membranes were probed with polyclonal antibodies against RASA1 (1:500), SPRED1 (1:300), ERK1/2 (1:800), or pERK1/2 (p44/42) (1:500) overnight at 4°C , and β -actin antibody (1:1,000) was used as a loading control. Then, the membranes were incubated with horseradish peroxidase-conjugated secondary antibody (1:5,000) for 1 hour. The signals were developed with enhanced chemiluminescence reagent, and digital images were captured using an LAS 4000 charge-coupled device camera system. The relative intensity of bands was analyzed with Image-Pro Plus 6.0 software.

miRNA transfection

The miR182 mimics (50 nM), negative control (NC; 50 nM), anti-miR182 oligonucleotides (100 nM), and anti-miR182 NC (100 nM) were synthesized for transfection. Anti-miR182 oligonucleotides are modified antisense oligonucleotides designed specifically to bind to and inhibit endogenous miR182 with rare off-target effect. The corresponding miRNA sequences were: miR-NC, 5'-AUAUGACGUACGUGUAACGUACUC-3'; miR182 mimics, 5'-UUUGGCAAUGGUAGAACUCACACU-3'; anti-miR NC, 5'-UCCGAGUGCUAUACGCUAGUAAAU-3'; and anti-miR182 oligonucleotides, 5'-AGUGUGAGUUCUACCAUUGCCAAA-3'. Transfections with miR182 mimics, anti-miR182 oligonucleotides, or controls were

performed using Lipofectamine 2000 reagent in Tca8113 according to the manufacturer's instructions.

Cell-proliferation assay

Cell-proliferation assays were conducted using CCK-8 reagent: 10^4 Tca8113 cells per well were seeded in 96-well plates and incubated overnight. Tca8113 cells were transfected with miR182 mimics, anti miR182 oligonucleotides, or controls using Lipofectamine 2000. The cells were incubated with 10 μ L of CCK-8 reagent for 2 hours at 24, 36, 48, 60, and 72 hours after transfection, and then absorbance at 450 nm was measured at each time point. Assays were conducted in three replicate wells for each sample, and the experiment was performed in triplicate.

Colony-formation assay

Tca8113 cells (500 cells per well) were collected and seeded in six-well plates after transfection for 48 hours, and continued to culture for 14 days. Cells were subsequently fixed with methanol (20% v/v) and stained with crystal violet (0.2% w/v) for 5 minutes. Colonies with more than 50 cells per colony were counted. The experiments were repeated in triplicate to get standard deviations.

Cell-invasion assay

The invasive capacity of cells was tested using BD transwell invasion chambers. Transwell chambers were covered with Matrigel beforehand, and Tca8113 cells (10^5 per chamber) suspended in 100 μ L serum-free medium were added to the upper chambers. The lower chambers were filled with complete growth medium (Roswell Park Memorial Institute 1640+10% fetal bovine serum). After incubation for 48 hours, cells on the upper surface of the membrane were wiped off using a cotton swab. Then, the invaded cells were fixed in 4% formaldehyde for 30 minutes and stained with 0.1% crystal violet for 5 minutes. Finally, the cells were counted and imaged under a microscope. The experiment was performed in triplicate.

Cell-cycle analysis

Cells (0.5×10^6 per well) were harvested from six-well plates and then fixed in 75% ice-cold ethanol. The fixed cells were subsequently treated with a cell-cycle assay kit according to the manufacturer's manual protocol. Briefly, the fixed cells were resuspended in phosphate-buffered saline at a concentration of 3×10^5 /mL. Then, the cells were treated with RNase A for 20 minutes, followed by incubation with PI for 15 minutes. Cell-cycle analysis was performed using Guava® EasyCyte flow cytometry with GuavaSoft 2.5. Data were

analyzed with FlowJo software, and cell-cycle distribution was shown as the percentage of cells in the G₁, S, and G₂ populations. The experiment was performed three times to get standard deviations.

Cell-apoptosis analysis

Apoptosis analysis was performed using an annexin V-FITC/PI kit according to the manufacturer's instructions on Guava flow cytometry with GuavaSoft 2.5. Cells (0.5×10^6) were harvested from six-well plates and resuspended in binding buffer at a concentration of 10^5 /mL. Cells were subsequently incubated with annexin V-FITC and PI for 15 minutes at room temperature. Then, cells were detected on flow cytometry and the data analyzed with the FlowJo software. The percentage of cells that were annexin V-FITC-positive but PI-negative was compared among different treatment groups. Each experiment was performed in triplicate.

Active Ras pull-down assay

Active Ras was determined using an Ras-assay reagent, a glutathione *S*-transferase fusion protein corresponding to the Ras-binding domain of Raf1. According to the manufacturer's instructions, Tca8113 cells were seeded in 100 mm dishes at 2×10^6 per dish. After transfection with miR182 mimics or anti-miR182 oligonucleotides for 48 hours, the cells were lysed with lysis buffer. Then, cell lysates were centrifuged at $14,000 \times g$ at 4°C, and the supernatant was used as the sample for subsequent Ras pull-down assay. A protein sample (1 mg) was mixed with 10 μ g Raf1 Ras-binding domain agarose beads, and incubated for 45 minutes at 4°C. The beads were collected by centrifugation and resuspended in 40 μ L reducing sample buffer, followed by boiling at 95°C for 5 minutes. Next, the samples of each pull-down reaction group were subjected to Western blot analysis using anti-Ras antibody (1:2,000 dilution).

Luciferase activity assay

Partial lengths of the 3'UTR of *RASAI* (GenBank NM_002890.2) and *SPRED1* (GenBank NM_152594.2) genes were synthesized and inserted downstream of the luciferase reporter gene in a PmirGLO vector. For luciferase reporter assays, cells were seeded in 24-well plates and incubated for 24 hours before transfection. Firefly luciferase constructs containing the 3'UTR of the potential miR182 target and pRL-TK *Renilla* luciferase-normalization control, miR182 mimics, or anti-miR182 oligonucleotides and NCs were cotransfected into Tca8113 cells using Lipofectamine 2000. Lysates were collected 48 hours after

transfection and measured using a dual-luciferase reporter system according to the manufacturer's protocol. Three independent experiments were performed, and the data are presented as mean $\pm$ standard deviation.

Statistical analysis

All experiments were done in triplicate, and results are presented as mean $\pm$ standard deviation. Comparisons of the data among different groups were performed by paired *t*-tests and one-way analysis of variance using SPSS software version 13.0. $P < 0.05$ was considered statistically significant.

Results

miR182 expression is elevated in OSCC patients and inversely correlated with the expression of *RASA1* and *SPRED1*

Since a single miRNA can affect hundreds of different protein-coding genes and one specific mRNA can be targeted by multiple miRNAs, we attempted to identify the miRNAs that activate Ras–MEK–ERK signaling pathway in OSCC through targeting *RASA1* and *SPRED1*. To gain new insight into the mechanisms of activation of the Ras–MEK–ERK cascade in OSCC, the prediction algorithm of TargetScan was used to identify the potential miRNAs regulating both *RASA1* and *SPRED1*. The 24 overlapped miRNAs were thought to be potential target miRNAs according to the prediction algorithm (Figure 1).

Then, expression of the 24 predicted miRNAs was screened in malignant tissues and paired nonmalignant tissues of ten OSCC patients through a TaqMan qPCR-based method (for patient information, refer to Table 1).

Figure 1 Shared putative microRNAs potentially regulating both *RASA1* and *SPRED1* are identified by the prediction algorithm of TargetScan.

Notes: Based on the algorithm of TargetScan, there were 80 miRNAs that were predicted to only regulate *SPRED1*, 14 miRNAs to only regulate *RASA1*, and overlapped 24 miRNAs to regulate both *SPRED1* and *RASA1*.

The results showed that four miRNAs – miR31, miR182, miR200a, and miR141 – were significantly upregulated, while three miRNAs – miR1, miR206, and miR212 – were downregulated in OSCC tumor tissues compared with paired noncancerous tissues (Figure 2A), but there was no significant differentiation in the expression of other miRNAs (containing four miRNAs undetected). Among these, miR31 was the most significantly upregulated (4.27-fold upregulation) in OSCC tumor samples, and miR182, miR200a, and miR141 were upregulated to 2.87-fold, 2.4-fold, and 1.7-fold, respectively (Table 2). Notably, the upregulation of miR31 has been intensively investigated in a variety of cancers, and thus miR182 was our research focus, and the expression differentiation of miR182 between the pairing tissues was matched (Figure 2B).

Based on the sequence alignment, it was found that there was a perfect base pairing between the seed of miR182 and 3'UTR of *RASA1* or *SPRED1*. To identify further correlations among miR182 and *RASA1* and *SPRED1*, Western blot was performed to compare the expression of *RASA1* and *SPRED1* in malignant OSCC tissues and adjacent normal tissues. The results showed that expression of *RASA1* and *SPRED1* was dramatically downregulated in OSCC tumor tissues (Figure 3A), which indicated that miR182 was inversely correlated with the expression of *RASA1* and *SPRED1* (Figure 3B). As such, we speculated that miR182 might be an oncogene and *RASA1* and *SPRED1* would be the potential targets of miR182 in regulating the tumorigenesis and progression of OSCC.

miR182 promotes OSCC cell proliferation and cell-cycle progression

To elucidate the oncogenic role of miR182 in OSCC, Tca8113 cells were used for further investigation. Overexpression of miR182 was achieved by transfection of miR182 mimics, whereas downregulation of miR182 was achieved by transfection of anti-miR182 oligonucleotides. Similarly, control experiments were performed by transfection of miR-NC and anti-miR-NC.

Cell proliferation was determined using CCK-8 reagent at 24, 36, 48, 60, and 72 hours after transfection. As shown in Figure 4A, the overexpression of miR182 significantly increased the growth rate of Tca8113 cells compared with the miR-NC group; however, downregulation of miR182 led to a decrease in cell viability. To prove further the proproliferating effect of miR182 in OSCC cells, a colony-formation assay was performed. The results showed that overexpression of miR182 increased the colony numbers of Tca8113

Figure 2 Expression differentiation of predicted miRNAs between malignant tissues and adjacent nonmalignant tissues from ten OSCC patients.

Notes: (A) Heat map generated by cluster analysis of the predicted miRNAs potentially regulating *RASA1* and *SPRED1* in OSCC patients based on the data of TaqMan qRT-PCR; (B) expression of miR182 matched between ten paired OSCC tissues (** $P < 0.01$).

Abbreviations: T, malignant tumor tissues; N, adjacent nonmalignant tissues; miRNAs, microRNAs; OSCC, oral cavity squamous cell carcinoma; qRT-PCR, quantitative reverse-transcription polymerase chain reaction.

cells, and the colonies were much larger than the NC, while downregulation of miR182 depressed colony formation (Figure 4B). To understand further the proproliferating mechanism, the effects of miR182 on the cell cycle were investigated using flow cytometry. The data showed that there was a dramatic decrease in the percentage of cells in the G_1/G_0 phase (6.9%) and an increase in cells in the S phase (7.1%) in miR182-overexpressing Tca8113 cells compared with NC cells. In contrast, there was a significant increase in the percentage of cells in the G_1/G_0 phase (7.2%) and a decrease in the percentage of cells in the S phase (6.3%) in

miR182 downregulation of Tca8113 cells compared with anti-miR-NC cells (Figure 4C). As such, overexpression of miR182 promoted proliferation and cell-cycle progression of OSCC cells in vitro.

miR182 inhibits OSCC cell apoptosis

Since the proproliferating role of miR182 had been proved in OSCC cells, its involvement in cell apoptosis was further investigated by annexin V-FITC/PI costaining. The results showed that miR182-overexpressing cells had a lower apoptosis rate than miR-NC cells (3.45% vs 7.7%). On the contrary, the apoptosis rate was dramatically increased by downregulation of miR182 by transfection of anti-miR182 oligonucleotides (10.24% vs 7.53%) (Figure 5). The results indicated that overexpression of miR182 inhibited the apoptosis of OSCC cells, whereas inhibition of miR182 enhanced Tca8113 apoptosis.

miR182 enhances invasive capacity of OSCC cells

To study further the effects of miR182 on the invasive capacity of OSCC cells, a Matrigel invasion assay was performed using transwell chambers. Figure 6 shows that overexpression of miR182 significantly increased the invasive potential of Tca8113 cells compared with miR-NC cells. In contrast, downregulation of miR182 resulted in dramatically decreased

Table 2 Deregulated miRNAs potentially regulating *RASA1* and *SPRED1* between ten paired OSCC malignant tissues and adjacent noncancerous tissues

miRNA	Fold-change	P-value
miR31	4.27	0.0018
miR182	2.87	0.0009
miR200a	2.4	0.0039
miR141	1.7	0.0246
miR1	-4.46	0.0020
miR206	-2.73	0.0092
miR212	-1.96	0.0237

Notes: Fold-change values represented by ratio of mean values of malignant groups and normal groups by the method of $2^{-\Delta\Delta Ct}$. Negative fold-change values indicate overexpression in adjacent normal tissues. P-values indicate the significance level for each miRNA analyzed using paired t-test.

Abbreviations: miRNAs, microRNAs; OSCC, oral cavity squamous cell carcinoma.

Figure 3 The reverse correlation of miR182 and RASA1 or SPRED1 protein levels.

Notes: (A) Expression of RASA1 and SPRED1 between paired OSCC tissues was examined by Western blot and was normalized by β -actin; (B) correlation between miR182 and RASA1 or SPRED1 represented by relative expression.

Abbreviations: T, malignant tumor tissues; N, adjacent nonmalignant tissues; OSCC, oral cavity squamous cell carcinoma.

invasive cell numbers. The results indicated that overexpression of miR182 enhanced invasion of OSCC cell in vitro.

miR182 promotes activation of the Ras–MEK–ERK signaling pathway by regulating RASA1 and SPRED1

Since miR182 was predicted to be a putative miRNA regulating *RASA1* and *SPRED1* by in silico analysis and was found to be inversely correlated with the expression of RASA1 and SPRED1 based on the data from ten OSCC patients, we attempted to investigate whether miR182 directly regulates the expression of RASA1 and SPRED1. Western blot was performed to examine the expression differentiations of RASA1 and SPRED1 by upregulating or downregulating miR182 in Tca8113 cells. The results showed that RASA1 and SPRED1 expression was significantly decreased in miR182-overexpressing cells compared with miR-NC cells, whereas they were significantly increased in miR182-downregulated cells (Figure 7). Since RASA1 and SPRED1 are two important negative regulators that directly influence

the activation of GTP-Ras in the Ras–MEK–ERK signaling pathway, GTP-Ras and phospho-ERK1/2 were determined by Western blot after transfection of miR182 mimics or anti-miR182 oligonucleotides. The results revealed a significant increase in GTP-Ras and phospho-ERK1/2 by upregulation of miR182 (Figure 7), which indicated that miR182 played a crucial role in activating the Ras–MEK–ERK signaling cascade. These results suggested that miR182 promotes activation of the Ras–MEK–ERK signaling pathway by suppressing RASA1 and SPRED1.

miR182 represses the translation of RASA1 and SPRED1 by directly targeting their 3'UTRs

Given that Western blot results had revealed that expression of RASA1 and SPRED1 was decreased significantly in miR182-overexpressing cells and increased in miR182-downregulated cells, we hypothesized that *RASA1* and *SPRED1* might be direct downstream targets of miR182. To confirm our hypothesis, we first aligned the sequence of

Figure 4 miR182 promotes OSCC cell proliferation and cell-cycle progression.

Notes: Tca8113 cells were transfected with miR-NC, miR182 mimics, anti-miR-NC, and anti-miR182 oligonucleotides, and divided into four groups. (A) Effects of miR182 on the proliferation of Tca8113 cells were examined at 24, 36, 48, 60, and 72 hours after transfection using a CCK-8 assay kit. (B) Representative micrograph of each group's Tca8113 cell colonies determined by colony-formation assay. (C) Effects of miR182 on cell-cycle progression analyzed by flow cytometry and quantification of cell percentages in G₀-G₁, S, and G₂-M phases. Different colors represented the cells in different phases of cell cycles: green, G₁ phase; yellow, S phase; blue, G₂ phase. **P*<0.05 vs miR-NC; ***P*<0.01 vs miR-NC; ****P*<0.01 vs anti-miR-NC.

Abbreviations: Oligos, oligonucleotides; OSCC, oral cavity squamous cell carcinoma; NC, negative control; OD, optical density.

Figure 5 miR182 inhibits the apoptosis of OSCC cells.

Notes: Tca8113 cells were transfected with miR182 mimics, miR-NC, anti-miR182 oligonucleotides, or anti-miR-NC for 48 hours. Cell apoptosis was detected by flow cytometry by annexin V-FITC/PI staining. The experiment was performed in triplicate. ** $P < 0.01$ vs miR-NC; # $P < 0.05$ vs anti-miR-NC.

Abbreviations: Oligos, oligonucleotides; OSCC, oral cavity squamous cell carcinoma; NC, negative control; FITC, fluorescein isothiocyanate; PI, propidium iodide.

miR182 with the 3'UTR of *RASA1* and *SPRED1*. As shown in Figure 8A, there were two miR182-binding sites in the *RASA1* mRNA 3'UTR and one binding site in the *SPRED1* mRNA 3'UTR, and miR182 was predicted to bind to them strongly, due to good base pairing. Next, luciferase-binding assays were conducted using the PmirGLO luciferase plasmid to determine whether miR182 suppressed *RASA1* and *SPRED1* through directly binding to their 3'UTRs. The partial length of wild-type *RASA1* or *SPRED1* 3'UTRs were

synthesized and cloned into the PmirGLO luciferase reporter plasmid. Then, the constructed luciferase reporter plasmids were cotransfected with either miR182 mimics, anti-miR182 oligonucleotides, or controls into Tca8113 cells. As shown in Figure 8B, the luciferase activity of wild-type *RASA1* and *SPRED1* decreased by 6.33-fold and 3.41-fold, respectively, by transfection of miR182 mimics compared with miR-NC. These results indicated that *RASA1* and *SPRED1* were direct targets of miR182 in OSCC.

Figure 6 miR182 promotes the invasion of OSCC cells.

Notes: Invasion assays were performed using Tca8113 cells transfected with miR182 mimics, anti-miR182 oligonucleotides, or controls. Representative images of Tca8113 cells for the four groups were captured. Magnification $\times 200$. ** $P < 0.01$ vs miR-NC; ## $P < 0.01$ vs anti-miR-NC.

Abbreviations: Oligos, oligonucleotides; OSCC, oral cavity squamous cell carcinoma; NC, negative control.

Figure 7 miR182 promotes the activation of the Ras–MEK–ERK signaling pathway by suppressing the expression of RASA1 and SPRED1.

Notes: Western blot was performed to examine the expression differentiations of RASA1, SPRED1, Ras-GTP, and phospho-ERK1/2 in Tca8113 cells after transfection with miR182 mimics or anti-miR182 oligonucleotides. Relative band intensities were analyzed by Image-Pro Plus 6.0, Ras was used for normalization of Ras-GTP, and β-actin was used for others. ** $P < 0.01$ vs miR-NC; # $P < 0.05$ vs anti-miR-NC; ## $P < 0.01$ vs anti-miR-NC.

Abbreviations: NC, negative control; p, phospho.

Discussion

Deregulation of miRNAs has been demonstrated to be involved in tumorigenesis and progression in various types of cancers; however, elucidation of their potential roles in OSCC remains relatively rare. In the present study, it was

found that miR182 was significantly upregulated in OSCC carcinoma tissues compared with adjacent noncancerous tissues, which was consistent with previous data on differential miRNA-expression profiles in OSCC.²³ Overexpression of miR182 in OSCC cell lines sustained Ras–MEK–ERK

Figure 8 miR182 regulates RASA1 and SPRED1 by directly targeting their 3'UTRs.

Notes: (A) Schematic representation of mature human miR182 sequence and miR182 target-binding sites in the 3'UTRs of RASA1 and SPRED1 mRNAs. (B) For luciferase-activity assays, 3'UTR of RASA1 or SPRED1 was cloned into the pMirGlo reporter plasmid separately, and the constructed plasmid was cotransfected with miR182 mimics, miR-NC, anti-miR182 oligonucleotides, or anti-miR-NC individually. The normalized activity of the reporters relative to Renilla luciferase was analyzed 48 hours after transfection. ** $P < 0.01$ vs miR-NC; # $P < 0.05$ vs anti-miR-NC.

Abbreviations: UTRs, untranslated regions; mRNAs, messenger RNAs; NC, negative control.

signaling pathway activation by suppressing *RASA1* and *SPRED1*. Moreover, introduction of anti-miR182 oligonucleotides strikingly repressed the oncogenic properties of miR182, represented by inhibiting cell proliferation, colony formation, invasion, and promoting cell apoptosis. Therefore, our results provide new insights into the important role of miR182 in activation of the Ras–MEK–ERK signaling pathway, and demonstrate that miR182 functions as an oncogenic miRNA in OSCC.

The highly conserved Ras–MEK–ERK signaling pathway is involved in a wide range of cellular processes, including cell growth, proliferation, differentiation, survival, and senescence. This pathway is aberrantly activated in a variety of human cancers. Previous research has mostly focused on the somatic mutations in genes encoding Ras/Erk members that frequently occur in cancers, of which the *RAS* mutation is the most prevalent event in human cancers. Aberrant *RAS* function is closely associated with a single mutation, typically at codon 12, 13, or 61.²⁴ Mutation at these conserved sites favors GTP binding, and leads to constitutive activation of the Ras–MEK–ERK pathway. *RAS* plays essential roles in controlling the activity of multiple downstream effectors of pathways responsible for cellular proliferation and survival.¹⁸ However, *RAS* mutation in oral carcinomas is not so common as in other cancers, such as lung cancer and colorectal cancer. *RAS* mutation has been reported in 20%–35% of cases from India,²⁵ whereas lower frequencies (4%) were reported from the UK and no mutation reported from the US.^{26,27} Therefore, it is likely that mechanisms other than *RAS* mutation contribute to the constitutive activation of the Ras–MEK–ERK cascade in oral carcinomas.

The Ras protein cycles between the inactive GDP-bound and the active GTP-bound state, and the GTPase activity of normal Ras p21 is mediated by two negative regulators: *RASA1* and *SPRED1*. A number of studies have shown that deregulation of *RASA1* has an oncogenic effect in multiple types of human cancer. For instance, the expression of *RASA1* was repressed by miR31, and the Ras–MEK–ERK signaling pathway was activated to stimulate tumorigenesis in human colorectal cancer.²⁸ Downregulation of *RASA1* by miR31 was also found to promote cellular proliferation and inhibit cellular apoptosis in intrahepatic cholangiocarcinoma by activating the Ras–MAPK signaling pathway.¹⁰ In addition, miR206/21 was illustrated to promote coordinately Ras–ERK signaling by suppressing the translation of *RASA1* and *SPRED1* in breast cancer.²⁹ Herein, we found that overexpression of miR182 significantly increased but downregulation of miR182 inhibited ERK activation by

targeting *RASA1* and *SPRED1* in OSCC. Consistently, overexpression of miR182 can enhance the aggressiveness of oral carcinoma cells. Therefore, our results indicate that miR182 plays important oncogenic roles in the tumorigenesis and progression of OSCC through activating the Ras–MEK–ERK signaling pathway.

Our results showed that the expression of miR182 was elevated in malignant tissues compared with adjacent normal tissues in OSCC. Consistent with our findings, other studies have correlated a high level of miR182 to increased pathological aggression of tumors and poor survival of patients in various human cancers. For example, miR182 was found to be frequently upregulated in human breast-tumor tissues and cell lines, and was also significantly elevated in the serum of breast cancer patients compared with healthy controls.^{30,31} Although the role of miR182 in lung cancer seems to be controversial, upregulated expression of miR182 was indeed found in lung cancer cells and tissues.^{32,33} In addition, miR182 has also proved to be upregulated in clinical tissue samples of melanoma, glioma tumors, ovarian cancer, colorectal cancer, and prostate cancer, which promoted survival, migration, invasion, and metastasis of tumor cells.^{34–38}

There are several possible causes for the deregulation of miR182 in human cancers. One possible cause is regional amplification. miR182 is a member of an miRNA cluster (miR183/96/182), a highly conserved cluster residing on an intergenic region of human chromosomes 7q31–34, and is frequently amplified in human cancers.³⁹ For example, miR182 has been confirmed to be amplified in advanced human melanoma and soft tissue sarcomas, which correlates with poor prognosis.^{34,40} Another possible mechanism may be the regulation of transcriptional factors. It has been reported that miR182 is transcriptionally activated by the Wnt-pathway effector β -catenin to promote colony formation and invasiveness of MDA-MB-231 breast cancer cells.⁴¹ However, overexpression of miR182 has been reported to inhibit lung cancer tumorigenicity both in vitro and in vivo. In addition, miR182 exerts antiproliferative activity and inhibits cell invasion by targeting the G protein GNA13 in prostate cancer-cell lines.^{42,43} Since each single miRNA can target multiple mRNAs and be involved in various signaling pathways to exert diversified functions, it is reasonable to speculate that the molecular routes through which miR182 exerts its regulatory effects are largely context-dependent, which helps in understanding the pluripotency of this miRNA in cancer.

Upregulation of miR182 was demonstrated to promote cell proliferation and colony formation in Tca8113 cells.

The cell-cycle analysis by flow cytometry showed that overexpression of miR182 led to a significant decrease in the percentage of cells in the G₀/G₁ phase and a significant increase in the percentage of cells in the S phase. It has been reported that Ras can regulate cell-cycle progression by modulating transcriptional factors in diverse cell types and act at different phases of the cell cycle.⁴⁴ Therefore, it is likely that miR182 controls the cell cycle through regulating Ras by targeting *RASA1* and *SPRED1*. Moreover, elevation of miR182 was also found to promote cell invasion in Tca8113. In fact, miR182 has been proved to be correlated with metastasis in melanoma and sarcomas,^{34,40,45} which establishes a novel regulating mechanism for tumor metastasis. miR182 is overexpressed in human metastatic melanoma-cell lines and metastatic melanoma, and its overexpression can enhance the metastatic behavior of melanoma cells in vivo by converging onto FOXO3 and MITF inactivation. Besides, miR182 was found to be markedly overexpressed in sarcomas with lung metastasis, and downregulation of miR182 substantially inhibited, while elevated miR182 considerably accelerated the rate of lung metastasis of sarcomas. Furthermore, miR182 has been identified as a promising predictor of aggressive phenotype and an independent prognostic biomarker to predict the survival rate of patients following surgery in colorectal cancer, prostate cancer, and metastatic melanoma. Nevertheless, larger cohorts of patients with OSCC with complete clinical data, including staging, therapy, and disease-free survival, should be studied to determine whether miR182 can be a predictor of diagnosis and prognosis in OSCC.

Conclusion

In conclusion, our study identified miR182 as an oncogenic miRNA that exerts its regulatory effects on the Ras–MEK–ERK signaling pathway by inhibiting *RASA1* and *SPRED1* in OSCC. Therefore, miR182, *RASA1*, and *SPRED1* might be used as potential therapeutic targets for the treatment of OSCC.

Acknowledgment

This work was supported by the Foundation of Harbin Talent Introduction Project (grant 2014SYRCYJ03).

Author contributions

All authors contributed toward data analysis, drafting and critically revising the paper, and agree to be accountable for all aspects of the work.

Disclosure

The authors report no conflicts of interest in this work.

References

1. Rapidis AD, Gullane P, Langdon JD, Lefebvre JL, Scully C, Shah JP. Major advances in the knowledge and understanding of the epidemiology, aetiopathogenesis, diagnosis, management and prognosis of oral cancer. *Oral Oncol.* 2009;45(4–5):299–300.
2. Scully C, Bagan J. Oral squamous cell carcinoma overview. *Oral Oncol.* 2009;45(4–5):301–308.
3. Bozec A, Peyrade F, Fischel JL, Milano G. Emerging molecular targeted therapies in the treatment of head and neck cancer. *Expert Opin Emerg Drugs.* 2009;14(2):299–310.
4. Scully C, Field JK, Tanzawa H. Genetic aberrations in oral or head and neck squamous cell carcinoma (SCCHN) – I: carcinogen metabolism, DNA repair and cell cycle control. *Oral Oncol.* 2000;36(3):256–263.
5. Bartel DP. MicroRNAs: genomics, biogenesis, mechanism, and function. *Cell.* 2004;116(2):281–297.
6. He L, Hannon GJ. MicroRNAs: small RNAs with a big role in gene regulation. *Nat Rev Genet.* 2004;5(7):522–531.
7. Gomes CC, Gomez RS. MicroRNA and oral cancer: future perspectives. *Oral Oncol.* 2008;44(10):910–914.
8. Garzon R, Calin GA, Croce CM. MicroRNAs in cancer. *Annu Rev Med.* 2009;60:167–179.
9. Jiang L, Yu L, Zhang X, et al. miR-892b silencing activates NF-κB and promotes aggressiveness in breast cancer. *Cancer Res.* 2016;76(5):1101–1111.
10. Hu C, Huang F, Deng G, Nie W, Huang W, Zeng X. miR-31 promotes oncogenesis in intrahepatic cholangiocarcinoma cells via the direct suppression of *RASA1*. *Exp Ther Med.* 2013;6(5):1265–1270.
11. Yan B, Guo Q, Nan XX, et al. Micro-ribonucleic acid 29b inhibits cell proliferation and invasion and enhances cell apoptosis and chemotherapy effects of cisplatin via targeting of DNMT3b and AKT3 in prostate cancer. *Onco Targets Ther.* 2015;8:557–565.
12. Zhan Y, Xiang F, Wu R, et al. MiRNA-149 modulates chemosensitivity of ovarian cancer A2780 cells to paclitaxel by targeting MyD88. *J Ovarian Res.* 2015;8:48.
13. Li J, Huang H, Sun L, et al. MiR-21 indicates poor prognosis in tongue squamous cell carcinomas as an apoptosis inhibitor. *Clin Cancer Res.* 2009;15(12):3998–4008.
14. Liu CJ, Kao SY, Tu HF, Tsai MM, Chang KW, Lin SC. Increase of microRNA miR-31 level in plasma could be a potential marker of oral cancer. *Oral Dis.* 2010;16(4):360–364.
15. Liu CJ, Tsai MM, Hung PS, et al. miR-31 ablates expression of the HIF regulatory factor FIH to activate the HIF pathway in head and neck carcinoma. *Cancer Res.* 2010;70(4):1635–1644.
16. Kozaki K, Imoto I, Mogi S, Omura K, Inazawa J. Exploration of tumor-suppressive microRNAs silenced by DNA hypermethylation in oral cancer. *Cancer Res.* 2008;68(7):2094–2105.
17. Santarpia L, Lippman SM, El-Naggar AK. Targeting the MAPK-RAS-RAF signaling pathway in cancer therapy. *Expert Opin Ther Targets.* 2012;16(1):103–119.
18. Downward J. Targeting RAS signalling pathways in cancer therapy. *Nat Rev Cancer.* 2003;3(1):11–22.
19. Clark GJ, Westwick JK, Der CJ. p120 GAP modulates Ras activation of Jun kinases and transformation. *J Biol Chem.* 1997;272(3):1677–1681.
20. Stowe IB, Mercado EL, Stowe TR, et al. A shared molecular mechanism underlies the human rasopathies Legius syndrome and neurofibromatosis-1. *Genes Dev.* 2012;26(13):1421–1426.
21. Coutinho CM, Bassini AS, Gutierrez LG, et al. Genetic alterations in Ki-Ras and Ha-Ras genes in juvenile nasopharyngeal angiofibromas and head and neck cancer. *Sao Paulo Med J.* 1999;117(3):113–120.

22. Sathyan KM, Nalinakumari KR, Kannan S. H-Ras mutation modulates the expression of major cell cycle regulatory proteins and disease prognosis in oral carcinoma. *Mod Pathol*. 2007;20(11):1141–1148.
23. Severino P, Brüggemann H, Andreghetto FM, et al. MicroRNA expression profile in head and neck cancer: HOX-cluster embedded microRNA-196a and microRNA-10b dysregulation implicated in cell proliferation. *BMC Cancer*. 2013;13:533.
24. Prior IA, Lewis PD, Mattos C. A comprehensive survey of Ras mutations in cancer. *Cancer Res*. 2012;72(10):2457–2467.
25. Das N, Majumder J, DasGupta UB. Ras gene mutations in oral cancer in eastern India. *Oral Oncol*. 2000;36(1):76–80.
26. Chang SE, Bhatia P, Johnson NW, et al. Ras mutations in United Kingdom examples of oral malignancies are infrequent. *Int J Cancer*. 1991;48(3):409–412.
27. Xu J, Gimenez-Conti IB, Cunningham JE, et al. Alterations of p53, cyclin D1, Rb, and H-Ras in human oral carcinomas related to tobacco use. *Cancer*. 1998;83(2):204–212.
28. Sun D, Yu F, Ma Y, et al. MicroRNA-31 activates the RAS pathway and functions as an oncogenic microRNA in human colorectal cancer by repressing RAS p21 GTPase activating protein 1 (RASA1). *J Biol Chem*. 2013;288(13):9508–9518.
29. Sharma SB, Lin CC, Farrugia MK, et al. MicroRNAs 206 and 21 cooperate to promote RAS-extracellular signal-regulated kinase signaling by suppressing the translation of RASA1 and SPRED1. *Mol Cell Biol*. 2014;34(22):4143–4164.
30. Guttilla IK, White BA. Coordinate regulation of FOXO1 by miR-27a, miR-96, and miR-182 in breast cancer cells. *J Biol Chem*. 2009;284(35):23204–23216.
31. Li P, Sheng C, Huang L, et al. MiR-183/-96/-182 cluster is up-regulated in most breast cancers and increases cell proliferation and migration. *Breast Cancer Res*. 2014;16(6):473.
32. Roa W, Brunet B, Guo L, et al. Identification of a new microRNA expression profile as a potential cancer screening tool. *Clin Invest Med*. 2010;33(2):E124.
33. Cho WC, Chow AS, Au JS. Restoration of tumour suppressor hsa-miR-145 inhibits cancer cell growth in lung adenocarcinoma patients with epidermal growth factor receptor mutation. *Eur J Cancer*. 2009;45(12):2197–2206.
34. Segura MF, Hanniford D, Menendez S, et al. Aberrant miR-182 expression promotes melanoma metastasis by repressing FOXO3 and microphthalmia-associated transcription factor. *Proc Natl Acad Sci U S A*. 2009;106(6):1814–1819.
35. Jiang L, Mao P, Song L, et al. miR-182 as a prognostic marker for glioma progression and patient survival. *Am J Pathol*. 2010;177(1):29–38.
36. Liu Z, Liu J, Segura MF, et al. MiR-182 overexpression in tumorigenesis of high-grade serous ovarian carcinoma. *J Pathol*. 2012;228(2):204–215.
37. Liu H, Du L, Wen Z, et al. Up-regulation of miR-182 expression in colorectal cancer tissues and its prognostic value. *Int J Colorectal Dis*. 2013;28(5):697–703.
38. Hirata H, Ueno K, Shahryari V, et al. MicroRNA-182-5p promotes cell invasion and proliferation by down regulating FOXF2, RECK and MTSS1 genes in human prostate cancer. *PLoS One*. 2013;8(1):e55502.
39. Weeraratne SD, Amani V, Teider N, et al. Pleiotropic effects of miR-183–96–182 converge to regulate cell survival, proliferation and migration in medulloblastoma. *Acta Neuropathol*. 2012;123(4):539–552.
40. Larramendy ML, Tarkkanen M, Blomqvist C, et al. Comparative genomic hybridization of malignant fibrous histiocytoma reveals a novel prognostic marker. *Am J Pathol*. 1997;151(4):1153–1161.
41. Chiang CH, Hou MF, Hung WC. Up-regulation of miR-182 by β -catenin in breast cancer increases tumorigenicity and invasiveness by targeting the matrix metalloproteinase inhibitor RECK. *Biochim Biophys Acta*. 2013;1830(4):3067–3076.
42. Peng X, Li W, Yuan L, Mehta RG, Kopelovich L, McCormick DL. Inhibition of proliferation and induction of autophagy by atorvastatin in PC3 prostate cancer cells correlate with downregulation of Bcl2 and upregulation of miR-182 and p21. *PLoS One*. 2013;8(8):e70442.
43. Rasheed SA, Teo CR, Beillard EJ, Voorhoeve PM, Casey PJ. MicroRNA-182 and microRNA-200a control G-protein subunit α -13 (GNA13) expression and cell invasion synergistically in prostate cancer cells. *J Biol Chem*. 2013;288(11):7986–7995.
44. Winston JT, Coats SR, Wang YZ, Pledger WJ. Regulation of the cell cycle machinery by oncogenic Ras. *Oncogene*. 1996;12(1):127–134.
45. Sachdeva M, Mito JK, Lee CL, et al. MicroRNA-182 drives metastasis of primary sarcomas by targeting multiple genes. *J Clin Invest*. 2014;124(10):4305–4319.

OncoTargets and Therapy

Publish your work in this journal

OncoTargets and Therapy is an international, peer-reviewed, open access journal focusing on the pathological basis of all cancers, potential targets for therapy and treatment protocols employed to improve the management of cancer patients. The journal also focuses on the impact of management programs and new therapeutic agents and protocols on

Submit your manuscript here: <http://www.dovepress.com/oncotargets-and-therapy-journal>

patient perspectives such as quality of life, adherence and satisfaction. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Dovepress