The role of KPN β_1 in neuro-oncology

Megan Lilley^{1,2}
Faris Farassati^{1,3}

¹Midwest Biomedical Research Foundation, Kansas City Veterans Affairs Medical Center, ²School of Medicine, University of Missouri, ³Saint Luke's Cancer Institute–Saint Luke's Marion Bloch Neuroscience Institute, Kansas City, MO, USA

KPN β_1 , also known as importin- β , is part of the karyopherin superfamily of nuclear transport proteins.¹ The classical importin- α/β system is believed to import up to half of the nuclear traffic.² Importins and exportins transport pro-oncogenic mediators across the nuclear membrane and are found to be overexpressed in a number of cancer types, including breast, colon, esophageal, gastric, lung, lymphoma, melanoma, pancreatic, and prostate cancer.³ The nuclear pore complex (NPC) is composed of approximately 30 proteins (nucleoporins) that are arranged octagonally around a central channel. KPN β_1 participates in the classical nuclear import pathway alongside its adaptor protein – importin- α . Importin- α first recognizes and binds the cytoplasmic cargo via its nuclear localization signals, and then associates with importin- β via the importin- β binding (IBB) domain. This complex (nuclear localization signals–importin- α –importin- β) then traverses the nuclear pore complex.⁴ This active transport is able to occur against the concentration gradient, due to varying levels of Ran, a GTPase.⁴

KPN β_1 has recently been shown to regulate proliferation of human glioma cells via the Wnt- β -catenin pathway.⁵ Glioblastoma multiforme is the most frequent brain cancer in adults and is highly infiltrative.⁶ Despite current treatments of neurosurgery and chemoradiotherapy, median survival remains around 14.6 months, and 5-year survival is <5%.^{5,7} Lu et al showed that the relative expression of KPN β_1 correlated with the World Health Organization (WHO) grades of human glioma, with higher expression of KPN β_1 correlating with more severe WHO glioma classification.⁵ Additionally, higher expression of KPN β_1 correlated with lower 5-year survival ratio on Kaplan–Meier survival curves.⁵ Wnts are glycoproteins that are involved in cell proliferation, differentiation, and oncogenesis, and regulate β -catenin in their pathway.⁵ KPN β_1 has recently been elucidated as a regulator of glioma-cell proliferation via the Wnt- β -catenin pathway.⁵ Down-regulation of KPN β_1 has been shown to inhibit glioma proliferation in vitro. Additionally, cells with lower levels of KPN β_1 showed decreased nuclear β -catenin, demonstrating that KPN β_1 played a role in the nuclear transport of β -catenin in the Wnt- β -catenin pathway.⁵ KPN β_1 has previously been shown to play a role in translocating β -catenin, which accelerates glioma proliferation.⁵

Another role of KPN β_1 and glioma is the transport of GLI1 into the nucleus.⁸ GLI1 was discovered in human gliomata,⁶ and is a nuclear regulator of the Hedgehog (Hh)-signaling pathway.⁸ Dysregulation of this pathway leads to aggressive tumorigenesis. Hh normally binds to and inactivates Patched (Ptc). When Ptc is inhibited, Smoothened (Smo) is released and triggers a signaling cascade that ends in nuclear localization of GLI.⁸ SuFu is an additional negative regulatory protein that anchors GLI in the cytoplasm during inactivation of the Hh-signaling pathway.⁸

KPN β_1 binds GLI1 with high affinity, and the GLI1-binding site on the N-terminus for SuFu overlaps with the GLI1-binding site for KPN β_1 . This results in competitive

Correspondence: Faris Farassati
Midwest Biomedical Research Foundation, Kansas City Veterans Affairs Medical Center, 4801 East Linwood Boulevard, Kansas City, MO 64128, USA
Email ffarassati@gmail.com

binding of GLI1 based on relative concentrations of KPN β_1 and SuFu.⁸ When KPN β_1 is bound to GLI1 (rather than SuFu bound to GLI1), GLI1 can undergo nuclear import and thereby play a role in tumorigenesis.⁸ Various studies have found different levels of GLI1 in malignant gliomata. Zhu and Lo performed genome-wide copy-number analysis on 31 glioma samples and found that 22.6% of these samples had amplified *GLI1*.⁶ Therefore, inhibition of nuclear import of GLI1 via KPN β_1 has the potential to inhibit oncogenesis in gliomata as a novel therapeutic strategy.⁸ Interestingly, KPN β_1 has also been indicated in the development of secondary brain tumors.⁹ Childhood acute lymphoblastic leukemia commonly results in treatment-related secondary brain tumors.⁹ This is due in part to cranial irradiation (and treatment with antimetabolites), though a genetic predisposition is also necessary.⁹ In a study by Edick et al, ~20% of patients developed secondary brain tumors, comprised of glioblastoma multiforme, anaplastic astrocytoma, primitive neuroectodermal tumors, and embryoplastic neuroepithelial tumors.⁹ Genetic analysis of pretreatment acute lymphoblastic leukemia blasts indicated the *KPNB1* gene, along with *STAT4*, *NFIC*, and *HNRPL* (all involved in tumor growth and trafficking), to have high significance in the development of secondary brain tumors.⁹ This supports the role of KPN β_1 in oncogenesis and cancer-cell viability and suggests its potential use as a predictive factor for secondary brain tumors. In conclusion, KPN β_1 is a promising target for

anticancer therapeutics, including a potential for inhibition of certain neurological malignancies.

Disclosure

The authors declare no conflicts of interest in this work.

References

1. van der Watt PJ, Ngarande E, Leaner VD. Overexpression of Kpn β_1 and Kpn α_2 importin proteins in cancer derives from deregulated E2F activity. *PLoS One*. 2011;6(11):e27723.
2. Chook YM, Süel KE. Nuclear import by karyopherin- β s: recognition and inhibition. *Biochim Biophys Acta*. 2011;1813(9):1593–1606.
3. Mahipal A, Malafa M. Importins and exportins as therapeutic targets in cancer. *Pharmacol Ther*. 2016;164:135–143.
4. Lott K, Cingolani G. The importin β binding domain as a master regulator of nucleocytoplasmic transport. *Biochim Biophys Acta*. 2011;1813(9):1578–1592.
5. Lu T, Bao Z, Wang Y, et al. Karyopherin- β_1 regulates proliferation of human glioma cells via Wnt/ β -catenin pathway. *Biochem Biophys Res Commun*. 2016;478(3):1189–1197.
6. Zhu H, Lo HW. The human glioma-associated oncogene homolog 1 (GLI1) family of transcription factors in gene regulation and diseases. *Curr Genomics*. 2010;11(4):238–245.
7. Wangaryattawanich P, Hatami M, Wang J, et al. Multicenter imaging outcomes study of the Cancer Genome Atlas glioblastoma patient cohort: imaging predictors of overall and progression-free survival. *Neuro Oncol*. 2015;17(11):1525–1537.
8. Szczepny A, Wagstaff KM, Dias M, et al. Overlapping binding sites for importin β_1 and suppressor of fused (SuFu) on glioma-associated oncogene homologue 1 (Gli1) regulate its nuclear localization. *Biochem J*. 2014;461(3):469–476.
9. Edick MJ, Cheng C, Yang W, et al. Lymphoid gene expression as a predictor of risk of secondary brain tumors. *Genes Chromosomes Cancer*. 2005;42(2):107–116.

Dove Medical Press encourages responsible, free and frank academic debate. The content of the OncoTargets and Therapy 'Editorial' section does not necessarily represent the views of Dove Medical Press, its officers, agents, employees, related entities or the OncoTargets and Therapy editors. While all reasonable steps have been taken to confirm the content of each Editorial, Dove Medical Press accepts no liability in respect of the content of any Editorial, nor is it responsible for the content and accuracy of any Editorial.

OncoTargets and Therapy

Publish your work in this journal

OncoTargets and Therapy is an international, peer-reviewed, open access journal focusing on the pathological basis of all cancers, potential targets for therapy and treatment protocols employed to improve the management of cancer patients. The journal also focuses on the impact of management programs and new therapeutic agents and protocols on

Submit your manuscript here: <http://www.dovepress.com/oncotargets-and-therapy-journal>

Dovepress

patient perspectives such as quality of life, adherence and satisfaction. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.