Thiopurine S-methyltransferase polymorphisms in acute lymphoblastic leukemia, inflammatory bowel disease and autoimmune disorders: influence on treatment response

Rachid Abaji¹
Maja Krajinovic²

¹Department of Pharmacology,
²Departments of Pediatrics and
Pharmacology, CHU Sainte-Justine
Research Center, University of
Montreal, Montreal, QC, Canada

Abstract: The *thiopurine S-methyltransferase (TPMT)* gene encodes for the TPMT enzyme that plays a crucial role in the metabolism of thiopurine drugs. Genetic polymorphisms in this gene can affect the activity of the TPMT enzyme and have been correlated with variability in response to treatment with thiopurines. Advances in the pharmacogenetics of *TPMT* allowed the development of dosing recommendations and treatment strategies to optimize and individualize prescribing thiopurine in an attempt to enhance treatment efficacy while minimizing toxicity. The influence of genetic polymorphisms in the *TPMT* gene on clinical outcome has been well-documented and replicated in many studies. In this review, we provide an overview of the evolution, results, conclusions and recommendations of selected studies that investigated the influence of *TPMT* pharmacogenetics on thiopurine treatment in acute lymphoblastic leukemia, inflammatory bowel disease and autoimmune disorders. We focus mainly on prospective studies that explored the impact of individualized TPMT-based dosing of thiopurines on clinical response. Together, these studies demonstrate the importance of preemptive *TPMT* genetic screening and subsequent dose adjustment in mitigating the toxicity associated with thiopurine treatment while maintaining treatment efficacy and favorable long-term outcomes. In addition, we briefly address the cost-effectiveness of this pharmacogenetics approach and its impact on clinical practice as well as the importance of recent breakthrough advances in sequencing and genotyping techniques in refining the *TPMT* genetic screening process.

Keywords: TPMT, pharmacogenetics, thiopurine, 6-mercaptopurine, azathioprine, ADRs.

Introduction

Thiopurine S-methyltransferase (TPMT) is an important cytoplasmic enzyme that catalyses the rate-limiting step in the metabolism of thiopurine drugs. It is coded by the *TPMT* gene and exerts its effect via S-adenosyl-L-methionine as the S-methyl donor and S-adenosyl-L-homocysteine as a by-product.¹⁻³ Thiopurine drugs, mainly 6-mercaptopurine (6-MP), and its prodrug azathioprine (AZA), are implicated as anti-metabolite cytotoxic and immunosuppressive agents in the treatment of malignancies such as acute lymphoblastic leukemia (ALL), inflammatory disorders like inflammatory bowel disease (IBD) and many autoimmune disorders, including rheumatoid arthritis (RA), systemic lupus erythematosus (SLE), autoimmune hepatitis (AIH) and generalized eczematous disorders.³⁻⁵ However, gastrointestinal disturbances (like nausea and vomiting), rashes, as well as more serious adverse drug reactions (ADRs) like bone marrow toxicity, hepatotoxicity and pancreatitis can lead to discontinuation of therapy in up to one-third of patients;⁶ these factors limit the use of these drugs.²

Correspondance: Maja Krajinovic
Centre de recherche, CHU
Sainte-Justine, 3175 Chemin de la
Côte-Ste-Catherine, Montréal, QC
H3T 1C5, Canada
Tel +1 514 345 4931 Ext 6259
Email maja.krajinovic@umontreal.ca

while *TPMT**3C is the most frequently found allele in African and Southeast Asian populations.^{14,34}

Other than variants in the coding region of *TPMT*, it is being increasingly acknowledged that variants in the non-coding regions such as the *TPMT*-promoter and introns can also affect the activity of the *TPMT* enzyme, possibly by influencing the transcription of its gene.³ One well-studied example of such polymorphisms is the variable number of tandem repeats (VNTR) region, which is a rare micro-satellite region in the *TPMT* gene promoter. Interestingly, studies have shown that the architecture of this region can modulate *TPMT* transcription and possibly enzyme activity. For example, higher *TPMT* promoter activity was shown to be associated with a region that contains five or seven GCC repeats rather than six. Thus, studies suggest the use of VNTR architecture as a pharmacogenomic biomarker to refine the *TPMT* genetic screening process currently used prior to the introduction of thiopurine therapy to enhance the treatment outcome in ALL.³ However, contrary to the results of ALL studies, the expression of the *TPMT* gene seems to decrease in IBD patients treated with thiopurine drugs and thus VNTR genotype cannot predict the *TPMT* activity, which seems to be influenced by the treated condition, the protocol used and the concomitant administration of other drugs.³

Across all ethnic groups, ~1 in 300 individuals are homozygous (or compound heterozygous) for a mutant *TPMT* allele and have very low or absent *TPMT* activity while around 4%–11% of individuals are heterozygous and are generally considered to have intermediate enzymatic activity.^{1,18,31} Nonetheless, such genotype-based classification is not always representative of the actual state of enzymatic activity. In literature, conflicting data were obtained by studies that addressed the concordance between the genetic and phenotypic tests, as results ranged from 100% or almost perfect match in the majority of studies to as low as 77% concordance in a few of them.^{1,3,9–11,19,33,35–38} This discordance was particularly observed in patients with intermediate activity in whom the estimated probability of obtaining matching results varied from 70% to 86%.^{11,33,36,39}

Many factors influence *TPMT* enzyme activity and eventually affect this genotype–phenotype concordance such as the age and gender of the patient, coadministration of drugs that could potentially interfere with the disease condition or *TPMT* activity (e.g., methotrexate),⁴⁰ levels of *TPMT* cofactor S-adenosyl-methionine,⁴¹ recent blood transfusion,⁴² life span of red blood cells,⁴³ as well as untested rare or novel variants in the coding and regulatory

regions of the *TPMT* gene (e.g., *TPMT**38 and the VNTR architecture).^{3,37,44} Furthermore, interethnic variability in the *TPMT* enzymatic activity levels have been observed in people of Afro-Caribbean descent having lower activity than Caucasians and South Asians.^{1,45,46} Taken together, there is always a risk of misclassifying patients if the decision was based on only one of the two abovementioned methods, but it is also unreasonable to perform both tests for all patients. A recent randomized clinical trial (RCT) concluded that there was no advantage or disadvantage of *TPMT* genotyping compared with phenotyping,¹⁶ whereas a more recent study concluded that genotyping was superior to phenotyping and should be considered as the primary choice for pretreatment evaluation of *TPMT* function.³⁵ Nonetheless, phenotype testing supplemented by genotyping can be a useful strategy in specific circumstances (e.g., after recent blood transfusion and for confirmation of intermediate activity in known high-risk patients).¹⁶

Recent advances in the pharmacogenetics of *TPMT* have allowed for the development of dosing recommendations and treatment strategies to optimize and individualize therapy with thiopurines to obtain maximum treatment benefit with minimal toxicity.⁴⁷ However, the implementation of pharmacogenetic tests in clinical practice is still somewhat limited due to the lack of robust evidence stemming directly from large-scale RCTs and proving the clinical utility of such strategy.^{1,16,22} Nonetheless, given the indisputable influence of pharmacogenetics on *TPMT* activity and the seriousness of thiopurines-induced toxicities, particularly myelosuppression, several regulatory agencies and clinical guidelines such as the US Food and Drug Administration, British National Formulary and Clinical Pharmacogenetics Implementation Consortium (CPIC) recommend pretreatment *TPMT* activity testing either by genotyping or phenotyping.^{11,47} In general, most guidelines suggest that the initial dose of thiopurines should be reduced to 10% of the standard dose when administered to homozygous carriers of *TPMT*-deficient alleles, as well as a reduction in administration frequency. For heterozygous patients, the recommendation differs slightly depending on the type of thiopurine used, as CPIC guidelines suggest an initial dose of AZA and 6-MP that is 30%–70% of the standard protocol dose while the recommendation for TG is 30%–50% of that dose.⁴⁷

As new data is being continuously generated by RCTs and studies about the long-term outcome of previous treatment protocols, the strength of the clinical evidence should be constantly revised and the recommendations of the guidelines should be reevaluated and modified when deemed necessary.

In this review, we provide an overview of the evolution, results, conclusions and recommendations of studies that investigated the influence of *TPMT* pharmacogenetics on clinical response to thiopurines in ALL, IBD and autoimmune disorders.

Acute lymphoblastic leukemia

Childhood ALL is the most frequent pediatric cancer. The survival rate currently exceeds 85% in favorable settings. 6-MP is coadministered with methotrexate as key components in the maintenance therapy for pediatric ALL and their use is associated with significant reduction in disease relapse.³³

An early study reported that *TPMT* activity was significantly higher in blood samples of ALL patients on long-term 6-MP treatment compared with controls.⁴⁸ They also noted a relationship between low *TPMT* activity and the risk of developing severe myelosuppression in patients treated with thiopurine drugs, plausibly due to elevated 6-TGN concentrations.^{48,49} Others reported that higher *TPMT* activity was linked to an elevated risk of relapse.⁵⁰ These findings led to the suggestion that genetic screening of *TPMT* activity could play a role in influencing treatment response to childhood ALL.^{48,50} Indeed, one study of childhood ALL suggested that prospectively screening for major *TPMT* coding region polymorphisms followed by selective administration of an initially reduced dose of 6-MP to heterozygous patients and a subsequent gradual increase to a target range of blood cell count allowed these patients to eventually achieve the full drug dose without experiencing any toxicity.⁵¹ Many clinical trials have investigated the impact of *TPMT* gene polymorphisms on treatment outcome, with most of them demonstrating the benefit of preemptive *TPMT* screening, but results were somewhat inconsistent (Table 1).

The Total Therapy Study XII explored the impact of 6-MP dose reduction from the standard protocol dose to a maximum tolerable dose subsequently to the development of myelosuppression and investigated the association between the maximum tolerable doses and *TPMT* genotypes in a total of 188 patients. The results showed that *TPMT* genotype was an important predictor of 6-MP toxicity in ALL patients, as the cumulative incidence of dose reduction or treatment interruption was significantly different across the 3 groups ($P < 0.001$); the results also showed that wild-type patients had the lowest incidence (7%) followed by heterozygous carriers (35%) and homozygous carriers of *TPMT*-deficient alleles (100%).^{10,23} They also concluded that administering lower doses of 6-MP in these patients was successful in maintaining adequately high levels of 6-TGN while allowing

the administration of other agents at full protocol doses.¹⁰ Furthermore, the investigators pointed out that the reduced activity patients tended to improve event-free survival (EFS) compared with wild-type patients ($P = 0.096$) and that higher dose intensity of 6-MP was the most significant predictor of that outcome ($P = 0.020$).²³ However, the authors also observed a nonsignificant trend for patients with low *TPMT* activity to have higher incidence and shorter onset of secondary acute myeloid leukemia as well as higher cumulative incidence of brain tumors when compared with patients with wild-type.^{52,53} In their later trial, Total Therapy Study XIII, which included 247 patients and pioneered the implementation of pharmacogenetics in leukemia therapy, they continued to administer a standard initial dose of 6-MP at the start of the continuation therapy but then selectively decreased the dose when deemed necessary based on a strategy that involved up-front knowledge of *TPMT* status combined with clinical tolerance and measurement of thiopurine metabolite levels.^{54,55} They eventually reported that *TPMT* genotype was not associated with the risk of hematologic relapse and that the long-term outcome showed no association with *TPMT* status (5-year cumulative incidences of $13.2\% \pm 2.3\%$ and $6.7\% \pm 6.7\%$ for wild-type and low-activity genotypes, respectively; $P = 0.46$), further confirming that considering pharmacogenetics of *TPMT* for dose adjustment of 6-MP dosage in ALL can help to reduce treatment-associated toxicity while not compromising its efficacy.^{54–57}

In the NOPHO-ALL-92 study of The Nordic Society of Paediatric Haematology and Oncology, a higher risk of relapse was observed in patients homozygous for wild-type ($P = 0.02$) and/or high *TPMT* activity ($P = 0.002$).^{36,58,59} However, the authors also observed that patients with low *TPMT* activity, although at lower risk of relapse, had a higher risk of developing second myeloid neoplasms (SMN) associated with high levels of 6-TGN and methylated metabolites, probably leading to DNA damage and subsequent malignancies. The authors believe that this theory explains why low *TPMT* activity patients did not have a superior overall survival (OS) compared with those with wild-type activity ($P = 0.82$) despite their lower risk of relapse.^{38,60} These observations, together with the ones from the Total Therapy Study XII, led the NOPHO to adopt *TPMT* genotype-dependent initial dosing of 6-MP in their later protocols, ALL-2000 and ALL-2008.^{36,58} Indeed, the long-term survival results from the ALL-2000 trial indicate that selecting the initial 6-MP dose based on *TPMT* genotype did reduce the risk of SMN in heterozygous patients, but at the expense of an increased risk of relapse. This explains

Table 1 Summary of selected studies that investigated the influence of TPMT pharmacogenetics on thiopurine treatment response in childhood acute lymphoblastic leukemia

Trial name	Design relative to TPMT genotype	6-MP treatment strategy	6-MP dosing (maintenance phase)	Genotype included	Finding	Conclusion	Author (reference no.)
Total Therapy Study XII	Observational/retrospective	Individualized therapy (dose reduction in patients experiencing myelosuppression to the highest tolerable dose)	75 mg/m ² /day with selective dose reduction in patients based on clinical tolerance	TPMT*2, *3A, *3B and *3C	Highest cumulative incidence of toxicity-induced reductions among patients homozygous for mutant TPMT alleles (100%), intermediate among heterozygous (35%) and lowest among wild-type patients (7%) Reduced TPMT activity tends to be associated with improved EFS. 6-MP dose intensity is a significant predictor of EFS	TPMT genotype is an important determinant of 6-MP toxicity and tolerance Treatment should aim to administer maximal tolerable protocol dose of 6-MP, particularly to patients with wild-type TPMT activity TPMT-deficient genotype may be associated with an increased risk of secondary AML.	Relling et al ¹⁰
Total Therapy Study XIII B	Individualized/prospective	Individualized therapy (dose reduction in patients experiencing myelosuppression to the highest tolerable dose) with pharmacogenetics compartment	75 mg/m ² /day with selective dose reduction in patients with low or intermediate TPMT activity based on a strategy that involves up-front knowledge of TPMT status combined with clinical tolerance and measurement of thiopurine metabolites	TPMT*2, *3A, *3B and *3C	Reduced TPMT activity tends to associate with higher risk of secondary AML Significantly higher cumulative risk of brain tumors among patients with reduced activity TPMT polymorphisms, especially when combined with high doses of antimetabolites No association between hematologic relapse (or other long-term outcomes) and TPMT status	TPMT-deficient genotype may be associated with an increased risk of radiation-associated brain tumors Considering pharmacogenetics of TPMT for dose adjustment of 6-MP can help reduce treatment toxicity while not compromising its efficacy	Relling et al ⁵³ Relling et al ⁵² Relling et al ⁵⁵ Rocha et al ⁵⁶

(Continued)

Table 1 (Continued)

Trial name	Design relative to TPMT genotype	6-MP treatment strategy	6-MP dosing (maintenance phase)	Genotype included	Finding	Conclusion	Author (reference no.)
NOPHO-ALL-92	Observational/retrospective	Individualized randomized maintenance therapy based on clinical response and levels of metabolites	75 mg/m ² /day with subsequent dose adjustment to a target WBC	TPMT*3A, *3B and *3C	Higher risk of relapse in patients with high TPMT activity. Increased leukemogenic risk and higher rate of SMN in patients with low activity	No difference in OS between low vs high TPMT activity groups as the improved EFS is offset by the higher risk of SMN	Thomsen et al ⁶⁰ Schmiegelow et al ⁵⁹ Schmiegelow et al ³⁶ Schmiegelow et al ³⁸ Schmiegelow et al ⁵⁸
NOPHO-ALL-2000	Individualized/prospective	Individualized randomized maintenance therapy based on clinical response, levels of metabolites and pharmacogenetics	75 mg/m ² /day for wild-type patients, 50 mg/m ² /day for heterozygous patients and 5–10 mg/m ² /day for TPMT deficient patients, with subsequent dose adjustment to a target WBC during the first year of maintenance therapy	TPMT*3A, *3B and *3C	Reduced risk of SMN in heterozygous patients compared to ALL-92. Similar relapse risk between low TPMT activity patients and wild-type patients. No difference in overall EFS (any event) or OS compared to previous protocol	Initial 6-MP dose adjustment based on TPMT genotype did reduce the risk of SMN in heterozygous patients but at the expense of an increased risk of relapse	Schmiegelow et al ⁵⁸ Levinson et al ⁶¹
UK-ALL97 and ALL97/99	Individualized	Individualized randomized maintenance therapy based on TPMT activity and clinical response	75 mg/m ² /day for both TPMT heterozygous and wild-type patients subsequently titrated to toxicity Patients with TPMT deficiency were titrated from a starting dose of 10% of protocol dose (75 mg/m ² /day)	TPMT*2, *3A, *3B and *3C	Patients with TPMT*1/*3A genotype had better EFS than both wild-type TPMT*1/*1 and TPMT*1/*3C patients. Heterozygosity was not associated with a higher rate of second cancers. Higher hematotoxicity in heterozygous patients compared to wild-type patients	While, heterozygous patients had significantly more cytopenias and required dose adjustments more often than wild-type patients, TPMT*1/*3A patients had better EFS.	Lennard et al ⁶²
UK-ALL-2003	Individualized/prospective	Risk stratification based on MRD and individualized randomized maintenance therapy based on clinical response, TPMT activity and pharmacogenetics	75 mg/m ² /day for both TPMT heterozygous and wild-type patients subsequently titrated to toxicity Patients with TPMT deficiency were titrated from a starting dose of 10% of protocol dose (75 mg/m ² /day)	TPMT*2, *3A, *3B, *3C and *9	Improved overall EFS (all TPMT genotypes confounded) compared to previous protocol. No difference in EFS, RFS or OS between TPMT genotypes. The only significant factor affecting EFS was MRD status	Refinements of the risk stratification process and treatment strategies have reduced the influence of TPMT genotype on treatment outcome	Lennard et al ³³

ALL-BFM-2000	Observational/ prospective	Measurement of MRD load before and after 6-MP treatment in heterozygous vs wild-type patients	During consolidation phase, on treatment day 78, heterozygous and wild-type homozygous patients received a 4-week cycle of (60 mg/ m ² /day of 6-MP). Patients homozygous for mutant <i>TPMT</i> alleles were excluded from the analyses.	<i>TPMT</i> *2, *3A, *3B, *3C and *3D + direct sequencing for individuals with an intermediate activity not having any mutation in exon 5, 7 and 10	Significant reduction in the risk of having detectable MRD in <i>TPMT</i> heterozygous patients after induction consolidation treatment compared with wild-type patients. No difference in the rate of hematopoietic toxicity between <i>TPMT</i> heterozygous carriers and homozygous wild-type carriers. No association between low <i>TPMT</i> activity and risk of SMN	<i>TPMT</i> genotype has a significant impact on MRD during induction consolidation treatment as heterozygous patients showed a better MRD response, indicating an increased clearance of disease likely due to higher intensity of 6-MP effect. <i>TPMT</i> polymorphisms do not influence the risk of developing SMN in ALL patients treated with BFM protocols	Stanulla et al ⁶³ Stanulla et al ⁶⁴
--------------	-------------------------------	---	--	--	--	--	--

Abbreviations: 6-MP, 6-mercaptopurine; ALL, acute lymphoblastic leukemia; AML, acute myeloid leukemia; BFM, Berlin-Frankfurt-Münster; EFS, event-free survival; MRD, minimal residual disease; NOPHO, Nordic Society of Paediatric Haematology and Oncology; OS, overall survival; RFS, relapse-free survival; SMN, secondary malignant neoplasms; *TPMT*, thiopurine S-methyltransferase; WBC, white blood cells.

why although a slight nonsignificant improvement in EFS was achieved by the new protocol, it had no difference in overall EFS or OS from its predecessor (5-year results: NOPHO-ALL-92: EFS=77.4%±1.0%, OS=87.6%±0.8%, n=1654; and NOPHO-ALL-2000: EFS=79.4%±1.5% and OS=89.1%±1.1%, n=1023).^{58,61}

In the UK, ALL97 and ALL97/99 trials, wild-type and heterozygous patients on the 6-MP arm initially received full dose of the drug, which was later adjusted to clinical hematologic toxicity whereas *TPMT*-deficient patients received 10% of the dose also adjusted for toxicity. While investigators observed a finding similar to the aforementioned studies in that patients with the *TPMT**1/*3A genotype (n=99, EFS=88%) had better outcome at 5 years compared with *TPMT* wild-type patients (n=1206, EFS=80%; *P*=0.05), paradoxically, patients with the *TPMT**1/*3C genotype also had lower EFS than those with *1/*3A genotype (n=17, EFS=53%; *P*=0.002). Furthermore, patients with heterozygous genotypes were found to experience more myelosuppression, accumulated higher 6-TGN concentrations and required dose reduction more frequently. However, no association between the risk of secondary malignancy and *TPMT* genotype was found.^{13,33,62} In their subsequent trial, ALL-2003, which used minimal residual disease (MRD) to guide risk stratification and treatment intensity, the protocol prospectively observed the influence of *TPMT* genotype on treatment outcome by applying pretreatment genetic screening of the most common *TPMT* polymorphisms to 2387 of the study patients. The dosing regimen for 6-MP was similar to that of ALL97 in the sense that *TPMT*-deficient patients received 10% of the dose while the others received a standard dose subsequently adjusted according to a target cell count. The results showed that overall EFS – all *TPMT* genotypes confounded – was significantly higher than that of the previous protocol, which was attributed to the improved survival in the *TPMT* wild-type and *TPMT**1/*3C genotype groups (EFS at 5 years=88%, 88% and 94% for *TPMT* wild-type, *1/*3A and *1/*3C, respectively). However, within this protocol, no significant differences in OS, EFS or relapse-free survival were observed with respect to *TPMT* genotypes. Thus, it was concluded that the improved risk-adapted protocol had reduced the influence of *TPMT* genotypes on treatment outcome and that the only factor that affected the outcome was MRD. Furthermore, there was no difference in survival within each MRD risk group with respect to *TPMT* genotypes.³³ Overall, the cumulative experience of the many UK ALL trials led to mandating preemptive *TPMT* screening for all children and young adults who start the ALL-2011 trial protocol.¹¹

In the Berlin-Frankfurt-Münster-2000 (BFM-2000) trial, the 6-MP dose was reduced by 10-fold from the standard starting dose for TPMT-deficient patients, but no dose adjustment was carried for heterozygous carriers who were given doses equal to the homozygous carriers of the wild-type. The investigators assessed the genotypes of 814 patients and used MRD for risk-stratification. The results showed no difference in the rate of hematopoietic toxicity between *TPMT* heterozygous variant carriers and homozygous wild-type carriers or between *TPMT* status and the risk of developing secondary cancers. Interestingly, it was observed that *TPMT* genotype had a significant impact on MRD during induction consolidation treatment as heterozygous patients had better MRD response (2.9-fold reduction), indicating an increased clearance of disease likely due to higher intensity of 6-MP effect (Relative risk=0.34; 95% confidence interval [CI], 0.13–0.86; $P=0.02$).^{63,64}

In summary, all these trials demonstrate the importance of preemptive *TPMT* genetic screening and subsequent dose adjustment in mitigating the toxicity associated with thiopurine treatment while maintaining, if not enhancing, treatment efficacy and favorable long-term outcome.

Inflammatory bowel disease

IBD is a polygenic chronic, relapsing and remitting disease of the gastrointestinal tract that can be divided into two major clinical subtypes, Crohn's disease and ulcerative colitis.^{2,65} Thiopurines, particularly AZA, are proven effective in inducing and maintaining long-term remission in IBD patients.⁵ More than 20% of patients experience severe ADRs that lead to dose modification, treatment interruption or cessation.^{66,67} Bone marrow toxicity represented by leukopenia is one of the most serious thiopurine-related ADRs. Many studies investigated the influence of *TPMT* genotype on the efficacy and toxicity of thiopurines and most suggested a significant impact on clinical response (Table 2).

A systematic review followed by a meta-analysis that eventually combined the results of 47 studies that investigated the risk of myelosuppression with respect to intermediate *TPMT* activity demonstrated a 4.19-fold increase in odd-ratio of leukopenia (95% CI: 3.20–5.48) in IBD patients with reduced *TPMT* activity compared with wild-type. One critic of this meta-analysis is that it combined rather smaller studies with sample sizes of <100 patients in most cases and the majority having retrospective cohort designs. However, in a sub-analysis of this study that combined 834 patients coming only from the 11 studies that had a prospective cohort design, the significant association of reduced *TPMT* activity with the risk of myelosuppression had an odd-ratio

of 4.3 (95% CI: 2.53–7.29).¹ Among these prospective studies, an observational study with preemptive *TPMT* genetic testing for all patients and a relatively large sample size of 207 participants found that heterozygous *TPMT* genotype strongly predicted treatment withdrawal due to early-onset of ADRs following a conventional fixed-dosing regimen (79% vs 35% in heterozygous and wild-type, respectively; $P<0.001$). They highlighted that gastric intolerance (GI) was the most frequent reason for withdrawal among this group of patients and that myelotoxicity and GI occurred significantly more frequently among heterozygous (26% and 37%, respectively) than wild-type patients (0.5% and 7%, respectively). Interestingly, they had a 100% concordance of genotype to phenotype activity and found that *TPMT* activity was strongly predictive of clinical response as it was significantly higher in nonresponders.⁶⁸ Other prospective studies with preemptive *TPMT* genetic screening component reported that overall thiopurine-related ADRs were significantly more common among patients with low-to-intermediate *TPMT* activity when doses were not adjusted;⁶⁷ particularly myelotoxicity, which was more profound in *TPMT*-deficient genotype.^{67,69}

Two independent meta-analysis further investigated the impact of pharmacogenetics on treatment response by exclusively combining studies (14 and 9 studies, respectively) that investigated the association between *TPMT* polymorphisms and ADRs in IBD patients, regardless of the study design (i.e., cross-sectional cohort, prospective cohort and case-control studies).^{2,65} They involved 2206 and 1309 patients respectively, and both concluded that *TPMT* polymorphisms were significantly associated with thiopurine-induced overall ADRs and bone marrow toxicity (around 3- and 6-fold increase in the odd-ratios, respectively) but not with hepatotoxicity, pancreatitis, flu-like symptoms, GI or skin reactions.^{2,65}

The *TPMT*: Azathioprine Response to Genotyping and Enzyme Testing (TARGET) trial is a pragmatic RCT that prospectively investigated the impact of genotype-guided initial dosing of AZA followed by upward-titration to the maximum tolerable dose of the full protocol dose compared to no genotyping and full standard dose administration to all participants. It included 333 patients with inflammatory diseases and the primary aim was to see if this strategy would result in a significant reduction in the rate of ADRs-induced treatment cessation. No differences were found between the conventional and pharmacogenetics arms with respect to the frequency of treatment interruption due to ADRs (frequency: 27.7% vs 28.8%; odds ratio [OR]: 1.1; 95% CI: 0.66–1.8; $P=0.74$). On the other hand, the study did not find any difference in the rate of remission between the intervention

Table 2 Summary of selected studies which investigated the influence of TPMT pharmacogenetics on thiopurine treatment response in inflammatory bowel disease

Author (reference no.)	Design relative to TPMT genotype	Treatment strategy	Genotypes included	Findings and conclusion
Derijks et al ⁶⁹	Observational	6-MP as a single oral 50 mg evening dose	TPMT*2, *3A, *3B and *3C	TPMT genotype correlated with 6-TGN concentrations. Patients with mutant alleles have higher risk of developing leukopenia
Hindorf et al ⁶⁷	Observational	Dose escalation schedule to the target dose of (2.5 mg/kg) for azathioprine and (1.25 mg/kg) for 6-MP by week 3	TPMT*2, *3A, *3B, *3C, *3D, *4, *5, *6, *7, *8, *10, *14 and *15	Overall, thiopurine-related ADRs were significantly more common among patients with low-to-intermediate TPMT activity; particularly myelotoxicity in TPMT-deficient patients
Ansari et al ⁶⁸	Observational	AZA was started at 2 mg/kg daily and without dose alteration	TPMT*3A, *3B and *3C	Heterozygous TPMT genotype strongly predicts treatment withdrawal due to early-onset of ADRs. Gastric intolerance was the most frequent reason for withdrawal among heterozygous patients. Myelotoxicity and gastric intolerance occurred significantly more frequently among heterozygous than with TPMT wild-type patients
Newman et al ¹⁶ , TARGET trial	Individualized/prospective	Arm 1: Standard dosing without genotyping vs Arm 2: Pretreatment TPMT genotyping and AZA dosing. Wild-type (1.5–3 mg/kg/day); heterozygous (25–50 mg/day) and titrate to the maintenance dose; homozygous for TPMT variant alleles were given alternative drugs	TPMT*2, *3A, *3B and *3C	No differences between the two study arms or between heterozygous and wild-type homozygous patients with respect to the rate of stopping azathioprine due to ADRs. No difference in the rate of remission between the intervention and control groups
Coenen et al ²² , TOPIC trial	Individualized/prospective	Arm 1: Control group. No genotyping + standard dosing: 2–2.5 mg/kg/day AZA or 1–1.5 mg/kg/day 6-MP. Arm 2: Intervention group. Genotyping + individualized dosing. TPMT wild-type: same as control; heterozygous: 50% of control and homozygous variant: 0%–10% of control	TPMT*2, *3A and *3C	No significant overall impact of TPMT genotype-guided dosing of thiopurines on treatment efficacy or on the risk of hematologic ADRs (i.e., leukopenia and thrombocytopenia) between the genotyped and nongenotyped arms. Carriers of at least one genetic variant in the pharmacogenetics arm had a significant reduction in the risk of hematologic ADRs compared with same group in the conventional arm

Abbreviations: 6-MP, 6-mercaptopurine; 6-TGN, 6-thioguanine nucleotides; ADRs: adverse drug reactions; AZA, azathioprine; TPMT, thiopurine S-methyltransferase; TARGET, TPMT: Azathioprine Response to Genotyping and Enzyme Testing; TOPIC, Thiopurine response Optimization by Pharmacogenetic testing in Inflammatory bowel disease Clinics.

and control groups indicating that the adjustment did not affect treatment efficacy. However, the investigators did not provide a stratified analysis addressing the differences in outcomes according to genotype groups within each study arm or between the two arms, probably due to small sample size. Moreover, it is worth mentioning that a single patient with homozygous variant genotype in the study who was on the nongenotyping arm and subsequently received the full dose of AZA developed severe neutropenia, which underlines the importance of genetic testing to identify this group of patients.¹⁶

A larger and more recent prospective RCT, which involved 783 IBD patients, the Thiopurine response Optimization by

Pharmacogenetic testing in Inflammatory bowel disease Clinics (TOPIC) trial, similarly showed no significant overall impact of TPMT genotype-guided dosing of thiopurines on treatment efficacy or on the risk of hematologic ADRs (i.e., leukopenia and thrombocytopenia) between the genotyped and nongenotyped arms (frequency: 7.4% vs 7.9%; relative risk: 0.93; 95% CI: 0.57–1.52). The efficacy results of this study further advocate that a reduced thiopurine dose does not result in under-treatment. Moreover, a subgroup analysis of this study, which compared only carriers of TPMT variants between the two arms, revealed that the pharmacogenetic approach was able to significantly decrease the risk of hematologic ADRs by 10-fold in carriers of at least one genetic

variant (frequency: 2.6% vs 22.9%; relative risk: 0.11; 95% CI: 0.01–0.85).²² The results of the secondary aim of this study excluded any significant association between *TPMT* genotypes and anemia, hepatotoxicity, pancreatitis, skin rash, GI and general malaise, which is consistent with the results of the aforementioned meta-analysis and other results in the literature.^{2,22,65} It also suggested that factors other than *TPMT* genotype play an important role in the development of thiopurine-induced ADRs.²⁴

Autoimmune disorders

Autoimmune diseases are a group of heterogeneous conditions that involve a destructive attack against the host's tissues launched by a deregulated immune system as in SLE, RA, AIH and generalized-eczematous disorders. Thus, treatment strategies are usually based on the use of immunosuppressants that act by modifying the activity of the immune system. AZA is widely used as an immunosuppressive agent in autoimmune diseases but its use is limited by its ADRs.¹⁴ Similar to the meta-analysis that focused on IBD, another meta-analysis, which included 651 patients with autoimmune diseases coming from 11 studies demonstrated that overall ADRs and AZA-induced bone marrow toxicity are significantly associated with *TPMT* polymorphisms with OR of 3.12 (95% CI: 1.48–6.56) and 3.76 (95% CI: 1.97–7.17), respectively. The results remained significant in two analysis, one that grouped the homozygous and heterozygous carriers into one reduced-activity group and the other that focused on heterozygous carriers only. The study also showed a significant association with GI with OR of 6.43 (95% CI: 2.04–20.25), but the authors suspect that the observed association might have been driven by a single study, since after excluding this study, the association was no longer significant with OR of 2.1 (95% CI: 0.36–12.42).¹⁴ The study also excludes the association of *TPMT* polymorphisms with hepatotoxicity. The sub-analysis that examined the association with myelosuppression according to the type of disease found significant results in SLE, RA and AIH subgroups. They also concluded that the risk prediction of bone marrow toxicity and overall ADRs based on *TPMT* variant-positive genotypes has high specificity (94.10% and 92.93%, respectively) but at the expense of sensitivity (16.30% and 14.85%, respectively).¹⁴ Furthermore, in a prospective study that investigated the impact of *TPMT*-activity guided AZA dosing on the treatment response in patients with atopic eczema, the investigators concluded that *TPMT*-based dosing was able to maintain the drug efficacy while reducing the predicted toxicity.⁷⁰

Cost-effectiveness

Most of the above studies concluded that *TPMT* testing could lead to improved prescribing of thiopurines, which would ultimately result in an increased treatment efficacy and a reduction in the rate and intensity of ADRs. Nonetheless, the cost-effectiveness of such an intervention is still open to debate. Only a few studies have addressed the cost-effectiveness of *TPMT* pharmacogenetics interventions. In an effort toward this evaluation, a case study examined the cost-effectiveness of prospective *TPMT* genotyping in children with ALL treated with thiopurines and suggested positive results manifested in financial savings and a gain in life-years in the most favorable settings of the sensitivity analysis.⁷¹ Similarly, another study established a model based on a theoretical IBD population treated with AZA and found that pretreatment screening for *TPMT* genotype would be cost-effective in avoiding patient mortality due to myelosuppression.²⁹ However, data coming from RCTs do not necessarily support this conclusion as one perspective study found that such a technique incurred excessive cost associated with genotyping but did not predict AZA-induced toxicity in IBD patients.⁷² Nonetheless, these studies were too small and not adequately powered to answer this question.¹⁶ A systematic review came to the conclusion that screening for *TPMT* activity either by genotyping or phenotyping was a cost-effective strategy that can be used to reduce health-care costs while improving clinical effectiveness.⁷³ Another study aimed at the evaluation of the added-value of genetic screening of *TPMT* followed by dose adjustment of AZA prior to the initiation of treatment found that genetic-based dosing dominated the standard dosing strategy in patients with rheumatological disease by reducing the treatment cost and the frequency of AZA-induced side effects.⁷⁴ In a more recent prospective economic evaluation that was conducted alongside the TARGET study, in which the study aim was to test the cost-effectiveness of the *TPMT* genotyping approach in autoimmune diseases, the researchers concluded that the genetic approach had up to 71% probability of being cost-effective depending on the cost of the genetic test. The results, however, were not conclusive as the observed economic advantage in the intervention group owing to lower use of resources was accompanied by a slight (almost negligible) reduction in the quality of life.⁷⁵

Impact on clinical practice

Over the past decade, *TPMT* enzyme testing has gained a lot of acceptance, as reflected by the rapid increase in the number of tests performed in clinical practice.^{16,76} This

sudden increase was the inevitable result of multiple factors supporting this approach, which include the increase in the available knowledge about the role of *TPMT* in treatment outcome, the stronger recommendations coming from clinical guidelines like the CPIC and the wider accessibility to genetic testing (i.e., larger availability, reduced cost, faster turnaround of results and shorter interpretation time).^{1,16} This shift in clinical practice was evaluated in the TARGET study, which observed that the physicians did follow the recommendations coming from British clinical guidelines (e.g., British Association of Dermatologists Therapy and British Society for Rheumatology) for *TPMT* heterozygous patients and chose a lower initial dose of AZA for those patients but the investigators also noted that the physicians used overall lower starting doses for wild-type patients as well.¹⁶ This “safe” practice reflects the physicians’ reservation regarding the sensitivity and specificity of this test, which stems from the fact that being homozygous carrier of *TPMT* wild-type, although predictive of a reduced risk of AZA-induced myelosuppression, does not completely eliminate the possibility. Indeed, it was mentioned earlier that a fraction of *TPMT* wild-type patients can still have intermediate *TPMT*-activity and that other factors play a role in the development of this ADR.^{16,33,36} Moreover, other side effects such as hepatotoxicity, pancreatitis, nausea and vomiting cannot be predicted by *TPMT* testing.^{1,14,65} The adoption of pretreatment *TPMT* screening seems to vary according to discipline, as reported by one survey, with 94% of dermatologists, 60% of gastroenterologists and only 47% rheumatologists requesting it.⁷⁶ This could be related to the level of evidence available in the domain of practice and the strength of the recommendations of the respective guidelines and protocols used by each specialist (e.g., UK guidelines in dermatology and gastroenterology recommend genetic screening while ALL-2011 protocol mandates it).^{1,11,76} However, from an evidence-based perspective, and besides the universally accepted association with hematotoxicity, the recommendations for preemptive genetic testing still have some margin to evolve. Plus, even in well-established scenarios, like in the case of myelosuppression in *TPMT*-deficient patients, strong evidence is still lacking to support that the pharmacogenetic approach would result in a significantly better outcome.^{14,65}

TPMT in the new era of sequencing

The influence of genetic polymorphisms in the *TPMT* gene on treatment outcome has been well-documented and replicated in many studies. However, studies have also concluded that the genetic-based screening for *TPMT* activity should be

interpreted with caution, as the activity of the *TPMT* enzyme can be co-influenced by other factors, and the development of thiopurine-induced ADRs is a multifactorial event.¹⁴ For instance, most of the presented studies inferred *TPMT* activity by genotyping the most common nonfunctional *TPMT* alleles while results of a recent study that explored the sequencing data suggest that in certain populations, the inferred activity can be refined by incorporating the genotypes of other alleles. The study also identified a new variant in the *TPMT* gene, *TPMT**38 (T514C), which had an allelic frequency of 0.11% and was predicted to be a damaging mutation.¹⁵ Moreover, as increasingly reported by different studies, genetic variants in other genes involved in thiopurines metabolism like *ITPA*, *HGPRT* and *MTHFR* as well as variants in genes independent of *TPMT* can influence thiopurines treatment outcome.^{14,20,25,41,70,77} For example, genome-wide association studies have identified variants in the *PACSLN2* gene that influence *TPMT* activity and were linked to 6-MP-related gastrointestinal toxicity in children with ALL, whereas variants in the *NUDT15* gene were associated with thiopurine-induced leukopenia.^{26,41} However, since a lot of genes have significant differences in the frequencies of polymorphisms across major ethnic groups, it is important to evaluate the genetic profiles of patients in a global frame that considers all of the genes involved in a specific pathway to better understand the impact of ethnic diversity on drug response. One particularly interesting example of the role of pharmacogenetics is the case of *NUDT15* in Japanese population, in whom polymorphisms of this gene were associated with higher risk of toxicity and were more frequent than *TPMT*-deficient variants.²⁶ Studies also suggest that combining the effects of such polymorphisms with variants in *TPMT* gene could strengthen the predictive power of the risk of developing thiopurines-related toxicity.^{14,22} This should soon become feasible with the breakthrough advances in sequencing and genotyping techniques. Indeed, in a recent study that tested the sensitivity, specificity and predictive values of the imputation of *TPMT*-alleles, most values were over 90%, indicating that imputation of *TPMT* alleles can be used as a screening method for individuals with high-risk of developing serious thiopurine-induced ADRs.⁴⁶ Furthermore, nongenetic factors should be taken into consideration before thiopurine initiation as they can have a big influence on the outcome and might interfere with the genotype-guided dosing.¹⁴

Conclusion

In conclusion, although it is currently well established that *TPMT* polymorphisms can explain a certain portion of

thiopurine-induced ADRs, particularly hematotoxicity, it is surely not capable of predicting all of them. Indeed, many studies^{2,14,22,65} have found that certain ADRs were not associated with a reduced TPMT activity such as pancreatitis and hepatotoxicity. This holds true in the context of ALL, IBD and the different types of autoimmune disorders. What is clear so far is that TPMT-deficient genotypes (homozygous variant carriers and compound heterozygous), and to lesser extent, heterozygous patients are predisposed to thiopurine-induced severe hematotoxicity.^{1,47} However, other factors such as disease progression and co-medications can also modulate the risk of myelosuppression, regardless of the genotype. While TPMT-deficient patients will definitely benefit from dose reduction of thiopurines, the validity of this approach for heterozygous carriers is still arguable since studies have shown that not all of these patients are intolerant to thiopurine; in fact, 30%–60% of heterozygous patients do tolerate it.^{33,47} Moreover, depending on the treated condition and treatment protocol used, *TPMT* wild-type patients also exhibit higher risks of worse outcome such as hematologic relapse in ALL and treatment failure in IBD, which adds an extra layer of complexity to the already troublesome process of finding the best therapeutic regimen that would ensure maximum efficacy and minimum toxicity.⁴⁷ Consequently, regular clinical testing and hematologic assessment remain the mainstay in the monitoring of thiopurine treatment while genetic testing adds the advantage of refining the initial dosing and patient-stratification processes, as well as suggesting customized monitoring for certain patient groups. One nice example backed with strong clinical evidence is the abovementioned scenario of myelosuppression. Preemptive *TPMT* genetic screening and tailored thiopurine initial dosing followed by upward/downward titration and hematological monitoring to a target level of myelosuppression can be considered a cost-effective approach which would allow the prevention and early detection of myelosuppression in this vulnerable population without compromising the efficacy of the treatment.⁴⁷

Perspective

While the goal of personalized medicine in general, and pharmacogenetics in particular, is to deliver patient-tailored treatments that would ensure maximum efficacy with minimum toxicity, the studies presented in this review make the argument that this is not an easy task. There is a balance to consider between treatment benefits and ADRs, which is controlled by multiple factors. This being said, what we can be sure of, for now, is that the more we get to know about the impact of pharmacogenetics on the variability of

treatment response, the better we will be able to control the outcome to the advantage of the patient. Moreover, most pharmacoeconomic analyses have indicated that screening for *TPMT* pharmacogenetics promises to be cost-effective. With the advent of next-generation sequencing and the many breakthroughs in bioinformatics, the cost of analyzing the entire human genome is bound to drop, which would allow for greater accessibility to genetic data and a larger understanding of how their interactions with each other and with other factors influence the treatment. In the meantime, it is very promising to see that most major institutions have already incorporated preemptive *TPMT* screening in their treatment protocols to enhance treatment outcome and the continuously emerging long-term data proving the utility of doing so, which should encourage other institutions to follow.

Acknowledgment

Rachid Abaji is a scholar of the Cole Foundation and the Network of Applied Medical Genetics and acknowledges the support of both organizations.

Disclosure

The authors report no conflicts of interest in this work.

References

- Higgs JE, Payne K, Roberts C, Newman WG. Are patients with intermediate TPMT activity at increased risk of myelosuppression when taking thiopurine medications? *Pharmacogenomics*. 2010;11(2):177–188.
- Dong XW, Zheng Q, Zhu MM, Tong JL, Ran ZH. Thiopurine S-methyltransferase polymorphisms and thiopurine toxicity in treatment of inflammatory bowel disease. *World J Gastroenterol*. 2010;16(25):3187–3195.
- Kotur N, Dokmanovic L, Janic D, et al. TPMT gene expression is increased during maintenance therapy in childhood acute lymphoblastic leukemia patients in a TPMT gene promoter variable number of tandem repeat-dependent manner. *Pharmacogenomics*. 2015;16(15):1701–1712.
- Chouchana L, Narjoz C, Beaune P, Lorient MA, Roblin X. Review article: the benefits of pharmacogenetics for improving thiopurine therapy in inflammatory bowel disease. *Aliment Pharmacol Ther*. 2012;35(1):15–36.
- Gisbert JP, Gomollon F. Thiopurine-induced myelotoxicity in patients with inflammatory bowel disease: a review. *Am J Gastroenterol*. 2008;103(7):1783–1800.
- Fraser AG, Orchard TR, Jewell DP. The efficacy of azathioprine for the treatment of inflammatory bowel disease: a 30 year review. *Gut*. 2002;50(4):485–489.
- Lennard L. The clinical pharmacology of 6-mercaptopurine. *Eur J Clin Pharmacol*. 1992;43(4):329–339.
- Lennard L. TPMT in the treatment of Crohn's disease with azathioprine. *Gut*. 2002;51(2):143–146.
- Chouchana L, Narjoz C, Roche D, et al. Interindividual variability in TPMT enzyme activity: 10 years of experience with thiopurine pharmacogenetics and therapeutic drug monitoring. *Pharmacogenomics*. 2014;15(6):745–757.
- Relling MV, Hancock ML, Rivera GK, et al. Mercaptopurine therapy intolerance and heterozygosity at the thiopurine S-methyltransferase gene locus. *J Natl Cancer Inst*. 1999;91(23):2001–2008.

11. Lennard L. Implementation of TPMT testing. *Br J Clin Pharmacol*. 2014;77(4):704–714.
12. Wang L, Pellemounter L, Weinshilboum R, et al. Very important pharmacogene summary: thiopurine S-methyltransferase. *Pharmacogenet Genomics*. 2010;20(6):401–405.
13. Vora A, Mitchell CD, Lennard L, et al. Toxicity and efficacy of 6-thioguanine vs. 6-mercaptopurine in childhood lymphoblastic leukaemia: a randomised trial. *Lancet*. 2006;368(9544):1339–1348.
14. Liu YP, Xu HQ, Li M, et al. Association between thiopurine S-methyltransferase polymorphisms and azathioprine-induced adverse drug reactions in patients with autoimmune diseases: a meta-analysis. *PLoS One*. 2015;10(12):e0144234.
15. Kim HY, Lee SH, Lee MN, et al. Complete sequence-based screening of TPMT variants in the Korean population. *Pharmacogenet Genomics*. 2015;25(3):143–146.
16. Newman WG, Payne K, Tricker K, et al. A pragmatic randomized controlled trial of thiopurinemethyltransferase genotyping prior to azathioprine treatment: the TARGET study. *Pharmacogenomics*. 2011;12(6):815–826.
17. Evans WE, Horner M, Chu YQ, Kalwinsky D, Roberts WM. Altered mercaptopurine metabolism, toxic effects, and dosage requirement in a thiopurinemethyltransferase-deficient child with acute lymphocytic leukemia. *J Pediatr*. 1991;119(6):985–989.
18. Yates CR, Krynetski EY, Loennechen T, et al. Molecular diagnosis of thiopurine S-methyltransferase deficiency: genetic basis for azathioprine and mercaptopurine intolerance. *Ann Intern Med*. 1997;126(8):608–614.
19. Evans WE, Hon YY, Bomgaars L, et al. Preponderance of thiopurine S-methyltransferase deficiency and heterozygosity among patients intolerant to mercaptopurine or azathioprine. *J Clin Oncol*. 2001;19(8):2293–2301.
20. Karas-Kuzelicki N, Jazbec J, Milek M, Mlinaric-Rascan I. Heterozygosity at the TPMT gene locus, augmented by mutated MTHFR gene, predisposes to 6-MP related toxicities in childhood ALL patients. *Leukemia*. 2009;23(5):971–974.
21. Peregud-Pogorzelski J, Tetera-Rudnicka E, Kurzawski M, et al. Thiopurine S-methyltransferase (TPMT) polymorphisms in children with acute lymphoblastic leukemia, and the need for reduction or cessation of 6-mercaptopurine doses during maintenance therapy: the Polish multicenter analysis. *Pediatr Blood Cancer*. 2011;57(4):578–582.
22. Coenen MJ, de Jong DJ, van Marrewijk CJ, et al. Identification of patients with variants in TPMT and dose reduction reduces hematologic events during thiopurine treatment of inflammatory bowel disease. *Gastroenterology*. 2015;149(4):907–917.e7.
23. Relling MV, Hancock ML, Boyett JM, Pui CH, Evans WE. Prognostic importance of 6-mercaptopurine dose intensity in acute lymphoblastic leukemia. *Blood*. 1999;93(9):2817–2823.
24. Colombel JF, Ferrari N, Debuysere H, et al. Genotypic analysis of thiopurine S-methyltransferase in patients with Crohn's disease and severe myelosuppression during azathioprine therapy. *Gastroenterology*. 2000;118(6):1025–1030.
25. Stocco G, Cheok MH, Crews KR, et al. Genetic polymorphism of inosine triphosphate pyrophosphatase is a determinant of mercaptopurine metabolism and toxicity during treatment for acute lymphoblastic leukemia. *Clin Pharmacol Ther*. 2009;85(2):164–172.
26. Moriyama T, Nishii R, Perez-Andreu V, et al. NUDT15 polymorphisms alter thiopurine metabolism and hematopoietic toxicity. *Nat Genet*. 2016;48(4):367–373.
27. Dubinsky MC, Yang H, Hassard PV, et al. 6-MP metabolite profiles provide a biochemical explanation for 6-MP resistance in patients with inflammatory bowel disease. *Gastroenterology*. 2002;122(4):904–915.
28. Ansari A, Hassan C, Duley J, et al. Thiopurinemethyltransferase activity and the use of azathioprine in inflammatory bowel disease. *Aliment Pharmacol Ther*. 2002;16(10):1743–1750.
29. Winter J, Walker A, Shapiro D, Gaffney D, Spooner RJ, Mills PR. Cost-effectiveness of thiopurinemethyltransferase genotype screening in patients about to commence azathioprine therapy for treatment of inflammatory bowel disease. *Aliment Pharmacol Ther*. 2004;20(6):593–599.
30. Appell ML, Berg J, Duley J, et al. Nomenclature for alleles of the thiopurinemethyltransferase gene. *Pharmacogenet Genomics*. 2013;23(4):242–248.
31. Collie-Duguid ES, Pritchard SC, Powrie RH, et al. The frequency and distribution of thiopurinemethyltransferase alleles in Caucasian and Asian populations. *Pharmacogenetics*. 1999;9(1):37–42.
32. Katara P, Kuntal H. TPMT polymorphism: when shield becomes weakness. *Interdiscip Sci*. 2016;8(2):150–155.
33. Lennard L, Cartwright CS, Wade R, Vora A. Thiopurinemethyltransferase and treatment outcome in the UK acute lymphoblastic leukaemia trial ALL2003. *Br J Haematol*. 2015;170(4):550–558.
34. Winter JW, Gaffney D, Shapiro D, et al. Assessment of thiopurinemethyltransferase enzyme activity is superior to genotype in predicting myelosuppression following azathioprine therapy in patients with inflammatory bowel disease. *Aliment Pharmacol Ther*. 2007;25(9):1069–1077.
35. Hindorf U, Appell ML. Genotyping should be considered the primary choice for pre-treatment evaluation of thiopurinemethyltransferase function. *J Crohns Colitis*. 2012;6(6):655–659.
36. Schmiegelow K, Forestier E, Kristinsson J, et al. Thiopurinemethyltransferase activity is related to the risk of relapse of childhood acute lymphoblastic leukemia: results from the NOPHO ALL-92 study. *Leukemia*. 2009;23(3):557–564.
37. Schaeffeler E, Fischer C, Brockmeier D, et al. Comprehensive analysis of thiopurine S-methyltransferase phenotype-genotype correlation in a large population of German-Caucasians and identification of novel TPMT variants. *Pharmacogenetics*. 2004;14(7):407–417.
38. Schmiegelow K, Al-Modhwhi I, Andersen MK, et al. Methotrexate/6-mercaptopurine maintenance therapy influences the risk of a second malignant neoplasm after childhood acute lymphoblastic leukemia: results from the NOPHO ALL-92 study. *Blood*. 2009;113(24):6077–6084.
39. Booth RA, Ansari MT, Loit E, et al. Assessment of thiopurine S-methyltransferase activity in patients prescribed thiopurines: a systematic review. *Ann Intern Med*. 2011;154(12):814–823, W-295–W-298.
40. Levinsen M, Rosthøj S, Nygaard U, et al. Myelotoxicity after high-dose methotrexate in childhood acute leukemia is influenced by 6-mercaptopurine dosing but not by intermediate thiopurinemethyltransferase activity. *Cancer Chemother Pharmacol*. 2015;75(1):59–66.
41. Tamm R, Magi R, Tremmel R, et al. Polymorphic variation in TPMT is the principal determinant of TPMT phenotype: a meta-analysis of three genome-wide association studies. *Clin Pharmacol Ther*. Epub 2016 Oct 22.
42. Ford L, Prout C, Gaffney D, Berg J. Whose TPMT activity is it anyway? *Ann Clin Biochem*. 2004;41(Pt 6):498–500.
43. Lennard L, Chew TS, Lilleyman JS. Human thiopurinemethyltransferase activity varies with red blood cell age. *Br J Clin Pharmacol*. 2001;52(5):539–546.
44. Yan L, Zhang S, Eiff B, et al. Thiopurinemethyltransferase polymorphic tandem repeat: genotype-phenotype correlation analysis. *Clin Pharmacol Ther*. 2000;68(2):210–219.
45. Lau J, Ioannidis JP, Terrin N, Schmid CH, Olkin I. The case of the misleading funnel plot. *BMJ*. 2006;333(7568):597–600.
46. Almoguera B, Vazquez L, Connolly JJ, et al. Imputation of TPMT defective alleles for the identification of patients with high-risk phenotypes. *Front Genet*. 2014;5:96.
47. Relling MV, Gardner EE, Sandborn WJ, et al. Clinical pharmacogenetics implementation consortium guidelines for thiopurinemethyltransferase genotype and thiopurine dosing: 2013 update. *Clin Pharmacol Ther*. 2013;93(4):324–325.
48. Lennard L, Van Loon JA, Lilleyman JS, Weinshilboum RM. Thiopurine pharmacogenetics in leukemia: correlation of erythrocyte thiopurinemethyltransferase activity and 6-thioguanine nucleotide concentrations. *Clin Pharmacol Ther*. 1987;41(1):18–25.
49. Lennard L, Gibson BE, Nicole T, Lilleyman JS. Congenital thiopurinemethyltransferase deficiency and 6-mercaptopurine toxicity during treatment for acute lymphoblastic leukaemia. *Arch Dis Child*. 1993;69(5):577–579.

50. Lennard L, Lilleyman JS, Van Loon J, Weinshilboum RM. Genetic variation in response to 6-mercaptopurine for childhood acute lymphoblastic leukaemia. *Lancet*. 1990;336(8709):225–229.
51. Dokmanovic L, Urosevic J, Janic D, et al. Analysis of thiopurine S-methyltransferase polymorphism in the population of Serbia and Montenegro and mercaptopurine therapy tolerance in childhood acute lymphoblastic leukemia. *Ther Drug Monit*. 2006;28(6):800–806.
52. Relling MV, Rubnitz JE, Rivera GK, et al. High incidence of secondary brain tumours after radiotherapy and antimetabolites. *Lancet*. 1999;354(9172):34–39.
53. Relling MV, Yanishevski Y, Nemec J, et al. Etoposide and antimetabolite pharmacology in patients who develop secondary acute myeloid leukemia. *Leukemia*. 1998;12(3):346–352.
54. Pui CH, Pei D, Sandlund JT, et al. Long-term results of St Jude Total Therapy Studies 11, 12, 13A, 13B, and 14 for childhood acute lymphoblastic leukemia. *Leukemia*. 2010;24(2):371–382.
55. Relling MV, Pui CH, Cheng C, Evans WE. Thiopurinomethyltransferase in acute lymphoblastic leukemia. *Blood*. 2006;107(2):843–844.
56. Rocha JC, Cheng C, Liu W, et al. Pharmacogenetics of outcome in children with acute lymphoblastic leukemia. *Blood*. 2005;105(12):4752–4758.
57. Moriyama T, Relling MV, Yang JJ. Inherited genetic variation in childhood acute lymphoblastic leukemia. *Blood*. 2015;125:3988–3995.
58. Schmiegelow K, Forestier E, Hellebostad M, et al. Long-term results of NOPHO ALL-92 and ALL-2000 studies of childhood acute lymphoblastic leukemia. *Leukemia*. 2010;24(2):345–354.
59. Schmiegelow K, Bjork O, Glomstein A, et al. Intensification of mercaptopurine/methotrexate maintenance chemotherapy may increase the risk of relapse for some children with acute lymphoblastic leukemia. *J Clin Oncol*. 2003;21(7):1332–1339.
60. Thomsen JB, Schröder H, Kristinsson J, et al. Possible carcinogenic effect of 6-mercaptopurine on bone marrow stem cells. *Cancer*. 1999;86(6):1080–1086.
61. Levinsen M, Rotevatn EØ, Rosthøj S, et al. Pharmacogenetically based dosing of thiopurines in childhood acute lymphoblastic leukemia: influence on cure rates and risk of second cancer. *Pediatric Blood Cancer*. 2014;61(5):797–802.
62. Lennard L, Cartwright CS, Wade R, Vora A. Thiopurine dose intensity and treatment outcome in childhood lymphoblastic leukaemia: the influence of thiopurine methyltransferase pharmacogenetics. *Br J Haematol*. 2015;169(2):228–240.
63. Stanulla M, Schaeffeler E, Flohr T, et al. Thiopurine methyltransferase (TPMT) genotype and early treatment response to mercaptopurine in childhood acute lymphoblastic leukemia. *JAMA*. 2005;293(12):1485–1489.
64. Stanulla M, Schaeffeler E, Moricke A, et al. Thiopurine methyltransferase genetics is not a major risk factor for secondary malignant neoplasms after treatment of childhood acute lymphoblastic leukemia on Berlin-Frankfurt-Munster protocols. *Blood*. 2009;114(7):1314–1318.
65. Liu YP, Wu HY, Yang X, et al. Association between thiopurine S-methyltransferase polymorphisms and thiopurine-induced adverse drug reactions in patients with inflammatory bowel disease: a meta-analysis. *PLoS One*. 2015;10(3):e0121745.
66. de Jong DJ, Derijks LJ, Naber AH, Hooymans PM, Mulder CJ. Safety of thiopurines in the treatment of inflammatory bowel disease. *Scand J Gastroenterol Suppl*. 2003;(239):69–72.
67. Hindorf U, Lindqvist M, Peterson C, et al. Pharmacogenetics during standardised initiation of thiopurine treatment in inflammatory bowel disease. *Gut*. 2006;55(10):1423–1431.
68. Ansari A, Arenas M, Greenfield SM, et al. Prospective evaluation of the pharmacogenetics of azathioprine in the treatment of inflammatory bowel disease. *Aliment Pharmacol Ther*. 2008;28(8):973–983.
69. Derijks LJ, Gilissen LP, Engels LG, et al. Pharmacokinetics of 6-mercaptopurine in patients with inflammatory bowel disease: implications for therapy. *Ther Drug Monit*. 2004;26(3):311–318.
70. Meggitt SJ, Gray JC, Reynolds NJ. Azathioprine dosed by thiopurine-methyltransferase activity for moderate-to-severe atopic eczema: a double-blind, randomised controlled trial. *Lancet*. 2006;367(9513):839–846.
71. van den Akker-van Marle ME, Gurwitz D, Detmar SB, et al. Cost-effectiveness of pharmacogenomics in clinical practice: a case study of thiopurinomethyltransferase genotyping in acute lymphoblastic leukemia in Europe. *Pharmacogenomics*. 2006;7(5):783–792.
72. Sayani FA, Prosser C, Bailey RJ, Jacobs P, Fedorak RN. Thiopurine methyltransferase enzyme activity determination before treatment of inflammatory bowel disease with azathioprine: effect on cost and adverse events. *Can J Gastroenterol*. 2005;19(3):147–151.
73. Payne K, Newman WG, Gurwitz D, Ibarreta D, Phillips KA. TPMT testing in azathioprine: a ‘cost-effective use of healthcare resources’? *Personal Med*. 2008;6(1):103–113.
74. Marra CA, Esdaile JM, Anis AH. Practical pharmacogenetics: the cost effectiveness of screening for thiopurine S-methyltransferase polymorphisms in patients with rheumatological conditions treated with azathioprine. *J Rheumatol*. 2002;29(12):2507–2512.
75. Thompson AJ, Newman WG, Elliott RA, Roberts SA, Tricker K, Payne K. The cost-effectiveness of a pharmacogenetic test: a trial-based evaluation of TPMT genotyping for azathioprine. *Value Health*. 2014;17(1):22–33.
76. Fargher EA, Tricker K, Newman W, et al. Current use of pharmacogenetic testing: a national survey of thiopurine methyltransferase testing prior to azathioprine prescription. *J Clin Pharm Ther*. 2007;32(2):187–195.
77. Zelinkova Z, Derijks LJ, Stokkers PC, et al. Inosine triphosphate pyrophosphatase and thiopurine s-methyltransferase genotypes relationship to azathioprine-induced myelosuppression. *Clin Gastroenterol Hepatol*. 2006;4(1):44–49.

Pharmacogenomics and Personalized Medicine

Publish your work in this journal

Pharmacogenomics and Personalized Medicine is an international, peer-reviewed, open access journal characterizing the influence of genotype on pharmacology leading to the development of personalized treatment programs and individualized drug selection for improved safety, efficacy and sustainability. This journal is indexed on the American Chemical

Submit your manuscript here: <https://www.dovepress.com/pharmacogenomics-and-personalized-medicine-journal>

Dovepress

Society's Chemical Abstracts Service (CAS). The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.