Stigma associated with mental illness: perspectives of university students in Qatar

Monica Zolezzi^{1,2}

Nawal Bensmail²

Farah Zahrah²

Salma Mawfek Khaled³

Tayseer El-Gaili⁴

¹Clinical Pharmacy and Practice,

²College of Pharmacy, ³Research Unit, Social and Economic Survey Research Institute, ⁴Student Counseling Center, Qatar University, Doha, Qatar

Background: Stigma in relation to mental illness is one of the main factors inhibiting people from seeking help. Studies have been undertaken looking into the knowledge, attitudes, and beliefs (KAB) about mental illness among residents in Qatar; however, none have looked specifically at students in higher education. The aim of this study was to understand the KAB toward mental illness among students at a Qatari university and determine if there are any differences based on gender, nationality, and college type.

Methods: A convenience sample of students from all genders, colleges, and nationalities was approached to participate in a survey that consisted of four sections: demographic, beliefs, attitudes, and help-seeking and treatment preferences associated with mental illness. Chi-square testing was performed to test for differences in the distribution of proportions of our primary outcomes (students' beliefs, attitudes, and help-seeking and treatment preferences).

Results: A total of 282 students completed the survey. The majority of the participating students were females (59.3%), non-Qataris (64.3%), and enrolled in science-based colleges (62.7%). Beliefs reflecting poor mental health literacy, such as "medications to treat mental illness can cause addiction", "mental illness is not like any other illness", or that "mental illness is a punishment from God", were reported by a majority of students (84.4%, 56.7%, and 50.2%, respectively). Stigmatizing attitudes that were endorsed by a majority of students included believing that people with mental illness cannot have regular jobs (60.2%), that people with mental illness are dangerous (65.7%), and that they would not marry someone with a mental illness (88.9%). Additionally, 33.6% of students indicated they would be ashamed to mention if someone in their family or they themselves, had a mental illness. A vast majority of students (86.3%) indicated to prefer family and friend's support as treatment options. Significant differences in KAB about mental illness between genders, colleges, and college type were found only for a few items.

Conclusion: The findings of this study suggest that a high percentage of students reported KAB about mental illness, that are considered stigmatizing. Students' KAB about mental illness was consistent with that reported by the general population in Qatar. Nevertheless, the sample was small and as such, larger studies from a randomly selected population are needed to confirm these findings.

Keywords: knowledge, attitude and behavior, university students, stigmatization, mental illness

Introduction

Worldwide reports indicate that one in every four people suffer from a mental illness throughout their life, which accounts for ~15% of the global disease burden.^{1,2} It is also known that many mental disorders can start at an early age, some as early as in childhood.² According to Qatar's National Mental Health Strategy 2013–2018, the point prevalence of mental illnesses in Qatari nationals is ~20%, with females and young adults (aged 18–34 years) being the most affected.³ In a study conducted in

Correspondence: Monica Zolezzi
Clinical Pharmacy and Practice, College of Pharmacy, Qatar University, Doha, Qatar
Tel +974 4403 5623
Fax +974 4403 5551
Email mzolezzi@qu.edu.qa

Qatar, it was reported that generalized anxiety disorders and major depression were the most prevalent among Qatari nationals.⁴ Despite the fact that these disorders are frequently encountered, significant underutilization of mental health care services by patients who suffer from mental illness has been reported in the international literature, with nearly two-thirds never seeking attention from a trained or qualified health care professional.¹

Stigma has been defined as a “sign of disgrace or discredit, which sets a person apart from others”.⁵ Stigma in mental illness can be described as a cluster of negative attitudes and beliefs that motivate fear, reject, avoid, and discriminate against people who have mental illnesses.⁵ Attitudes toward mental illness are also shaped by the individuals’ mental health literacy; that is, the knowledge and beliefs about mental disorders which aid their recognition, management, or prevention.⁶ Beliefs about mental illness are also shaped by knowing and interacting with someone living with mental illness, the media, and cultural stereotypes.^{5–8}

As such, beliefs and attitudes about mental illness influence how people interact with those with a mental illness, as well as how they experience and express their own emotional problems and psychological distress and whether they disclose these symptoms and seek care. In mental illness, stigma is one of the main factors that inhibit people from seeking help.⁹ Stigmatization occurs globally and is prevalent not only among the general public, but even among health care professionals.^{10,11} Although there are well-known manuals with validated criteria which are well known and used internationally for diagnosing mental illnesses (such as the *Diagnostic and Statistical Manual for Mental Disorders*¹² and the International Classification of Diseases),¹³ treatment seeking is influenced by the individual’s mental health literacy which is also culturally shaped, as the manifestations of any particular mental disorder may vary with the culture in which it presents.¹⁴ For example, in Malaysia, belief of psychiatric patients in supernatural causes was associated with greater use of traditional healers and poorer compliance with medication.¹⁵ Understanding the role of these influential factors is important in order to implement culturally appropriate strategies to manage mental disorders.

Qatar is a small country and its population comprises mostly non-Qatari expatriate workers of diverse ethnicity, predominantly from the Indian subcontinent and Arabs from various nationalities (mostly Egyptians, Syrians, Jordanians, Lebanese, Palestinians, Tunisians, and people from other Gulf countries).¹⁶ Considering this demographic pattern, studies to explore differences in the knowledge, attitudes,

and beliefs (KAB) about mental illness among residents in Qatar have been undertaken.^{17–19} In one of these studies in the general population, authors found that Qataris had poorer knowledge and more negative attitudes toward mental illnesses and their treatment, compared to non-Qataris.¹⁹

There are no current published studies examining the KAB toward mental illness among university students in Qatar. Considering that in Qatar, students in higher education comprise mostly young adults (the majority being females) in the age range of 18–34 years,⁴ who are also the most vulnerable for mental illness,³ it is important to examine if findings similar to those reported by the general population in Qatar are seen among university students. We hypothesized that university students in Qatar were better informed about mental illness than the general population and aimed at establishing if, within this population, there were any differences in KAB between Qataris and non-Qataris, females and males, and those attending science-based and non-science-based colleges.

Methods

A cross-sectional survey of students at Qatar University (QU) was carried out during the fall semester in 2015. The study was approved by the Institutional Review Board at QU (reference number: QU-IRB 501-E/15).

Sample

All undergraduate and graduate students at QU were eligible for inclusion, provided they were able to understand English. Researchers visited all colleges at QU and explained the study and its objectives to prospective participants. Informed consent was obtained from each student before the survey was completed.

Questionnaire

An adapted version of a questionnaire used to survey beliefs and attitudes toward mental illness in a sample of the general population in Qatar was used.¹⁹ Questions were changed considering the aims of this study and the student population that was being targeted. The final questionnaire consisted of four parts (Supplementary material):

1. Participants’ demographic characteristics (such as age, nationality, marital/living/financial status, educational background, and some information about the students’ parents).
2. Participants’ beliefs associated with mental illness. Questions in this section focused on the etiology of mental illness in addition to beliefs related to treatment.

3. Participants' attitudes toward individuals with mental illness. Questions in this section focused on the students' willingness to interact with individuals with mental illness, in order to identify any stigmatizing behavior on their part.
4. Participants' help-seeking preferences when it comes to management of mental illness. Questions in this section focused on students' approaches to treatment for mental illness, including treatment preferences.

The adapted version of the questionnaire was piloted on a sample of 20 randomly selected QU students from both genders in order to establish face validity, readability, clarity of questions, as well as time needed to complete the survey.

Data collection and analyses

A convenience (nonrandom sampling) approach was used. Researchers approached students from all genders, colleges, and nationalities at their respective colleges during the fall and spring semesters of the 2015–2016 academic year (November 2015–May 2016). An information sheet on the study was provided to the students, together with a consent form to give students more information about the study and its objectives. Consenting students were offered the choice for completing the questionnaire online (through SurveyMonkey® accessible via an iPad® computer) or were given paper-based questionnaires. Data from the paper-based questionnaires were later entered into SurveyMonkey by the researchers. Once data from all surveys were available, these were exported into SPSS® (version 21) for analysis.

Descriptive statistics were performed and reported as frequencies and percentages. The proportions (and percentages) were calculated based on valid responses, that is, nonmissing values and responses other than refusals. The latter were treated as missing and were not counted as part of the denominator for the corresponding proportion. "I don't know" answers were treated as valid responses when calculating the proportions of those endorsing stigmatizing statements with respect to beliefs and attitudes toward mental illness. Chi-square testing was performed to test for differences in the distribution of proportions of our primary outcomes (students' beliefs, attitudes, help-seeking and treatment preferences) between groups based on the following.

Nationality

Given the diversity in nationalities among non-Qatari students at QU (which for the most part include Egyptians, Syrians, Jordanians, Persians, Pakistanis, Sudanese, Indians, and Omanis, among others), these were grouped together in

order to provide more robust results when analyzing differences in KAB among Qatari and non-Qatari students.

Gender

Considering that the majority of students at QU comprise females, it was important to analyze differences in KAB among males and females.

College type

Assuming that students in science programs would be better informed about mental illness, QU colleges were classified as science-based (SB; colleges of medicine, engineering, pharmacy, and general sciences) and non-science-based (NSB; colleges of law, business, education, arts, and Islamic studies) to explore if any differences in KAB about mental illness exist among the students attending these two types of colleges.

Statements about beliefs were classified further to provide an indication of the students' level of awareness about mental health. Depending on the statement, a "yes" or "I don't know" answer was considered as reflecting poor awareness about mental illness in the part of the student (eg, the statement "do you think mental illness is due to possession by evil spirits?"); whereas for other statements, a "no" or "I don't know" answer would also be considered as being poorly aware about mental illness (eg, the statement "genetic inheritance can be the cause of mental illness"). Statements reflecting poor awareness were also considered to be potentially stigmatizing. A similar classification was used for the statements about attitudes toward people with mental illness. Depending on the statement, a "yes" or "I don't know" answer was considered as stigmatizing (eg, the statement "people with mental illness are dangerous"); whereas for other statements, a "no" or "I don't know" answer would also be considered as stigmatizing (eg, the statement "would you be willing to maintain friendship with someone with a mental illness?").

Multiple answers were allowed only for the section exploring students' help-seeking and treatment preferences. The level of significance for statistical tests was set a priori at $P < 0.05$.

Results

A total of 282 students completed the survey. A summary of the characteristics of the study sample is presented in Table 1. The majority of the student participants were females (59.3%), non-Qataris (64.3%), aged between 18 and 24 years (85.8%), single (92.5%), and living with their

Table 1 Demographic characteristics of the studied sample (N=282)

Students' characteristics	n (%)
Gender	
Male	114 (40.7)
Female	116 (59.3)
Nationality	
Qatari	100 (35.7)
Non-Qatari	180 (64.3)
Age group, years	
<18	17 (6.0)
18–24	241 (85.8)
25–30	20 (7.1)
>30	3 (1.1)
Social status	
Married	21 (7.5)
Single	259 (92.5)
Living status	
Alone	7 (2.5)
With family	271 (96.8)
With friends	2 (0.7)
Academic year	
First	69 (24.6)
Second	76 (27.0)
Third	64 (22.8)
Fourth	37 (13.2)
Fifth	14 (5.0)
Post-graduate	21 (7.5)
College	
Science based	177 (62.7)
Arts and sciences	37 (13.1)
Pharmacy	21 (7.4)
Engineering	79 (28.0)
Medicine	40 (14.2)
Non-science based	105 (37.2)
Business	57 (20.2)
Law	15 (5.3)
Education	25 (8.9)
Islamic studies	8 (2.8)
History of mental illness	
Yes	45 (18.9)
No	169 (71.0)
I don't know	24 (10.1)

Notes: N= total number of students completing the survey, n= number of valid responses, %= percent of valid responses.

families (96.8%). The majority of participants were enrolled in SB programs (62.7%), most of them at the College of Engineering (28.0%). The majority (71.0%) of the students who completed the survey indicated that they did not have a history of mental illness or had ever been told they have a mental illness.

Students' knowledge and beliefs associated with mental illness

The percent responses of the students' beliefs regarding mental illness are summarized in Table 2. Students' beliefs

Table 2 Stigmatizing beliefs about mental illness among different student cohorts

Beliefs about mental illness	Male (n=114)		Female (n=116)		χ^2	P-value	Qataris (n=100)		Non-Qataris (n=180)		χ^2	P-value	Science based (n=177)		Non-science based (n=105)		Total sample (N=282)	
	% stigmatizing response ^a (n/total)	% stigmatizing response ^a (n/total)	% stigmatizing response ^a (n/total)	% stigmatizing response ^a (n/total)			% stigmatizing response ^a (n/total)	% stigmatizing response ^a (n/total)	% stigmatizing response ^a (n/total)	% stigmatizing response ^a (n/total)			% stigmatizing response ^a (n/total)	% stigmatizing response ^a (n/total)	% stigmatizing response ^a (n/total)	% stigmatizing response ^a (n/total)	% stigmatizing response ^a (n/total)	% stigmatizing response ^a (n/total)
Mental illness is like any other illness ^a	64.2 (70/109)	52.4 (77/147)	58.3 (53/91)	57.0 (94/165)	0.01*	0.20	55.9 (90/161)	60.4 (58/96)	0.02*	57.4 (147/256)								
Mental illness is due to possession by evil spirits ^b	28.5 (31/109)	33.0 (50/148)	39.6 (36/91)	26.5% (44/166)	0.46	0.02*	28.4 (46/162)	36.5 (35/96)	0.39	31.5 (81/257)								
Chemical imbalances in the brain can result in mental illness ^a	34.0 (37/109)	15.6 (23/147)	24.2 (22/91)	22.4 (37/165)	<0.01***	0.20	16.7 (27/162)	34.7 (33/95)	<0.01**	23.4 (60/256)								
Mental illness is a punishment from God ^b	58.7 (64/109)	43.8 (64/146)	60.0 (54/90)	44.8 (74/165)	0.06	0.06	46.2 (74/160)	57.3 (55/96)	<0.01**	50.2 (128/255)								
Stress or traumatic events can result in mental illness ^a	4.6 (5/109)	10.9 (16/147)	17.8 (16/90)	3.6 (6/166)	0.11	<0.01***	5.6 (9/161)	13.5 (13/96)	0.01*	8.2 (21/256)								
Genetic inheritance can be the cause of mental illness ^a	36.7 (40/109)	34.0 (50/147)	38.9 (35/90)	33.2 (55/166)	0.33	0.65	31.6 (51/161)	41.7 (40/96)	0.11	35.2 (90/256)								
Mental illness can be treated ^d	28.4 (29/102)	25.2 (34/135)	26.5 (22/83)	26.8 (41/153)	0.03*	0.73	25.6 (38/148)	28.1 (25/89)	0.89	23.1 (63/237)								
Medications used to treat mental illness can cause addiction ^b	88.1 (96/109)	81.6 (120/147)	88.9 (80/90)	82.0 (136/166)	0.03*	0.10	84.5 (136/161)	82.6 (81/96)	0.23	84.0 (216/256)								

Notes: n= number of observations; total = total of valid responses; χ^2 = chi-square test; * $p<0.05$, ** $p<0.01$, *** $p<0.001$. ^aStigmatizing responses, those associated with poor awareness about mental illness, the aggregate of either "yes" and "do not know" or "yes" responses, depending on the statement about beliefs associated with mental illness.

reflecting poor awareness of mental illness included: “mental illness is like any other illness” (57.6% of students responding “no” and “I don’t know”), “mental illness is a punishment from God” (50.2% of students responding “yes” and “I don’t know”), and “medications to treat mental illness can cause addiction” (84.4% of students responding “yes” and “I don’t know”). Over a quarter of the students (26.6%) indicated that they believed mental illness cannot be treated.

Students’ attitudes toward mental illness

Table 3 shows the percentage of students endorsing stigmatizing attitudes toward the mentally ill. The majority of students indicated believing that people with mental illness cannot have regular jobs (60.2% answering “no” or “I don’t know”), that people with mental illness are dangerous (65.7% answering “yes” or “I don’t know”), and that they would not marry someone with a mental illness (88.9% answering “no” or “I don’t know”). Additionally, 33.6% indicated they would be ashamed to mention if someone in their families or themselves had a mental illness.

Students’ help-seeking and treatment preferences in relation to mental illness

As summarized in Table 4, the vast majority of students (84.5%) indicated that they would ask help from family members and 81.7% would consult a psychiatrist. Close to half of the students would consider seeking help from a pharmacist (44.8%) and only about one-third (37.3%) from a nurse. With respect to mental health treatments, the majority of students preferred support from family and friends (83.6%). Individual counseling (68.6%) and medications (68.1%) were among the top preferred treatments. Less than a quarter of the students indicated that marriage could be a treatment for mental illness (20.5%). Only 35.9% of the students were aware of the services provided by the university’s Student Counseling Center (SCC).

Gender differences

As summarized in Tables 2–4, significantly more male (64.2%) than female (52.4%) students believed that “mental illness is like any other illness” ($P=0.01$), that “chemical imbalances can result in mental illness” (male =34.0%, female =15.6%; $P<0.01$), that “mental illness can be treated” (male =28.4%, female =25.2%; $P=0.03$), and that “medications to treat mental illness can cause addiction” (male =88.1%, female =81.6%; $P=0.03$). Significantly more male than female students reported to be “willing to marry someone with a mental illness” (male =93.4%,

Table 3 Stigmatizing attitudes about mental illness among different student cohorts

Attitudes toward mental illness	Male (n=114)		Female (n=116)		χ^2 P-value		Qataris (n=100)		Non-Qataris (n=180)		χ^2 P-value		Science based (n=177)		Non-science based (n=105)		χ^2 P-value		Total sample (N=282)	
	% stigmatizing response ^a (n/total)	% stigmatizing response ^a (n/total)	% stigmatizing response ^a (n/total)	% stigmatizing response ^a (n/total)			% stigmatizing response ^a (n/total)	% stigmatizing response ^a (n/total)	% stigmatizing response ^a (n/total)	% stigmatizing response ^a (n/total)			% stigmatizing response ^a (n/total)	% stigmatizing response ^a (n/total)	% stigmatizing response ^a (n/total)	% stigmatizing response ^a (n/total)			% stigmatizing response ^a (n/total)	% stigmatizing response ^a (n/total)
People with mental illness can study at university ^a	41.5 (44/106)	52.1 (73/140)	0.10	51.1 (44/86)	45.9 (73/159)	0.42	43.1 (66/153)	54.8 (51/93)	0.16	47.6 (117/246)										
People with mental illness can work in regular jobs ^a	59.4 (63/106)	60.7 (85/140)	0.46	64.4 (56/87)	58.5 (93/159)	0.14	57.8 (89/154)	64.5 (60/93)	0.10	60.2 (148/246)										
People with mental illness can get married ^a	51.9 (55/106)	52.1 (73/140)	0.87	52.4 (45/86)	51.6 (82/159)	0.07	47.7 (73/153)	59.1 (55/93)	0.14	52.0 (128/246)										
People with mental illness can have children ^a	50.9 (54/106)	55.0 (77/140)	0.79	54.6 (47/86)	52.2 (83/159)	0.04*	50.9 (78/153)	57.0 (53/93)	0.20	53.3 (131/246)										
People with mental illness are dangerous ^b	73.6 (77/106)	60.4 (84/139)	0.14	70.1 (61/87)	62.7 (99/158)	0.34	64.7 (99/153)	66.7 (62/93)	0.03	65.7 (161/245)										
Willing to maintain friendship with someone with mental illness ^a	46.2 (49/106)	41.4 (59/140)	0.75	47.8 (42/88)	40.5 (64/158)	0.52	37.9 (58/153)	52.1 (49/94)	0.01*	43.5 (107/246)										
Willing to study/work with someone with a mental illness ^a	58.5 (62/106)	46.4 (65/140)	0.12	47.1 (41/87)	54.7 (87/159)	0.52	50.4 (77/153)	54.3 (51/94)	0.77	51.6 (127/246)										
Willing to marry someone with mental illness ^a	93.4 (99/106)	85.5 (118/138)	0.05	87.4 (76/87)	89.8 (141/157)	0.64	88.8 (135/152)	89.3 (83/93)	0.84	88.9 (217/244)										
Ashamed if someone in the family or themselves have a mental illness ^b	36.8 (39/106)	31.4 (44/140)	0.67	31.8 (28/88)	34.6 (55/159)	0.10	33.1 (51/154)	34.4 (32/93)	0.45	33.7 (83/246)										

Notes: n= number of observations; total = total of valid responses; χ^2 = chi-square test; * $p<0.05$. ^aStigmatizing responses, those associated with poor awareness about mental illness, the aggregate of either “no” and “do not know” or “yes” and “do not know” responses, depending on the statement about beliefs associated with mental illness.

Table 4 Help-seeking and treatment preferences

Sources of help ^a	Male (n=114)		Female (n=116)		χ^2 P-value		Qataris (n=100)		Non-Qataris (n=180)		χ^2 P-value		Science based (n=177)		Non-science based (n=105)		χ^2 P-value	
	% yes/maybe responses (n/total)		% yes/maybe responses (n/total)				% yes/maybe responses (n/total)		% yes/maybe responses (n/total)				% yes/maybe responses (n/total)		% yes/maybe responses (n/total)			
Family member	84.3 (86/102)		84.6 (110/130)		0.53		83.8 (67/80)		84.7 (128/151)		0.38		84.1 (122/145)		85.0 (74/87)		0.78	
Friend	85.3 (87/102)		70.8 (92/130)		<0.01**		72.8 (59/81)		79.3 (119/150)		0.46		76.6 (111/145)		78.2 (68/87)		0.91	
Psychiatric doctor	80.4 (82/102)		82.7 (105/127)		0.88		77.6 (62/80)		83.8 (124/148)		0.39		86.6 (123/142)		73.5 (64/87)		<0.01**	
Counselor	66.3 (67/101)		66.1 (84/127)		0.89		69.6 (55/79)		64.2 (95/148)		0.67		68.4 (97/142)		62.8 (54/86)		0.48	
Spiritual healer	63.7 (65/102)		59.8 (76/127)		0.31		60.7 (48/79)		62.4 (93/149)		0.17		59.5 (85/143)		65.2 (56/86)		<0.01***	
Pharmacist	42.6 (43/101)		46.7 (57/122)		0.82		45.5 (35/77)		44.9 (65/145)		0.35		45.7 (63/138)		43.5 (37/85)		0.95	
General physician	45.6 (46/101)		49.2 (62/126)		0.30		48.7 (38/78)		47.3 (70/148)		0.92		47.6 (67/141)		47.7 (41/86)		0.64	
Nurse	34.6 (35/101)		39.5 (49/124)		0.37		41.6 (32/77)		35.4 (52/147)		0.66		34.3 (48/140)		42.3 (36/85)		0.35	
Internet	62.4 (63/101)		68.6 (85/124)		0.49		64.5 (51/79)		66.9 (97/145)		0.80		68.6 (96/140)		61.2 (52/85)		0.41	
Treatment preferences^a																		
Medication	62.0 (62/100)		72.7 (96/132)		0.04*		67.1 (55/82)		68.5 (102/149)		0.86		71.2 (104/146)		62.8 (54/86)		0.41	
Natural product	50.0 (50/100)		50.4 (64/127)		0.94		54.3 (44/81)		47.6 (69/145)		0.14		47.1 (66/140)		55.2 (48/87)		0.44	
Support group	65.0 (65/100)		56.9 (74/130)		0.43		57.3 (47/82)		61.9 (91/147)		0.35		62.9 (90/143)		56.3 (49/87)		0.41	
Individual counseling	72.0 (72/100)		65.9 (85/129)		0.10		68.3 (56/82)		68.5 (100/146)		0.89		73.4 (105/143)		60.5 (52/86)		0.08	
Spiritual healing	54.0 (54/100)		48.0 (60/125)		0.30		45.0 (36/80)		54.2 (78/144)		0.09		60.0 (84/140)		35.3 (30/85)		<0.01***	
Family or friend's support	83.0 (83/100)		84.1 (111/132)		0.94		85.4 (70/82)		82.6 (123/149)		0.78		84.2 (123/146)		82.6 (71/86)		0.49	
Marriage	27.3 (27/99)		15.2 (19/125)		0.04*		17.3 (14/81)		22.5 (32/142)		0.64		21.0 (29/138)		19.8 (17/86)		0.82	
Cope by self	36.0 (36/100)		33.3 (42/126)		0.66		38.8 (31/80)		32.4 (47/145)		0.57		37.1 (52/140)		30.2 (26/86)		0.51	

Notes: n= number of observations; total = total of valid responses; *P<0.05, **P<0.01, ***P<0.001. ^aParticipants were able to select more than one source of help and treatment preferences.

female =85.5%; $P=0.05$), would ask for help from friends when confronted with a mental health problem (male =85.3%, female =70.8%; $P<0.01$), but significantly more females (72.7%) than males (62.0%) would use medications to treat mental illness ($P=0.04$). Significantly more male (27.3%) than female (15.2%) students reported that they would consider marriage as a treatment option for dealing with mental health problems ($P=0.04$).

Qatari versus non-Qatari differences

Significantly more Qatari (39.6%) than non-Qatari (26.5%) students believed that “mental illness is the result of possession of evil spirits” ($P=0.02$). Significantly more Qatari (17.8%) than non-Qatari (3.6%) students believed that “stress or traumatic events can result in mental illness” ($P<0.01$), and that “people with mental illness can have children” (Qatari =54.6%, non-Qatari =52.2%; $P=0.04$). No significant differences were found in help-seeking or treatment preferences for mental health problems between Qatari versus non-Qatari students.

Differences among students in SB versus NSB colleges

Significantly more students attending SB (44.1%) versus NSB (39.6%) colleges believed that “mental illness is like any other illness” ($P=0.02$), that “chemical imbalances in the brain can result in mental illness” (SB =83.3%, NSB =65.3%; $P<0.01$), and that “stress or traumatic events can result in mental illness” (SB =94.4%, NSB =86.5%; $P=0.01$). Significantly more students in NSB colleges indicated they perceive people with mental illness as dangerous (NSB =66.7%, SB =64.7%; $P=0.03$), but were also significantly more willing to maintain friendship with someone with mental illness (NSB =52.1%, SB =37.9%; $P=0.01$). Significantly more students in SB colleges reported they would seek help from a psychiatric doctor (SB =86.6%, NSB =73.5%; $P<0.01$), and significantly more students in NSB colleges reported they would seek help from a spiritual healer (NSB =65.2%, SB =59.5%; $P<0.01$).

Discussion

Although similar studies in the general population in Qatar have been published,^{17–19} to the best of our knowledge, this is the first exploring KAB about mental illness among university students. In contrast to what was hypothesized, university students’ stigmatizing beliefs were similar to those reported in the study by Bener and Ghuloum in the general population.¹⁹ Even though university students may be considered to be

well educated, several beliefs reflecting poor mental health literacy were highly prevalent, such as “medications to treat mental illness can cause addiction”, “mental illness is not like any other illness”, and “mental illness is a punishment from God”. Similar views were reported by medical students and the general public in Oman, who rejected a genetic factor as the cause of mental illness and rather favored the role of spiritual factors.²⁰ These findings suggest that other factors, such as some cultural and religious teachings, possibly influence beliefs about the origins and nature of mental illness.

The sample of university students in our study also shared similar negative attitudes toward people with mental illness as those reported by the general population in Qatar.¹⁹ Stigmatizing attitudes, such as that mentally ill individuals cannot hold regular jobs, or that they would not be willing to work with mentally ill individuals, were also highly prevalent in our study. In the Omani study, the majority of students also reported other negative attitudes such as that mentally ill individuals should not have children, get married, or maintain contact with the public.²⁰ Interestingly, more than half of QU students were comfortable with maintaining a friendship with someone who is mentally ill, an attitude that was reported by only less than a quarter of participants in the Qatar-based study in the general population.¹⁹ It is possible that this positive attitude by students compared to the general population is related to age. Other authors have found that older people had more negative attitudes (specifically social distancing) toward the mentally ill than their younger counterparts.²¹

Another important finding in our study was in relation to nationality differences in beliefs and attitudes toward people with mental illness. Qatari students in our study sample were less aware of the causes of mental illness and showed less favorable attitudes toward individuals with mental illness, compared to non-Qatari students. A possible explanation for these differences could be related to non-Qatari students having been exposed to other sociocultural environments, which may in turn contribute to an increased awareness about mental illness. Other authors with similar findings have also suggested that individuals who have lived or have had upbringing in other more “westernized” Arab societies may become more aware of mental illness and its treatments.²² Our results suggests that ethnic backgrounds can influence people’s knowledge and attitudes toward mental illness, and support existing literature that mental illness is a concept deeply tied to culture.^{7,19,22}

The majority of QU students prefer seeking the support from family and friends when facing psychological problems. Most students also held misconceptions regarding treatment,

such as believing that medications to treat mental illness can cause addiction, and mostly also appeared to underestimate the importance of seeking professional help for mental or psychological problems. Many studies in the Middle East have suggested that individuals may rather seek social support from family members or friends when faced with mental illness rather than seeking professional help.^{22–25} This help-seeking attitude has been attributed to the Arab culture and among individuals of Muslim faith, who have a strong family bond, and thus, involving non-family members in mental health issues, which implicates revealing familial or personal issues, is less acceptable than in western societies.^{23,26–28} Misconceptions (such as believing that medications used to treat mental illnesses are associated with addiction reported by 84.5% of students in our study) and lack of awareness (such as being unaware of the services provided by the university's Student Counseling Center) may be influencing students to rely more on friends and family for mental health support. These findings are consistent with those from another study conducted in the United Arab Emirates, in which most students were unaware of both individual and group counseling and psychological consultations provided on campus.²³ These findings highlight the need for educational interventions in this population to increase awareness about the importance of pharmacologic treatment and for improving access to mental health services within the university.²⁹

Differences among different student cohorts

Overall, male QU students appear to be better informed about the causes of mental illness than QU female students. However, overall attitudes toward mental illness did not appear to be significantly affected by gender. An exception to this finding was that QU male students appeared to be more likely than QU female students to marry someone with a mental illness, and also more males than females would consider marriage as a preferred treatment. Other studies in the Middle East have reported that females believe more in modern treatments as they were more inclined to consult professional mental health services for treatment.^{22,25,30} In our study, significantly more female students also preferred medication to treat mental illness. Combined, these results suggest female students may have a stronger inclination toward more modern treatments in comparison to their male counterparts.

Surprisingly, students in NSB colleges showed an overall favorable KAB about mental illness compared to those in SB colleges, except for believing that mental illness is a punishment from God. Results of studies in other universities in the

Middle East have shown conflicting results in this regard. For example, in a study in Jordan, medical students reported more favorable attitudes toward people with mental illness,²⁴ while a study in Oman reported that both medical students and the public rejected a genetic factor as the cause of mental illness and rather favored the role of spirits.²⁰ Combined, these results suggest that social and cultural underpinnings are more likely to influence students' KAB about mental illness than the college where these students receive their education.

Study limitations

The nationality variable was Qatari and non-Qatari. However, the non-Qataris represent a heterogeneous group from various nationalities, both from Arabic and non-Arabic speaking countries. Hence, there is a definite diversity within this group in their cultural beliefs and attitudes such that lumping them into one category may not be appropriate. Findings from this study are based on a convenience sample. Therefore, the proportion estimates and confidence intervals may not have their presumed frequency properties. Furthermore, the nonrandom nature of the sample may impact the validity and generalizability of our findings. In fact, the distributional properties of the study sample differed substantially from what is known about QU students' population, including that 65% of the actual population are Qataris, while the reverse was found to be true in our sample with approximately two-thirds reported being non-Qataris. It is also known that females make up ~70% of the students at QU, but were underrepresented in our sample at 53.3%. With this in mind, it is possible that our estimates of the proportions related to lack of knowledge about the roots of mental illness and corresponding negative attitudes and beliefs about people with mental illness may be an underestimate of the actual proportions in the larger QU student population. Multiple statistical tests were conducted simultaneously, which increases the type I error. This may explain why some results were statistically significant, though the differences between the groups were small. Therefore, caution must be taken when interpreting the results. Multiple responses were allowed for the main questions of interest in this study. This approach has the advantage of approximating real-life experiences by allowing participants to express multiple points of view toward the subject matter. Nevertheless, this type of question design poses analytical challenges. Specifically, the use of traditional Pearson's chi-squared tests is limited in the presence of multiple-response questions because of within-subject dependence among responses.³¹ While sophisticated statistical approaches that deal with this issue are beyond the scope of the paper, we acknowledge that

allowing multiple answers makes the results more difficult to interpret, and is thus a limitation of this study. Finally, due to the self-reported nature of the data, it is impossible to rule out social desirability bias. This type of bias can lead to participants choosing answers that they perceive as being more socially desirable and are not an actual representation of their perceptions. For example, they might have reported to be more willing to interact with mentally ill individuals, when in reality, they are not comfortable with such behavior; hence, this would result in falsely elevated nonstigmatizing attitudes toward people with mental illness.

Conclusion

The results of this study indicate that the students' beliefs about mental illness were interrelated to their attitudes, both of which set the stage for how they interact with people with mental illness as well as how they seek help and support when faced with a psychological problem themselves. The findings of this study were similar and consistent with those found in the general population in Qatar, supporting existing literature that attitudes and beliefs about mental illness are shaped by knowledge and cultural stereotypes. Nevertheless, the sample was small and as such, larger studies from a randomly selected population are needed in order to make more conclusive remarks. Because knowledge and attitudes are fundamental for reducing stigmatization, the study findings are important for determining the need for culturally specific targeted stigma intervention programs at the university level.

Treatment preferences were variable among students, although most students were not aware of the psychiatric help services provided by the university. These results suggest that strategies at the university to promote help-seeking in the part of the students are also needed. This could be done through awareness campaigns promoted by the SCC to educate students about the causes and the available treatment options for mental illnesses. Such a strategy is likely to also increase awareness of students about the services provided by the SCC. A collaborative strategy between the College of Pharmacy at QU and the SCC is also being planned to help support the SCC in educating QU students about medications. Future studies can find if awareness campaigns can have positive outcomes in terms of decreasing misperceptions among QU students in relation to mental illness.

Acknowledgments

Many thanks to all the students who accepted to participate in our study. We also acknowledge the help from Ahmed Al-Kaddour in survey distribution and data collection.

Disclosure

The authors report no conflicts of interest in this work.

References

1. World Health Organization (WHO). Mental disorders affect one in four people [Internet]. WHO; 2015 [cited March 21, 2017]. Available from: http://www.who.int/whr/2001/media_centre/press_release/en/. Accessed March 22, 2017.
2. National Alliance on Mental Illness (NAMI). Mental Health Conditions [Internet]. NAMI; 2015 [cited March 21, 2017]. Available from: <https://www.nami.org/Learn-More/Mental-Health-Conditions>. Accessed March 22, 2017.
3. Supreme Council of Health (SCH). Qatar National Mental Health Strategy Qatar [Internet]. SCH; 2015 [cited March 21, 2017]. Available from: <http://nhsq.info/app/media/1166>. Accessed March 22, 2017.
4. Bener A, Ghuloum S, Dafeeah E, Al-Yazidi T, Mustapha A, Zakareia A. Lifetime prevalence of common mental disorders in Qatar: using WHO composite international diagnostic interview (WHO-CIDI). *Int J Clin Psychiat Ment Health*. 2014;2(1):38–46.
5. Byrne P. Stigma of mental illness and ways of diminishing it. *Adv Psychiatr Treat*. 2000;6(1):65–72.
6. Jorm AF. Mental health literacy. Public knowledge and beliefs about mental disorders. *Br J Psychiatry*. 2000;177:396–401.
7. Rao D, Feinglass J, Corrigan P. Racial and ethnic disparities in mental illness stigma. *J Nerv Ment Dis*. 2007;195(12):1020–1023.
8. Corrigan PW, Markowitz FE, Watson AC. Structural levels of mental illness: stigma and discrimination. *Schizophr Bull*. 2004;20(3):481–491.
9. Thornicroft G. Stigma and discrimination limit access to mental health care. *Epidemiol Psychiatr Soc*. 2008;17(1):14–19.
10. Thornicroft G, Brohan E, Rose D, Sartorius N, Leese M. Global pattern of experienced and anticipated discrimination against people with schizophrenia: a cross-sectional survey. *Lancet*. 2009;373(9661):408–415.
11. Corrigan PW, Watson AC. Understanding the impact of stigma on people with mental illness. *World Psychiatry*. 2002;1(1):16–20.
12. American Psychiatric Association (APA). *The Diagnostic and Statistical Manual of Mental Disorders*. 5th ed. Washington, DC: APA; 2013.
13. World Health Organization (WHO). *International Classification of Diseases and Related Health Problems*. 10th revision. Geneva: WHO; 1992.
14. Kirmayer LJ. Cultural variations in the response to psychiatric disorders and psychological distress. *J Clin Psychiatry*. 2001;62(Suppl 13):22–28.
15. Razali SM, Khan UA, Hasanah CI. Belief in supernatural causes of mental illness among Malay patients: impact on treatment. *Acta Psychiatr Scand*. 1996;94(4):229–233.
16. Snoj J. Population of Qatar by nationality – 2017 report. [Internet, Feb 7, 2017]. [cited March 21, 2017]. Available from: <http://priyad-souza.com/population-of-qatar-by-nationality-in-2017/>.
17. Bener A, Ghuloum S. Gender differences in the knowledge, attitude and practice towards mental health illness in a rapidly developing Arab society. *Int J Social Psychiatry*. 2010;57(5):480–486.
18. Ghuloum S, Bener A, Burgut F. Epidemiological survey of knowledge, attitudes, and health literacy concerning mental illness in a national community sample: a global burden. *J Prim Care Community Health*. 2010;1(2):111–118.
19. Bener A, Ghuloum S. Ethnic differences in the knowledge, attitude and beliefs towards mental illness in a traditional fast developing country. *Psychiatria Danubina*. 2011;23(2):157–164.
20. Al-Adawi S, Dorvlo A, Al-Ismaïly S, et al. Perception of attitude towards mental illness in Oman. *Int J Soc Psychiatry*. 2002;48(4):305–317.
21. Yuan Q, Abidin E, Picco L, et al. Attitudes to mental illness and its demographic correlates among general population in Singapore. *PLoS One*. 2016;11(11):e0167297.
22. Al-Krenawi A, Graham JR, Al-Bedah EA, Kadri HM, Schwail MA. Cross-national comparison of Middle Eastern university students: help-seeking behaviors, attitudes toward helping professionals, and cultural beliefs about mental health problems. *Community Ment Health J*. 2009;45(1):26–36.

23. Al-Darmaki F. Needs, attitudes toward seeking professional help, and preferred sources of help among Emirati College students. *J Int Couns Educ*. 2011;3(1):39–57.
24. Rayan A, Jaradat A. Stigma of mental illness and attitudes toward psychological help-seeking in Jordanian university students. *Res Psychol Beha Sci*. 2016;4(1):7–14.
25. Bilican F. Help-seeking attitudes and behaviors regarding mental health among Turkish college students. *Int J Ment Health*. 2013; 42(2):43–59.
26. Dardas LA, Simmons LA. The stigma of mental illness in Arab families: a concept analysis. *J Psychiatr Ment Health Nurs*. 2015;22(9): 668–679.
27. Karam EG, Mneimneh ZN, Karam AN, et al. Prevalence and treatment of mental disorders in Lebanon: a national epidemiological survey. *Lancet*. 2006;367(9515):1000–1006.
28. Karim S, Saeed K, Rana MH, Mubhashar MH, Jenkins R. Pakistan mental health country profile. *Int Rev Psychiatry*. 2004;16(1–2):83–92.
29. Wright A, McGorry PD, Harris MG, Jorm AF, Pennell K. Development and evaluation of a youth mental health community awareness campaign. The Compass Strategy. *BMC Public Health*. 2006;6:215.
30. Al-Krenawi A, Graham JR, Kandah J. Gendered utilization differences of mental health services in Jordan. *Community Ment Health J*. 2000; 36(5):501–511.
31. Thomas RD, Decady YJ. Analyzing categorical data with multiple responses per subject. Statistics Society of Canada: proceedings of the Survey Methods Section; 121–130, 2000.

Supplementary material

Qatar University students' beliefs and preferences for the management of mental illness and psychological problems

Dear participant,

Thank you very much for agreeing to participate in this survey. The questionnaire is divided into 3 parts. Please try to answer ALL the questions to the best of your ability and as completely as possible.

Part 1: Demographics

1. Gender:

☐ Male
☐ Female
2. Nationality:

☐ Qatari
☐ Non-Qatari
3. Age group (years):

☐ <18
☐ 18–24
☐ 25–30
☐ >30
4. Social status:

☐ Married
☐ Single
5. If married: do you have children:

☐ Yes
☐ No
6. Academic year:

☐ First year
☐ Second year
☐ Third year
☐ Fourth year
☐ Fifth year
☐ Post-graduate
7. College:

☐ Arts & sciences
☐ Pharmacy

☐ Business
☐ Law

☐ Engineering
☐ Medicine

☐ Education
☐ Sharia
8. Parent's working status:

Father: ☐ Working in Qatar
☐ Working abroad

☐ Not working

Mother: ☐ Working in Qatar
☐ Working abroad

☐ Not working
9. Parent's highest education level:

Father: ☐ Primary school
☐ High school

☐ College/university

Mother: ☐ Primary school
☐ High school

☐ College/university
10. Living status:

☐ Alone
☐ With parents/family
☐ With friends
11. Do you have a scholarship:

☐ Yes
☐ No

Part 2: Beliefs associated with mental illness (select only ONE answer, "Yes", "No", or "I don't know")

Questions	Yes	No	I don't know
1. Do you think that mental illnesses are like any other illness?			
2. Do you think mental illness is due to possession by evil spirits?			
3. Do you think chemical imbalances in the brain can cause mental illness?			
4. Do you think mental illness can be punishment from God?			
5. Do you think stress in daily life, traumatic event, or shock can cause mental illness or psychological problems?			
6. Do you think genetic inheritance may be the cause of mental illness?			
7. Do you think medications used to treat mental illness cause addiction?			
8. Do you think mental illness can be treated (Select only ONE answer: "Yes", "No", or "I don't know")			

Part 3: Attitudes associated with mental illness (select only ONE answer, “Yes”, “No”, or “I don’t know”)

Question:	Yes	No	I don’t know
1. Do you think people with mental illness can live normal lives like:			
a. Study at University			
b. Work in regular jobs			
c. Get married			
d. Have children			
2. Do you think people with mental illness are dangerous?			
3. Would you be willing to:			
a. Maintain a friendship with someone with mental illness			
b. Study or work with someone with mental illness			
c. Marry someone with mental illness			
4. Are you ashamed to mention that someone in your family or yourself have a mental illness and/or experienced psychological problems?			

Part 4: Help-seeking and treatment preferences

Questions	Yes	No	I don’t know
1. During the course of your studies, have you ever had psychological problems or been told you have a mental illness? (Select only ONE answer: “Yes”, “No”, or “I don’t know”)			
2. Are you aware of the services provided by the Student Counseling Center at QU to help students with psychological problems? (Select only ONE answer: “Yes”, “No”, or “I don’t know”)			
3. Do you think pharmacists can be trusted to educate individuals with psychological problems on how to use their medications? (Select only ONE answer: “Yes”, “No”, or “I don’t know”)			
4. Do you think that QU Pharmacy students could have a role in helping fellow QU students with psychological problems by providing medication counseling services in collaboration with the QU Student Counseling Center? (Select only ONE answer: “Yes”, “No”, or “I don’t know”)			
5. If you are faced with a psychological problem who would you approach for help: (Select only ONE answer, “Yes”, “No”, or “Maybe”)			
Would you approach:	Yes	No	Maybe
a. Family members?			
b. Friends?			
c. Psychiatrist Doctor?			
d. Counselor?			
e. Spiritual/religious healer?			
f. Pharmacist?			
g. General Physician?			
h. Nurse?			
i. Internet?			
6. If you had a mental illness which treatment(s) would you prefer: (Select only ONE answer, “Yes” or “No”)			
a. Medication			
b. Support groups/group counseling			
c. Individual counseling			
d. Spiritual healing			
e. Family or friends support			
f. Natural product, like herbs			
g. Marriage			
h. I can get over the problem on my own			

Thank you for completing the survey.

Neuropsychiatric Disease and Treatment

Dovepress

Publish your work in this journal

Neuropsychiatric Disease and Treatment is an international, peer-reviewed journal of clinical therapeutics and pharmacology focusing on concise rapid reporting of clinical or pre-clinical studies on a range of neuropsychiatric and neurological disorders. This journal is indexed on PubMed Central, the 'PsycINFO' database and CAS,

and is the official journal of The International Neuropsychiatric Association (INA). The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <http://www.dovepress.com/neuropsychiatric-disease-and-treatment-journal>