Should physicians always rely on estimated glomerular filtration rate without knowing the equation?

Fabio Fabbian
Dario Priori
Alfredo De Giorgi

Department of Medical Sciences,
Clinica Medica Unit, University
Hospital St Anna, University of
Ferrara, Ferrara, Italy

Dear editor

We read with interest the paper written by Deskur-Smielecka et al¹ who investigated the performance of three equations for valuating renal function in 174 patients, aged 78 years, with different cancer diagnosis. They used Cockcroft–Gault (C–G), Modification of Diet in Renal Disease (MDRD), and Berlin Initiative Study1 (BIS1) equations and found that there is a considerable disagreement between renal function estimation formulas. We also performed two studies evaluating different formulas for the calculation of glomerular filtration rate (GFR) in patients with eating disorders² and in those with type 2 diabetes mellitus (DM).³ In the first study, GFR was evaluated in 55 patients with anorexia nervosa and 44 patients with bulimia nervosa using C–G, Mayo Clinic Quadratic (MAYO), and MDRD equations. Bland–Altman analysis suggested that the three formulae were not in agreement C–G formula was inaccurate when it was applied to obese or cachectic subjects, MDRD equation underestimated renal function in subjects with normal–high GFR, while MAYO equation seemed to be a good alternative to the other equations suggesting correct classification of patients. Subsequently, we evaluated 1,686 type 2 DM patients aged 68 years. GFR was estimated by MDRD, MAYO, Chronic Kidney Disease Epidemiology Collaboration (CKD-EPI), and C–G formulae. The whole population showed a better renal function when it was evaluated with MAYO and C–G formulae, and a worse renal function when it was calculated with MDRD formula. Bland–Altman analysis confirmed that formulae results were not in agreement, especially in the case of high GFR. Deskur-Smielecka et al¹ adjusted C–G equation using total body weight in underweight patients, ideal body weight in patients with normal body weight, and adjusted body weight in overweight and obese patients. They found that GFR estimated with MDRD equation was the highest, and the mean difference with the other two equations varied from 18 to 12 mL/min/1.73 m², confirming our results on the lack of agreement between different equations. However, they found a strong negative correlation between difference in GFR calculated with MDRD and C–G, and BIS1 and serum creatinine values. The correlation explained the worse performance of MDRD formula in subjects with normal serum creatinine levels.

MDRD and C–G equations' performances were compared with measured (125) I-iothalamate GFR in 828 patients with CKD and in 457 potential kidney donors. In CKD patients with GFR <60 mL/min/1.73 m², the MDRD equation performed better than the C–G formula. In the kidney donor group, the MDRD equation significantly

Correspondence: Fabio Fabbian
Department of Medical Sciences, Clinica
Medica Unit, University of Ferrara,
Via L. Ariosto 25–44121, Ferrara, Italy
Tel +39 532 23 7071
Fax +39 532 23 6816
Email f.fabbian@ospfe.it

underestimated the measured GFR when compared with the C–G formula. Moreover, both the MDRD and C–G equations overestimated the strength of the association of GFR with measured serum creatinine. The authors concluded that results derived from MDRD equation could be problematic in healthy individuals.⁴

All data suggest that sensitivity of C–G and MDRD equations in diagnosing CKD is lower when GFR is >60 mL/min/1.73 m².

Disclosure

The authors report no other conflicts of interest in this communication.

References

1. Deskur-Smielecka E, Kotlinska-Lemieszek A, Chdek J, Wieczorowska-Tobis K. Assessment of renal function in geriatric palliative care patients – comparison of creatinine-based estimation equations. *Clin Interv Aging*. 2017;12:977–983.
2. Fabbian F, Pala M, Scanelli G, Manzato E, Longhini C, Portaluppi F. Estimation of renal function in patients with eating disorders. *Int J Eat Disord*. 2011;44(3):233–237.
3. Fabbian F, Pala M, Monesi M, et al. The estimation of glomerular filtration rate in type 2 diabetic patients may depend on the equation used. *Eur Rev Med Pharmacol Sci*. 2013;17(20):2791–2797.
4. Poggio ED, Wang X, Greene T, Van Lente F, Hall PM. Performance of the modification of diet in renal disease and Cockcroft-Gault equations in the estimation of GFR in health and in chronic kidney disease. *J Am Soc Nephrol*. 2005;16(2):459–466.

Authors' reply

Ewa Deskur-Smielecka^{1,2}

Aleksandra Kotlinska-Lemieszek^{1,2}

Jerzy Chudek^{3,4}

Katarzyna Wieczorowska-Tobis^{1,2}

¹Department of Palliative Medicine, Poznan University of Medical Sciences, ²Palliative Medicine Unit, University Hospital of Lord's Transfiguration, Poznan, Poland; ³Pathophysiology Unit, Department of Pathophysiology, ⁴Department of Internal Medicine and Oncological Chemotherapy, Medical Faculty in Katowice, Medical University of Silesia, Katowice, Poland

Correspondence: Ewa Deskur-Smielecka
Department of Palliative Medicine, Poznan University of Medical Sciences, Hospicjum Palium, os. Rusa 55, 61-245 Poznan, Poland
Tel/fax +48 61 8738 303
Email edeskur@ump.edu.pl

Dear editor

The letter from Fabbian et al, in general, supports our findings concerning the disagreement between currently used equations for the estimation of glomerular filtration rate (GFR). We agree that disagreement in the estimation of kidney function was previously found by a number of authors. However, in our paper¹ we describe this aspect in very specific population – geriatric, palliative care patients – with high prevalence of cachexia and substantial muscle mass loss. Our group has shown some similarities (low creatinine generation) to the group of patients with anorexia nervosa and bulimia nervosa studied by Fabbian et al.² However, it has to be stressed that patients with anorexia nervosa and bulimia nervosa were much younger, and therefore had much better, mostly normal kidney function. The second population analyzed by Fabbian et al³ – diabetic patients – also differ from our cohort, as they were better nourished. Regardless of these differences, the conclusions concerning the disagreement of different

methods in the estimation of kidney function are quite similar. Importantly, the disagreement is more clinically significant in older population, including geriatric, palliative care patients, due to much greater prevalence of chronic kidney disease (CKD)⁴ that affects adjusting drug dosage.

Fabbian et al in their letter conclude that sensitivity of Cockcroft–Gault (C–G) and Modification of Diet in Renal Disease (MDRD) equations in diagnosing CKD is lower in subjects with GFR >60 mL/min/1.73 m². This statement, however, is controversial, as MDRD formula should not be used for the estimation of GFR >60 mL/min/1.73 m². Moreover, the process of kidney aging that results in the GFR decline moves the threshold level below 60 mL/min/1.73 m², and even 45 mL/min/1.73 m² in 80 years old and above.⁵

Disclosure

The authors report no other conflicts of interest in this communication.

References

1. Deskur-Smielecka E, Kotlinska-Lemieszek A, Chudek J, Wieczorowska-Tobis K. Assessment of renal function in geriatric palliative care patients – comparison of creatinine-based estimation equations. *Clin Interv Aging*. 2017;12:977–983.
2. Fabbian F, Pala M, Scanello G, Manzato E, Longhini C, Portaluppi F. Estimation of renal function in patients with eating disorders. *Int J Eat Disord*. 2011;44(3):233–237.
3. Fabbian F, Pala M, Monesi M, et al. The estimation of glomerular filtration rate in type 2 diabetic patients may depend on the equation used. *Eur Rev Med Pharmacol Sci*. 2013;17(20):2791–2797.
4. Chudek J, Wieczorowska-Tobis K, Zejda J, et al. The prevalence of chronic kidney disease and its relation to socioeconomic conditions in an elderly Polish population: results from the national population-based study PolSenior. *Nephrol Dial Transplant*. 2014;29(5):1073–1082.
5. KDOQI clinical practice guidelines for chronic kidney disease: evaluation, classification, and stratification. Available from: www.kidney.org/professionals/KDOQI/guidelines_ckd/toc.htm. Accessed July 12, 2017.

Dove Medical Press encourages responsible, free and frank academic debate. The content of the Clinical Interventions in Aging 'letters to the editor' section does not necessarily represent the views of Dove Medical Press, its officers, agents, employees, related entities or the Clinical Interventions in Aging editors. While all reasonable steps have been taken to confirm the content of each letter, Dove Medical Press accepts no liability in respect of the content of any letter, nor is it responsible for the content and accuracy of any letter to the editor.

Clinical Interventions in Aging

Publish your work in this journal

Clinical Interventions in Aging is an international, peer-reviewed journal focusing on evidence-based reports on the value or lack thereof of treatments intended to prevent or delay the onset of maladaptive correlates of aging in human beings. This journal is indexed on PubMed Central, MedLine,

CAS, Scopus and the Elsevier Bibliographic databases. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <http://www.dovepress.com/clinical-interventions-in-aging-journal>

Dovepress