

Pharmacogenomic testing: aiding in the management of psychotropic therapy for adolescents with autism spectrum disorders

Seuli Bose-Brill¹
Jinming Xing²
Debra J Barnette^{1,2}
Christopher Hanks¹

¹Internal Medicine and Pediatrics at Grandview, Wexner Medical Center,

²Department of Practice and Science, College of Pharmacy, The Ohio State University, Columbus, OH, USA

Abstract: Adolescents with autism have higher rates of anxiety than the general adolescent population. They often struggle to express psychological symptoms verbally where their symptoms may manifest as withdrawal and agitation. Adolescent patients with autism have higher rates of polypharmacy and high-risk psychiatric medication use (eg, atypical antipsychotics) than other patients with psychiatric illness. Primary care pediatricians are at the front lines of psychiatric management for patients with autism. Yet, they have inadequate access to pediatric psychiatry for complex medication management. Pharmacogenomic testing can provide personalized drug metabolism profiles for a majority of psychotropic medications. Primary care based pharmacogenomic testing for adolescents with autism on one or more psychiatric medications may help individualize and optimize complex medication regimens, while promoting drug safety.

Keywords: autism, pharmacogenomic, pharmacogenetic, pharmacotherapy, anxiety, depression, psychiatry, pediatrics

Introduction

Pharmacogenomic testing is a genetic test that helps assess and provide possible explanations for a patient's response or lack thereof, therefore aiding drug selection and dosing for improved outcomes. Although pharmacogenomics testing has been a crucial part in the personalized medicine movement, there is little clarity on how much it will benefit patients with autism spectrum disorder (ASD).

Pharmacogenomics

"Pharmacogenomics," (also referred to as pharmacogenetics) is a relatively new field, which studies how a patient's genes influence drug therapy response. Clinical pharmacogenomics testing has evolved from benchwork to direct patient care testing. It has received increased public attention since President Obama's call for personalized medicine in his 2015 State of the Union Address. Gene testing to understand a patient's genetic makeup permits tailoring the choice of medication therapy and/or dosing to an individual's determined drug metabolism. The patients will benefit through having fewer side effects and lower drug-related costs as a result of shorter hospital stays, reduced medical visits, and reduced medical absence days.¹ Genetic testing assessing common polymorphisms affecting selective serotonin reuptake inhibitors (SSRIs) metabolism may provide guidance to prescribers about individual drug metabolizing patterns for specific psychotropic medications.² Genetic differences have increasingly been shown to play a vital role in the variability of treatment response, optimal dosage,

Correspondence: Seuli Bose-Brill
Internal Medicine and Pediatrics at Grandview, Wexner Medical Center, The Ohio State University, 895 Yard Street, Columbus, OH 43212, USA
Email Seuli.Brill@osumc.edu

and tolerability.³ For example, an ultrarapid metabolizer phenotype to a particular drug can be a cause of therapeutic inefficacy, whereas a poor metabolizer phenotype to a drug carries an increased risk of toxicity.²

The use of pharmacogenomics in ASD

Side effects of psychotropic medications (some of which mimic anxiety-related symptoms such as irritability and insomnia), are related to individual pharmacogenetics influencing drug metabolism.⁴ With the profound side effect profile of atypical antipsychotics, pharmacogenomic testing could serve as a beneficial tool for reducing polypharmacy and negative side effects related to their use among adolescents with ASD. In adult medicine, it is likely that this testing modality will be ordered preemptively to aid in the selection and dosing of psychotropic medications in the near future.⁵ For adults, the Clinical Pharmacogenetics Implementation Consortium Guidelines from 2015 strongly recommends SSRI dosing adjustment based on the rate of metabolism through the CYP2C19 pathway (sertraline, citalopram, and escitalopram) and the CYP2D6 pathway (fluvoxamine).⁶ While specific pharmacogenomic guidelines have not been established for pediatrics or autism, there is precedence for use in these populations. For example, CYP2C19 testing in patients with autism identified a clear disparity in tolerability of SSRI dose titration between ultrarapid and normal metabolizers.⁷ At the same time, studies looking at how drug metabolism rates affect dosing at different stages of pediatric and adolescent development are needed to further refine their clinical application in adolescents with and without autism.⁸ Pharmacogenomic testing has already been specifically applied to the pediatric population for medications other than SSRIs. It has been used to identify patients who are at higher risk of respiratory depression from codeine,² predict toxicity in pediatric patients receiving 6-mercaptopurine and azathioprine, and predict risk for Stevens–Johnson syndrome in patients receiving carbamazepine treatment.^{9,10} There have been case reports where pharmacogenomics testing results were successfully utilized resulting in improved symptom management and overall quality of life among children and adolescents with complex behavior health issues.^{11,12} There is, therefore, precedence in pediatric care for pharmacogenomic testing in high-risk populations. While pharmacogenomic testing must be carefully researched and monitored prior to broad generalization to general adolescent populations, implementing pharmacogenomic testing in adolescents with

autism on one or more psychotropic medications may be warranted because of their high-risk profile.^{12,13}

Complex mental health issues

Data from the Center for Disease Control and Prevention in 2012 estimate 1 in 68 children have been diagnosed with ASD.¹⁴ Comparing to typical adolescents, young patients with ASD have more complex mental health concerns and frequently suffer from comorbid psychiatric disorders such as anxiety, attention deficit hyperactivity disorder, and obsessive compulsive disorder.¹⁵ Agitation, self-injury, and temper outbursts are often manifestations of anxiety in patients with autism who have difficulty verbally describing their symptoms.¹⁵ A practice parameter published in the Journal of the American Academy of Child and Adolescent Psychiatry suggests that pharmacotherapy may be offered to children with ASD when there is a specific target symptom or comorbid condition.¹⁶ While adolescents without ASD often have a hard time expressing psychiatric symptoms,¹⁷ adolescents with ASD may face behavioral and verbal challenges that make it even more difficult to describe symptoms. Adolescents with autism may also have difficulty expressing experienced medication side effects.¹⁸ Moreover, presenting symptoms may also vary by sex in ASD, with females tending to show more internalizing symptoms, such as withdrawal and males showing more externalizing behaviors such as aggression.¹⁹ This combination of complex manifestations of psychiatric symptoms and unrecognized side effects requires high levels of medication management expertise.²⁰

Complicated pharmacotherapy

Almost 1 in 10 patients aged 10–19 years are on some psychotropic medication.²¹ Patients with ASD are commonly diagnosed with anxiety and refractory depression.²² Several randomized clinical trials have been conducted indicating that cognitive behavioral therapy (CBT) is associated with positive treatment responses such as reduced anxiety in children with ASD.^{24,25} SSRIs are first line pharmacotherapy for patients with uncontrolled symptoms after behavioral interventions alone.¹⁶ Along with behavioral therapy, SSRIs have been indicated to reduce repetitive behavior among patients with ASD.²⁶ However, only half of pediatric patients with depression or anxiety who are prescribed a single SSRI have full remission of symptoms after 6 months.²⁷ In comparison, only one third of pediatric patients with autism who have anxiety have remission of symptoms with a single SSRI at 6 months. The remainder of patients either cycle through

numerous psychotropic medications due to side effects or are classified as having SSRI failure.²⁸ The side effect profile of SSRIs (such as increased nervousness) may exacerbate anxiety and depression symptoms.²⁹ Patients with ASD are more likely to be on psychotropic therapy for depression or other mental health conditions.³⁰ At the same time, they are less equipped to express potential side effects of medications or have full remission of symptoms with medication.^{18,31} This results in a higher risk of polypharmacy. Adolescents with ASD have higher prevalence of psychotropic use. Analysis of a national registry reviewing children with ASD indicated that >35% of this patient population use at least one psychotropic medication, and ~10% use psychotropic medications in ≥3 major classes concurrently.²³

Antipsychotics, stimulants, anticonvulsants, and antidepressants are often prescribed in combination for symptom control in these patients at much higher rates than the general population.³⁰ Over the past two decades, these psychotropic medications are increasingly being prescribed to children and adolescents with ASD, making polypharmacy among these patients a concerning issue.³² Teens with ASD are more likely to be on an atypical antipsychotic than those without autism.³³ These medications are often prescribed to manage symptoms related to resistant depression and anxiety, as well as behavioral problems in this population.³³ As superior efficacy of combination therapy has been demonstrated among pediatric patients in past clinical studies, pharmacological interventions are often added on top of behavioral therapy to help patients improve their daily function.^{34,35} Guidelines for adolescent depression in primary care suggest that treatment with antidepressants or psychotherapy such as CBT should be offered to adolescent patients with depression.³⁶ According to a 2011 report published by the Agency for Healthcare Research and Quality, 90% of children taking antipsychotic medications would receive an atypical antipsychotic for mostly off-label indications.³⁷ However, limited clinical data evaluating acute and long-term risks place these adolescents at a greater chance of undertreatment or overtreatment, in addition to long-term side effects.^{30,33} While regulatory approval for new indications depends on large randomized controlled clinical trials, studies in adolescents with autism have been limited in size and scope.³³ Other factors that intensify polypharmacy in this patient population include adolescence, handicap, and group home environment.³⁸ Because it may be very challenging for patients with ASD to express anxiety symptoms resulting intrinsically or from medication, a vicious cycle of chasing behavioral symptoms (unknowingly resulting from medication side effects) further

exacerbates psychotropic polypharmacy. With psychotropic medication's profound side effect profile (such as sedation, weight gain, metabolic disorders, hyperprolactinemia, movement disorders, and QTc prolongation),³⁹ it is imperative to maintain doses as low as possible while optimizing therapy for the best outcome.

Struggles of providers

Primary care providers are at the front lines of ASD management in the pediatric population. Families often have longstanding trusting relationships with them and often seek guidance from them both pre- and post-diagnosis.^{40,41} Pediatricians are encouraged to refer more complex patients, including those diagnosed with behavioral health disorders such as ASD, to mental health providers who practice in a multidisciplinary care team; the child and adolescent mental health workforce can currently care for only 10% of these patients, and they have limited tools to guide management during the often long waiting period to establish psychiatric care.^{27,40,42} In a recent study, 63% of pediatric patients with autism reported unmet needs, frequently lacking family services such as mental health and respite care.⁴³ About half of all pediatric patients who are provided a psychiatric referral actually see a mental health provider within 6 months.²⁷ These delays are due to an acute shortage of psychiatric providers who are comfortable providing care to adolescents with autism.⁴⁴

Pharmacogenomic testing in primary care setting

There are positive and negative logistical aspects to ordering pharmacogenomic testing in a primary care pediatrician's office. Over the past few years, the cost associated with pharmacogenomic testing has decreased drastically.⁵ With increased public awareness of such tests, parents and patients may specifically request testing from their provider. Such requests can help caregivers advocate for personalized psychotropic prescribing patterns, but require primary care providers to be familiar with testing. Many pharmacogenomic samples can be collected using a minimally invasive buccal swab, negating the need for venipuncture. Most samples can be processed within 7 days, allowing for real-time medication adjustments. Companies offering these testing services provide a comprehensive decision support analysis to assist in the genomic results interpretation and application to a wide array of medications within the selected therapeutic drug classes. The report is meant to provide guidance in selecting drug therapy based on that patient's specific results. Medicare and Medicaid provide insurance coverage for pharmacogenomic

testing, but many private insurance companies do not currently cover the cost of these tests.⁴⁵ Patients and parents may need to communicate directly with testing companies to determine out-of-pocket costs. These administrative burdens may dissuade patients with autism and their parents from pursuing this testing. Practices may have to facilitate such communication in addition to collecting insurance information for the testing vendor. In addition, outpatient clinics generally need to establish a relationship with specific pharmacogenomic testing vendors to obtain supplies and establish a practice workflow. Individual clinics and providers need to weigh these factors in deciding whether or not to offer these services for psychotropic medication management.

Proposed practice model

While pharmacogenomic testing is an emerging resource, it can be an effective tool for primary care pediatricians to assist in the management of adolescent patients with autism who are on psychotropic medication.¹¹ In our proposed practice, non-invasive gene testings can be offered to patients with autism who are on one or more of these medications. Patients who can potentially benefit from such testings are identified by our clinical staff using electronic health record based population management. Clinical pharmacists then will perform comprehensive medication review and pre-visit planning to further stratify for best candidates. Samples can be collected via buccal swab and are then sent for genetic testing. This testing is used to guide medication changes in patients who have suboptimal control of behavioral symptoms. Standardized behavioral symptom assessments in conjunction with medication monitoring can be used to monitor the impact of medication changes on behavior over time. Psychology practices are already using these tools as part of disease management. Practice guidelines are available, guiding clinicians on the use of pharmacogenomic testing with antidepressants and antipsychotics metabolized through CYP450 2D6 and CYP450 2C19.⁴⁶ Past study on major depression in an outpatient psychiatric practice also demonstrated improved outcomes with the use of pharmacogenomic testing.⁴⁷ While pharmacogenomic testings have been increasingly used for management of psychiatric symptoms in psychiatric practices, they are rarely used in primary care settings. If practices choose to incorporate pharmacogenomic testing, they should collect data on what impact it has on providers and patients so this information can be disseminated. Moreover, concurrent research and policy discussions should be pursued to ensure that this clinical application is optimized for pediatric patients with autism.

Conclusion

Adolescent patients with ASD often suffer from comorbid psychiatric disorders that may be more challenging to manage. They are frequently treated with complex drug regimens and are at increased risk of undertreatment and overtreatment, side effects, and polypharmacy. Although primary care providers are at the front lines of ASD management in this population, they are often overwhelmed and lack the right tools to better care for these patients. Through aiding in drug selection and dose optimization, pharmacogenomic testing may be beneficial in aiding better management of adolescent patients with ASD. Our proposed practice model should be individualized to best meet the needs of each practice.

Acknowledgments

The project described was supported by grant award number UL1TR001070 from the National Center for Advancing Translational Sciences. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Center for Advancing Translational Sciences or the National Institutes of Health.

Disclosure

The authors report no conflicts of interest in this work.

References

1. Perlis RH. Pharmacogenomic testing and personalized treatment of depression. *Clin Chem*. 2014;60(1):53–59.
2. Samer CF, Lorenzini KI, Rollason V, Daali Y, Desmeules JA. Applications of CYP450 testing in the clinical setting. *Mol Diagn Ther*. 2013;17(3):165–184.
3. Mrazek DA. Psychiatric pharmacogenomic testing in clinical practice. *Dialogues Clin Neurosci*. 2010;12(1):69–76.
4. Murphy GM, Kremer C, Rodrigues HE, Schatzberg AF. Pharmacogenetics of antidepressant medication intolerance. *Am J Psychiatry*. 2003;160(10):1830–1835.
5. Kieling C, Genro JP, Hutz MH, Rohde LA. A current update on ADHD pharmacogenomics. *Pharmacogenomics*. 2010;11(3):407–419.
6. Hicks JK, Bishop JR, Sangkuhl K, et al. Clinical pharmacogenetics implementation consortium (CPIC) guideline for CYP2D6 and CYP2C19 genotypes and dosing of selective serotonin reuptake inhibitors. *Clin Pharmacol Ther*. 2015;98(2):127–134.
7. Bishop JR, Najjar F, Rubin LH, et al. Escitalopram pharmacogenetics: CYP2C19 relationships with dosing and clinical outcomes in autism spectrum disorder. *Pharmacogenet Genomics*. 2015;25(11):548–554.
8. Stevens A, De Leonibus C, Hanson D, et al. Pediatric perspective on pharmacogenomics. *Pharmacogenomics*. 2013;14(15):1889–1905.
9. Gazouli M, Pachoula I, Panayotou I, et al. Thiopurine methyltransferase genotype and thiopurine S-methyltransferase activity in Greek children with inflammatory bowel disease. *Ann Gastroenterol*. 2012;25(3):249–253.
10. Wu XT, Hu FY, An DM, et al. Association between carbamazepine-induced cutaneous adverse drug reactions and the HLA-B*1502 allele among patients in central China. *Epilepsy Behav*. 2010;19(3):405–408.

11. Smith T, Sharp S, Manzardo AM, Butler MG. Pharmacogenetics informed decision making in adolescent psychiatric treatment: a clinical case report. *Int J Mol Sci.* 2015;16(3):4416–4428.
12. Kronenberg S, Apter A, Brent D, et al. Serotonin transporter polymorphism (5-HTTLPR) and citalopram effectiveness and side effects in children with depression and/or anxiety disorders. *J Child Adolesc Psychopharmacol.* 2007;17(6):741–750.
13. Bowers K, Lin PI, Erickson C. Pharmacogenomic medicine in autism: challenges and opportunities. *Paediatr Drugs.* 2015;17(2):115–124.
14. Christensen DL, Baio J, Braun KV, et al. Prevalence and characteristics of autism spectrum disorder among children aged 8 years – autism and developmental disabilities monitoring network, 11 sites, United States, 2012. *MMWR Surveill Summ.* 2016;65(3):1–23.
15. Leyfer OT, Folstein SE, Bacalman S, et al. Comorbid psychiatric disorders in children with autism: interview development and rates of disorders. *J Autism Dev Disord.* 2006;36(7):849–861.
16. Volkmar F, Siegel M, Woodbury-smith M, King B, McCracken J, State M. Practice parameter for the assessment and treatment of children and adolescents with autism spectrum disorder. *J Am Acad Child Adolesc Psychiatry.* 2014;53(2):237–257.
17. Achenbach TM, McConaughy SH, Howell CT. Child/adolescent behavioral and emotional problems: implications of cross-informant correlations for situational specificity. *Psychol Bull.* 1987;101(2):213–232.
18. Posey DJ, Stigler KA, Erickson CA, McDougle CJ. Antipsychotics in the treatment of autism. *J Clin Invest.* 2008;118(1):6–14.
19. Øien RA, Hart L, Schjølberg S, et al. Parent-endorsed sex differences in toddlers with and without ASD: utilizing the M-CHAT. *J Autism Dev Disord.* 2017;47(1):126–134.
20. Hepburn SL, Stern JA, Blakeley-Smith A, Kimel LK, Reaven JA. Complex psychiatric comorbidity of treatment-seeking youth with autism spectrum disorder and anxiety symptoms. *J Ment Health Res Intellect Disabil.* 2014;7(4):359–378.
21. Zito J, Safer DJ, Lolkje TW, et al. A three-country comparison of psychotropic medication prevalence in youth. *Child Adolesc Psychiatry Ment Health.* 2008;2(1):26.
22. Soorya L, Kiarashi J, Hollander E. Psychopharmacologic interventions for repetitive behaviors in autism spectrum disorders. *Child Adolesc Psychiatr Clin N Am.* 2008;17(4):753–771.
23. Rosenberg RE, Mandell DS, Farmer JE, Law JK, Marvin AR, Law PA. Psychotropic medication use among children with autism spectrum disorders enrolled in a national registry, 2007–2008. *J Autism Dev Disord.* 2010;40(3):342–351.
24. Reaven J, Blakeley-Smith A, Culhane-Shelburne K, Hepburn S. Group cognitive behavior therapy for children with high-functioning autism spectrum disorders and anxiety: a randomized trial. *J Child Psychol Psychiatry.* 2012;53(4):410–419.
25. Storch EA, Arnold EB, Lewin AB, et al. The effect of cognitive-behavioral therapy versus treatment as usual for anxiety in children with autism spectrum disorders: a randomized, controlled trial. *J Am Acad Child Adolesc Psychiatry.* 2013;52(2):132–142.
26. McDougle CJ, Kresch LE, Posey DJ. Repetitive thoughts and behavior in pervasive developmental disorders: treatment with serotonin reuptake inhibitors. *J Autism Dev Disord.* 2000;30(5):427–435.
27. Van Cleave J, Le TT, Perrin JM. Point-of-care child psychiatry expertise: the Massachusetts Child Psychiatry Access Project. *Pediatrics.* 2015;135(5):834–841.
28. Kennard BD, Silva SG, Tonev S, et al. Remission and recovery in the treatment for adolescents with depression study (TADS): acute and long-term outcomes. *J Am Acad Child Adolesc Psychiatry.* 2009;48(2):186–195.
29. Baldwin D. Generalized anxiety disorder. In: *Encyclopedia of Psychopharmacology.* Berlin, Heidelberg: Springer; 2010: 551–553.
30. Baribeau DA, Anagnostou E. An update on medication management of behavioral disorders in autism. *Curr Psychiatry Rep.* 2014;16(3):437.
31. Johnco CJ, De Nadai AS, Lewin AB, Ehrenreich-May J, Wood JJ, Storch EA. Defining treatment response and symptom remission for anxiety disorders in pediatric autism spectrum disorders using the Pediatric Anxiety Rating Scale. *J Autism Dev Disord.* 2015;45(10):3232–3242.
32. Coury DL, Anagnostou E, Manning-courtney P, et al. Use of psychotropic medication in children and adolescents with autism spectrum disorders. *Pediatrics.* 2012;130(Suppl 2):S69–S76.
33. Schneider C, Taylor D, Zalsman G, Frangou S, Kyriakopoulos M. Antipsychotics use in children and adolescents: an on-going challenge in clinical practice. *J Psychopharmacol.* 2014;28(7):615–623.
34. Wolraich M, Brown L, Brown RT, et al. ADHD: clinical practice guideline for the diagnosis, evaluation, and treatment of attention-deficit/hyperactivity disorder in children and adolescents. *Pediatrics.* 2011;128(5):1007–1022.
35. Autism Speaks. What is autism? 2015. Available from: <http://www.autismspeaks.org/what-autism>. Accessed May 13, 2017.
36. Cheung AH, Zuckerbrot RA, Jensen PS, Ghalib K, Laraque D, Stein REK. Guidelines for adolescent depression in primary care (GLAD-PC): II. Treatment and ongoing management. *Pediatrics.* 2007;120(5):e1313–e1326.
37. Centers for Medicare and Medicaid Services. Atypical antipsychotic medications: use in pediatric patients. 2013. Available from: <https://www.cms.gov/medicare-medicare-coordination/fraud-prevention/medicaid-integrity-education/pharmacy-education-materials/downloads/atyp-antipsych-pediatric-factsheet.pdf>. Accessed May 13, 2017.
38. Aman MG, Lam KS, Van bourgondien ME. Medication patterns in patients with autism: temporal, regional, and demographic influences. *J Child Adolesc Psychopharmacol.* 2005;15(1):116–126.
39. Wink LK, Pedapati EV, Horn PS, McDougle CJ, Erickson CA. Multiple antipsychotic medication use in autism spectrum disorder. *J Child Adolesc Psychopharmacol.* 2017;27(1):91–94.
40. Garg P, Lillystone D, Dossetor D, et al. A framework for developing a curriculum regarding autism spectrum disorders for primary care providers. *J Clin Diagn Res.* 2015;9(10):SC01–SC06.
41. Carbone PS, Behl DD, Azor V, Murphy NA. The medical home for children with autism spectrum disorders: parent and pediatrician perspectives. *J Autism Dev Disord.* 2010;40:317–324.
42. Thomas CR, Holzer CE. The continuing shortage of child and adolescent psychiatrists. *J Am Acad Child Adolesc Psychiatry.* 2006;45(9):1023–1031.
43. Farmer JE, Clark MJ, Mayfield WA, et al. The relationship between the medical home and unmet needs for children with autism spectrum disorders. *Matern Child Health J.* 2014;18(3):672–680.
44. Lake JK, Vogan V, Sawyer A, Weiss JA, Lunsky Y. Psychotropic medication use among adolescents and young adults with an autism spectrum disorder: parent views about medication use and healthcare services. *J Child Adolesc Psychopharmacol.* 2015;25(3):260–268.
45. Dougherty MJ, Wicklund C, Johansen Taber KA. Challenges and opportunities for genomics education: insights from an Institute of Medicine roundtable activity. *J Contin Educ Health Prof.* 2016;36(1):82–85.
46. De Leon J, Armstrong SC, Cozza KL. Clinical guidelines for psychiatrists for the use of pharmacogenetic testing for CYP450 2D6 and CYP450 2C19. *Psychosomatics.* 2006;47(1):75–85.
47. Hall-Flavin DK, Winner JG, Allen JD, et al. Utility of integrated pharmacogenomic testing to support the treatment of major depressive disorder in a psychiatric outpatient setting. *Pharmacogenet Genomics.* 2013;23(10):535–548.

Pharmacogenomics and Personalized Medicine

Dovepress

Publish your work in this journal

Pharmacogenomics and Personalized Medicine is an international, peer-reviewed, open access journal characterizing the influence of genotype on pharmacology leading to the development of personalized treatment programs and individualized drug selection for improved safety, efficacy and sustainability. This journal is indexed on the American Chemical

Society's Chemical Abstracts Service (CAS). The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/pharmacogenomics-and-personalized-medicine-journal>