Investigating a rare methicillin-resistant *Staphylococcus aureus* strain: first description of genome sequencing and molecular characterization of CC15-MRSA

Abiola C Senok¹Ali M Somily²Peter Slickers^{3,4}Muhabat A Raji⁵Ghada Garaween⁵Atef Shibl⁵Stefan Monecke^{3,4,6}Ralf Ehrlich^{3,4}

¹Department of Basic Science, College of Medicine, Mohammed Bin Rashid University of Medicine and Health Sciences, Dubai, United Arab Emirates; ²Department of Pathology and Laboratory Medicine, College of Medicine, King Khalid University Hospital and King Saud University, Riyadh, Saudi Arabia; ³Alere Technologies GmbH, Jena, Germany; ⁴InfectoGnostics Research Campus, Jena, Germany; ⁵Department of Microbiology and Immunology, College of Medicine, Alfaisal University, Riyadh, Saudi Arabia; ⁶Institute for Medical Microbiology and Hygiene (IMMH), Technische Universität Dresden, Dresden, Germany

Purpose: Methicillin resistant *Staphylococcus aureus* CC15 strains (CC15-MRSA) have only been sporadically described in literature. This study was carried out to describe the genetic make-up for this rare MRSA strain.

Methods: Four CC15-MRSA isolates collected in Riyadh, Saudi Arabia, between 2013 and 2014 were studied. Two isolates were from clinical infection and 2 from retail meat products. Whole genome sequencing was carried out using Illumina HiSeq2500 genome analyzer.

Results: All the CC15-MRSA isolates had the multilocus sequence typing profile ST1535, 13–13–1–1–81–11–13, which is a single locus variant of ST15. Of the 6 contigs related to the SCC element, one comprised a recombinase gene *ccrAA*, *ccrC-PM1*, *fusC* and a helicase, another one included *mvaS*, *dru*, *mecA* and 1 had *yobV* and *Q4LAG7*. The SCC element had 5 transposase genes, namely 3 identical paralogs of *tnpIS431* and 2 identical paralogs of *tnpIS256*. Two identical copies of a *tnpIS256*-based insertion element flank the *aacA-aphD* gene. Two copies of this insertion element were present with 1 located in the SCC element and another inserted into the *sasC* gene. A short 3 kb region, which lacks any bacteriophage structural genes and site-specific DNA integrase, was inserted into the *hbl* gene. The *hsdM* and the 5'-part of the *hsdS* gene are replaced by a copy of the *hsdM/hsdS* paralogs from *vSaa* giving rise to a new chimeric paralog of *hsdS* in *vSaa*.

Conclusion: CC15-MRSA shows a novel SCCmecV/SCCfus composite element. Its variant of *hsdM/hsdS* probably facilitated uptake of foreign mobile genetic elements that promoted emergence of CC15-MRSA. Close surveillance is needed to monitor spread and emergence of further CC15 MRSA strains.

Keywords: whole genome sequencing, MRSA, MLST, clonal complex, SCCmec, Saudi Arabia

Introduction

In recent years, the landscape of the molecular epidemiology of methicillin resistant *Staphylococcus aureus* (MRSA) has been characterized by the emergence and dissemination of new strains. Clonal complex 15 (CC15) is ubiquitous and widely described in the literature, but these isolates are mostly methicillin susceptible *S. aureus* (MSSA).¹ CC15-MSSA was recently identified as a predominant nasal colonizer in a report from Saudi Arabia.² Previously, methicillin resistant CC15 strains (CC15-MRSA) have only been sporadically described in literature.^{3–5} In a large scale genotyping study of MRSA isolates, no CC15-MRSA was identified.¹ Two isolates of CC15-MRSA associated with nasal colonization have been reported in Iran and Saudi Arabia.^{3,5} While whole genome sequencing data are available for CC15-MSSA, there are, to the best of our knowledge, no publications on the genomic data for the rare CC15-MRSA. Recently, we reported the first

Correspondence: Abiola C Senok
Mohammed Bin Rashid University of
Medicine and Health Sciences, Building
14, Dubai Healthcare City, PO Box
505055, Dubai, United Arab Emirates
Tel +971 4 383 8717
Email abiola.senok@mbru.ac.ae

identification of CC15-MRSA from clinical infections and retail meat products in the Middle East.^{6,7} In light of the emergence of CC15-MRSA in our setting and to provide much-needed insight into the genetic make-up of this rare MRSA clone, we have carried out whole genome sequencing of these isolates.

Materials and methods

The human isolates were identified as part of a larger MRSA study for which ethical approval was obtained from the Institutional Review Board, King Khalid University Hospital, Riyadh, Kingdom of Saudi Arabia. Patient consent was waived as the study involved use of archived isolates from specimens submitted for routine diagnostic tests and without use of patient identifiers. Four CC15-MRSA isolates collected in Riyadh, Saudi Arabia between 2013 and 2014 were studied. Two isolates (RUH-2 and RUH-71) were from patients with sepsis and wound infection, respectively, while the other 2 (RUH-98 and RUH-99) were from retail camel meat. *S. aureus* identification and confirmation of methicillin resistance was performed as previously described.^{6,7} Genomic DNA was extracted using Qiagen DNA isolation kit (Qiagen, Hilden, Germany) in accordance with manufacturer's instructions. Whole genome sequencing was carried out using the Illumina HiSeq2500 genome analyzer.

Sequencing reads were assembled de-novo with SPAdes and the final assembly was done with SPAdes version 3.10.1 (<http://bioinf.spbau.ru/spades>).⁸ Contigs shorter than 500 nt were dropped. Reads were mapped to the SPAdes contigs but also to the reference sequence (ST15-MSSA strain ST20130938, GenBank: CP012972.1⁹) with the Burrow–Wheeler aligner “bwa” using the local aligning algorithm “mem” (“bwa” version 0.7.12-r1039, <https://github.com/lh3/bwa>).¹⁰ We also used “bwa-mem” to map the whole SPAdes contigs on the reference sequence (ST15-MSSA strain ST20130938, GenBank: CP012972.1). Read mappings and coverage were visually inspected with “tablet” (“tablet” version 1.14.10.21, <https://ics.hutton.ac.uk/tablet/>).¹¹ We manually scaffolded and annotated the contigs from isolate RUH-2, which cover the genomic islands of *vSaβ* and *vSaa*, a 3 kb element inserted into the *hly* gene and the SCC element. We used the GenomeDiagram module from Biopython to draw sketches from the manually annotated sequences.¹² The reads and the SPAdes contigs were submitted to NCBI sequence database. The manually scaffolded and annotated regions were submitted to Genbank as short sequences.

Results

For each isolate, a de-novo assembly of the genomic sequence was carried out. The assemblies comprised 73 and 71 contigs

for the human isolates RUH-2 and RUH-71, respectively. Isolates RUH-98 and RUH-99 from camel meat had 72 and 66 contigs, respectively. The overall G/C content for the chromosomal contigs was 33%. All the CC15-MRSA isolates had the MLST profile 13–13–1–1–81–11–13. All of the 4 isolates sequenced carried a 30-kb plasmid harboring additional antibiotic resistance genes, namely *cadD*, *cadX*, *blaI*, *blaR*, *blaZ*, *lnuA*, *aadD*. In addition, isolate (RUH-71, from human wound infection) harbored another putative plasmidic contig encoding *tetK*. A comparison of the genomic features of the 4 CC15-MRSA isolates reported in this study (RUH-2, RUH71, RUH-98, RUH-99), with CC15-MSSA sequences in the NCBI GenBank (VCU006, MPROS1797, 08–02119, ST20130938, ST20130940, ST20130941) is given in Table S1.

The 6 contigs related to the SCC element (Figure 1) were identical in all isolates. One contig comprised a recombinase gene “*ccrAA*”, *ccrC-PM1*, *fusC* and a helicase; another contig included *mvaS*, *dru*, *mecA* and 1 contig had *yobV* and *Q4LAG7* (putative protein associated with SCCmec V/VT, GenBank AM990992.1: 52112 to 54100) (Figure 1). The SCC element presumably comprises 5 transposase genes, namely 3 identical paralogs of *tnpIS431* (size 675 nt) and 2 identical paralogs of *tnpIS256* (size 1173 nt) (Figure 1). Two identical copies of a *tnpIS256*-based insertion element flank the bifunctional kanamycin resistance determinant *aacA-aphD*. Two copies of this insertion element were present in the genome with 1 copy located in the SCC element and another copy inserted into the *sasC* gene encoding a surface protein.

The CC15-MRSA isolates had a short 3 kb region inserted into the *hly* gene (Figure 2). The 3 kb insertion element lacks any bacteriophage structural genes and site-specific DNA integrase. The insertion element comprised 5 genes, including *scn* (staphylococcal complement inhibitor) and *chp* (chemotaxis inhibitor), but *sak* (staphylokinase) was absent (Figure 2).

The CC15-MRSA isolates showed a variant of *hsdM/hsdS* at the major pathogenicity island *vSaa* compared with the reference CC15-MSSA genome (Figure 3). The *hsdM* and the 5'-part of the *hsdS* gene were replaced by a copy of the *hsdM/hsdS* paralogs from *vSaβ*. This gives rise to a new chimeric paralog of *hsdS* in *vSaa* (Figure 3). The chimeric *hsdS* has an intact reading frame. We can see this recombination in all of the 4 CC15-MRSA isolates. Furthermore, a *Sau3AI* restriction system is present in all of the CC15 isolates analyzed, while the type IV restriction system *SauUSI* is absent (Table S1).

Discussion

All the CC15-MRSA isolates had the MLST profile 13–13–1–1–81–11–13, which is a single locus variant of ST15. This

Figure 1 CC15-MRSA SCC element.

Notes: Six contigs related to the SCC element (MF185204 to MF185209). The first contig comprises the flanking *orfX* gene, and the last contig the flanking *dusC* gene. In between, four contigs are shown carrying genes typically found in SCC elements. Contig 2 has the fusidic acid resistance gene *fusC* and the SCC recombinase genes *ccrAA* and *ccrC*. Contig 3 comprises the *mecA* gene cluster with *mecR1* truncated by *tnpIS431*. Contig 5 constitutes a true insertion element, where the bifunctional kanamycin resistance determinant *aacAaphD* is flanked by two copies of *tnpIS256*.

Abbreviation: MRSA, methicillin-resistant *Staphylococcus aureus*.

MLST profile has been assigned to ST1535 (https://pubmlst.org/bigsdb?db=pubmlst_s aureus_isolates&page=profiles) and comprises pta-81 instead of pta-12 in canonical ST15. This pta-81 differs from pta-12 by only 1 single-nucleotide

polymorphism, which was present in all our isolates. Three of the 4 isolates assigned to ST1535 in the PUBMLST database are MSSA (https://pubmlst.org/bigsdb?db=pubmlst_s aureus_isolates&page=profiles). The fourth is MRSA isolate

Figure 2 The *hlb*-3kb-insert in CC15-MRSA.

Notes: The hemolysin beta gene (*hlb*) is interrupted by a 3 kb insertion element in CC15-MRSA genomes.

Abbreviation: MRSA, methicillin-resistant *Staphylococcus aureus*.

Figure 3 *hsdM/hsdS* recombination in CC15-MRSA.

Notes: The Figure shows the contents of genomic islands *vSaα* and *vSaβ* in isolate RUH-2 (ST1535/CC15, MFI85202, MFI85203) and in ST20130938 (ST15/CC15, Genbank accession CP012972.1). The reference genome CP012972.1 comprises two distinct paralogs of *hsdM/hsdS* in genomic islands alpha and beta. The mapping of the sequencing reads from isolate RUH-2 onto the reference sequence CP012972.1 reveals that *hsdM-alpha* and the 5'-end of *hsdS-alpha* are missing in RUH-2, while the coverage of *hsdM-beta* and the 5'-end of *hsdS-beta* is doubled with respect to other chromosomal genes, indicating that this stretch of DNA is duplicated in RUH-2. We extracted the duplicated region of *vSaβ* from the SPAdes contigs and were able to link it to contigs mapping to *vSaα*.

Abbreviation: MRSA, methicillin-resistant *Staphylococcus aureus*.

MPROS1797, which has a similar SCC element as the CC15 MRSA in this study (<https://www.ncbi.nlm.nih.gov/biosample/SAMEA2664415>; Table S1). Due to the presence of repeats, the SCC element could not be scaffolded into a single contiguous sequence. The overall constellation of the SCC element as shown in Figure 1 was interpreted as a novel SCCmecV/SCCfus composite element. A very similar element has also been found by microarray hybridization in CC97-MRSA from Saudi Arabia.¹³ Furthermore, reports from Saudi Arabia have described MRSA isolates from other lineages that also harbored SCCfus in addition to SCCmec IV

or V elements.^{6,13} Insertion elements flanked by 2 antiparallel copies of a transposase are common in bacteria, and often found in association with antibiotic resistance genes. The *sasC* gene, which is interrupted by insertion of another copy of the *tnpIS256*-based insertion element, has been linked with biofilm production in *S. aureus*.¹⁴

In *S. aureus*, an insert in the *hlb* gene is typically a prophage comprising several structural genes encoding the capsule, head and tail of the phage alongside an integrase at the terminus. It also frequently carries virulence associated genes like *sea*, *sep* (N315), *see*, *chp*, *sak* and *scn* in

various combinations.¹⁵ The Riyadh CC15-MRSA isolates (as well as other published CC15 genomes) had a short 3 kb region inserted into the *hly* gene (Figure 2). The absence of bacteriophage structural genes and site-specific DNA integrase suggests that it is no longer mobile but blocking the *hly* insertion site. This insertion element seems to be a remnant of a bacteriophage as it is homologous to the terminus of the *hly* converting phage in MRSA252 (Genbank accession BX571856.1, genes SAR2032 to SAR2036). In MRSA252, the *hly* converting phage has a size of about 44 kb. However, we did not find any resemblance between the putative SCC elements in our CC15-MRSA isolates and the SCC element in MRSA252. MRSA252 has a full *mec* gene cluster comprising *mecA/mecR1/mecI/mecR2*, while the *mec* cluster is truncated in *mecR1* in the CC15-MRSA isolates. MRSA252 has cassette recombinase *ccrA-2/ccrB-2*, while we find *ccrC* in the CC15-MRSA isolates. The *sak* gene is associated with tissue invasion.^{16,17} Its absence could be the reason why most CC15 isolates are associated with carriage rather than invasive infection.

Two distinct copies of the *hsdM* and *hsdS* genes are present in most genomes of *S. aureus*. These genes encode components of a type I restriction-modification (R-M) system (*hsdM* encodes a DNA methylase, *hsdS* encodes the specificity determinate). One pair of *hsdM/hsdS* is typically located in the genomic island *vSaa* between the superantigen-like genes *ssl10* and *ssl11*. A second pair of *hsdM/hsdS* genes resides in genomic island *vSaβ*. The CC15-MRSA isolates showed a different variant of *hsdM/hsdS* at the major pathogenicity island *vSaa* compared with the reference CC15-MSSA genome. This has arisen presumably due to intrachromosomal recombination, resulting in a repertoire of *hsdM/hsdS* restriction enzymes that deviate from the CC15 parent. This might have played a role in acquiring an SCC element as bacteria use the type I R-M system to control uptake of foreign DNA. In the type I R-M system, the *hsdM/hsdS* gene products are required in this process.¹⁸ Usually, the composition of the genomic island *vSaa* and *vSaβ* is highly conserved within clonal complexes. It has been shown that the type I R-M system has facilitated the evolution of distinct *S. aureus* lineages and controls the horizontal transfer of mobile genetic elements.¹⁸ The *Sau3AI* is a type II system that digests DNA at GATC sites and that was first described by Seeber *et al.*¹⁹ Interestingly, *Sau3AI* is present in all CC15 isolates, while *SauUSI* is absent (Table S1). This is rather unusual as *Sau3AI* is uncommon in *S. aureus*. Indeed, most *S. aureus* isolates harbor the *SauUSI* (type IV) at the DNA locus for

Sau3AI.^{19,20} Therefore, based on our findings, we suggest that changes in the *hsdM/hsdS* system and the type II R-M locus facilitated uptake of foreign mobile genetic elements, that is, of *SCCmec/SCCfus* by the ancestral CC15-MSSA promoting emergence of CC15-MRSA.

The limitation of our work is that the gaps between the contigs, which are presumably caused by repeated sequence elements, could not be resolved since the average fragment size of the Illumina library was only about 250 nt. Also, we were unable to determine the *spa* type reliably from our assembly since *spa* is a highly repetitive locus of a variable number of imperfect repeats. This genomic arrangement typically provokes artifacts in read assembly.

Accession numbers

The raw read sequences have been deposited in the Sequence Read Archive database (Bioproject PRJNA386092) with accession numbers: SAMN06925301, SAMN06925302, SAMN06925303, SAMN06925304.

De-novo assembled contigs have been deposited at DDBJ/ENA/GenBank under the accession NHZU000000,000, NHZV000000000, NHZW000000000, NHZX000000000. The version described in this paper is version NHZU01000000, NHZV01000000, NHZW01000000, NHZX01000000.

The manually scaffolded sequences for *hly_3kb_insert*, *vSaa*, *vSaβ* and the 6 SCC element contigs have been submitted to the GenBank under the following accession numbers: MF185201, MF185202, MF185203, MF185204, MF185205, MF185206, MF185207, MF185208, MF185209

Conclusion

We provide the molecular characterization of a MRSA strain from a common lineage that until recently gave rise only to very few MRSA. The findings indicate that CC15-MRSA has a novel *SCCmecV/SCCfus* composite element. Changes in the *hsdM/hsdS* system and the type II R-M locus probably played role in the emergence of this rare MRSA strain. Close surveillance is needed, especially with regard to spread among humans and livestock in the Middle East and emergence of further CC15 MRSA strains.

Acknowledgment

This work was presented in part at the 27th European Congress of Clinical Microbiology and Infectious Diseases (22–25 April 2017, Vienna, Austria).

Disclosure

The authors report no conflicts of interest in this work.

References

1. Monecke S, Coombs G, Shore AC, et al. A field guide to pandemic, epidemic and sporadic clones of methicillin-resistant *Staphylococcus aureus*. *PLoS One*. 2011;6(4):e17936.
2. Sarkar A, Raji A, Garaween G, et al. Antimicrobial resistance and virulence markers in methicillin sensitive *Staphylococcus aureus* isolates associated with nasal colonization. *Microb Pathog*. 2016;93:8–12.
3. Abou Shady HM, Bakr AE, Hashad ME, Alzohairy MA. *Staphylococcus aureus* nasal carriage among outpatients attending primary health care centers: a comparative study of two cities in Saudi Arabia and Egypt. *Braz J Infect Dis*. 2015;19(1):68–76.
4. Campanile F, Bongiorno D, Borbone S, Stefani S. Hospital-associated methicillin-resistant *Staphylococcus aureus* (HA-MRSA) in Italy. *Ann Clin Microbiol Antimicrob*. 2009;8:22.
5. Japoni-Nejad A, Rezazadeh M, Kazemian H, Fardmousavi N, van Belkum A, Ghaznavi-Rad E. Molecular characterization of the first community-acquired methicillin-resistant *Staphylococcus aureus* strains from Central Iran. *Int J Infect Dis*. 2013;17(11):e949–e954.
6. Senok A, Ehricht R, Monecke S, Al-Saedan R, Somily A. Molecular characterization of methicillin-resistant *Staphylococcus aureus* in nosocomial infections in a tertiary-care facility: emergence of new clonal complexes in Saudi Arabia. *New Microbes New Infect*. 2016;14:13–18.
7. Raji MA, Garaween G, Ehricht R, Monecke S, Shibl AM, Senok A. Genetic Characterization of *Staphylococcus aureus* Isolated from Retail Meat in Riyadh, Saudi Arabia. *Front Microbiol*. 2016;7:911.
8. Bankevich A, Nurk S, Antipov D, et al. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol*. 2012;19(5):455–477.
9. Trouillet-Assant S, Lelievre L, Martins-Simoes P, et al. Adaptive processes of *Staphylococcus aureus* isolates during the progression from acute to chronic bone and joint infections in patients. *Cell Microbiol*. 2016;18(10):1405–1414.
10. Li H. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM. 2013; arXiv:1303.3997v2. Available from: <https://arxiv.org/abs/1303.3997>. Accessed August 24, 2017.
11. Milne I, Stephen G, Bayer M, et al. Using Tablet for visual exploration of second-generation sequencing data. *Brief Bioinform*. 2013;14(2):193–202.
12. Pritchard L, White JA, Birch PR, Toth IK. GenomeDiagram: a python package for the visualization of large-scale genomic data. *Bioinformatics*. 2006;22(5):616–617.
13. Monecke S, Skakni L, Hasan R, et al. Characterization of MRSA strains isolated from patients in a hospital in Riyadh, Kingdom of Saudi Arabia. *BMC Microbiol*. 2012;12:146.
14. Speziale P, Pietrocola G, Foster TJ, Geoghegan JA. Protein-based biofilm matrices in *Staphylococci*. *Front Cell Infect Microbiol*. 2014;4:171.
15. van Wamel WJ, Rooijackers SH, Ruyken M, van Kessel KP, van Strijp JA. The innate immune modulators staphylococcal complement inhibitor and chemotaxis inhibitory protein of *Staphylococcus aureus* are located on beta-hemolysin-converting bacteriophages. *J Bacteriol*. 2006;188(4):1310–1315.
16. Monecke S, Luedicke C, Slickers P, Ehricht R. Molecular epidemiology of *Staphylococcus aureus* in asymptomatic carriers. *Eur J Clin Microbiol Infect Dis*. 2009;28(9):1159–1165.
17. Wieckowska-Szakiel M, Sadowska B, Rozalska B. Staphylokinase production by clinical *Staphylococcus aureus* strains. *Pol J Microbiol*. 2007;56(2):97–102.
18. Waldron DE, Lindsay JA. *SauI*: a novel lineage-specific type I restriction-modification system that blocks horizontal gene transfer into *Staphylococcus aureus* and between *S. aureus* isolates of different lineages. *J Bacteriol*. 2006;188(15):5578–5585.
19. Seeber S, Kessler C, Gotz F. Cloning, expression and characterization of the *Sau3AI* restriction and modification genes in *Staphylococcus carnosus* TM300. *Gene*. 1990;94(1):37–43.
20. Xu SY, Corvaglia AR, Chan SH, Zheng Y, Linder P. A type IV modification-dependent restriction enzyme *SauUSI* from *Staphylococcus aureus* subsp. *aureus* USA300. *Nucleic Acids Res*. 2011;39(13):5597–5610.

Supplementary material

Table S1 Comparison of CC15-MRSA and CC15-MSSA

Isolate name	RUH-2	RUH-71	RUH-98	RUH-99	VCU006	MPROS1797	08-02,119	ST2,01,30,938	ST2,01,30,940	ST2,01,30,941
Biosample accession	SAMN06925302	SAMN06925301	SAMN06925303	SAMN06925304	SAMN00138234	SAMEA2664415	SAMN04939716	SAMN04166246	SAMN04166494	SAMN04166543
Collection date	07-Nov-2013	04-Apr-2014	26-Oct-2014	26-Oct-2014						
Collection place	Saudi Arabia: Riyadh	Saudi Arabia: Riyadh	Saudi Arabia: Riyadh	Saudi Arabia: Riyadh						
Host	<i>Homo sapiens</i>	<i>Homo sapiens</i>	<i>Camelus dromedarius</i>	<i>Camelus dromedarius</i>						
Host disease	Sepsis	Wound infection								
Isolation source			Retail meat, neighborhood meat shop	Retail meat, neighborhood meat shop						
MLST	I535	I535	I535	I535	I5	I535	582	I5	I5	I5
Clonal complex	I5	I5	I5	I5	I5	I5	CC15/CC20 chimera	I5	I5	I5
SCC element	SCCmecV / SCCfus	SCCmecV / SCCfus	SCCmecV / SCCfus	SCCmecV / SCCfus	None	SCCmecV / SCCfus	None	None		
Paired end sequencing	2x51	2x51	2x51	2x51		2x100				
Average insert size	260	270	290	310		260				
Fragments sequenced	10461449	9646853	8433245	7667462		2852668				
Total number of bases	1067067798	983979006	860190990	782081124		570533600				
Estimated coverage	360	330	290	260		190				
WGS accession	SAMN06925302	SAMN06925301	SAMN06925303	SAMN06925304	AGTZ000000000.1		CP0156451	CP012972.1	CP012979.1	CP012978.1
Number of contigs	NHZU000000000	NHZV000000000	NHZW000000000	NHZX000000000						
capsular genotype (assembly)	73 contigs	71 contigs	72 contigs	66 contigs						
agr type (assembly)	8	8	8	8						
RIDOM spa type (assembly)	II	II	II	II						
RIDOM spa profile	t328, uneven coverage	t328, uneven coverage	t328, uneven coverage	t328, uneven coverage						
	07-23-12-34-	07-23-12-34-	07-23-12-34-	07-23-12-34-						
	34-12-12-23-	34-12-12-23-	34-12-12-23-	34-12-12-23-						
	02-12-23-02-	02-12-23-02-	23-02-12-23	02-12-23-02-						
	12-23	12-23		12-23						
RIDOM spa repeat count	14	14	14	14						

(Continued)

Table S1 Comparison of CC15-MRSA and CC15-MSSA

Isolate name	RUH-2	RUH-71	RUH-98	RUH-99	VCU006	MPROSI797	08-02,119	ST2,01,30,938	ST2,01,30,940	ST2,01,30,941
<i>cna</i> (assembly)	Missing	Missing	Missing	Missing	Missing	Missing	Missing	Missing	Missing	Missing
<i>sarT/sarU</i> (assembly)	Present	Present	Present	Present	Present	Present	Present	Present	Present	Present
<i>sasC</i>	Truncated	Truncated	Truncated	Truncated	Intact	Truncated	Intact	Intact	Intact	Intact
<i>tetK</i>	Missing	Present	Missing	Missing	Missing	Missing	Missing	Missing	Missing	Missing
<i>blaZ</i>	Present	Present	Present	Present	Present	Present	Present	Present	Present	Present
<i>fusC</i>	Present	Present	Present	Present	Missing	Present	Missing	Missing	Missing	Missing
<i>mecA</i>	Present	Present	Present	Present	Missing	Present	Missing	Missing	Missing	Missing
<i>hly</i>	Truncated	Truncated	Truncated	Truncated	Truncated	Truncated	Truncated	Truncated	Truncated	Truncated
<i>scn</i>	Present	Present	Present	Present	Present	Present	Present	Present	Present	Present
<i>chp</i>	Present	Present	Present	Present	Present	Present	Present	Present	Present	Present
3 kb <i>hly</i> insert	Present	Present	Present	Present	Present	Present	Present	Present	Present	Present
<i>sau3AI</i>	Present	Present	Present	Present	Present	Present	Present	Present	Present	Present
<i>sauUSI</i>	Missing	Missing	Missing	Missing	Missing	Missing	Missing	Missing	Missing	Missing

Notes: Comparison genome properties of CC 15-MRSA from this study with those of CC15-MSSA/MRSA from the NCBI GenBank.

Abbreviations: MRSA, methicillin-resistant *Staphylococcus aureus*; MSSA, methicillin susceptible *S. aureus*; MLST, multilocus sequence typing; WGS, whole genome shotgun.

Infection and Drug Resistance

Dovepress

Publish your work in this journal

Infection and Drug Resistance is an international, peer-reviewed open-access journal that focuses on the optimal treatment of infection (bacterial, fungal and viral) and the development and institution of preventive strategies to minimize the development and spread of resistance. The journal is specifically concerned with the epidemiology of antibiotic

resistance and the mechanisms of resistance development and diffusion in both hospitals and the community. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/infection-and-drug-resistance-journal>