

Role of poly (ADP-ribose) polymerase-1 and cytokines in acute renal injury in elderly Chinese patients with diabetes mellitus

Li Cao
Yan Liu
Li Zhang
Yanfei Wei

Intensive Medicine, Nanxishan
Hospital of Guangxi Zhuang
Autonomous Region, Guilin,
Guangxi, China

Objective: The primary objective of this study was to evaluate whether the polymorphism of poly (ADP-ribose) polymerase-1 (PARP-1) is involved as a potential risk factor in the development of acute renal injury in elderly Chinese patients with diabetes mellitus.

Subjects and methods: In this pilot study, diabetic patients of either gender (aged ≥ 65 years) with a confirmed diagnosis of acute renal injury and individuals with no clinical symptoms of acute renal injury were enrolled at Nanxishan Hospital of Guangxi Zhuang Autonomous Region, China. Genetic polymorphism of PARP-1 was assessed using polymerase chain reaction-restriction fragment length polymorphism assay. Cytokine levels (interleukin [IL]-6, IL-1 β and tumor necrosis factor- α) were measured in the serum samples by sandwich enzyme-linked immunosorbent assay technique.

Results: A total of 130 Chinese patients with acute renal injury and 130 Chinese individuals with no clinical symptoms of acute renal injury were included. We found that the patients with GG genotype and carriers of the G and C alleles of PARP-1 were at high risk of developing acute renal injury. Moreover, del/ins polymorphism of the NF- κ B1 gene was also found to be associated with acute renal injury. In addition, the levels of IL-6, IL-1 β and tumor necrosis factor- α were significantly increased in patients with acute renal injury ($p < 0.05$).

Conclusion: Our findings showed the involvement of PARP-1 polymorphisms in the development of acute renal injury in Chinese individuals. This study identified the involvement of two SNPs of PARP-1 (C410T and G1672A) in development of acute renal injury among Chinese diabetic patients. Also, increased expression of C and G alleles of PARP-1 can be considered as one of the potential risk factors for developing acute renal injury. Increased serum cytokine levels can be considered as one of the potential risk factors for developing acute renal injury.

Keywords: acute renal injury, poly (ADP-ribose) polymerase-1, diabetes mellitus, polymorphism

Introduction

Diabetes-induced renal injury is one of the most common chronic microvascular complications in patient of type 2 diabetes mellitus. In patients with uncontrolled diabetes, diabetic nephropathy is known to cause end-stage renal disease.¹ The key manifestations of diabetic nephropathy are albuminuria and high blood pressure, eventually lead to renal failure which is one of the most common causes of death in patients with diabetes.² Sequist et al showed the involvement of genetic susceptibility in diabetes-induced renal injury.^{3,4} Hur et al⁵ found the involvement of polymorphisms of poly (ADP-ribose) polymerase (single nucleotide polymorphisms [SNPs] 1963AG and β 28077GA) in development of nephritis among Korean patients.

Correspondence: Yanfei Wei
Intensive Medicine, Nanxishan Hospital
of Guangxi Zhuang Autonomous Region,
No 46 Chonxin Road, Xiangshan District,
Guilin, Guangxi 541002, China
Tel +86 1800 783 5532
Email yuearllg@163.com

It has been reported that poly (ADP-ribose) polymerase-1 (PARP-1)-mediated signaling can trigger an outburst of proinflammatory regulators and the cell death cascade is also well known.⁶ The PARP-1 catalyzes the process of PARYlation by attaching the polymers of ADP-ribose to target protein motifs via ester linkage and triggers an array of vital cell functions including chromatin structure, DNA repair, transcriptional regulation, apoptosis, necrosis, cell separation and differentiation.⁷ PARP-1 maintains the genomic integrity by PARYlation of histones and other key enzymes, the DNA repair mechanism, interprotein interactions and gene expression to ensure optimum cellular homeostasis.^{8–13} However, through cellular and/or genotoxic stress, prolonged activation of PARP-1 leads to a decrease in β -nicotinamide adenine dinucleotide and adenosine triphosphate and causes release of apoptosis-inducing factor, a mitochondrial proapoptotic protein.¹³ Such stress-induced dysregulation in cellular homeostasis mediated by PARP-1 triggers the downstream necrosis cascade of cell death.¹²

Hur et al¹⁵ suggested the involvement of PARP-1 polymorphisms in the development of nephritis among Korean patients and identified the involvement of two SNPs (1963AG and β 28077GA). However, there was no study evaluating the involvement of PARP-1 polymorphisms in development of acute renal injury in diabetic patients. The primary objective of this study was to evaluate whether the polymorphism of PARP-1 is involved as a potential risk factor in the development of acute renal injury in elderly Chinese patients with diabetes mellitus. Our pilot study was designed to assess involvement of PARP-1 alleles and SNPs in development of acute renal injury in diabetic patients. We also investigated the cytokine levels among Chinese patients with acute kidney injury. Our study results can serve the basis for conducting large multicentric randomized clinical study.

Subjects and methods

In this pilot study, the diabetic patients of both genders (age ≥ 65 years) with a confirmed diagnosis of acute renal injury and individuals with no acute renal injury (control group) were enrolled at Nanxishan Hospital of Guangxi Zhuang Autonomous Region, China. The study was approved by the institutional ethics committee of Nanxishan Hospital of Guangxi Zhuang Autonomous Region, and written consent was obtained from each study participant. All participants were informed about the study procedures and the potential benefits to society. All participants underwent laboratory tests to confirm their eligibility.

Plasma samples were collected from each participant, and DNA from leukocytes was extracted by the high salt

DNA extraction method. All isolated DNA samples were kept at -80°C for further assessment. Polymorphisms of the PARP-1, NF- κB1 and NF- κBIA genes were studied in DNA samples by polymerase chain reaction-restriction fragment length polymorphism analysis. For the PARP-1 C410T and G1672A promoter regions, 297 and 187 base pair polymerase chain reaction (PCR) fragments were amplified in a 25 μL reaction volume at pH 8.3 consisting of 50 mM KCl, 100 ng genomic DNA, 200 pM deoxynucleotide, 200 pM of 10 mM Tris-HCl, 1 U Taq polymerase (Sigma) and 2 mM MgCl_2 (Fermentas). The upstream sequences of PARP-1 promoter regions C410T and G1672A were 5'-TCCAGTGGCACTA TCAT-3' and 5'-GCGAGACCCTGTCCCTAA-3', respectively. The downstream sequences of PARP-1 promoter regions C410T and G1672A were 5'-GTTGTGA GACATAGGCCGAAT-3' and 5'-TCCCCCTTTTATTT TTGAGACTG-3', respectively. The polymerase chain reaction-restriction fragment length polymorphism was performed using an optimized sequence of cycles: a denaturation phase at 95°C (2 min) followed by 30 cycles at 95°C (30 s), 60°C (30 s), 72°C (1 min) and a final incubation phase at 72°C (5 min). The PCR products were treated with 5 U of HpyF3I (DdeI; Fermentas) and Bsh1236I (Fermentas) at 37°C overnight for digestion. The resulting fragments were run on ethidium bromide-stained 3% agarose gel for 45 min at 90 V and quantified through direct detection under ultraviolet light (Figures 1 and 2). Likewise, the NF- κB1 and NF- κBIA genes were amplified by 285 and 424 base pair PCR fragments using the same reagents. The PCR running sequence included a denaturing step at 95°C (1 min) followed by 35 cycles at 95°C (30 s), 61°C (30 s), 72°C (1 min) and a final incubation at 72°C (5 min). The PCR products were treated with 5 U PfuMI (Van91I) and HaeIII (BsuRI; Fermentas) at 37°C overnight for digestion and the digested products were run on ethidium bromide-stained 3% agarose gel for 45 min at 90 V and quantified directly under ultraviolet light. Cytokine levels (interleukin [IL]-6, IL-1 β and tumor necrosis factor alpha [TNF- α]) were measured in the serum samples by a specific sandwich enzyme-linked immunosorbent assay technique using commercial kits (Assaypro Human IL-6, IL-1 β and TNF- α) coated with antibodies specific to human leptin and adiponectin antibodies, according to the manufacturer's protocols (St Charles).

Statistical analysis

The study was designed as a preliminary pilot study to assess the polymorphism association of PARP-1 with acute renal injury in Chinese diabetic patients. Hence, no formal sample




Figure 1 PARP-I C410T polymorphism (A) and G1672A polymorphism (B).

Notes: (A) Sample 1 shows TT genotype, samples 2 and 3 show CC genotype and samples 4 and 5 show CT genotype of PARP-I C410T. (B) Samples 1 and 4 show GA genotype, samples 2 and 3 show GG genotype and sample 5 shows AA genotype of PARP-I G1672A. Patient-related data were masked in the figure.

Abbreviation: PARP-I, poly (ADP-ribose) polymerase-I.

size calculation was performed. We planned to enroll at least 100 individuals in each group (patients with acute renal injury and patients without acute renal injury). Our study can serve as a basis for large genetic clinical studies to assess PARP-I polymorphism in patients with acute renal injury. Quantitative variables were presented as mean (\pm SD) and analyzed by parametric and nonparametric statistical tests depending on the number of groups for comparison and the distribution of data, using two-sided statistical tests. Categorical variables were presented as absolute numbers and/or percentages of subjects in each category and analyzed by the chi-square or Fisher's exact test depending on the size of the data, using two-sided statistical test. Demographic characteristics were presented by descriptive statistics, using mean (SD) (for numerical data) and as absolute values (for categorical data). Genotype involved in gene polymorphism in patients with acute renal injury was analyzed using univariate analysis. Odds ratios (ORs) were calculated based on univariate analysis using the chi-square or Fisher's exact test depending

on the size of the data. Plasma levels of cytokines in individuals with acute renal injury and without acute renal injury by different genotypes were analyzed either using unpaired *t*-test or Mann–Whitney test, based on normality of data. Kolmogorov–Smirnov test or Shapiro–Wilks test was used to check the distribution of numerical data. Unpaired *t*-test was applied if the data followed normal distribution (parametric data), whereas Mann–Whitney test was applied if data followed non-normal distribution (nonparametric data). All the statistical tests were two-sided statistical tests. The data of each study participant were coded, and analyzed using the GraphPad Prism statistical software (version 6.0).

Results

A total of 130 Chinese patients with acute renal injury and 130 individuals without acute renal injury (healthy groups) were enrolled and completed the study. The data of all the enrolled individuals were analyzed. Demographic characteristics were found to be similar in both the groups (Table 1).




Figure 2 NF-κB1 polymorphism (A) and NF-κB1A polymorphism (B).

Notes: (A) Samples 2 and 6 show del/del genotype, samples 4, 5, 7, 8 and 10 show del/ins genotype and sample 9 shows ins/ins genotype. (B) Samples 3 and 9 show AA genotype, samples 2, 4, 7 and 8 show GG genotype and samples 5, 6 and 10 show GA genotype. Patient-related data were masked in the figure.

Table 1 Demographic characteristics of healthy individuals and acute renal injury patients

Variables	Individuals with acute renal injury, n=130	Healthy individuals, n=130
Age (years)	68.7 (±5.3)	66.7 (±4.6)
Gender (male/female)	80/50	78/52
BMI	27.13 (3.45)	26.82 (3.43)

Note: Values are expressed as mean (±SD) for age and BMI, and absolute values are presented for gender.

Abbreviation: BMI, body mass index.

For the G1672A genotype, we found that the individuals with a G allele in PARP-1 were also at very high risk of developing acute renal injury (OR 3.96, 95% CI 2.33–6.74, $p<0.0001$). Also, we observed that the individuals with a GG genotype in the PARP-1 gene were at very high risk of developing acute renal injury (OR 4.09, 95% CI 2.42–6.90, $p<0.001$). There was no involvement of other genotypes such as GA or AA among patients with acute renal injury. None of the other genotypes of NF- κ B1, NF- κ B1A and PARP-1 were found to be involved (Table 2). Moreover, no evidence

of PARP-1 polymorphisms was observed in individuals with any acute renal injury.

For the C410T genotype, we found that the individuals with a C allele in the PARP-1 gene were at high risk of developing acute renal injury (OR 3.77, 95% CI 1.99–7.12, $p<0.0001$). Among patients with acute renal injury, del/ins of the NF- κ B1 gene was also noted (OR 3.32, 95% CI 1.96–5.61, $p<0.001$). This suggests the involvement of del/ins of NF- κ B1 in the development of acute renal injury. No statistically significant association of other NF- κ B1 genotypes with acute renal injury was observed (Table 2). We also studied the role of cytokines in acute renal injury in relation to different genes. The results showed a significantly higher level of cytokines (IL-6, IL-1 β and TNF- α) in acute renal injury patients as compared to patients without acute kidney injury (Table 3).

Discussion

This was the first study to assess involvement of PARP-1 alleles and SNPs in development of acute renal injury among

Table 2 Genotypes involved in gene polymorphism in patients with acute renal injury

Genotype	Individuals with acute renal injury, n=130, n (%)	Healthy individuals, n=130, n (%)	Odds ratio (95% CI)	p-value
G1672A genotype				
GG	88 (67.7)	42 (32.3)	4.09 (2.42–6.90)	<0.001
GA	43 (33.1)	41 (31.5)	1.07 (0.63–1.80)	0.83
AA	22 (16.9)	21 (16.2)	1.05 (0.54–2.03)	0.86
Allele frequency in G1672A genotype				
G allele	72 (55.4)	31 (23.8)	3.96 (2.33–6.74)	<0.001
A allele	34 (26.2)	32 (24.6)	1.08 (0.62–1.89)	0.88
C410T genotype				
CC	86 (66.2)	78 (60.0)	1.15 (0.88–1.75)	0.08
CT	34 (26.2)	32 (24.6)	1.08 (0.71–1.31)	0.21
TT	18 (13.8)	14 (10.8)	1.05 (0.80–1.21)	0.12
Allele frequency in C410T				
C allele	114 (87.7)	85 (65.4)	3.77 (1.99–7.12)	<0.0001
T allele	33 (25.4)	31 (23.8)	1.08 (0.61–1.91)	0.88
NF- κ B1 gene				
Ins/ins	84 (64.6)	82 (63.1)	1.06 (0.64–1.77)	0.79
Del/ins	69 (53.1)	33 (25.4)	3.32 (1.96–5.61)	<0.001
Del/del	15 (11.5)	11 (8.5)	1.41 (0.62–3.20)	0.68
Allele frequency in NF- κ B1				
Ins	81 (62.3)	78 (60.0)	1.13 (0.64–1.93)	0.77
NF- κ B1A gene				
AA	91 (70.0)	84 (64.6)	0.71 (0.84–1.10)	0.33
AG	25 (19.2)	21 (16.2)	1.05 (0.74–1.09)	0.79
GG	13 (10.0)	21 (16.2)	1.34 (0.98–1.92)	0.27
Allele frequency in NF- κ B1A				
A allele	64 (49.2)	67 (51.5)	1.23 (0.71–1.87)	0.70
G allele	52 (40.0)	48 (36.9)	1.13 (0.66–1.83)	0.62

Notes: Values are expressed as the absolute number (%) of subjects. p-value was calculated by univariate analysis using the chi-square.

Table 3 Plasma level of cytokines in individuals with acute renal injury and in healthy individuals by different genotypes

IL-6, ^a mean \pm SD			IL-1 β , ^a mean \pm SD		TNF- α , ^a mean \pm SD	
Acute renal injury patients (n=130)	Healthy people (n=130)		Acute renal injury patients (n=130)	Healthy people (n=130)	Acute renal injury patients (n=130)	Healthy people (n=130)
G1672A						
GG	131 \pm 24.2	12.6 \pm 4.2	84.2 \pm 2.4	12.2 \pm 1.8	188.1 \pm 2.5	17.3 \pm 8.3
GA	129 \pm 12.1	15.4 \pm 5.3	16.9 \pm 1.3	185.1 \pm 8.3	15.1 \pm 7.2	25.1 \pm 1.4
AA	142 \pm 18.2	18.2 \pm 3.3	87.3 \pm 2.2	19.1 \pm 1.4	191.1 \pm 9.1	23.9 \pm 8.9
C410T						
CC	115 \pm 12.3	17.6 \pm 1.2	88.3 \pm 2.1	17.4 \pm 1.1	96.8 \pm 1.3	17.4 \pm 1.1
CT	134 \pm 14.4	19.1 \pm 5.3	93.5 \pm 2.2	19.2 \pm 2.4	97.3 \pm 2.1	19.2 \pm 2.4
TT	124 \pm 12.4	18.2 \pm 3.3	92.3 \pm 2.1	12.8 \pm 1.2	171.1 \pm 12.1	23.2 \pm 6.2
NF- κ B1						
Ins/ins	134 \pm 12.1	18.1 \pm 2.2	83.2 \pm 3.1	134 \pm 14.4	19.1 \pm 5.3	134 \pm 4.4
Del/ins	117 \pm 15.6	17.4 \pm 1.1	96.8 \pm 1.3	124 \pm 12.4	18.2 \pm 3.3	124 \pm 12.4
Del/del	111 \pm 14.4	19.2 \pm 2.4	97.3 \pm 2.1	19.1 \pm 1.4	191.1 \pm 9.1	23.9 \pm 8.9
NF- κ BIA						
AA	134 \pm 14.4	19.6 \pm 3.2	98.1 \pm 2.1	18.5 \pm 1.2	117 \pm 5.6	17.4 \pm 1.1
AG	134 \pm 11.1	18.4 \pm 3.2	94.5 \pm 2.7	17.1 \pm 1.8	111 \pm 4.4	19.2 \pm 2.4
GG	180 \pm 12.2	12 \pm 3.8	99.7 \pm 1.3	21.1 \pm 1.1	195.1 \pm 5.1	22.5 \pm 9.2

Note: ^ap<0.05 for between-group comparisons (acute renal injury patients vs healthy controls) by unpaired t-test.

Abbreviations: IL, interleukin; TNF- α , tumor necrosis factor alpha.

Chinese diabetic patients. We found that individuals with the GG genotype were at very high risk of developing acute renal injury. Our finding in Chinese patients was consistent with the findings of Hur et al⁵ in Korean patients in terms of involvement of polymorphisms of PARP-1 in development of kidney-related diseases. Hur et al have identified two SNPs (1963AG and β 28077GA) of PARP-1 in development of nephritis among Korean patients. Our study identified the involvement of two new SNPs of PARP-1 (C410T and G1672A) in development of acute renal injury in Chinese diabetic patients. In this Chinese study, we also observed that individuals carrying C and G alleles of PARP-1 were at high risk of developing acute renal injury. Additionally, we observed the involvement of a del/ins polymorphism in NF- κ B1 in acute renal injury development among Chinese individuals.

In our study, the individuals with a GG genotype and G allele were at very high risk of developing acute renal injury. The possible reason for involvement of the GG genotype as a risk factor for acute renal injury could be explained by molecular heterosis, which is observed in ~50% of cases of gene associations.¹⁴ We also observed the involvement of an NF- κ B1 polymorphism in developing acute renal injury.

It has been reported that cytokines that are released by renal tubular cells into the injured kidney play an important role in the initiation and extension of inflammation in acute kidney injury.^{15,16} Among the cytokines, interleukin IL-6, IL-1 β and TNF- α play an important role in development

of kidney injury.^{15,16} Studies have shown a close correlation between IL-6 expression and acute kidney injury.¹⁷ Also, IL-6 contributes to the progression of renal disease and associated complications.¹⁸ IL-1 β is one of the cytokines most commonly present in development of kidney diseases.^{15,16} Also, IL-1 β is associated with the development and progression of acute kidney injury.¹⁹ TNF- α is a potent proinflammatory cytokine and an important mediator of inflammatory tissue damage,¹⁶ and has been implicated in the pathobiology of acute kidney injury.²⁰ A role for TNF- α in mediating the inflammatory injury in cisplatin-induced acute renal failure has recently been established.²¹ Based on the above facts, we measured the cytokine levels in Chinese subjects with and without acute kidney injury. In our study, we noted high levels of cytokines among Chinese patients with kidney injury; our finding was consistent with the previous reports that showed the involvement of cytokines in acute kidney injury.^{15–21}

This was the first study to suggest the association of two SNPs of PARP-1 (C410T and G1672A) with acute renal injury in Chinese patients. We also identified the association of C and G alleles of PARP-1 in acute kidney injury. Since the study was designed as pilot study and conducted at single study center in China (limitation of study). Therefore, the present findings cannot be generalized to the overall Chinese population. Based on the study results, we suggest for conducting large multi-centric randomized clinical study in future to generalize our findings. We also suggest investigating ethnic difference in involvement of PARP-1

polymorphism in development of acute renal injury among Chinese diabetic patients, which was not possible in our study. Our study is the first pilot study which identified the involvement of SNPs and allele of PARP-1 in diabetic patient with acute renal injury. The involvement of identified SNPs in other kidney disorders subtypes has not yet been established. Our finding encourages the researchers to focus on these two SNPs (C410T and G1672A) in other disease conditions associated with chronic diabetes mellitus. Since the objective of our pilot study was to assess the involvement of PARP-1 alleles and SNPs in development of acute renal diseases in diabetic patients; we therefore have not measured the sirtuin and PARP activity in relation with development of diabetes and associated kidney problems. We also encourage measuring the sirtuin and PARP activity in relation with the development of diabetes and associated kidney problems in future studies.

Conclusion

Our findings showed the involvement of PARP-1 polymorphisms in the development of acute renal injury in Chinese individuals. Our study identified the involvement of two SNPs of PARP-1 (C410T and G1672A) in development of acute renal injury among Chinese diabetic patients. Also, increased expression of C and G alleles of PARP-1 can be considered as one of the potential risk factors for developing acute renal injury. We also observed the involvement of an NF- κ B1 polymorphism in the increased risk of acute renal injury. Moreover, the increased serum cytokine levels can be considered as one of the potential risk factors for developing acute renal injury. Our study results can serve as a basis for conducting large multicenter, multicountry, genetic clinical studies to assess the involvement of PARP-1 and NF- κ B1 polymorphism in acute renal injury in elderly Chinese patients with diabetes mellitus.

Acknowledgments

This study was supported by a grant from the Health and Family Planning Commission of Guangxi Zhuang Autonomous Region Self-financing Funds in Clinical Application Plan of Scientific Research Subject (NO Z2016261). The authors thank all patients for their participation in the study. They would also like to thank Dr Rakesh Ojha (senior consultant medical writer) for writing the manuscript and for the editorial support provided in the preparation of this manuscript.

Disclosure

The authors report no conflicts of interest in this work.

References

1. Dronavalli S, Duka I, Bakris GL. The pathogenesis of diabetic nephropathy. *Nat Clin Pract Endocrinol Metab.* 2008;4(8):444–452.
2. Nakai S, Iseki K, Itami N, et al. An overview of regular dialysis treatment in Japan (as of 31 December 2010). *Ther Apher Dial.* 2012;16(6):483–521.
3. Seaquist ER, Goetz FC, Rich S, Barbosa J. Familial clustering of diabetic kidney disease. Evidence for genetic susceptibility to diabetic nephropathy. *N Engl J Med.* 1989;320(18):1161–1165.
4. Freedman BI, Bostrom M, Daeihagh P, Bowden DW. Genetic factors in diabetic nephropathy. *Clin J Am Soc Nephrol.* 2007;2(6):1306–1316.
5. Hur JW, Sung YK, Shin HD, Park BL, Cheong HS, Bae SC. Poly(ADP-ribose) polymerase (PARP) polymorphisms associated with nephritis and arthritis in systemic lupus erythematosus. *Rheumatology (Oxford).* 2006;45(6):711–717.
6. Smulson ME, Simbulan-Rosenthal CM, Boulares AH, et al. Roles of poly(ADP-ribosylation) and PARP in apoptosis, DNA repair, genomic stability and functions of p53 and E2F-1. *Adv Enzyme Regul.* 2000;40(1):183–215.
7. Corcoran NM, Clarkson MJ, Stuchbery R, Hovens CM. Molecular pathways: targeting DNA repair pathway defects enriched in metastasis. *Clin Cancer Res.* 2016;22(13):3132–3137.
8. Hassa PO, Hottiger MO. The diverse biological roles of mammalian PARPs, a small but powerful family of poly-ADP-ribose polymerases. *Front Biosci.* 2008;13:3046–3082.
9. Phulwani NK, Kielian T. Poly (ADP-ribose) polymerases (PARPs) 13 regulate astrocyte activation. *J Neurochem.* 2008;106(2):578–590.
10. Ba X, Garg NJ. Signaling mechanism of poly(ADP-ribose) polymerase-1 (PARP-1) in inflammatory diseases. *Am J Pathol.* 2011;178(3):946–955.
11. Chiarugi A, Moskowitz MA. Poly(ADP-ribose) polymerase-1 activity promotes NF- κ B-driven transcription and microglial activation: implication for neurodegenerative disorders. *J Neurochem.* 2003;85(2):306–317.
12. Yu SW, Wang H, Poitras MF, et al. Mediation of poly(ADP-ribose) polymerase-1-dependent cell death by apoptosis-inducing factor. *Science.* 2002;297(5579):259–263.
13. Albulescu R, Codrici E, Popescu ID, et al. Cytokine patterns in brain tumour progression. *Mediators Inflamm.* 2013;2013:979748.
14. Comings DE, MacMurray JP. Molecular heterosis: a review. *Mol Genet Metab.* 2000;71(1–2):19–31.
15. John DI, Michael JR. Immune and inflammatory role in renal disease. *Compr Physiol.* 2013;3(2):957–976.
16. Ali A, Quocan N, Charles L. Edelstein mediators of inflammation in acute kidney injury. *Mediators of Inflamm.* 2009;2009:12.
17. Grigoryev DN, Liu M, Hassoun HT, Cheadle C, Barnes KC, Rabb H. The local and systemic inflammatory transcriptome after acute kidney injury. *J Am Soc Nephrol.* 2008;19(3):547–558.
18. Jones SA, Fraser DJ, Fielding CA, Jones GW. Interleukin-6 in renal disease and therapy. *Nephrol Dial Transplant.* 2015;30(4):564–574.
19. Tadayoshi K, Rei N, Ryo M, et al. Expression and function of interleukin-1 β -induced neutrophil gelatinase-associated lipocalin in renal tubular cells. *PLoS One.* 2016;11(11):e0166707.
20. Susantitaphong P, Perianayagam MC, Tighiouart H, Liangos O, Bonventre JV, Jaber BL. Tumor necrosis factor alpha promoter polymorphism and severity of acute kidney injury. *Nephron Clin Pract.* 2013;123(1–2):67–73.
21. Ramesh G, Reeves WB. Inflammatory cytokines in acute renal failure. *Kidney Int Suppl.* 2004;91:S56–S61.

Clinical Interventions in Aging

Dovepress

Publish your work in this journal

Clinical Interventions in Aging is an international, peer-reviewed journal focusing on evidence-based reports on the value or lack thereof of treatments intended to prevent or delay the onset of maladaptive correlates of aging in human beings. This journal is indexed on PubMed Central, MedLine,

CAS, Scopus and the Elsevier Bibliographic databases. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <http://www.dovepress.com/clinical-interventions-in-aging-journal>