Prognostic role of CD82/KAI1 in multiple human malignant neoplasms: a meta-analysis of 31 studies

Jundong Zhu*
Chenkui Miao*
Shouyong Liu*
Ye Tian
Chao Zhang
Chao Liang
Aiming Xu
Qiang Cao
Zengjun Wang

Department of Urology, The First
Affiliated Hospital of Nanjing Medical
University, Nanjing, China

*These authors contributed equally
to this work

Abstract: Tetraspanin CD82, also known as KAI1, was revealed as an attractive prognostic tumor biomarker in recent studies. However, some results of these studies remained debatable and inconclusive. Therefore, we conducted a meta-analysis to clarify the precise predictive value of CD82 in various neoplasms. Qualified studies were identified up to April 27, 2017, by searching PubMed, EMBASE, and the Web of Science. In total, 29 eligible studies were ultimately enrolled in this meta-analysis. Pooled hazard ratios (HRs) with 95% CIs of overall survival and disease/recurrence/progression-free survival were calculated to evaluate the correct prognostic role of CD82. Statistical analysis demonstrated that high expression of CD82 was significantly associated with enhanced overall survival (HR =0.56, 95% CI: 0.47–0.67) and disease/recurrence/progression-free survival (HR =0.42, 95% CI: 0.30–0.59) in cancer patients. Furthermore, we also conducted the subgroup analysis and the results revealed that CD82 was associated with favorable outcomes in cancer patients. Taken together, CD82 could be a promising biomarker for predicting the prognosis of patients with malignant neoplasms, and the biological functions of CD82 are of great research value of the subject.

Keywords: CD82, KAI1, prognosis, meta-analysis

Introduction

Tetraspanins are a family of 34 proteins, which are involved in diverse functions such as B- and T-cell activation, platelet aggregation, migration, proliferation, morphogenesis, and tumor cell progression.¹ The key feature of tetraspanins is their potential to associate with each other and with a multitude of molecules from other protein families.^{2,3} CD82, also known as KAI1, belongs to tetraspanin family associated not only with extensive physiological processes, but also in pathological situations such as cancer invasion and metastasis,^{4,5} and its differential expressions are found in various normal and malignant tissues,^{6–8} which indicate that CD82 may play a pivotal role in cancer growth, progression, motility, invasion, and metastasis.⁹

A number of studies have demonstrated that decreased CD82 expression in tumor tissues was associated with unfavorable survival in cancer patients.^{10–15} Whereas in some individual studies focused on gastric carcinoma,¹⁶ osteosarcoma,¹⁷ colorectal carcinoma,¹⁸ and clear cell renal cell carcinoma,¹⁹ increased expression of CD82 might predict diverse, even opposing outcome. The discrepancies between these studies highlight the importance of evaluating the prognostic significance of CD82 in human malignant neoplasms. Therefore, we conducted this systematic review using meta-analysis to shed light on the relationship between CD82 expression and the prognosis of patients with carcinoma.

Correspondence: Qiang Cao;
Zengjun Wang
Department of Urology, The First
Affiliated Hospital of Nanjing Medical
University, 300 Guangzhou Road,
Nanjing, Jiangsu 210029, China
Tel/fax +86 25 8378 0079
Email qiang_cao@126.com;
zengjunwang@njmu.edu.cn

Methods

Search strategy

We searched online databases, including PubMed, EMBASE, and Web of Science, to identify relevant literature published until April 27, 2017. For the literature retrieval, combinations of the keywords were used as follows: (“cancer” or “carcinoma” or “neoplasm” or “tumor” or “tumour”) and (“KAI1” or “CD82”) and (“prognostic” or “prognosis” or “survival” or “outcome” or “recurrence” or “relapse”). The following criteria should be considered to select the published studies: 1) human, English publications; 2) a relationship of CD82 expression with cancer prognosis. In addition, we searched for studies published in Chinese to comprehensively understand the role of miRNA-205 in cancer. In order to supplement our literature search, the bibliographies in these studies were also carefully scanned.

Quality assessment

To evaluate the retrieved studies, they should include clear definitions of the following: 1) the study population and country; 2) the study design; 3) assay method to determine CD82 expression: immunohistochemistry (IHC), quantitative reverse transcription-polymerase chain reaction (qRT-PCR), or Western-blot; 4) the prognosis or survival assessment; 5) the detected tumor and pathology information; 6) the cutoff point of CD82; and 7) the follow-up duration (Table 1). Sensitivity analyses and published bias were performed to promote the quality of this meta-analysis. A flow diagram of the study selection process is presented in Figure 1.

Data extraction

All data from eligible studies were extracted independently; ambiguous data were reviewed in detail. Parameters of these literatures were extracted from each single paper, including the first author's surname, publication year, patients' median or mean age, nationality, dominant ethnicity, number of patients, investigating method, cutoff value, follow-up time, and hazard ratios (HRs) for prognostic outcomes (overall survival [OS] and disease/recurrence/progression-free survival [DFS/RFS/PFS]) along with their 95% CI and *p*-values. If only Kaplan–Meier curves were available, data were extracted from graphical survival curves to extrapolate HRs with 95% CIs using previously described methods.^{20–22} All of the aforementioned data are comprehensively detailed in Tables 1 and 2.

Statistical analysis

We quantified the effect of heterogeneity via $I^2=100\% \times (Q - df)/Q$. A random-effects model (DerSimonian–Laird

method) was applied if $p < 0.10$ or $I^2 > 50\%$; otherwise, a fixed-effects model (Mantel–Haenszel method) was used instead.²³ In addition, we classified the enrolled studies into subgroups to reduce the influence of heterogeneity. The publication bias was evaluated by the Begg's funnel plot and Egger linear regression test with a funnel plot.²⁴ Sensitivity analysis was also tested. All *p*-values were calculated using a two-sided test, and $p < 0.05$ was considered statistically significant. Statistical analyses were performed via the Stata 12.0 (StataCorp, College Station, TX, USA) and Microsoft Excel (V.2007, Microsoft Corporation, Redmond, WA, USA).

Results

Summary of enrolled studies

A total of 157 studies were retrieved from PubMed, EMBASE, and Web of Science. After initial scanning of titles and abstracts, 93 studies were excluded because they were review articles/letters or non-English publications, were not associated with CD82 or prognosis/survival. The remaining 64 records were downloaded as full text and accessed very carefully. Among them, 35 potentially suitable studies were excluded because they lacked sufficient survival data (HRs and 95% CIs), did not report comprehensive data, or extracted their survival data from an existing database. Ultimately, 29 studies were considered eligible for our meta-analysis. The inclusion and exclusion reasons for candidate studies are presented in detail in Figure 1.

The main features of the 29 enrolled studies are systematically summarized in Tables 1 and 2. Twenty-two studies reported patient OS, one focused on RFS, and six investigated OS as well as DFS or PFS. Eight of the studies focused on Caucasian populations, which mainly came from European countries, and 21 focused on Asian populations, of which 10 were from China, 9 from Japan, 1 from India, and 1 from Korea. As for cancer type, malignant neoplasms assessed in these studies included colorectal carcinoma, gastric carcinoma, breast cancer, laryngeal squamous cell carcinoma (LSCC), esophageal squamous cell carcinoma (ESCC), and oral squamous cell carcinoma (OSCC). Among the 29 studies, the pathological types of adenocarcinoma (AdenoCa), squamous carcinoma (SqCa), transitional cell carcinoma (TCC), sarcoma, and melanoma were covered. All of these studies were retrospective in design and determined CD82 expression using tissue samples. IHC and qRT-PCR were used in the majority of all eligible studies to detect CD82 expression, and Western-blot analysis was conducted in one study. When we analyzed the HR and 95% CI in each study, we extracted two sets of data from the same article

Table 1 Main characteristics of studies included in the meta-analysis

First author, publication year	Case nationality	Dominant ethnicity	Median or mean age	Study design	Malignant disease	Main type of pathology	Detected sample	Assay method	Survival analysis	Source of HR	Maximum months of follow-up
Zhu, 2017 ¹⁰	China	Asian	59.4	R	Colorectal carcinoma	AdenoCa	Tissue	IHC	OS	Reported	96
Lu, 2016 ¹²	China	Asian	57.7	R	Gastric carcinoma	AdenoCa	Tissue	IHC	OS	Reported	95
Singh, 2016 ¹¹	India	Asian	49	R	Breast cancer	AdenoCa	Tissue	qRT-PCR	OS	SC	36
Guo, 2015 ^{16a}	China	Asian	48	R	Gastric carcinoma	AdenoCa	Tissue	IHC	OS	SC	60
Guo, 2015 ^{16b}	China	Asian	48	R	Gastric carcinoma	AdenoCa	Tissue	qRT-PCR	OS	SC	60
Wu, 2015 ³²	China	Asian	62.1	R	Colorectal carcinoma	AdenoCa	Tissue	IHC	OS	Reported	108
Han, 2015 ³⁶	China	Asian	45.6	R	Breast cancer	AdenoCa	Tissue	IHC	OS	Reported	NM
Yu, 2014 ¹³	China	Asian	62.1	R	LSCC	SqCa	Tissue	IHC	OS/DFS	Reported	NM
Kwon, 2014 ¹⁹	Korea	Asian	55.8	R	CCRCC	AdenoCa	Tissue	IHC	OS	Reported	223
Tang, 2014 ^{14c}	Canada	Caucasian	60	R	Melanoma	Melanoma	Tissue	IHC	OS	Reported	120
Tang, 2014 ^{14d}	Canada	Caucasian	60	R	Melanoma	Melanoma	Tissue	IHC	OS	Reported	60
Zhang, 2013 ³⁷	China	Asian	58.7	R	LSCC	SqCa	Tissue	IHC	OS	SC	NM
Zhang, 2013 ³⁸	China	Asian	59.6	R	LSCC	SqCa	Tissue	WB	OS	SC	84
Wu, 2012 ¹⁵	China	Asian	58.6	R	NSCLC	AdenoCa	Tissue	IHC	OS	Reported	88
Knoener, 2012 ²³	Germany	Caucasian	NM	R	Gastric carcinoma	AdenoCa	Tissue	IHC	OS	SC	163
Guo, 2009 ³⁹	China	Asian	60	R	Hepatocellular carcinoma	AdenoCa	Tissue	IHC	OS	SC	24
Protzel, 2008 ¹⁴	Germany	Caucasian	66.5	R	Penile carcinoma	SqCa	Tissue	IHC	OS	SC	125
Miyazaki, 2005 ⁴⁰	Japan	Asian	61	R	ESCC	SqCa	Tissue	IHC	OS	SC	60
Leavey, 2005 ¹⁷	America	Caucasian	11	R	Osteosarcoma	Sarcoma	Tissue	IHC	OS/PFS	SC	NM
Farhadi, 2004 ⁴¹	Australia	Caucasian	65	R	OSCC	SqCa	Tissue	IHC	OS/DFS	Reported	258
Goncharuk, 2004 ⁴²	America	Caucasian	65	R	NSCLC	AdenoCa	Tissue	IHC	OS	SC	NM
Su, 2004 ⁵⁰	Japan	Asian	64	R	Bladder cancer	TCC	Tissue	IHC	RFS	Reported	78
Hashida, 2003 ¹⁸	Japan	Asian	62.8	R	Colorectal carcinoma	AdenoCa	Tissue	IHC	OS/DFS	Reported	85.9
Imai, 2002 ⁴³	Japan	Asian	62.6	R	OSCC	SqCa	Tissue	qRT-PCR	OS	SC	60
Schindl, 2001 ⁴⁴	Austria	Caucasian	56.8	R	Epithelial ovarian cancer	AdenoCa	Tissue	IHC	OS/DFS	Reported	130
Miyazaki, 2000 ⁴⁵	Japan	Asian	61.8	R	ESCC	SqCa	Tissue	IHC	OS	SC	195.2
Yang, 2000 ⁴⁶	America	Caucasian	NM	R	Breast cancer	AdenoCa	Tissue	IHC	OS	SC	NM
Sho, 1998 ⁴⁷	Japan	Asian	66	R	Pancreatic cancer	AdenoCa	Tissue	IHC	OS	SC	62
Huang, 1998 ⁴⁸	Japan	Asian	50	R	Breast cancer	AdenoCa	Tissue	IHC	OS/DFS	Reported	NM
Higashiyama, 1997 ³⁵	Japan	Asian	63.7	R	NSCLC	AdenoCa	Tissue	IHC	OS	Reported	61.3
Adachi, 1996 ⁴⁹	Japan	Asian	60	R	NSCLC	AdenoCa	Tissue	qRT-PCR	OS	Reported	58

Notes: ^aData extracted from one study due to different assay methods (IHC and qRT-PCR). ^bData extracted from one study due to different follow-up times (120 and 60 months). Study design is described as retrospective.

Abbreviations: AdenoCa, adenocarcinoma; CCRCC, clear cell renal cell carcinoma; DFS, disease-free survival; ESCC, esophageal squamous cell carcinoma; HR, hazard ratio; IHC, immunohistochemistry; LSCC, laryngeal squamous cell carcinoma; NM, not mentioned; NSCLC, non-small-cell lung cancer; OS, overall survival; OSCC, oral squamous cell carcinoma; PFS, progression-free survival; qRT-PCR, quantitative reverse transcription-polymerase chain reaction; R, retrospective; RFS, recurrence-free survival; SC, survival curve; SqCa, squamous carcinoma; TCC, transitional cell carcinoma; WB, Western-blot.

Figure 1 Flow diagram of the study selection process.

Abbreviation: HRs, hazard ratios.

due to different assay methods or different follow-up times in two studies (Guo et al¹⁶ and Tang et al¹⁴). The source of HR and 95% CI was extracted from survival curves or article reports.

OS associated with CD82 expression

A total of 28 original studies were included to analyze the OS, with a random-effects model on account of the moderate heterogeneity ($p=0.000$, $I^2=54.5\%$). The results indicated that CD82-positive expression was significantly associated with favorable OS in cancer patients (pooled HR = 0.56, 95% CI: 0.47–0.67, $p<0.05$; Figure 2A). Subgroup study was then performed; increased CD82 expression was significantly associated with enhanced OS in the Asian patients (pooled HR = 0.55, 95% CI: 0.43–0.70) as well in Caucasian (pooled HR = 0.57, 95% CI: 0.47–0.70; Figure 3A). In tumor subgroup analysis, we found high expression of CD82 correlated with longer OS in colorectal carcinoma, gastric carcinoma, breast cancer, LSCC, and non-small-cell lung cancer (NSCLC) (Figure 3B). Due to insufficient studies, correlation between CD82 and OS in other tumor types have not been further analyzed. Stratification analyses for other subgroups are presented in detail in Figure 3C and D.

DFS/RFS/PFS associated with CD82 expression

Seven of the studies analyzed DFS/RFS/PFS. The heterogeneity between these studies was low ($p=0.153$, $I^2=36.0\%$);

thus, a fixed-effects model was applied to calculate a pooled HR of 0.42 (95% CI: 0.30–0.59). This result demonstrated that CD82 overexpression predicted low risk of cancer progression (Figure 2B). Ethnic and pathological subgroup analysis for DFS/RFS/PFS also demonstrated the protective effect of CD82 and that it may play a key role of progression in cancer patients (Figure 4).

Sensitivity analyses

In order to determine the robustness of the above results and evaluate the stability of results, a sensitivity analysis was performed by Stata12.0 software. Individual data that could affect the final conclusions have been deleted in advance (see footnote “f” in Table 2). The analyzed result from a fixed model suggested that our results are comparatively credible and stable (Figure 5A and B).

Publication bias

Begg’s funnel and the Egger’s test were used to evaluate the possible publication bias in this meta-analysis. The funnel plots of the publication bias are presented in Figure 5C and D. p -values calculated from Egger’s test with higher detection effectiveness were 0.135 for OS, 0.610 for DFS/RFS/PFS, respectively, indicating no significant publication bias in the meta-analysis.

Discussion

In recent years, elaborate efforts have been invested to detect promising biomarkers for patients with multiple carcinomas.

Table 2 HRs and 95% CIs of patient survival or cancer progression relating to CD82 expression in eligible studies

First author, publication year	Assay method	Cutoff point	Case number		OS	DFS/RS/PFS		p-value
			High expression	Low expression		HR (95% CI) (U/M)	HR (95% CI) (U/M)	
Zhu, 2017 ¹⁰	IHC	IRS scores ≥ 3 (range of 0–12)	65	139		0.302 (0.184–0.497) ^M	NM	<0.001
Lu, 2016 ¹²	IHC	IRS scores ≥ 3 (range of 0–12)	134	191		0.648 (0.492–0.854) ^M	NM	0.002
Singh, 2016 ¹¹	qRT-PCR	NM	46	29		0.66 (0.26–1.68) ^U	NM	0.047
Guo, 2015 ^{6a}	IHC	> 10% of tumor cells stained	28	100		1.07 (0.22–5.24) ^U	NM	<0.05
Guo, 2015 ^{6b}	qRT-PCR	NM	40	88		0.62 (0.19–1.98) ^U	NM	<0.05
Wu, 2015 ³²	IHC	IRS scores ≥ 3 (range of 0–12)	56	118		0.430 (0.269–0.687) ^M	NM	<0.001
Han, 2015 ⁶	IHC	IRS scores ≥ 3 (range of 0–12)	137	188		0.617 (0.462–0.823) ^M	NM	0.001
Yu, 2014 ¹³	IHC	IRS scores ≥ 3 (range of 0–12)	34	49		0.226 (0.094–0.545) ^M	0.278 (0.119–0.647) ^M	0.003
Kwon, 2014 ¹⁹	IHC	>0% of tumor cells stained	98	546		1.513 (1.057–2.165) ^M	^f	0.024
Tang, 2014 ^{14c}	IHC	IRS scores ≥ 8 (range of 0–12)	114	303		0.42 (0.29–0.63) ^M	NM	0.000147
Tang, 2014 ^{14d}	IHC	IRS scores ≥ 8 (range of 0–12)	114	303		0.59 (0.36–0.96) ^M	NM	0.029
Zhang, 2013 ³⁷	IHC	Accumulated points ≥ 80	52	48		0.54 (0.27–1.05) ^U	NM	NM
Zhang, 2013 ³⁸	WB	NM	43	43		0.36 (0.17–0.77) ^U	NM	NM
Wu, 2012 ¹⁵	IHC	IRS scores > 1 (range of 0–12)	19	31		0.039 (0.007–0.236) ^M	NM	0.00
Knoener, 2012 ³³	IHC	IRS scores ≥ 3 (range of 0–12)	168	103		0.77 (0.55–1.07) ^U	NM	0.2305
Guo, 2009 ⁹	IHC	> 10% of tumor cells stained	21	59		0.81 (0.28–2.32) ^U	NM	0.022
Protzel, 2008 ³⁴	IHC	>50% of tumor cells stained	17	13		0.34 (0.02–4.90) ^U	NM	0.0042
Miyazaki, 2005 ⁴⁰	IHC	Ki index ≥ 39	48	43		0.48 (0.22–1.03) ^U	NM	0.0023
Leavey, 2005 ¹⁷	IHC	NM	16	31		1.14 (0.27–4.81) ^U	0.99 (0.35–2.85) ^U	0.28
Farhadi, 2004 ⁴¹	IHC	> 10% of tumor cells stained	15	42		0.52 (0.27–1.01) ^M	0.4 (0.2–0.8) ^M	0.053
Goncharuk, 2004 ⁴²	IHC	Median	77	27		0.47 (0.23–0.97) ^U	NM	0.034
Su, 2004 ⁴⁰	IHC	> 50% of tumor cells stained	54	33		NM	0.255 (0.11–0.588) ^M	NM
Hashida, 2003 ¹⁸	IHC	Scores ≥ 120 (range of 0–300)	63	83		0.903 (0.297–2.747) ^M	1.054 (0.431–2.577) ^M	0.0013
Imai, 2002 ⁴³	qRT-PCR	NM	25	18		0.51 (0.06–4.38) ^U	NM	0.858
Schindl, 2001 ⁴⁴	IHC	Scores ≥ 4 (range of 3–7)	48	59		0.45 (0.22–0.92) ^M	0.3 (0.09–0.93) ^M	NM
Miyazaki, 2000 ⁴⁵	IHC	> 10% of tumor cells stained	36	19		0.41 (0.17–1.00) ^U	0.3 (0.09–0.93) ^M	0.0282
Yang, 2000 ⁴⁶	IHC	>5% of tumor cells stained	36	36		0.72 (0.20–2.61) ^U	NM	0.024
Sho, 1998 ⁴⁷	IHC	>50% of tumor cells stained	15	25		1.16 (0.29–4.64) ^U	NM	NM
Huang, 1998 ⁴⁸	IHC	HSCORE ≥ 50	44	65		0.442 (0.109–1.789) ^M	NM	0.018
Higashiyama, 1997 ³⁵	IHC	>50% of tumor cells stained	65	135		0.700 (0.502–0.976) ^M	0.360 (0.149–0.870) ^M	0.2528
Adachi, 1996 ⁴⁹	qRT-PCR	Gene conservation rate value > 1.2	35	116		0.416 (0.175–0.986) ^M	NM	0.037

Notes: The source of HR and 95% CI was extracted from survival curves or article reports. ^aData extracted from one study due to its heterogeneity and publication bias. ^bData extracted from one study due to different assay method (IHC and qRT-PCR). ^cData extracted from one study due to different follow-up time (120 and 60 months). ^dHR calculated from survival curves. ^eExcluded due to its heterogeneity and publication bias.

Abbreviations: IHC, immunohistochemistry; qRT-PCR, quantitative reverse transcription-polymerase chain reaction; WB, western-blot; NM, not mentioned; IRS, immunoreactivity score; HSCORE, the intensity and respective percentage cells that stain at each intensity were multiplied to reach a HSCORE that ranged from 0 to 300; OS, overall survival; DFS, disease-free survival; RFS, recurrence-free survival; PFS, progression-free survival; SC, survival curve; HR, hazard ratio; CI, confidence interval; U, univariate analysis; M, multivariate analysis.

Figure 2 Forest plots of combined analyses associated with CD82 expression.

Notes: (A) OS and (B) DFS/RFS/PFS. Weights are from random effects analysis.

Abbreviations: DFS, disease-free survival; HR, hazard ratio; ID, identifier; OS, overall survival; PFS, progression-free survival; RFS, recurrence-free survival.

Figure 3 (Continued)

Figure 3 Forest plots of stratified analysis of the OS.

Notes: (A) Stratified by ethnic subgroup, (B) stratified by tumor subgroup, (C) stratified by pathological subgroup, and (D) stratified by assay method subgroup. Weights are from random effects analysis.

Abbreviations: AdenoCa, adenocarcinoma; HR, hazard ratio; ID, identifier; IHC, immunohistochemistry; LSCC, laryngeal squamous cell carcinoma; NSCLC, non-small-cell lung cancer; OS, overall survival; qRT-PCR, quantitative reverse transcription-polymerase chain reaction; SqCa, squamous carcinoma.

Figure 4 Forest plots of stratified analysis of the DFS/RFS/PFS.

Notes: (A) Stratified by ethnic subgroup and (B) stratified by pathological subgroup. Weights are from random effects analysis.

Abbreviations: AdenoCa, adenocarcinoma; DFS, disease-free survival; HR, hazard ratio; ID, identifier; OS, overall survival; PFS, progression-free survival; RFS, recurrence-free survival; SqCa, squamous carcinoma.

Tetraspanins are a family of integral membrane proteins with four transmembrane helices, a small extracellular loop, and a large extracellular loop. Recent studies have revealed the importance of tetraspanins in solid tumors and hematologic

malignancies, and the expression of tetraspanins is associated with the tumor biologic characteristics.⁴ Some members of this family are known as metastasis suppressor genes, while others are supposed to promote tumor progression.¹

Figure 5 Sensitivity analysis under specific model and Begg's funnel plots of publication bias test.

Notes: (A) Effect of individual studies on the combined HR for OS, (B) effect of individual studies on the combined HR for DFS/RFS/PFS, (C) Begg's funnel plots of OS, and (D) Begg's funnel plots of DFS/RFS/PFS.

Abbreviations: DFS, disease-free survival; HR, hazard ratio; OS, overall survival; PFS, progression-free survival; RFS, recurrence-free survival.

CD82 is considered to be important in tetraspanin family given its differential expression between cancer and normal tissues. CD82 is downregulated in many types of cancers and loss of CD82, both protein and mRNA, is strongly correlated with poor prognosis in many malignancies.^{9,25} Current understanding of CD82 function indicates that it is likely to be involved in detachment, motility/invasion, and cell survival, which are associated with various adhesion receptors (eg, integrins), receptor tyrosine kinases (eg, epithelial growth factor receptor [EGFR] and c-Met), and other signaling pathway molecules.^{4,9,26} CD82 can interact with other tetraspanin proteins (eg, CD151 and CD81), integrins (eg, $\alpha_3\beta_1$, $\alpha_4\beta_1$, and $\alpha_5\beta_1$), and chemokines to regulate the migration, adhesion, and signaling of cells.^{9,27,28} Integrins are not the only molecules that CD82 regulate, and studies have revealed that tetraspanins play a critical role in regulating receptor tyrosine kinase signaling in immune cells.^{29,30} Odintsova et al reported that CD82 was a regulator of epithelial growth

factor (EGF)-induced signaling and showed that the association of EGFR with the tetraspanin is critical in EGFR desensitization.³¹ The diverse biological activities and molecular mechanisms of CD82 may partially contribute to the tumor prognosis in cancer patients.

In this meta-analysis, we first evaluated the correlation of CD82 with prognostic outcomes (OS, DFS/RFS/PFS) of patients with various cancers systemically. We also performed subgroup, sensitivity, and heterogeneity analyses to explore the effects of dominant characteristics from available studies. Many studies have reported that decreased expression of CD82 often predicts unfavorable outcome in cancers and the level of CD82 was negatively correlated with invasion of depth, vessel invasion, lymph node metastasis, distant metastasis, and TNM stages.^{10,16,32–34} However, there are also some individual studies that came to a diverse, even opposing, conclusion in our included literature. Our prognosis analysis revealed the pooled HR of 0.56 (OS)

and 0.42 (DFS/RFS/PFS), demonstrating that increased CD82 expression could be a favorable prognostic factor of various tumors. Similarly, in subgroup analysis based on the characteristics of the individual studies, we observed the statistically significant outcomes when data were stratified according to ethnicity, pathology, assay method, and tumor types. Although AdenoCa and squamous cell carcinoma have oncologically different characteristics, according to our study, high expression of CD82 correlated with longer OS or DFS/RFS/PFS in both types. Due to few relevant studies and small sample size, results of some tumors and pathologic types were not presented in the forest plot when we conducted the subgroup analysis. This means the conclusion should be considered with caution when we come across these neoplasms, including hepatocellular carcinoma, clear cell renal cell carcinoma, melanoma, osteosarcoma, and pancreatic cancer. Therefore, more existing investigations toward these tumors are needed for further evidence.

After extracting the data from studies, sensitivity analysis was conducted to check which individual data could affect the final conclusions. When we analyzed the stability of results, we found that if the data toward DFS/RFS/PFS from Higashiyama et al³⁵ and Kwon et al¹⁹ were included in our review, they might have significant impact on the pooled significance. Therefore, we deleted these two sets of data in advance to ensure the stability of our analysis. The final results were assured that exclusion of any individual study alters little change of the pooled significance. We attributed this to the different ethnicity, tumor type, and pathological type, as well as the different source of HR. No obvious publication bias was detected in this meta-analysis, indicating our analysis was stable.

Despite the meta-analysis was performed with rigorous statistics, our conclusion still has several limitations for the following reasons. First, heterogeneity existed in the OS analyses and it was likely due to the different characteristics of the patients, such as age, ethnicity, tumor type, and pathological type, as well as the different source of HR. Second, most studies established their own varied expression cutoff, and a standard cutoff value was hard to define so that the pooled outcome may be different with the actual value. This may cause a bias in the results of the effectiveness of CD82 as a prognostic factor. Third, no independent investigation on Negroid was included in this meta-analysis, which might undermine the comprehensiveness to some extent. Fourth, the validity of results might be impaired due to the lack of prospective studies. Taking these limitations into consideration, our results should be interpreted rigorously, and more well-designed studies are needed to verify the function of CD82 in various carcinomas.

Conclusion

In summary, the significant relationship between high CD82 expression and favorable prognostic in various neoplasms was clearly revealed in this meta-analysis. The results indicated that CD82 could be a promising biomarker for predicting the prognosis of patients with malignant neoplasms, and the biological functions of CD82 are of great research value of the subject.

Acknowledgments

This study was funded by National Natural Science Foundation of China (grant numbers 81270685 and 81402321), and Project of Nanjing Science and Technology Committee (201605001).

Disclosure

The authors report no conflicts of interest in this work.

References

1. Zoller M. Gastrointestinal tumors: metastasis and tetraspanins. *Z Gastroenterol*. 2006;44(7):573–586.
2. Yunta M, Lazo PA. Tetraspanin proteins as organisers of membrane microdomains and signalling complexes. *Cell Signal*. 2003;15(6):559–564.
3. Hemler ME. Tetraspanin functions and associated microdomains. *Nat Rev Mol Cell Biol*. 2005;6(10):801–811.
4. Yang YG, Sari IN, Zia MF, Lee SR, Song SJ, Kwon HY. Tetraspanins: spanning from solid tumors to hematologic malignancies. *Exp Hematol*. 2016;44(5):322–328.
5. Richardson MM, Jennings LK, Zhang XA. Tetraspanins and tumor progression. *Clin Exp Metastasis*. 2011;28(3):261–270.
6. You J, Madigan MC, Rowe A, Sajinovic M, Russell PJ, Jackson P. An inverse relationship between KAI1 expression, invasive ability, and MMP-2 expression and activity in bladder cancer cell lines. *Urol Oncol*. 2012;30(4):502–508.
7. Liu W, Iizumi-Gairani M, Okuda H, et al. KAI1 gene is engaged in NDRG1 gene-mediated metastasis suppression through the ATF3-NFkappaB complex in human prostate cancer. *J Biol Chem*. 2011;286(21):18949–18959.
8. Christgen M, Christgen H, Heil C, et al. Expression of KAI1/CD82 in distant metastases from estrogen receptor-negative breast cancer. *Cancer Sci*. 2009;100(9):1767–1771.
9. Miranti CK. Controlling cell surface dynamics and signaling: how CD82/KAI1 suppresses metastasis. *Cell Signal*. 2009;21(2):196–211.
10. Zhu B, Zhou L, Yu L, et al. Evaluation of the correlation of vasculogenic mimicry, ALDH1, KAI1 and microvessel density in the prediction of metastasis and prognosis in colorectal carcinoma. *BMC Surg*. 2017;17(1):47.
11. Singh R, Bhatt ML, Singh SP, et al. Expression levels of Tetraspanin KAI1/CD82 in breast cancers in North Indian females. *Asian Pac J Cancer Prev*. 2016;17(7):3431–3436.
12. Lu G, Zhou L, Zhang X, et al. The expression of metastasis-associated in colon cancer-1 and KAI1 in gastric adenocarcinoma and their clinical significance. *World J Surg Oncol*. 2016;14(1):276.
13. Yu L, Zhou L, Wu S, et al. Clinicopathological significance of cancer stem cells marked by CD133 and KAI1/CD82 expression in laryngeal squamous cell carcinoma. *World J Surg Oncol*. 2014;12:118.
14. Tang Y, Cheng Y, Martinka M, Ong CJ, Li G. Prognostic significance of KAI1/CD82 in human melanoma and its role in cell migration and invasion through the regulation of ING4. *Carcinogenesis*. 2014;35(1):86–95.

15. Shiwu WU, Lan Y, Wenqing S, Lei Z, Yisheng T. Expression and clinical significance of CD82/KAI1 and E-cadherin in non-small cell lung cancer. *Arch Iran Med*. 2012;15(11):707–712.
16. Guo J, Fan KX, Xie LI, et al. Effect and prognostic significance of the KAI1 gene in human gastric carcinoma. *Oncol Lett*. 2015;10(4):2035–2042.
17. Leavey PJ, Timmons C, Frawley W, Lombardi D, Ashfaq R. KAI-1 expression in pediatric high-grade osteosarcoma. *Pediatr Dev Pathol*. 2006;9(3):219–224.
18. Hashida H, Takabayashi A, Tokuhara T, et al. Clinical significance of transmembrane 4 superfamily in colon cancer. *Br J Cancer*. 2003;89(1):158–167.
19. Kwon HJ, Min SY, Park MJ, et al. Expression of CD9 and CD82 in clear cell renal cell carcinoma and its clinical significance. *Pathol Res Pract*. 2014;210(5):285–290.
20. Williamson PR, Smith CT, Hutton JL, Marson AG. Aggregate data meta-analysis with time-to-event outcomes. *Stat Med*. 2002;21(22):3337–3351.
21. DerSimonian R, Laird N. Meta-analysis in clinical trials revisited. *Contemp Clin Trials*. 2015;45(Pt A):139–145.
22. Tierney JF, Stewart LA, Ghersi D, Burdett S, Sydes MR. Practical methods for incorporating summary time-to-event data into meta-analysis. *Trials*. 2007;8:16.
23. DerSimonian R, Laird N. Meta-analysis in clinical trials. *Control Clin Trials*. 1986;7(3):177–188.
24. Egger M, Davey SG, Schneider M, Minder C. Bias in meta-analysis detected by a simple, graphical test. *BMJ*. 1997;315(7109):629–634.
25. Feng J, Huang C, Wren JD, et al. Tetraspanin CD82: a suppressor of solid tumors and a modulator of membrane heterogeneity. *Cancer Metastasis Rev*. 2015;34(4):619–633.
26. Romanska HM, Berditchevski F. Tetraspanins in human epithelial malignancies. *J Pathol*. 2011;223(1):4–14.
27. Malik FA, Sanders AJ, Jiang WG. KAI-1/CD82, the molecule and clinical implication in cancer and cancer metastasis. *Histol Histopathol*. 2009;24(4):519–530.
28. Lee J, Byun HJ, Lee MS, et al. The metastasis suppressor CD82/KAI1 inhibits fibronectin adhesion-induced epithelial-to-mesenchymal transition in prostate cancer cells by repressing the associated integrin signaling. *Oncotarget*. 2017;8(1):1641–1654.
29. Delaguillaumie A, Harriague J, Kohanna S, et al. Tetraspanin CD82 controls the association of cholesterol-dependent microdomains with the actin cytoskeleton in T lymphocytes: relevance to co-stimulation. *J Cell Sci*. 2004;117(Pt 22):5269–5282.
30. Odintsova E, Voortman J, Gilbert E, Berditchevski F. Tetraspanin CD82 regulates compartmentalisation and ligand-induced dimerization of EGFR. *J Cell Sci*. 2003;116(Pt 22):4557–4566.
31. Odintsova E, Sugiura T, Berditchevski F. Attenuation of EGF receptor signaling by a metastasis suppressor, the tetraspanin CD82/KAI-1. *Curr Biol*. 2000;10(16):1009–1012.
32. Wu Q, Yang Y, Wu S, et al. Evaluation of the correlation of KAI1/CD82, CD44, MMP7 and beta-catenin in the prediction of prognosis and metastasis in colorectal carcinoma. *Diagn Pathol*. 2015;10:176.
33. Pal'Tseva EM, Samofalova O, Gorbacheva I, Tsar'Kov PV. [Expression of some biomarkers in primary colon adenocarcinomas and their lymph node metastases]. *Arkh Patol*. 2012;74(4):12–18.
34. Protzel C, Kakies C, Kleist B, Poetsch M, Giebel J. Down-regulation of the metastasis suppressor protein KAI1/CD82 correlates with occurrence of metastasis, prognosis and presence of HPV DNA in human penile squamous cell carcinoma. *Virchows Arch*. 2008;452(4):369–375.
35. Higashiyama M, Kodama K, Yokouchi H, et al. KAI1/CD82 expression in nonsmall cell lung carcinoma is a novel, favorable prognostic factor: an immunohistochemical analysis. *Cancer-Am Cancer Soc*. 1998;83(3):466–474.
36. Han Z, Chen Z, Zheng R, Cheng Z, Gong X, Wang D. Clinicopathological significance of CD133 and CD44 expression in infiltrating ductal carcinoma and their relationship to angiogenesis. *World J Surg Oncol*. 2015;13:56.
37. Zhang BH, Liu W, Li L, et al. KAI1/CD82 and MRP1/CD9 serve as markers of infiltration, metastasis, and prognosis in laryngeal squamous cell carcinomas. *Asian Pac J Cancer Prev*. 2013;14(6):3521–3526.
38. Zhang B, Liu W, Li L, et al. KAI1/CD82 and cyclin D1 as biomarkers of invasion, metastasis and prognosis of laryngeal squamous cell carcinoma. *Int J Clin Exp Pathol*. 2013;6(6):1060–1067.
39. Guo C, Liu QG, Zhang L, Song T, Yang X. Expression and clinical significance of p53, JunB and KAI1/CD82 in human hepatocellular carcinoma. *Hepatobiliary Pancreat Dis Int*. 2009;8(4):389–396.
40. Miyazaki T, Kato H, Kimura H, et al. Evaluation of tumor malignancy in esophageal squamous cell carcinoma using different characteristic factors. *Anticancer Res*. 2005;25(6B):4005–4011.
41. Farhadieh RD, Smee R, Ow K, et al. Down-regulation of KAI1/CD82 protein expression in oral cancer correlates with reduced disease free survival and overall patient survival. *Cancer Lett*. 2004;213(1):91–98.
42. Goncharuk VN, Del-Rosario A, Kren L, et al. Co-downregulation of PTEN, KAI-1, and nm23-H1 tumor/metastasis suppressor proteins in non-small cell lung cancer. *Ann Diagn Pathol*. 2004;8(1):6–16.
43. Imai Y, Sasaki T, Shinagawa Y, Akimoto K, Fujibayashi T. Expression of metastasis suppressor gene (KAI1/CD82) in oral squamous cell carcinoma and its clinico-pathological significance. *Oral Oncol*. 2002;38(6):557–561.
44. Schindl M, Birner P, Breitenacker G, Oberhuber G. Downregulation of KAI1 metastasis suppressor protein is associated with a dismal prognosis in epithelial ovarian cancer. *Gynecol Oncol*. 2001;83(2):244–248.
45. Miyazaki T, Kato H, Shitara Y, et al. Mutation and expression of the metastasis suppressor gene KAI1 in esophageal squamous cell carcinoma. *Cancer-Am Cancer Soc*. 2000;89(5):955–962.
46. Yang X, Wei L, Tang C, Slack R, Montgomery E, Lippman M. KAI1 protein is down-regulated during the progression of human breast cancer. *Clin Cancer Res*. 2000;6(9):3424–3429.
47. Sho M, Adachi M, Taki T, et al. Transmembrane 4 superfamily as a prognostic factor in pancreatic cancer. *Int J Cancer*. 1998;79(5):509–516.
48. Huang CI, Kohno N, Ogawa E, Adachi M, Taki T, Miyake M. Correlation of reduction in MRP-1/CD9 and KAI1/CD82 expression with recurrences in breast cancer patients. *Am J Pathol*. 1998;153(3):973–983.
49. Adachi M, Taki T, Ieki Y, Huang CL, Higashiyama M, Miyake M. Correlation of KAI1/CD82 gene expression with good prognosis in patients with non-small cell lung cancer. *Cancer Res*. 1996;56(8):1751–1755.
50. Su JS, Arima K, Hasegawa M, et al. Decreased expression of KAI1 metastasis suppressor gene is a recurrence predictor in primary pTa and pT1 urothelial bladder carcinoma. *Int J Urol*. 2004;11(2):74–82.

OncoTargets and Therapy

Publish your work in this journal

OncoTargets and Therapy is an international, peer-reviewed, open access journal focusing on the pathological basis of all cancers, potential targets for therapy and treatment protocols employed to improve the management of cancer patients. The journal also focuses on the impact of management programs and new therapeutic agents and protocols on

Submit your manuscript here: <http://www.dovepress.com/oncotargets-and-therapy-journal>

patient perspectives such as quality of life, adherence and satisfaction. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Dovepress