

Perirenal fat stranding is not a powerful diagnostic tool for acute pyelonephritis [Corrigendum]

Fukami H, Takeuchi Y, Kagaya S, et al. *Int J Gen Med.* 2017;10:137–144.

Page 137, Abstract, Results, the text “The frequency of PFS was 72% in the pyelonephritis group vs 39% in the control group” should read “The frequency of PFS was 72% in the pyelonephritis group vs 29% in the control group”.

Page 140, Table 1, Control (PRB) column, last row, the data “93 (39%)” should read “93 (29%)”.

Page 141, Discussion, line 5, the text “PFS was found in 39% of patients” should read “PFS was found in 72% of patients”.

International Journal of General Medicine

Dovepress

Publish your work in this journal

The International Journal of General Medicine is an international, peer-reviewed open-access journal that focuses on general and internal medicine, pathogenesis, epidemiology, diagnosis, monitoring and treatment protocols. The journal is characterized by the rapid reporting of reviews, original research and clinical studies across all disease areas.

The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/international-journal-of-general-medicine-journal>