Response to crizotinib in a non-small-cell lung cancer patient harboring an *EML4-ALK* fusion with an atypical *LTBP1* insertion

Cristina Aguado^{1,*}
 Maria-de-los-Llanos Gil^{2,*}
 Zaira Yeste¹
 Ana Giménez-Capitán¹
 Cristina Teixidó¹
 Niki Karachaliou²
 Santiago Viteri²
 Rafael Rosell^{1,2}
 Miguel A Molina-Vila¹

¹Laboratory of Oncology, Pangaea Oncology, ²Dr Rosell Oncology Institute, Quirón Dexeus University Hospital, Barcelona, Spain

*These authors contributed equally to this work

Abstract: Fusion of the anaplastic lymphoma receptor tyrosine kinase gene (*ALK*) with the echinoderm microtubule-associated protein 4 gene (*EML4*) is the second most common actionable alteration in non-small-cell lung cancer, with a frequency of 5%. Here, we present a case of an *EML4-ALK*-positive patient with an atypical in-frame insertion from the *LTBP1* gene in the canonical junction of *variant 1*. The patient was a 39-year-old never-smoker female diagnosed with Stage IV lung adenocarcinoma. A core biopsy was negative for *EGFR* and *KRAS* mutations but positive for *ALK* immunohistochemistry and fluorescence in situ hybridization. When submitted to nCounter, the sample showed a 3'/5' imbalance indicative of an *ALK* rearrangement, but failed to give a positive signal for any of the variants tested. Finally, a band with a molecular weight higher than expected appeared after reverse transcriptase-polymerase chain reaction analysis. When Sanger sequencing was performed, the band revealed an atypical *EML4-ALK* fusion gene with an in-frame 129 bp insertion. A 115 bp segment of the insertion corresponded to an intronic region of *LTBP1*, a gene located in the short arm of chromosome 2, between *ALK* and *EML4*. The patient received crizotinib and showed a good therapeutic response that is still ongoing after 12 months. Our result suggests that short in-frame insertions of other genes in the *EML4-ALK* junction do not affect the sensitivity of the *EML4-ALK* fusion protein to crizotinib.

Keywords: lung cancer, NSCLC, *EML4-ALK*, *LTBP1*, crizotinib, targeted therapy

Introduction

Fusions of the *ALK* gene with the *EML4* gene is the second most common actionable alteration in non-small-cell lung cancer, with a frequency of 5%.¹ More than 15 *EML4-ALK* fusion variants with various breakpoints on the *EML4* and *ALK* genes have been reported. The most frequent is *variant 1* (*v1*, 33%), followed by *v3a/3b* (29%) and *v2* (10%) but other breakpoints and 5' fusion gene partners different from *EML4* have also been described.¹⁻³ Here, we report the first case of a tumor harboring an *EML4-ALK* fusion with an atypical in-frame insertion from the *LTBP1* gene. Together with previous reports, our result suggests that in-frame insertions of other genes in the *EML4-ALK* junction might be associated with good responses to crizotinib.

Case presentation

A 39-year-old never-smoker female without prior relevant medical history was admitted to the hospital with progressive symptoms of abdominal pain, dyspnea, and bilateral

Correspondence: Miguel A Molina-Vila
 Laboratory of Oncology, Pangaea Oncology, Quirón Dexeus University Hospital, C Sabino Arana 5-19, 08023 Barcelona, Spain
 Tel +34 93 546 0140
 Fax +34 93 546 0142
 Email mamolina@panoncology.com

of mutations. Immunostaining with IHC VENTANA clone DF53 identified overexpression of ALK, and fluorescence in situ hybridization using Vysis LSI ALK dual-color break-apart probe demonstrated an *ALK* rearrangement. When the sample was analyzed by nCounter, it showed a 3'/5' imbalance indicative of an *ALK* rearrangement, but failed to give a positive signal for the EML4-ALK *v1*, *v2*, *v3*, or *v5*; TFG-ALK_T5:A20; KIF5B-ALK_K17:A20; or KIF5B-ALK_K24:A20 fusions.⁴ Finally, a band with a molecular weight higher than expected (239 bp) appeared after reverse transcriptase-polymerase chain reaction analysis (RT-PCR) with primers specific for EML4-ALK *v1*. No additional bands were apparent. The 239 bp band was submitted to Sanger sequencing revealing an atypical *EML4-ALK* fusion gene with a 129 bp insertion in the canonical junction of *v1*. A 115 bp segment of the insertion corresponded to an intronic region of *LTBP1*, a gene located in the short arm of chromosome 2, between the *ALK* and *EML4* genes. The in silico translation of this new variant, which will be referred to as *v1insLTBP1*, showed an in-frame insertion of 43 aminoacids (Figure 1).

The patient started crizotinib with good tolerance. The computed tomography scan performed a month later showed a reduction of the primary lesion, disappearance of hilary and reroperitoneal lymphadenopathies, and a reduction of the mediastinal lymph nodes. After 14 months, the patient continues to demonstrate partial response.

Written informed consent has been provided by the patient to have the case details and any accompanying images published.

Discussion

A majority of laboratories determine *ALK* rearrangements by the two US Food and Drug Administration-approved techniques, fluorescence in situ hybridization and immunohistochemistry, and do not test for specific variants due to cost-effectiveness considerations. Consequently, it is difficult to estimate the real frequency of new variants such as the *v1insLTBP1*, described in this paper. Using an nCounter methodology, we have recently reported that 6/32 (18.8%) *ALK* rearrangements in a retrospective cohort of positive cases were not *EML4-ALK v1*, *v2*, *v3*, or *v5*; TFG-ALK_T5:A20 KIF5B-ALK_K17:A20; or KIF5B-ALK_K24:A20.⁴ The exact variant of those cases could not be identified either by nCounter or RT-PCR. In addition, as a part of our routine clinical practice, we prospectively test advanced non-small-cell lung cancer patients for *ALK*

translocations by an RT-PCR technique that can identify *v1*, *v2*, and *v3*. We have found 38 positive cases, including the patient with the new variant, suggesting that the frequency of the *v1insLTBP1* could be as high as 2.7% (1/38).

The clinical relevance of the different *ALK* fusion partners and variants is poorly understood, and inconsistent results have been reported. A retrospective study including 55 *ALK*-positive patients found an association of *v1* with a longer progression-free survival (PFS) to crizotinib, while a second study reported a shorter PFS for those carrying *v3a/b*.^{3,5} Regarding rare variants, the recently described E6:A18 was intrinsically refractory to crizotinib,² while a patient with an uncommon 138 bp in-frame insertion from the *ATRNL1* gene in *v3* derived benefit from this drug.⁶ The partial response we also observed in the patient with the *v1insLTBP1* suggests that in-frame, atypical insertions do not affect the sensitivity of the EML4-ALK fusion protein to crizotinib.

Acknowledgment

The present address for Cristina Teixidó is the Department of Medical Oncology, Hospital Clínic, Barcelona, Spain.

Disclosure

Dr Santiago Viteri reports speaker honoraria from BMS and Roche, advisory board fees from Roche and Boehringer Ingelheim, and meeting inscription/travel expenses fees from Merck Serono. The authors report no other conflicts of interest in this work.

References

1. Sasaki T, Rodig SJ, Chirieac LR, Janne PA. The biology and treatment of EML4-ALK non-small cell lung cancer. *Eur J Cancer*. 2010;46(10):1773–1780.
2. Anai S, Takeshita M, Ando N, et al. A case of lung adenocarcinoma resistant to crizotinib harboring a novel EML4-ALK variant, exon 6 of EML4 fused to exon 18 of ALK. *J Thorac Oncol*. 2016;11(10):e126–e128.
3. Woo CG, Seo S, Kim SW, et al. Differential protein stability and clinical responses of EML4-ALK fusion variants to various ALK inhibitors in advanced ALK-rearranged non-small cell lung cancer. *Ann Oncol*. 2017;28(4):791–797.
4. Reguart N, Teixido C, Gimenez-Capitan A, et al. Identification of ALK, ROS1, and RET fusions by a multiplexed mRNA-based assay in formalin-fixed, paraffin-embedded samples from advanced non-small-cell lung cancer patients. *Clin Chem*. 2017;63(3):751–760.
5. Yoshida T, Oya Y, Tanaka K, et al. Differential crizotinib response duration among ALK fusion variants in ALK-positive non-small-cell lung cancer. *J Clin Oncol*. 2016;34(28):3383–3389.
6. Robesova B, Bajero M, Hausnerova J, Skrickova J, Tomiskova M, Dvorakova D. Identification of atypical ATRNL1 insertion to EML4-ALK fusion gene in NSCLC. *Lung Cancer*. 2015;87(3):318–320.

OncoTargets and Therapy**Dovepress****Publish your work in this journal**

OncoTargets and Therapy is an international, peer-reviewed, open access journal focusing on the pathological basis of all cancers, potential targets for therapy and treatment protocols employed to improve the management of cancer patients. The journal also focuses on the impact of management programs and new therapeutic agents and protocols on

patient perspectives such as quality of life, adherence and satisfaction. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <http://www.dovepress.com/oncotargets-and-therapy-journal>