Association of expression of *DRD2* rs1800497 polymorphism with migraine risk in Han Chinese individuals

Yingfeng Deng
Jianping Huang
Huijun Zhang
Xueqin Zhu
Qin Gong

Department of Anesthesiology, The
Affiliated Hospital of Medical School,
Ningbo University, Ningbo, China

Background: Previous studies suggested that single-nucleotide polymorphisms in dopamine receptor D2 (*DRD2*) are the susceptibility loci for migraine. This study was aimed at evaluating the contribution of *DRD2* rs1800497 and its expression to migraine risk in Han Chinese subjects.

Methods: In total, 250 patients with migraine and 250 age- and sex-matched control subjects were included in this study. TaqMan allelic discrimination assay was used for *DRD2* rs1800497 genotyping. Plasma *DRD2* concentration was determined using enzyme-linked immunosorbent assay.

Results: Significant associations were observed for the rs1800497 genotype ($\chi^2=6.37, p=0.041$) and allele ($\chi^2=4.69, p=0.03$; odds ratio [OR]=1.33, 95% CI=1.03–1.72, power=58%) frequencies between the migraine and control groups. Sex analysis indicated a positive association for rs1800497 between female patients with migraine and control individuals (genotype: $\chi^2=7.84, p=0.019$; allele: $\chi^2=6.60, p=0.010$; OR=1.61, 95% CI=1.12–2.30, power=73.4%). Furthermore, a significant association was observed only in female patients with migraine without aura (MO) (genotype: $\chi^2=6.88, p=0.032$; allele: $\chi^2=5.65, p=0.017$; OR=1.59, 95% CI=1.08–2.36, power=65.1%). The mean plasma *DRD2* levels in the control group (mean $\pm$ SD: 24.20 $\pm$ 2.78) were significantly lower than those in the migraine with aura (MA) (30.86 $\pm$ 3.69, $p<0.0001$) and MO groups (31.88 $\pm$ 4.99, $p<0.0001$). Additionally, there was a sex-based difference in *DRD2* expression in the MA (male vs female: 29.46 $\pm$ 3.59 vs 32.27 $\pm$ 3.27, $p<0.01$) and MO groups (male vs female: 29.18 $\pm$ 3.50 vs 34.58 $\pm$ 4.84, $p<0.0001$). Moreover, plasma *DRD2* levels in patients were significantly different among the three genotypes (CC vs CT vs TT: 24.76 $\pm$ 3.76 vs 30.93 $\pm$ 3.85 vs 37.06 $\pm$ 3.95, $p<0.0001$). Similar results were observed both in the MA (CC vs CT vs TT: 25.09 $\pm$ 3.84 vs 28.57 $\pm$ 2.84 vs 33.37 $\pm$ 1.58, $p<0.0001$) and MO groups (CC vs CT vs TT: 24.65 $\pm$ 3.79 vs 31.65 $\pm$ 3.86 vs 38.29 $\pm$ 3.74, $p<0.0001$).

Conclusion: Our case-control study suggested that the *DRD2* polymorphism rs1800497 was significantly associated with the risk of migraine in Han Chinese females. Additionally, the plasma *DRD2* level was high in patients with migraine. Females with migraine had considerably higher *DRD2* levels than males with migraine. *DRD2* expression may be regulated by *DRD2* rs1800497 genotype in patients with migraine.

Keywords: dopamine receptor D2, female, rs1800497, ELISA, migraine

Correspondence: Yingfeng Deng
Department of Anesthesiology, The
Affiliated Hospital of Medical School,
Ningbo University, No. 247 Renming
Road, Ningbo 315020, Zhejiang, China
Email dengyin4507@163.com

Introduction

Migraine is a common neurologic disorder that has become a major public health concern worldwide.¹ This disorder is characterized by recurrent, moderate-to-severe headaches. Clinical manifestations typically include unilateral headache with nausea, emesis, phonophobia, and visual sensory disturbance.² The two major types of migraine

are migraine with aura (MA) and migraine without aura (MO), as defined by the International Headache Society.³ Migraine is a complex disorder, and involves the influence of various environmental and genetic factors.^{4,5} While scientists have obtained valuable information concerning the development and clinical treatment of migraine, the mechanisms underlying the pathogenesis of migraine remain unclear.⁶

Dopamine has been suggested to be the second putative protagonist in headache.⁷ Among the proposed mechanisms, the dopaminergic system is responsible for maintaining the dopamine–norepinephrine ratio.⁸ An imbalance in this ratio makes individuals susceptible to migraine.⁹ Dopamine receptors are believed to play specific roles in the pathogenesis of migraine.¹⁰ Dopamine receptor D2 (DRD2) has been widely studied in central nervous system disorders, including schizophrenia, posttraumatic stress disorder, movement disorders, and migraine.¹¹ Both *DRD2* genotyping and monitoring of plasma drug concentrations may be useful for alleviating clinically dominant symptoms in patients.¹² Previous studies suggested that single-nucleotide polymorphisms (SNPs) in *DRD2* are susceptibility loci for migraine.¹³ *DRD2* polymorphism was observed to be significantly associated with the risk of migraine in an Indian population.¹⁴ *DRD2* haplotype may be a risk factor in migraine susceptibility. In Japanese subjects, Onaya et al showed that *DRD2* polymorphism significantly and independently contributed to the development of migraine from medication overuse headache.¹⁵ However, other studies did not support the contribution of *DRD2* toward the genetic predisposition to migraine in a Spanish population and a North Indian population.^{13,16} The SNP rs1800497 on chromosome 11q23.2 is located in exon 8 of the ankyrin repeat domain containing one gene downstream of *DRD2*.¹⁷ *DRD2* rs1800497 was observed to be associated with different conditions such as autism,¹⁸ cramps,¹⁹ migraine,¹⁴ memory loss,²⁰ and schizophrenia.²¹ *DRD2* rs1800497-T-carriers faced a significantly reduced risk of schizophrenia susceptibility in Asian populations.²² Gluskin and Mickey suggested that *DRD2* rs1800497 probably contributes to important individual differences in human striatal function, neuropsychiatric disease risk, and pharmacologic response.²³ *DRD2* rs1800497 has been consistently reported to be linked with reduced striatal D2 receptor expression in both postmortem expression studies and in vivo radioligand binding studies.^{24,25} However, this link between rs1800497 and D2 receptor expression was not apparent in patients with neuropsychiatric disorders.²³ The D2 receptor expression in these subjects tended to be higher among rs1800497-T-carriers than among C homozygotes.²³ Significant associations between *DRD2*

rs1800497 and migraine risk were observed in a primary cohort study (208 patients with migraine and 200 healthy control subjects).¹⁴ However, no significant associations were observed in a secondary cohort (127 patients with migraine and 200 healthy control individuals) of Indian subjects.¹⁴ Importantly, no studies have investigated the putative relationship between *DRD2* rs1800497 and migraine risk in Han Chinese populations.

In this study, we enrolled 500 age- and sex-matched volunteers and conducted a case–control study to validate the relationship between the *DRD2* polymorphism rs1800497 and migraine risk in a Chinese Han population. Additionally, we examined the role of *DRD2* in the pathogenesis of migraine by determining plasma *DRD2* levels.

Materials and methods

Case–control subjects

This study was approved by the Ethics Committees of the Affiliated Hospital of Ningbo University, and written informed consent was obtained from all participants. This study included 250 unrelated patients with migraine and 250 age- and sex-matched healthy control subjects. All subjects were enrolled randomly between January 2014 and October 2017 from the Affiliated Hospital of Ningbo University, Ningbo city, Zhejiang province, China. Patients were diagnosed by at least two independent pain specialists and neurologists based on their responses provided in a validated medical questionnaire prepared according to the International Headache Society.²⁶ Patients with migraine were assigned to two subgroups: MA and MO. The control subjects had no history of personal or familial migraine, and were selected randomly while undergoing a health examination in the hospital. Individuals with obesity, hypertension, diabetes, and other neurologic or psychiatric disorders were excluded from this study.

Laboratory analysis

A venous blood sample was collected from each subject within 12 h after fasting in an EDTA-coated vacutainer. Biochemical indices, including levels of triglyceride (TG), total cholesterol (TC), high-density lipoprotein cholesterol, low-density lipoprotein cholesterol, apolipoprotein A-I, and apolipoprotein B, were measured using an automated clinical chemistry analyzer (Beckman Coulter, Brea, CA, USA). Plasma was separated from the blood samples by centrifugation at 1000×g for 10 min. Totally, 168 sex- and age-matched individuals, including 56 patients with MA, 56 patients with MO, and 56 control individuals, were enrolled

for protein expression analysis. Plasma DRD2 concentration was determined using enzyme-linked immunosorbent assay kits (R&D Systems, Minneapolis, MN, USA).

Genotype analysis

Lymphocyte DNA was isolated from peripheral blood using the salting-out procedure. The TaqMan Allelic Discrimination Assay System (assay ID: C_7486676_10) was used for *DRD2* rs1800497 genotyping. All assays were optimized to a total volume of 10 μ L, containing 10 ng of genomic DNA, 0.125 μ L of primer-TaqMan Probe mixture, and 5 μ L of TaqMan Universal PCR Master Mix 2X (Thermo Fisher Scientific, Waltham, MA, USA). Polymerase chain reaction amplification was conducted using a 7500 Real-Time PCR System (Thermo Fisher Scientific). Thermal cycling conditions included 95°C for 10 min, followed by 50 cycles at 95°C for 15 s, and 60°C for 1 min. Genotypes were determined using the algorithm and software supplied by the manufacturer (Thermo Fisher Scientific). To determine the random genotyping error rate, 5% DNA samples were examined in duplicate, showing 100% consistency in genotype.

Statistical analysis

Continuous variables were expressed as the mean and SD and compared using the independent samples *t*-test. Categorical variables were compared using Pearson's chi-squared test. Quantitative data were compared using one-way analysis of variance or the Kruskal–Wallis test. The odds ratio (OR) and 95% CI were calculated by logistic regression to determine the relative risk of rs1800497 genotypes for migraine development. All statistical analyses were conducted with SPSS statistical software (version 16.0; SPSS, Inc., Chicago, IL, USA). Power analysis was performed using Power and Sample Size Calculation software (version 3.0.43).²⁷ Statistical significance was defined as a *p*-value <0.05.

Results

Clinical characteristics of subjects

The clinical characteristics of both groups are listed in Table 1. The case group comprised 250 patients (125 males and 125 females) experiencing migraine with a mean age of 43.02±16.70 years, while the control group comprised 250 healthy subjects (125 males and 125 females) with a mean age of 41.98±11.23 years. Among the patients, 68 had MA (39 males and 29 females) with a mean age of 44.00±17.33 years and 182 had MO (86 males and 96 females) with a mean age of 41.04±9.70 years. No significant differences were observed in the biochemical indices (levels of TG, TC, high-density lipoprotein, low-density lipoprotein, apolipoprotein A-I, and apolipoprotein B) between the patients with migraine and controls (*p*>0.05).

Comparison of genotype and allelic distributions

The genotype distributions of *DRD2* rs1800497 in the control subjects and patients were within the Hardy–Weinberg equilibrium model (*p*>0.05). As shown in Table 2, significant differences were observed in the genotype ($\chi^2=6.37$, *p*=0.041) and allele ($\chi^2=4.69$, *p*=0.03; OR=1.33, 95% CI=1.03–1.72, power=58%) frequencies between the migraine and the control groups. Sex analysis indicated a positive association for rs1800497 between female patients with migraine and control subjects (genotype: $\chi^2=7.84$, *p*=0.019; allele: $\chi^2=6.60$, *p*=0.010; OR=1.61, 95% CI=1.12–2.30, power=73.4%). Furthermore, a significant association was observed only in female patients with MO (genotype: $\chi^2=6.88$, *p*=0.032; allele: $\chi^2=5.65$, *p*=0.017; OR=1.59, 95% CI=1.08–2.36, power=65.1%). However, no significant difference was observed in the male groups (*p*>0.05). The genetic models analyzed are listed in Table 3, and an increased risk was observed between the case and control groups under the

Table 1 Comparison of characteristics between cases and controls

Characteristics	Control (250)	Migraine (250)	MA (68)	MO (182)	<i>p</i> ₁	<i>p</i> ₂	<i>p</i> ₃
Age, years, mean±SD	41.98±11.23	43.02±16.70	44.00±17.33	41.04±9.70	0.414	0.248	0.364
Male (n)	125	125	39	86	1.000	0.549	0.741
TG (mmol/L), mean±SD	1.48±0.88	1.50±1.11	1.58±0.99	1.48±1.04	0.823	0.419	1.000
TC (mmol/L), mean±SD	4.38±1.01	4.41±1.18	4.49±1.08	4.39±1.04	0.760	0.433	0.920
HDL (mmol/L), mean±SD	1.12±0.31	1.16±0.69	1.10±0.78	1.18±0.39	0.403	0.747	0.076
LDL (mmol/L), mean±SD	2.55±0.86	2.44±0.89	2.49±0.91	2.39±0.99	0.160	0.615	0.074
ApoA (g/L), mean±SD	1.09±0.21	1.07±0.22	1.09±0.52	1.06±0.72	0.299	1.000	0.533
ApoB (g/L), mean±SD	0.81±0.41	0.79±0.25	0.89±0.23	0.75±0.45	0.511	0.124	0.150

Note: *p*₁, the *p*-value for control vs migraine; *p*₂, the *p*-value for control vs MA; *p*₃, the *p*-value for control vs MO.

Abbreviations: ApoA, apolipoprotein A; ApoB, apolipoprotein B; HDL, high-density lipoprotein; LDL, low-density lipoprotein; MA, migraine with aura; MO, migraine without aura; TC, total cholesterol; TG, triglyceride.

Table 2 Distribution of rs1800497 genotypes and alleles in both groups

Gender	Group	Genotype			χ^2	p (df=2)	Allele		χ^2	p (df=1)	OR (95% CI)
		CC, n (%)	CT, n (%)	TT, n (%)			C, n (%)	T, n (%)			
All	Control	99 (0.40)	135 (0.54)	16 (0.06)			333 (0.67)	167 (0.33)			
	Case	80 (0.32)	140 (0.56)	30 (0.12)	6.37	0.041	300 (0.60)	200 (0.40)	4.69	0.03	1.33 (1.03–1.72)
	MO	59 (0.32)	102 (0.56)	21 (0.12)	4.81	0.09	220 (0.60)	144 (0.40)	3.47	0.062	1.31 (0.98–1.73)
	MA	21 (0.31)	38 (0.56)	9 (0.13)	4.29	0.117	80 (0.59)	56 (0.41)	2.84	0.092	1.39 (0.95–2.06)
Male	Control	48 (0.38)	70 (0.56)	7 (0.06)			166 (0.66)	84 (0.34)			
	Case	46 (0.37)	69 (0.55)	10 (0.08)	0.58	0.748	161 (0.64)	89 (0.36)	0.22	0.639	1.09 (0.76–1.58)
	MO	32 (0.37)	49 (0.57)	5 (0.06)	0.03	0.984	113 (0.66)	59 (0.34)	0.02	0.887	1.03 (0.68–1.55)
	MA	14 (0.36)	20 (0.51)	5 (0.13)	2.29	0.318	48 (0.62)	30 (0.39)	0.62	0.431	1.23 (0.73–2.09)
Female	Control	51 (0.41)	65 (0.52)	9 (0.07)			167 (0.67)	83 (0.33)			
	Case	34 (0.27)	71 (0.57)	20 (0.16)	7.84	0.019	139 (0.56)	111 (0.44)	6.60	0.01	1.61 (1.12–2.30)
	MO	27 (0.28)	53 (0.55)	16 (0.17)	6.88	0.032	107 (0.56)	85 (0.44)	5.65	0.017	1.59 (1.08–2.36)
	MA	7 (0.24)	18 (0.62)	4 (0.14)	3.39	0.184	32 (0.55)	26 (0.45)	2.78	0.095	1.63 (0.91–2.92)

Note: Results with p -values less than 0.05 are shown in bold font. The minor allele frequency of rs1800497-T was 0.19 in HapMap-CEU; 0.41 in -JPT; 0.35 in -CHB; 0.44 in -CHD.

Abbreviations: df, degrees of freedom; MA, migraine with aura; MO, migraine without aura; OR, odds ratio.

Table 3 Comparison of the dominant model and recessive model in rs1800497 between cases and controls by gender

Gender	Group	Dominant model		χ^2	p	OR (95% CI)	Recessive model		χ^2	p	OR (95% CI)
		CC, n (%)	CT+TT, n (%)				CC+CT, n (%)	TT, n (%)			
All	Control	99 (0.40)	151 (0.60)				234 (0.94)	16 (0.06)			
	Case	80 (0.32)	170 (0.68)	3.14	0.076	1.39 (0.97–2.01)	220 (0.88)	30 (0.12)	4.69	0.03	1.99 (1.06–3.76)
	MO	59 (0.32)	123 (0.68)	2.34	0.126	1.37 (0.92–2.04)	161 (0.88)	21 (0.12)	3.55	0.059	1.91 (0.96–3.77)
	MA	21 (0.31)	47 (0.69)	1.73	0.188	1.47 (0.83–2.60)	59 (0.87)	9 (0.13)	3.45	0.063	2.23 (0.94–5.29)
Male	Control	48 (0.38)	77 (0.62)				118 (0.94)	7 (0.06)			
	Case	46 (0.37)	79 (0.63)	0.07	0.791	1.07 (0.64–1.78)	115 (0.92)	10 (0.08)	0.57	0.45	1.47 (0.54–3.98)
	MO	32 (0.37)	54 (0.63)	0.03	0.862	1.05 (0.59–1.85)	81 (0.94)	5 (0.06)	NA	NA	1.04 (0.32–3.39)
	MA	14 (0.36)	25 (0.64)	0.08	0.777	1.11 (0.53–2.35)	34 (0.87)	5 (0.13)	NA	NA	2.48 (0.74–8.31)
Female	Control	51 (0.41)	74 (0.59)				116 (0.92)	9 (0.08)			
	Case	34 (0.27)	91 (0.73)	5.15	0.023	1.84 (1.08–3.14)	105 (0.84)	20 (0.16)	4.72	0.029	2.45 (1.07–5.63)
	MO	27 (0.28)	69 (0.62)	3.82	0.05	1.76 (1.00–3.11)	80 (0.83)	16 (0.17)	4.85	0.027	2.58 (1.09–6.12)
	MA	7 (0.24)	22 (0.76)	2.78	0.095	2.17 (0.86–5.45)	25 (0.86)	4 (0.14)	NA	NA	2.06 (0.59–7.23)

Abbreviations: MA, migraine with aura; MO, migraine without aura; OR, odds ratio; NA, not available.

recessive model (CC+CT vs TT: $\chi^2=4.69$, $p=0.03$; OR=1.99, 95% CI=1.06–3.76, power=57.9%). In the sex analysis, positive associations were observed in female patients with migraine under the dominant ($\chi^2=5.15$, $p=0.023$; OR=1.84, 95% CI=1.08–3.14, power=61.9%) and recessive models ($\chi^2=4.72$, $p=0.029$; OR=2.45, 95% CI=1.07–5.63, power=58.3%). Significant collections were likely to be present in the patients with MO under the dominant ($\chi^2=3.82$, $p=0.05$; OR=1.76, 95% CI=1.00–3.11, power=49.7%) and recessive models ($\chi^2=4.85$, $p=0.027$; OR=2.58, 95% CI=1.09–6.12, power=59.3%).

DRD2 expression and genotype

The mean plasma DRD2 levels in the control individuals (mean±SD: 24.20±2.78 ng/L) were significantly lower than those in the patients with MA (30.86±3.69 ng/L, $p<0.0001$) and MO (31.88±4.99 ng/L, $p<0.0001$; Figure 1). Addition-

Figure 1 Comparison of DRD2 expression between cases and controls.

Notes: The plasma level of DRD2 (mean±SD): controls (24.20±2.78 ng/L), MA patients (30.86±3.69 ng/L), MO patients (31.88±4.99 ng/L), male controls (24.45±3.14 ng/L), female controls (23.96±2.39 ng/L), male MA patients (29.46±3.59 ng/L), female MA patients (32.27±3.27 ng/L), male MO patients (29.18±3.50 ng/L), female MO patients (34.58±4.84 ng/L). ** $p<0.01$, *** $p<0.0001$.

Abbreviations: DRD2, dopamine receptor D2; MA, migraine with aura; MO, migraine without aura.

ally, there was a sex-based difference in DRD2 expression in the patients with MA (male vs female: 29.46 ± 3.59 vs 32.27 ± 3.27 ng/L, $p < 0.01$) and MO (male vs female: 29.18 ± 3.50 vs 34.58 ± 4.84 ng/L, $p < 0.0001$; Figure 1). No significant difference was observed between the male (24.45 ± 3.14 ng/L) and female (23.96 ± 2.39 ng/L, $p > 0.05$; Figure 1) control individuals. Plasma DRD2 levels for different genotypes are shown in Figure 2. There were no differences in plasma DRD2 levels among three genotypes in the control group (CC vs CT vs TT: 23.41 ± 2.40 vs 24.51 ± 2.69 vs 25.52 ± 3.99 ng/L, $p > 0.05$). The DRD2 concentration in patients with migraine with genotype CC (24.76 ± 3.76 ng/L) was much lower than that in patients with genotypes CT (30.93 ± 3.85 ng/L, $p < 0.0001$) and TT (37.06 ± 3.95 ng/L, $p < 0.0001$). In the subsequent subgroup analysis, similar results were obtained for the patients with MA (CC vs CT vs TT: 25.09 ± 3.84 vs 28.57 ± 2.84 vs 33.37 ± 1.58 ng/L, $p < 0.0001$) and MO (CC vs CT vs TT: 24.65 ± 3.79 vs 31.65 ± 3.86 vs 38.29 ± 3.74 ng/L, $p < 0.0001$).

Discussion

In this case-control study, 250 patients with migraine (68 patients with MA and 182 with MO) and 250 unrelated healthy control individuals were analyzed. Our results showed that the *DRD2* polymorphism rs1800497 was strongly related to the risk of MO in Han Chinese subjects. Furthermore, through sex-stratified comparison, rs1800497 was observed to be associated with the risk of MO in females. DRD2 was highly expressed in the patients with MA and MO, and in the patients with migraine, its expression was regulated by the *DRD2* rs1800497 genotypes.

Growing evidence suggests that high lipid concentration constitutes a risk factor for the development of migraine.^{28,29}

Migraine frequency was shown to be associated with dyslipidemia in women.^{30,31} Plasma TC and TG levels may influence migraine severity.³⁰ Cinzia et al suggested that lipoprotein (a) levels differed significantly between female Italian patients and female Italian control subjects.³² However, Goulart et al reported that different lipid and lipoprotein subfractions were positively associated with MO in both sexes in Brazilians.³³ Rist et al showed that TG and TC levels were associated with the risk of MA development, although not with other forms of headache in the French elderly.²⁹ Chorazka et al observed that the levels of TC and low-density lipoprotein cholesterol may contribute to an increased risk of migraine in Polish subjects.³⁴ In this study, we observed no differences in lipid concentrations between the patients with migraine and control individuals. These conflicting findings may be related to the different dietary habits of the Chinese population.

Genome-wide association studies have revealed numerous genetic variants responsible for the risk factors for migraine development.^{35,36} Dopaminergic gene variants are the most studied entities in migraine studies.³⁷ For example, the dopamine beta-hydroxylase gene polymorphism rs6271 was suggested to be a genetic factor influencing migraine susceptibility in the Turkish population.³⁸ The dopamine transporter gene variant *SLC6A3* rs40184 contributed to the risk of MA development in Germans.¹³ An imbalance in dopamine receptors D2/D3 was shown to be associated with migraine attacks and ictal cutaneous allodynia.³⁹ *DRD2* polymorphisms may be the genetic determinants of detoxification outcome in patients with medication overuse headache.^{15,40} Although several studies reported that *DRD2* polymorphisms are associated with the risk of migraine, many other studies

Figure 2 Comparison of DRD2 expression among three genotypes in different groups.

Notes: The plasma level of DRD2 (mean ± SD): controls CC (23.41 ± 2.40 ng/L), controls CT (24.51 ± 2.69 ng/L), controls TT (25.52 ± 3.99 ng/L), migraines CC (24.76 ± 3.76 ng/L), migraines CT (30.93 ± 3.85 ng/L), migraines TT (37.06 ± 3.95 ng/L), MA patients CC (25.09 ± 3.84 ng/L), MA patients CT (28.57 ± 2.84 ng/L), MA patients TT (33.37 ± 1.58 ng/L), MO patients CC (24.65 ± 3.79 ng/L), MO patients CT (31.65 ± 3.86 ng/L), MO patients TT (38.29 ± 3.74 ng/L). * $p < 0.05$, *** $p < 0.0001$.

Abbreviations: DRD2, dopamine receptor D2; MA, migraine with aura; MO, migraine without aura.

showed both positive and negative relationships with *DRD2*. Significant associations of a functional polymorphism at the dopamine beta-hydroxylase locus with migraine risk were observed in Australian and German subjects,^{13,41} although not in Spanish subjects.⁴² The variant rs1800497 was located 9.5 kb downstream of *DRD2*. Ghosh et al observed a significant association between rs1800497 and migraine risk in a meta-analysis of an Indian population.¹⁴ However, the results of the meta-analysis involved in the two independent studies were contradictory.¹⁴ Our study including 500 age- and sex-matched individuals revealed that *DRD2* rs1800497 contributed to the risk of migraine in Han Chinese. This could be partly explained by the specific genetic background and dietary habits of the Chinese.

The SNP rs1800497 was reported to mediate protein–protein interactions.¹⁷ Previous studies suggested that rs1800497 robustly influenced the availability of striatal D2 dopamine receptor.²³ rs1800497 is located in the 3′ untranslated region of *DRD2*, near the promoter, which is an inhibitory domain for the binding of nuclear factor-kappa B (NF-κB)-regulated genes. There are two NF-κB binding sites in the *DRD2* promoter, and *DRD2* expression is regulated by NF-κB.⁴³ rs1800497 variants may influence NF-κB-mediated regulation of *DRD2* and upregulate *DRD2* transcription.¹⁴ *DRD2* expression may be involved in the mechanism of migraine pathogenesis, and *DRD2* antagonists have been proposed as antimigraine drugs.⁴⁴ Our study showed similar results. Plasma *DRD2* levels were significantly higher in the patients with migraine than in the control subjects. Additionally, the plasma *DRD2* level was regulated by different rs1800497 genotypes.

Sexual dimorphism has been observed to influence the prevalence and severity of neurologic disorders.⁴⁵ Females show a much higher prevalence of migraine worldwide than males.⁴⁶ Previous studies revealed that female sex hormones are the major factors determining the risk and characteristics of migraine, accounting for the sex-based differences observed in this study; however, there was evidence supporting the underlying genetic variance as well.¹⁹ Sazci et al reported a significant association of the *NNMT* rs694539 variant with female patients with migraine, but no association with male patients.⁴⁷ CoSkun et al suggested that the genotypes rs2229741-GG and rs726281-GG are responsible for reduced risk of migraine in female Turkish patients.⁴⁸ Dopamine signaling was demonstrated to be linked to the female hormones, namely, progesterone and prolactin, in the migraineurs.⁴⁹ Serum prolactin level was considerably high in the female migraineurs.⁵⁰ Our results showed that

rs1800497 was associated with the risk of migraine in female patients among the three genotypes. Plasma *DRD2* levels were significantly higher in female patients than in male patients. Our findings provide insight into the prediction of the risk of headache in females.

Conclusion

Our case–control study showed that the *DRD2* polymorphism rs1800497 was significantly associated with the risk of migraine in Han Chinese females. Additionally, plasma *DRD2* level was higher in patients with migraine than in control subjects. Female patients with migraine had much higher *DRD2* levels than male patients with migraine. *DRD2* expression may be regulated by *DRD2* rs1800497 genotypes in patients with migraine.

Author contributions

All authors contributed toward data analysis, drafting and revising the paper, and agree to be accountable for all aspects of the work.

Disclosure

The authors report no conflicts of interest in this work.

References

1. Raval AD, Shah A. National trends in direct health care expenditures among US adults with migraine: 2004 to 2013. *J Pain*. 2017;18(1):96–107.
2. Jacobs B, Dussor G. Neurovascular contributions to migraine: moving beyond vasodilation. *Neuroscience*. 2016;338:130–144.
3. Steiner TJ, Stovner LJ, Birbeck GL. Migraine: the seventh disabling. *J Headache Pain*. 2013;14:1.
4. Gasparini CF, Smith RA, Griffiths LR. Genetic and biochemical changes of the serotonergic system in migraine pathobiology. *J Headache Pain*. 2017;18(1):20.
5. Kowalska M, Predecki M, Kozubski W, Lianeri M, Dorszewska J. Molecular factors in migraine. *Oncotarget*. 2016;7(31):50708–50718.
6. Eising E, Huisman SM, Mahfouz A, et al. Gene co-expression analysis identifies brain regions and cell types involved in migraine pathophysiology: a GWAS-based study using the Allen Human Brain Atlas. *Hum Genet*. 2016;135(4):425–439.
7. Sicuteri F. Dopamine, the second putative protagonist in headache. *Headache*. 1977;17(3):129–131.
8. Barbanti P, Fofi L, Aurilia C, Egeo G. Dopaminergic symptoms in migraine. *Neurol Sci*. 2013;34(Suppl 1):S67–S70.
9. Gasparini CF, Sutherland HG, Griffiths LR. Studies on the pathophysiology and genetic basis of migraine. *Curr Genomics*. 2013;14(5):300–315.
10. Marmura MJ. Use of dopamine antagonists in treatment of migraine. *Curr Treat Options Neurol*. 2012;14(1):27–35.
11. Noble EP. D2 dopamine receptor gene in psychiatric and neurologic disorders and its phenotypes. *Am J Med Genet B Neuropsychiatr Genet*. 2003;116B(1):103–125.
12. Yasui-Furukori N, Tsuchimine S, Saito M, et al. Comparing the influence of dopamine D(2) polymorphisms and plasma drug concentrations on the clinical response to risperidone. *J Clin Psychopharmacol*. 2011;31(5):633–637.

13. Todt U, Netzer C, Toliat M, et al. New genetic evidence for involvement of the dopamine system in migraine with aura. *Hum Genet.* 2009;125(3):265–279.
14. Ghosh J, Pradhan S, Mittal B. Identification of a novel ANKK1 and other dopaminergic (DRD2 and DBH) gene variants in migraine susceptibility. *Neuromol Med.* 2013;15(1):61–73.
15. Onaya T, Ishii M, Katoh H, et al. Predictive index for the onset of medication overuse headache in migraine patients. *Neurol Sci.* 2013;34(1):85–92.
16. Ghosh J, Pradhan S, Mittal B. Role of dopaminergic gene polymorphisms (DBH 19 bp indel and DRD2 Nco I) in genetic susceptibility to migraine in North Indian population. *Pain Med.* 2011;12(7):1109–1111.
17. Neville MJ, Johnstone EC, Walton RT. Identification and characterization of ANKK1: a novel kinase gene closely linked to DRD2 on chromosome band 11q23.1. *Hum Mutat.* 2004;23(6):540–545.
18. Nuntamool N, Ngamsamut N, Vanwong N, et al. Pharmacogenomics and efficacy of risperidone long-term treatment in Thai autistic children and adolescents. *Basic Clin Pharmacol Toxicol.* 2017;121(4):316–324.
19. Zeuner KE, Acewicz A, Knutzen A, Dressler D, Lohmann K, Witt K. Dopamine DRD2 polymorphism (DRD2/ANKK1-Taq1A) is not a significant risk factor in writer's cramp. *J Neurogenet.* 2016;30(3–4):276–279.
20. Papenberg G, Becker N, Ferencz B, et al. Dopamine receptor genes modulate associative memory in old age. *J Cogn Neurosci.* 2017;29(2):245–253.
21. Miura I, Zhang JP, Hagi K, et al. Variants in the DRD2 locus and antipsychotic-related prolactin levels: a meta-analysis. *Psychoneuroendocrinology.* 2016;72:1–10.
22. Wang Y, Liu L, Xin L, et al. The -141C Ins/Del and Taq1A polymorphism in the dopamine D2 receptor gene may confer susceptibility to schizophrenia in Asian populations. *J Clin Neurosci.* 2016;30:1–7.
23. Gluskin BS, Mickey BJ. Genetic variation and dopamine D2 receptor availability: a systematic review and meta-analysis of human in vivo molecular imaging studies. *Transl Psychiatry.* 2016;6:e747.
24. Ritchie T, Noble EP. Association of seven polymorphisms of the D2 dopamine receptor gene with brain receptor-binding characteristics. *Neurochem Res.* 2003;28(1):73–82.
25. Hirvonen MM, Laakso A, Nagren K, Rinne JO, Pohjalainen T, Hietala J. C957T polymorphism of dopamine D2 receptor gene affects striatal DRD2 in vivo availability by changing the receptor affinity. *Synapse.* 2009;63(10):907–912.
26. Headache Classification Committee of the International Headache Society (IHS). The International Classification of Headache Disorders, 3rd edition (beta version). *Cephalalgia.* 2013;33(9):629–808.
27. Huang Y, Ye H, Hong Q, et al. Association of CDKN2BAS polymorphism rs4977574 with coronary heart disease: a case-control study and a meta-analysis. *Int J Mol Sci.* 2014;15(10):17478–17492.
28. Teber S, Bektas O, Yilmaz A, Aksoy E, Akar N, Deda G. Lipoprotein levels in pediatric migraine. *Pediatr Neurol.* 2011;45(4):225–228.
29. Rist PM, Tzourio C, Kurth T. Associations between lipid levels and migraine: cross-sectional analysis in the epidemiology of vascular ageing study. *Cephalalgia.* 2011;31(14):1459–1465.
30. Janoska M, Chorazka K, Domitrz I. Migraine frequency and its association with dyslipidemia in women. *Neurol Neurochir Pol.* 2015;49(2):95–98.
31. Kurth T, Ridker PM, Buring JE. Migraine and biomarkers of cardiovascular disease in women. *Cephalalgia.* 2008;28(1):49–56.
32. Cinzia F, Daniela P, Elena S, et al. Lipoprotein (a) [Lp(a)]: a possible link between migraine and stroke. *Transl Res.* 2009;153(1):44–47.
33. Goulart AC, Lotufo PA, Santos IS, et al. The relationship between migraine and lipid sub-fractions among individuals without cardiovascular disease: a cross-sectional evaluation in the Brazilian Longitudinal Study of Adult Health (ELSA-Brasil). *Cephalalgia.* Epub 2017 Jan 1.
34. Chorazka K, Janoska M, Swic P, Domitrz I. Body mass index and serum lipid levels in effect on the incidence and course of migraine. *Neurol Neurochir Pol.* 2013;47(6):572–576.
35. Freilinger T, Anttila V, de Vries B, et al. Genome-wide association analysis identifies susceptibility loci for migraine without aura. *Nat Genet.* 2012;44(7):777–782.
36. Gormley P, Anttila V, Winsvold BS, et al. Meta-analysis of 375,000 individuals identifies 38 susceptibility loci for migraine. *Nat Genet.* 2016;48(8):856–866.
37. Ducros A. Genetics of migraine. *Rev Neurol.* 2013;169(5):360–371.
38. Sezer S, Kurt S, Ates O. Analysis of dopamine beta hydroxylase gene polymorphisms in migraine. *Clin Neurol Neurosurg.* 2016;145:96–100.
39. DaSilva AF, Nascimento TD, Jassar H, et al. Dopamine D2/D3 imbalance during migraine attack and allodynia in vivo. *Neurology.* 2017;88(17):1634–1641.
40. Cargnin S, Viana M, Sances G, et al. Combined effect of common gene variants on response to drug withdrawal therapy in medication overuse headache. *Eur J Clin Pharmacol.* 2014;70(10):1195–1202.
41. Fernandez F, Colson N, Quinlan S, MacMillan J, Lea RA, Griffiths LR. Association between migraine and a functional polymorphism at the dopamine beta-hydroxylase locus. *Neurogenetics.* 2009;10(3):199–208.
42. Corominas R, Ribases M, Camina M, et al. Two-stage case-control association study of dopamine-related genes and migraine. *BMC Med Genet.* 2009;10:95.
43. Bontempi S, Fiorentini C, Busi C, Guerra N, Spano P, Missale C. Identification and characterization of two nuclear factor-kappaB sites in the regulatory region of the dopamine D2 receptor. *Endocrinology.* 2007;148(5):2563–2570.
44. Peroutka SJ. Dopamine and migraine. *Neurology.* 1997;49(3):650–656.
45. Clayton JA. Sex influences in neurological disorders: case studies and perspectives. *Dialogues Clin Neurosci.* 2016;18(4):357–360.
46. Woldeamanuel YW, Cowan RP. Migraine affects 1 in 10 people worldwide featuring recent rise: a systematic review and meta-analysis of community-based studies involving 6 million participants. *J Neurol Sci.* 2017;372:307–315.
47. Sazci A, Sazci G, Sazci B, Ergul E, Idrisoglu HA. Nicotinamide-N-Methyltransferase gene rs694539 variant and migraine risk. *J Headache Pain.* 2016;17(1):93.
48. CoSkun S, Yucel Y, Cim A, et al. Contribution of polymorphisms in ESR1, ESR2, FSHR, CYP19A1, SHBG, and NR1P1 genes to migraine susceptibility in Turkish population. *J Genet.* 2016;95(1):131–140.
49. Charbit AR, Akerman S, Goadsby PJ. Dopamine: what's new in migraine? *Curr Opin Neurol.* 2010;23(3):275–281.
50. Cavestro C, Rosatello A, Marino MP, Micca G, Asteggiano G. High prolactin levels as a worsening factor for migraine. *J Headache Pain.* 2006;7(2):83–89.

Journal of Pain Research

Publish your work in this journal

The Journal of Pain Research is an international, peer reviewed, open access, online journal that welcomes laboratory and clinical findings in the fields of pain research and the prevention and management of pain. Original research, reviews, symposium reports, hypothesis formation and commentaries are all considered for publication.

Submit your manuscript here: <https://www.dovepress.com/journal-of-pain-research-journal>

Dovepress

The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.