The cytochrome P450 isoenzyme and some new opportunities for the prediction of negative drug interaction in vivo

Dmitrij A Sychev,¹ Ghulam Md Ashraf,² Andrey A Svistunov,³ Maksim L Maksimov,⁴ Vadim V Tarasov,³ Vladimir N Chubarev,³ Vitalij A Otdelenov,¹ Natal'ja P Denisenko,¹ George E Barreto,^{5,6} Gjurmakch Aliev⁷⁻⁹

¹Russian Medical Academy of Postgraduate Education Studies, Moscow, Russia; ²King Fahd Medical Research Center, King Abdulaziz University, Jeddah, Saudi Arabia; ³Sechenov First Moscow State Medical University, Moscow, Russia; ⁴Branch Campus of the Federal State Budgetary Educational Institution of Further Professional Education «Russian Medical Academy of Continuous Professional Education» of the Ministry of Healthcare of the Russian Federation, Kazan State Medical Academy, Volga Region, Kazan, Russia; ⁵Departamento de Nutrición y Bioquímica, Facultad de Ciencias, Pontificia Universidad Javeriana, Bogotá D.C., Colombia; ⁶Instituto de Ciencias Biomédicas, Universidad Autónoma de Chile, Santiago, Chile; ⁷GALLY International Biomedical Research Consulting LLC, San Antonio, TX, USA; ⁸School of Health Science and Healthcare Administration, University of Atlanta, Johns Creek, GA, USA; ⁹Institute of Physiologically Active Compounds Russian Academy of Sciences, Chernogolovka, Russia

Correspondence: Gjurmakch Aliev
GALLY International Biomedical
Research Consulting LLC, 7733 Louis
Pasteur Drive, #330, San Antonio,
TX 78229, USA
Tel +1 440 263 7461
Email aliev03@gmail.com

Abstract: Cytochrome (CYP) 450 isoenzymes are the basic enzymes involved in Phase I biotransformation. The most important role in biotransformation belongs to CYP3A4, CYP2D6, CYP2C9, CYP2C19 and CYP1A2. Inhibition and induction of CYP isoenzymes caused by drugs are important and clinically relevant pharmacokinetic mechanisms of drug interaction. Investigation of the activity of CYP isoenzymes by using phenotyping methods (such as the determination of the concentration of specific substrates and metabolites in biological fluids) during drug administration provides the prediction of negative side effects caused by drug interaction. In clinical practice, the process of phenotyping of CYP isoenzymes and some endogenous substrates in the ratio of cortisol to 6 β -hydroxycortisol in urine for the evaluation of CYP3A4 activity has been deemed to be a quite promising, safe and minimally invasive method for patients nowadays.

Keywords: cytochrome CYP450, drug interaction, drug metabolism, phenotyping

Introduction

It is well known that efficacy and safety of drugs used in clinical practice mainly depend on drug concentration near target molecules. This concentration usually correlates with plasma drug concentration, which depends on absorption, distribution and elimination processes. Elimination includes biotransformation and/or excretion of a drug (Figure 1). The main enzymes of oxidation reaction in Phase I are cytochrome (CYP) 450 isoenzymes, alcohol dehydrogenase, aldehyde dehydrogenase and aminooxidase.¹ Recently, a number of studies devoted to endogenous and exogenous markers of CYP expression and activity were published. The results of those investigations could be promising in the clinical practice for the prediction of the patient's personal reaction to medication and negative drug interactions. Diczfalusy et al² reported that 4 β -hydroxycholesterol (4 β -OHC), which is formed by CYP3A4 from cholesterol, is a promising endogenous marker for the prediction of negative drug interactions with inducers of CYP3A enzymes or CYP3A inhibitors such as ritonavir or itraconazole. In this review, the authors also established a relationship between the concentration of 4 β -OHC and the number of active CYP3A5*1 alleles showing that 4 β -OHC was formed not only by CYP3A4 but also by CYP3A5 and thus proved that the above-mentioned correlations depend on gender as well (women had higher concentration of 4 β -OHC in comparison with men). 4 β -OHC has half-life of 17 days, and this is the reason why, in short-term studies, exogenous markers such as midazolam or quinine probes may be superior; but in long-term studies, 4 β -OHC is a sensitive marker of

Figure 1 Phases of biotransformation.

CYP3A activity especially to assess drug interactions caused by induction and inhibition of CYP.² CYP3A4 converts cortisol CYP3A4 to 6 β -hydroxycortisol (6 β -OHC), which is used as an endogenous marker in urine for the determination of CYP3A activity. CYP3A4 also converts cholesterol to another marker 4 β -OHC.² Mårde Arrhén et al³ compared the ratio of 4 β -OHC to cholesterol in plasma with the ratio of 6 β -OHC to cortisol in urine after the administration of rifampicin for 2 weeks, and after the analysis of obtained data, they concluded that the abovementioned markers give similar results about CYP activity. The concentration of cortisol and its metabolites in the body is flexible and depends on chronobiological rhythms; that is why the use of the ratio of 6 β -OHC to cortisol is more rational for the precision prediction of CYP3A4 induction by rifampicin.³ In their comparison of endogenous 4 β -OHC with midazolam as markers for CYP3A4 induction by rifampicin, Björkhem-Bergman et al⁴ showed that the 4 β -OHC ratio is comparable with midazolam clearance as a marker of CYP3A4 induction, and each may be used to evaluate CYP3A4 induction in clinical trials evaluating drug–drug interactions for new drugs. These data are in good agreement with the data of Kasichayanula et al,⁵

which showed that changes in the plasma of 4 β -OHC can be utilized as a surrogate for midazolam pharmacokinetics after the administration of multiple doses of potent CYP3A inducers (rifampicin). There is a more limited dynamic range for 4 β HCH for the assessment of potential CYP3A inhibitors. This review also confirmed the opinion of previous authors that 4 β -OHC is a valuable marker for the assessment of potential CYP3A inducers in early drug development and particularly in drug interaction.^{5,6} This review is devoted to the recent data about CYP isoenzyme and some new opportunities for the prediction of negative drug interaction in vivo.

Role of CYP isoenzymes in drug metabolism

The major biotransformation enzyme is CYP, which has more than 1,000 isoenzymes, of which five (CYP3A4, CYP2D6, CYP2C9, CYP2C19 and CYP1A2) metabolize 90% of all drugs (Table 1).⁷ CYP is a group of enzymes first isolated from liver microsomes of rats separately by Klingenberg⁸ and Garfinkel.⁹ Nomenclature of CYP includes family (CYP2, CYP3, etc.), subfamily (CYP2D, CYP3A, etc.) and the name of definite isoenzyme (CYP2D6, CYP3A4, etc.).¹⁰ CYP can be found in hepatocytes as well as in other cells of the body. According to the analysis of 200 most commonly prescribed drugs in the US (2002), about 73% of drugs are subject to metabolism, of which about 75% of the drugs are metabolized by CYP.¹¹ Type CYP3A isoenzyme metabolizes 46% of drugs, type CYP2C9 isoenzyme metabolizes 16% of drugs, type CYP2C19 + CYP2D6 isozymes metabolize 12% of drugs, type CYP1A isoenzyme metabolizes 9% of drugs and type CYP2B6 + CYP2E1 isoenzymes metabolize 2% of drugs.¹² Substrate specificity is an inherent characteristic of CYP isozymes, ie, they have the ability to bind and transform molecules with specific shape, charge and hydrophilic/hydrophobic characteristics.^{13,14} Some of the CYP isozymes have substrate stereospecificity, for example, CYP2C9 metabolizes S-warfarin (more active enantiomer of warfarin) and R-warfarin is metabolized by CYP1A2 and CYP3A4.^{15,16}

Table 1 The relative content of CYP isoenzymes in the liver, their basic share and their participation in the metabolism of drugs

CYP isoenzyme	Content in the liver (%)	Contribution to metabolism (% metabolizing drugs)
1A2	~13	8.2
2C	~18	15.8 (2C8, 2C9)
2D6	Up to 2.5	18.8 (2C18, 2C19)
3A4	Up to 28	34.1

Note: Data from Kasichayanula et al.⁵

Abbreviation: CYP, cytochrome.

Table 2 Typical substrates of basic CYP isozymes

CYP isozyme	Substrates
CYP1A2	Clozapine, caffeine, paracetamol, theophylline, phenacetin, R-warfarin
CYP2C9	Hexobarbital, zidovudine, losartan, paracetamol, testosterone, tolbutamide, phenytoin, celecoxib, S-warfarin
CYP2C19	Hexobarbital, diazepam, zidovudine, omeprazole, pantoprazole, testosterone, phenytoin, R-warfarin, S-warfarin
CYP2D6	Haloperidol, dextromethorphan, codeine, metoprolol, nortriptyline, paracetamol, pravastatin, propafenone
CYP3A4	Alprazolam, atorvastatin, vincristine, halothane, hydrocortisone, zidovudine, carbamazepine, codeine, cortisol, caffeine, lidocaine, lovastatin, midazolam, nifedipine, paracetamol, tacrolimus, tamoxifen, testosterone, phenytoin, cyclosporine, cyclophosphamide, erythromycin, R-warfarin, S-warfarin

Note: Data from Wadelius et al.⁵⁷ and Kasichayanula et al.⁵

Abbreviation: CYP, cytochrome.

Most isoenzymes demonstrate broad substrate specificity, ie, each isozyme is able to metabolize a wide spectrum of xenobiotics, including drugs¹ (Table 2). The activity of CYP isoenzymes may vary over a wide range after the exposure of inducers or inhibitors, thereby altering the metabolism of substrates of isoenzymes, which could result in drug–drug interactions.¹⁷ Isoenzyme CYP3A4 metabolizes approximately 40%–50% of the drugs used in clinical practice, including calcium slow channel blockers,¹⁸ macrolide antibiotics,¹⁹ statins (simvastatin, lovastatin, atorvastatin)²⁰ and some of the “new” oral anticoagulants from the group of direct factor Xa inhibitors (rivaroxaban, apixaban).²¹ The most important inducers of CYP3A4 are carbamazepine,²² phenobarbital,²³ phenytoin,²⁴ rifampicin²⁵ and St John’s wort extract.²⁶ Some antifungal agents from the azole group (ketoconazole, itraconazole),²⁷ protease inhibitors (indinavir, nelfinavir, ritonavir)²⁸ and clarithromycin²⁹ are the most important CYP3A4 inhibitors.

CYP2C9 is the main enzyme involved in the metabolism of many nonsteroidal anti-inflammatory drugs (NSAIDs): celecoxib,³⁰ diclofenac,³¹ ibuprofen,³² lornoxicam,³³ meloxicam,³⁴ naproxen,³⁵ S-warfarin,²² many antidiabetic sulfonylureas (glibenclamide,³⁶ glimepiride,³⁷ glipizide,³⁸ tolbutamide³⁹), angiotensin II receptor blockers (irbesartan,⁴⁰ losartan⁴¹), fluvastatin,⁴² phenytoin,^{38,43} tamoxifen⁴⁴ and cyclophosphamide.^{45,46} The most important inducer of CYP2C9 is rifampicin.⁴⁷ Fluconazole and amiodarone are significant inhibitors of CYP2C9. CYP2C19 is the main metabolic enzyme for proton pump inhibitors, which in turn are the inhibitors of this isoenzyme (the so-called autoinhibitors).^{48,49}

This enzyme is induced by carbamazepine, prednisolone and rifampicin.^{50,51} CYP2D6 metabolizes up to 20% of drugs, including tricyclic antidepressant amitriptyline, neuroleptics and β -adrenoblockers. Codeine is metabolized by CYP2D6 into the active metabolite of morphine.⁵² CYP2D6 is inhibited by fluoxetine,⁵³ quinidine⁵⁴ and bupropion.⁵⁵ Unlike other isozymes, CYP2D6 has no verified inducers,⁵⁶ but isoenzyme activity increases in the gestation period.⁵⁷ However, there are controversial data about its weak induction by dexamethasone and rifampicin.⁵⁸ CYP1A2 has no endogenous substrates. It metabolizes mainly xenobiotics, including theophylline, caffeine and acetaminophen. Autoinducers of CYP1A2 are polycyclic aromatic hydrocarbons (the main component of inhaled tobacco smoke), which by means of CYP1A2 are transformed into carcinogenic compounds.⁵⁹ Besides, components of tobacco, smoke, grilled meal⁶⁰ and broccoli⁶¹ are inducers of CYP1A2. Ciprofloxacin and fluvoxamine inhibit CYP1A2.⁶²

Gene polymorphism in many CYP isozymes may be responsible for the interindividual differences in the speed of biotransformation of drugs and certain drug–drug interactions.⁶³ The presence of single-nucleotide polymorphism (SNP) in the gene encoding a particular isozyme may lead to the synthesis of enzymes with altered activity, which in turn changes the pharmacokinetics of drugs metabolized by this isoenzyme. In clinical practice, it is possible to conduct pharmacogenetic testing of some drugs to identify genetic polymorphisms.⁴² This allows to predict the pharmacological response to these drugs, which in turn increases the efficacy and safety of drug therapy.⁵⁰ CYP isoenzymes play a pivotal role in the metabolism of many drugs used in clinical practice. The changes in isoenzymes activity are the basis of drug–drug interaction at the level of biotransformation; thus, it is important to study the effect of drugs on CYP isoenzymes for safe pharmacotherapy.⁵⁰

Inhibition of CYP isoenzymes

The inhibition of CYP isoenzymes is a particular mechanism of clinically significant drug interaction (Figure 2). A decrease in the metabolism of drugs which are substrates for the abovementioned enzyme leads to increased concentration of these drugs in plasma and exerts toxic effects.^{50,64} Inhibition of isoenzymes could be reversible as well as irreversible. The reversible mechanisms of inhibition can be competitive, noncompetitive and uncompetitive. In the type of competitive reversible inhibition, the drug inhibitor and drug substrate compete for the active site of isoenzyme; therefore, this type of inhibition could be overcome by increasing the concentration of drug substrate.⁶⁵ Mechanism of noncompetitive reversible

Figure 2 Schematic of drug–drug interaction between substrate and inhibitor of CYP isoenzyme.

Note: Data from Ritter et al.⁷²

Abbreviation: CYP, cytochrome.

inhibition is due to the binding of inhibitor with the nonfunctional part of CYP isoenzyme, which changes the conformation of active site and prevents its binding to the drug substrate; therefore, this type of inhibition cannot be reversed by increasing the concentration of drug substrate. Noncompetitive inhibition of CYP isoenzymes is observed upon binding of the inhibitor to isoenzyme–substrate complex.⁶⁶

There are two types of irreversible inhibition: true irreversible inhibition and quasi-irreversible inhibition. Under true irreversible inhibition, the inhibitor or its intermediate metabolite covalently binds to the heme of CYP isoenzyme, thereby inactivating it. A strong noncovalent bond is formed between inhibitor and CYP isoenzyme in the case of quasi-irreversible inhibition. During irreversible inhibition, the recovery time of isoenzyme activity depends on the time needed for the synthesis of a new isoenzyme. Inhibitors of CYP isoenzymes (Table 3) may be classified in vivo

according to the degree of inhibitory activity of drug substrates of the isoenzymes. The use of potent inhibitors result in more than five times increase in area under the curve (AUC) of drug substrate (a decrease in clearance of >80%); moderate inhibitors increase the AUC of drug substrate by two to five times (a decrease in clearance up to 50%–80%) and weak inhibitors are able to increase the AUC of drug substrate by 1.25–2 times (a decrease in clearance up to 20%–50%).⁶⁷

Induction of CYP isoenzymes

The absolute increase in the number and/or catalytic activity of isoenzymes and the associated reduction in the concentration of drug, which are substrate for CYP isoenzymes, are a result of the induction of isoenzymes. In most cases, the induction of isozymes is clinically expressed with the decrease in the pharmacological effects. In some cases, the induction of CYP isoenzymes results in the increase

Table 3 Inhibitors of major CYP isozymes

CYP isozyme	Potent inhibitors	Moderate inhibitors	Weak inhibitors
1A2	Ciprofloxacin, enoxacin, fluvoxamine	Zileuton, mexiletine, methoxsalen, felbamate, thiabendazole, phenylpropanolamine	Allopurinol, acyclovir, verapamil, disulfiram, caffeine, norfloxacin, propafenone, propranolol, ticlopidine, famotidine, cimetidine, echinacea extract
2C9		Amiodarone, miconazole, oxandrolone, fluconazole	Voriconazole, zafirlukast, capecitabine, co-trimoxazole (sulfamethoxazole + trimethoprim), metronidazole, sulfapyrazone, tigecycline, fluvastatin, fluvoxamine, etravirine
2C19	Fluconazole, fluvoxamine, ticlopidine	Fluoxetine, moclobemide, omeprazole, voriconazole	Armodafinil, carbamazepine, cimetidine, ethinyl estradiol, etravirine, somatotropin, felbamate, ketoconazole
2D6	Bupropion, paroxetine, fluoxetine, quinidine	Duloxetine, terbinafine, cinacalcet	Amiodarone, vemurafenib, verapamil, gefitinib, hydralazine, hydroxychloroquine, desvenlafaxine, diltiazem, diphenhydramine, imatinib, methadone, oral contraceptives, propafenone, ranitidine, ritonavir, sertraline, telithromycin, febuxostat, celecoxib, cimetidine, echinacea extract, escitalopram
3A4	Voriconazole, grapefruit juice (high concentration), itraconazole, ketoconazole, clarithromycin, lopinavir, nefazodone, posaconazole, ritonavir, saquinavir, telaprevir, telithromycin, conivaptan	Aprepitant, verapamil, grapefruit juice (normal concentration), darunavir, diltiazem, imatinib, ritonavir, fluconazole, fosamprenavir, ciprofloxacin, erythromycin	Alprazolam, amiodarone, amlodipine, atorvastatin, bicalutamide, zileuton, isoniazid, nilotinib, oral contraceptives, ranitidine, ranolazine, ticagrelor, tipranavir, fluvoxamine, fluoxetine, cyclosporin, cilostazol, cimetidine, an extract of goldenseal Canadian, Ginkgo biloba extract

Note: Data from Wu et al.⁵²

Abbreviation: CYP, cytochrome.

in pharmacological effects (in the case of the formation of active metabolites) and even in toxic effects. For example, the induction of CYP2E1 resulted in increased metabolism of acetaminophen and increased formation of hepatotoxic metabolite *N*-acetyl-*p*-benzoquinone imine.⁶⁸

The most important mechanism for the induction of CYP isozymes is the interaction of inducer with specific intracellular receptors that in fact are transcription regulator proteins (pregnane-X-receptor, androstane constitutive receptor, aryl-hydrocarbonate receptor, etc.),⁶⁹ thus forming a receptor-inducor complex. This complex penetrates into the cell nucleus and acts on the regulatory region of the gene that results in increased expression of the gene encoding CYP. There are mechanisms of induction, which are not associated with exposure to specific receptors. For example, the induction of CYP2E1 is associated with posttranscriptional stabilization of molecules of this isoenzyme.⁷⁰ Based on the degree of induction of biotransformation of drug substrates in vivo, inducers are classified into three types: potent ($\geq 80\%$ reduction in AUC), moderate (50%–80% reduction in AUC) and weak (20%–50% reduction in AUC).⁶⁷ Typical inductors are summarized in Table 4. Usually, it takes several days for the development of drug–drug interactions between drug inducer of CYP isoenzyme and drug substrate of this isozyme, because the mechanism of induction of CYP isoenzymes includes the induction of gene transcription encoding and subsequent synthesis.

Legal status of studies on the effect of drugs on biotransformation enzymes

The most important step in the safety assessment of new and already registered drugs is a comprehensive evaluation of their potential side effects.⁷¹ The demographic aging of the population and increase in the number of drugs used by patients (including nonprescription drugs) have enhanced the risk of drug–drug interactions.⁶⁰ Extensive study is being conducted worldwide to identify potential risk of negative

drug–drug interaction between new and registered drugs, and effective measures have been introduced to reduce the risk of side effects (dose adjustment, additional therapeutic monitoring, etc.).⁷² Usually, in vitro studies are conducted before in vivo studies. The aim of screening investigation in vitro is to determinate the usefulness of studies in vivo. The Food and Drug Administration (FDA) has recommended the inclusion of instructions for medical use of drugs with information on the possible inhibitory or inducing effect on the biotransformation enzymes in the section describing pharmacokinetics. In the European Union also, pharmacokinetic studies have been designed to assess the influence of new drug(s) on the effects of well-known drug(s).⁶⁷ The results of studies of side effects of new drug(s) could be used to predict their interaction with other drug(s) on the basis of the identified mechanisms of interaction.

Prediction of impact of drugs on CYP isoenzymes in vivo

The study of potential side effects of new drugs includes conducting studies on the effects of new drugs on the pharmacokinetics of marker substrates of CYP isoenzymes, extrapolation of the results of such study to other drugs, substrates of isozymes and, in some cases, the study of specific combinations of drugs to be able to work out specific recommendations on combined use of drugs.⁶⁰ Marker substrates are substances that are metabolized primarily by one CYP isoenzyme (Figure 3). The purpose of the review for the investigation of influence of drugs on the marker substrates of CYP isoenzymes was to determine the presence or absence of any effect on CYP isoenzymes as well as to observe the magnitude of such influence, so as to classify the drugs that exhibit inductive/inhibitory properties in three groups: potent, moderate and weak. It is noted that Table 5 does not list all substrates that could be used to assess the effects of new drugs on the activity of CYP isoenzymes in vivo. For some CYP isozymes, methods for assessing the

Table 4 Inductors of major CYP isozymes

CYP isozyme	Potent inducers	Moderate inducers	Weak inducers
1A2		Montelukast, tobacco smoke, phenytoin	Moricizine, omeprazole, phenobarbital
2C9		Arbamazepine, rifampicin	Aprepitant, bosentan, phenobarbital, St John's wort extract
2C19		Rifampicin, S-mephenytoin	Artemisinin
2D6	Inducers are not identified		Inducers are not identified
3A4	Avasimibe, carbamazepine, rifampicin, phenytoin, St John's wort extract	Bosentan, modafinil, nafcillin, etravirine, efavirenz, lurasidone	Aprepitant, armodafinil, pioglitazone, prednisolone, refinamid, echinacea extract

Note: Data from Wu et al.⁵²

Abbreviation: CYP, cytochrome.

Figure 3 Schematic of drug–drug interaction between the inducer and the substrate of CYP isoenzyme.

Note: Data from Ritter et al.⁷²

Abbreviation: CYP, cytochrome.

activity were developed. The special feature of those methods is that they do not require the administration of any xenobiotics. For example, in one of the guidelines for pharmaceutical companies, it was recommended to study biotransformation and transporters of new drugs for the evaluation of CYP3A4 activity in vivo by using the ratio of endogenous cortisol to one of its metabolites, 6 β -OHC, which is formed exclusively as a result of CYP3A4 activity⁷³ (Figure 4).

The past study showed that the dynamics of endogenous activity of CYP3A4 metabolic markers, including the ratio of 6 β -OHC to cortisol in urine, correlate with midazolam clearance dynamics under the induction or inhibition of CYP3A4 in healthy volunteers; therefore, the ratio of 6 β -OHC to cortisol in urine could be used to evaluate the activity of CYP3A4 in vivo.⁷⁴ Formation clearance of the sum of 6 β -OHC and 6 β -hydroxycortisone was shown to detect moderate and potent inhibition of CYP3A4 in vivo.⁷⁵ CYP2D6 activity may be evaluated using pinoline urinary metabolic ratio

(6-methoxy-1,2,3,4-tetrahydro- β -carboline/6-hydroxy-1,2,3,4-tetrahydro- β -carboline).⁷⁶ To estimate CYP2C19 activity in vivo, omeprazole plasma hydroxylation index (omeprazole/5-hydroxyomeprazole) was introduced.⁷⁷ Losartan test could be used to assess the activity of CYP2C9. This test is based on the determination of concentration of losartan and its active metabolite, E-3174, which is produced mainly by CYP2C9 activity.⁷⁸ According to in vitro studies, E-3174 is also formed by the action of CYP3A4,^{79,80} but in vivo studies with therapeutic doses of losartan showed no significant contribution to losartan CYP3A4 metabolism.⁸¹ The predominant contribution of CYP2C9 to losartan metabolism was confirmed indirectly by a decrease in the AUC of E-3174 when it was used in combination with moderate inhibitor CYP2C9 (fluconazole), while no change in AUC of E-3174 was observed when losartan was used in combination with a strong CYP3A4 inhibitor (itraconazole).⁴⁷ In many studies, losartan has been safely used as a marker substrate for

Table 5 Examples of sensitive substrates of basic CYP isoenzymes and substrates with a narrow therapeutic range

CYP isoenzyme	Sensitive substrate ^a	Substrates with a narrow therapeutic range
1A2	Alosetron, duloxetine, caffeine, melatonin, ramelteon, tacrine, tizanidine	Theophylline, tizanidine
2C9	Celecoxib	Warfarin, phenytoin
2C19	S-mephenytoin, clobazam, lansoprazole, omeprazole	S-mephenytoin
2D6	Atomoxetine, venlafaxine, desipramine, dextromethorphan, metoprolol, nebivolol, perphenazine, tolterodine	Pimozide, thioridazine
3A4	Alfentanil, aprepitant, budesonide, buspirone, vardenafil, dasatinib, darifenacin, darunavir, dronedarone, indinavir, quetiapine, conivaptan, lovastatin, lopinavir, lurasidone, maraviroc, midazolam, nisoldipine, saquinavir, sildenafil, simvastatin, sirolimus, ticagrelor, tipranavir, tolvaptan, triazolam, felodipine, fluticasone, everolimus, eletriptan, eplerenone	Alfentanil, astemizole, dihydroergotamine, pimozide, sirolimus, tacrolimus, terfenadine, fentanyl, quinidine, cisapride, cyclosporine, ergotamine

Notes: ^aThe sensitive substrate is a substrate with AUC values more than fivefold increase in the combined use with an inhibitor of CYP isoenzymes. Data from Wadelius et al.⁵⁷

Abbreviations: AUC, area under the curve; CYP, cytochrome.

Figure 4 Metabolic conversion of cortisol.

Note: Data from Gray et al.⁸⁵

Abbreviations: DHF, 20β-DHF (20β-dihydrocortisol); DHE, 20β-DHE(20β-dihydrocortisone); THF, 3α,5β-THF (3α,5β-tetrahydrocortisol); THE, 3α,5β-THE (3α,5β-tetrahydrocortisone); OHC, 6 β-OHC(6 β-hydrocortisol); OHE, 6 β-OHE(6 β-hydrocortisone).

phenotyping CYP2C9 activity.^{82–84} Relative clinical safety of losartan and reliability of losartan test allow its usage in clinical studies dedicated to the investigations of the influence of new drugs on CYP2C9 activity in vivo.

Conclusion and future perspectives

Prediction of negative drug interaction is one of the most important and difficult objectives of clinical pharmacology.

CYP isoenzymes are the basic enzymes involved in Phase I biotransformation. The use of CYP isoenzymes, cortisol to 6β-OHC ratio, losartan test and other tests used for the estimation of the activity of CYP so as to investigate the influence of new drugs can yield new and prospective findings related to the prediction of negative drug interaction, which in turn can provide new insights for researchers and clinicians.

Acknowledgments

Gjumrakch Aliev's work was supported by a grant from the Russian Science Foundation (directed design, synthesis and study of biological activity of multitarget compounds as innovative drugs for the treatment of neurodegenerative diseases: PHΦ 14-23-00160Π). Gjumrakch Aliev is also grateful for the equipment of Center for Collective Use, Institute of Physiologically Active Compounds of Russian Academy of Sciences Chernogolovka, Russia, that was used for this review.

Disclosure

The authors report no conflicts of interest in this work.

References

- Zanger UM, Schwab M. Cytochrome P450 enzymes in drug metabolism: regulation of gene expression, enzyme activities, and impact of genetic variation. *Pharmacol Ther*. 2013;138(1):103–141.
- Diczfalusy U, Nylén H, Elander P, Bertilsson L. 4β-Hydroxycholesterol, an endogenous marker of CYP3A4/5 activity in humans. *Br J Clin Pharmacol*. 2011;71(2):183–189.
- Mårde Arrhen Y, Nylén H, Lövgren-Sandblom A, Kanebratt KP, Wide K, Diczfalusy U. A comparison of 4β-hydroxycholesterol: cholesterol and 6β-hydroxycortisol: cortisol as markers of CYP3A4 induction. *Br J Clin Pharmacol*. 2013;75(6):1536–1540.
- Björkhem-Bergman L, Bäckström T, Nylén H, et al. Comparison of endogenous 4β-hydroxycholesterol with midazolam as markers for CYP3A4 induction by rifampicin. *Drug Metab Dispos*. 2013;41(8):1488–1493.
- Kasichayanula S, Boulton DW, Luo W-L, et al. Validation of 4β-hydroxycholesterol and evaluation of other endogenous biomarkers for the assessment of CYP3A activity in healthy subjects. *Br J Clin Pharmacol*. 2014;78(5):1122–1134.
- Leil TA, Kasichayanula S, Boulton DW, LaCreta F. Evaluation of 4β-hydroxycholesterol as a clinical biomarker of CYP3A4 drug interactions using a Bayesian mechanism-based pharmacometric model. *CPT Pharmacometrics Syst Pharmacol*. 2014;3(6):e120.
- Pinto N, Dolan ME. Clinically relevant genetic variations in drug metabolizing enzymes. *Curr Drug Metab*. 2011;12(5):487–497.
- Klingenberg M. Pigments of rat liver microsomes. *Arch Biochem Biophys*. 1958;75(2):376–386.
- Garfinkel D. Studies on pig liver microsomes. I. Enzymic and pigment composition of different microsomal fractions. *Arch Biochem Biophys*. 1958;77(2):493–509.
- Amal AIO, Murry DJ. Pharmacogenetics of the cytochrome P450 enzyme system: review of current knowledge and clinical significance. *J Pharm Pract*. 2007;20(3):206–218.
- Thunell S, Pomp E, Brun A. Guide to drug porphyrinogenicity prediction and drug prescription in the acute porphyrias. *Br J Clin Pharmacol*. 2007;64(5):668–679.
- Williams JA, Hyland R, Jones BC, et al. Drug–drug interactions for UDP-glucuronosyltransferase substrates: a pharmacokinetic explanation for typically observed low exposure (AUCi/AUC) ratios. *Drug Metab Dispos*. 2004;32(11):1201–1208.
- Roy J. Introduction to the Pharmaceutical Sciences. Elsevier, 2011.
- Sirim D, Widmann M, Wagner F, Pleiss J. Prediction and analysis of the modular structure of cytochrome P450 monooxygenases. *BMC Struct Biol*. 2010;10:34.
- Kaminsky LS, Zhang ZY. Human P450 metabolism of warfarin. *Pharmacol Ther*. 1997;73(1):67–74.
- Hosokawa M. Structure and catalytic properties of carboxylesterase isozymes involved in metabolic activation of prodrugs. *Molecules*. 2008;13(2):412–431.
- Teo YL, Ho HK, Chan A. Metabolism-related pharmacokinetic drug–drug interactions with tyrosine kinase inhibitors: current understanding, challenges and recommendations. *Br J Clin Pharmacol*. 2015;79(2):241–253.
- Katoh M, Nakajima M, Yamazaki H, Yokoi T. Inhibitory potencies of 1,4-dihydropyridine calcium antagonists to P-glycoprotein-mediated transport: comparison with the effects on CYP3A4. *Pharm Res*. 2000;17(10):1189–1197.
- Rodrigues AD, Roberts EM, Mulford DJ, Yao Y, Ouellet D. Oxidative metabolism of clarithromycin in the presence of human liver microsomes. Major role for the cytochrome P4503A (CYP3A) subfamily. *Drug Metab Dispos*. 1997;25(5):623–630.
- Corsini A, Bellista S, Baetta R, Fumagalli R, Paoletti R, Bernini F. New insights into the pharmacodynamic and pharmacokinetic properties of statins. *Pharmacol Ther*. 1999;84(3):413–428.
- Heidbuchel H, Verhamme P, Alings M, et al. Updated European Heart Rhythm Association Practical Guide on the use of non-vitamin K antagonist anticoagulants in patients with non-valvular atrial fibrillation. *Europace*. 2015;17(10):1467–1507.
- Bertilsson L, Tybring G, Widén J, Chang M, Tomson T. Carbamazepine treatment induces the CYP3A4 catalysed sulfoxidation of omeprazole, but has no or less effect on hydroxylation via CYP2C19. *Br J Clin Pharmacol*. 1997;44(2):186–189.
- Ohno M, Motojima K, Okano T, Taniguchi A. Induction of drug-metabolizing enzymes by phenobarbital in layered co-culture of a human liver cell line and endothelial cells. *Biol Pharm Bull*. 2009;32(5):813–817.
- Fleishaker JC, Pearson LK, Peters GR. Phenytoin causes a rapid increase in 6 beta-hydroxycortisol urinary excretion in humans – a putative measure of CYP3A induction. *J Pharm Sci*. 1995;84(3):292–294.
- Backman JT, Olkkola KT, Neuvonen PJ. Rifampin drastically reduces plasma concentrations and effects of oral midazolam. *Clin Pharmacol Ther*. 1996;59(1):7–13.
- Rahimi R, Abdollahi M. An update on the ability of St. John's wort to affect the metabolism of other drugs. *Expert Opin Drug Metab Toxicol*. 2012;8(6):691–708.
- Varhe A, Olkkola KT, Neuvonen PJ. Oral triazolam is potentially hazardous to patients receiving systemic antimycotics ketoconazole or itraconazole. *Clin Pharmacol Ther*. 1994;56(6 pt 1):601–607.
- Eagling VA, Back DJ, Barry MG. Differential inhibition of cytochrome P450 isoforms by the protease inhibitors, ritonavir, saquinavir and indinavir. *Br J Clin Pharmacol*. 1997;44(2):190–194.
- Akiyoshi T, Ito M, Murase S, et al. Mechanism-based inhibition profiles of erythromycin and clarithromycin with cytochrome P450 3A4 genetic variants. *Drug Metab Pharmacokinet*. 2013;28(5):411–415.
- Chan AT, Zauber AG, Hsu M, et al. Cytochrome P450 2C9 variants influence response to celecoxib for prevention of colorectal adenoma. *Gastroenterology*. 2009;136(7):2127–2136.e2121.
- Morin S, Lorient MA, Poirier JM, et al. Is diclofenac a valuable CYP2C9 probe in humans? *Eur J Clin Pharmacol*. 2001;56(11):793–797.
- Berka K, Hendrychová T, Anzenbacher P, Otyepka M. Membrane position of ibuprofen agrees with suggested access path entrance to cytochrome P450 2C9 active site. *J Phys Chem A*. 2011;115(41):11248–11255.
- Guo Y, Zhang Y, Wang Y, et al. Role of CYP2C9 and its variants (CYP2C9*3 and CYP2C9*13) in the metabolism of lornoxicam in humans. *Drug Metab Dispos*. 2005;33(6):749–753.
- Chesné C, Guyomard C, Guillozo A, Schmid J, Ludwig E, Sauter T. Metabolism of meloxicam in human liver involves cytochromes P4502C9 and 3A4. *Xenobiotica*. 1998;28(1):1–13.
- Bae J-W, Kim J-H, Choi C-I, et al. Effect of CYP2C9*3 allele on the pharmacokinetics of naproxen in Korean subjects. *Arch Pharm Res*. 2009;32(2):269–273.
- Zhang Y-F, Chen X-Y, Guo Y-J, Si D-Y, Zhou H, Zhong D-F. [Impact of cytochrome P450 CYP2C9 variant allele CYP2C9*3 on the pharmacokinetics of glibenclamide and lornoxicam in Chinese subjects]. *Acta Pharm Sin*. 2005;40(9):796–799.

37. Niemi M, Cascorbi I, Timm R, Kroemer HK, Neuvonen PJ, Kivistö KT. Glyburide and glimepiride pharmacokinetics in subjects with different CYP2C9 genotypes. *Clin Pharmacol Ther.* 2002;72(3):326–332.
38. Kidd RS, Straughn AB, Meyer MC, Blaisdell J, Goldstein JA, Dalton JT. Pharmacokinetics of chlorpheniramine, phenytoin, glipizide and nifedipine in an individual homozygous for the CYP2C9*3 allele. *Pharmacogenetics.* 1999;9(1):71–80.
39. Kirchheiner J, Bauer S, Meineke I, et al. Impact of CYP2C9 and CYP2C19 polymorphisms on tolbutamide kinetics and the insulin and glucose response in healthy volunteers. *Pharmacogenetics.* 2002; 12(2):101–109.
40. Chen G, Jiang S, Mao G, et al. CYP2C9 Ile359Leu polymorphism, plasma irbesartan concentration and acute blood pressure reductions in response to irbesartan treatment in Chinese hypertensive patients. *Methods Find Exp Clin Pharmacol.* 2006;28(1):19–24.
41. McCrear JB, Cribb A, Rushmore T, et al. Phenotypic and genotypic investigations of a healthy volunteer deficient in the conversion of losartan to its active metabolite E-3174. *Clin Pharmacol Ther.* 1999;65(3): 348–352.
42. Tutton R. Pharmacogenomic biomarkers in drug labels: what do they tell us? *Pharmacogenomics.* 2014;15(3):297–304.
43. Veronese ME, Mackenzie PI, Doecke CJ, McManus ME, Miners JO, Birkett DJ. Tolbutamide and phenytoin hydroxylations by cDNA-expressed human liver cytochrome P4502C9. *Biochem Biophys Res Commun.* 1991;175(3):1112–1118.
44. Collier JK, Krebsfaenger N, Klein K, et al. The influence of CYP2B6, CYP2C9 and CYP2D6 genotypes on the formation of the potent anti-oestrogen Z-4-hydroxy-tamoxifen in human liver. *Br J Clin Pharmacol.* 2002;54(2):157–167.
45. Ekhardt C, Doodeman VD, Rodenhuis S, Smits PHM, Beijnen JH, Huitema ADR. Influence of polymorphisms of drug metabolizing enzymes (CYP2B6, CYP2C9, CYP2C19, CYP3A4, CYP3A5, GSTA1, GSTP1, ALDH1A1 and ALDH3A1) on the pharmacokinetics of cyclophosphamide and 4-hydroxycyclophosphamide. *Pharmacogenet Genomics.* 2008;18(6):515–523.
46. Griskevicius L, Yasar U, Sandberg M, et al. Bioactivation of cyclophosphamide: the role of polymorphic CYP2C enzymes. *Eur J Clin Pharmacol.* 2003;59(2):103–109.
47. Rana R, Chen Y, Ferguson SS, Kissling GE, Surapureddi S, Goldstein JA. Hepatocyte nuclear factor 4{alpha} regulates rifampicin-mediated induction of CYP2C genes in primary cultures of human hepatocytes. *Drug Metab Dispos.* 2010;38(4):591–599.
48. Kaukonen KM, Olkkola KT, Neuvonen PJ. Fluconazole but not itraconazole decreases the metabolism of losartan to E-3174. *Eur J Clin Pharmacol.* 1998;53(6):445–449.
49. Wedemeyer R-S, Blume H. Pharmacokinetic drug interaction profiles of proton pump inhibitors: an update. *Drug Saf.* 2014;37(4):201–211.
50. Hodgson E [webpage on the Internet]. *A Textbook of Modern Toxicology*. 4th ed. Wiley. Available from: <http://eu.wiley.com/Wiley-CDA/WileyTitle/productCd-047046206X.html>. Accessed March 27, 2018.
51. Bibi Z. Role of cytochrome P450 in drug interactions. *Nutr Metab.* 2008;5:27.
52. Wu X, Yuan L, Zuo J, Lv J, Guo T. The impact of CYP2D6 polymorphisms on the pharmacokinetics of codeine and its metabolites in Mongolian Chinese subjects. *Eur J Clin Pharmacol.* 2014;70(1): 57–63.
53. Preskorn SH, Shah R, Neff M, Golbeck AL, Choi J. The potential for clinically significant drug–drug interactions involving the CYP 2D6 system: effects with fluoxetine and paroxetine versus sertraline. *J Psychiatr Pract.* 2007;13(1):5–12.
54. O'Hara GE, Philippon F, Gilbert M, et al. Combined administration of quinidine and propafenone for atrial fibrillation: the CAQ-PAF study. *J Clin Pharmacol.* 2012;52(2):171–179.
55. Spina E, Santoro V, D'Arrigo C. Clinically relevant pharmacokinetic drug interactions with second-generation antidepressants: an update. *Clin Ther.* 2008;30(7):1206–1227.
56. Haertter S. Recent examples on the clinical relevance of the CYP2D6 polymorphism and endogenous functionality of CYP2D6. *Drug Metabol Drug Interact.* 2013;28(4):209–216.
57. Wadelius M, Darj E, Frenne G, Rane A. Induction of CYP2D6 in pregnancy. *Clin Pharmacol Ther.* 1997;62(4):400–407.
58. Rae JM, Johnson MD, Lippman ME, Flockhart DA. Rifampin is a selective, pleiotropic inducer of drug metabolism genes in human hepatocytes: studies with cDNA and oligonucleotide expression arrays. *J Pharmacol Exp Ther.* 2001;299(3):849–857.
59. Ayari I, Fedeli U, Saguem S, Hidar S, Khelifi S, Pavanello S. Role of CYP1A2 polymorphisms in breast cancer risk in women. *Mol Med Rep.* 2013;7(1):280–286.
60. Fontana RJ, Lown KS, Paine MF, et al. Effects of a chargrilled meat diet on expression of CYP3A, CYP1A, and P-glycoprotein levels in healthy volunteers. *Gastroenterology.* 1999;117(1):89–98.
61. Hakooz N, Hamdan I. Effects of dietary broccoli on human in vivo caffeine metabolism: a pilot study on a group of Jordanian volunteers. *Curr Drug Metab.* 2007;8(1):9–15.
62. Karjalainen MJ, Neuvonen PJ, Backman JT. In vitro inhibition of CYP1A2 by model inhibitors, anti-inflammatory analgesics and female sex steroids: predictability of in vivo interactions. *Basic Clin Pharmacol Toxicol.* 2008;103(2):157–165.
63. Ingelman-Sundberg M, Sim SC, Gomez A, Rodriguez-Antona C. Influence of cytochrome P450 polymorphisms on drug therapies: pharmacogenetic, pharmacoeconomic and clinical aspects. *Pharmacol Ther.* 2007;116(3):496–526.
64. Feghali M, Venkataramanan R, Caritis S. Pharmacokinetics of drugs in pregnancy. *Semin Perinatol.* 2015;39(7):512–519.
65. Ogu CC, Maxa JL. Drug interactions due to cytochrome P450. *Proc (Bayl Univ Med Cent).* 2000;13(4):421–423.
66. Fowler S, Zhang H. In vitro evaluation of reversible and irreversible cytochrome P450 inhibition: current status on methodologies and their utility for predicting drug–drug interactions. *AAPS J.* 2008;10(2): 410–424.
67. Prueksaritanont T, Chu X, Gibson C, et al. Drug–drug interaction studies: regulatory guidance and an industry perspective. *AAPS J.* 2013;15(3):629–645.
68. Michaut A, Moreau C, Robin M-A, Fromenty B. Acetaminophen-induced liver injury in obesity and nonalcoholic fatty liver disease. *Liver Int.* 2014;34(7):e171–e179.
69. Tompkins LM, Wallace AD. Mechanisms of cytochrome P450 induction. *J Biochem Mol Toxicol.* 2007;21(4):176–181.
70. Azzalis LA, Fonseca FLA, Simon KA, et al. Effects of ethanol on CYP2E1 levels and related oxidative stress using a standard balanced diet. *Drug Chem Toxicol.* 2012;35(3):324–329.
71. Brewer L, Williams D. Clinically relevant drug–drug and drug–food interactions. *Pharm Med.* 2013;27(1):9–23.
72. Ritter J, Lewis L, Mant T, Ferro A. *A Textbook of Clinical Pharmacology and Therapeutics*. 5th ed. Boca Raton, Florida: CRC Press; 2008.
73. Рекомендации от 05.02.2009 – regmed.ru [webpage on the Internet]. Available from: <http://www.regmed.ru/Content/Doc.aspx?id=26a9128c-ee32-4469-9c64-5c666339049e>. Accessed March 27, 2018.
74. Shin KH, Choi MH, Lim KS, Yu KS, Jang IJ, Cho JY. Evaluation of endogenous metabolic markers of hepatic CYP3A activity using metabolic profiling and midazolam clearance. *Clin Pharmacol Ther.* 2013;94(5):601–609.
75. Peng CC, Templeton I, Thummel KE, Davis C, Kunze KL, Isoherranen N. Evaluation of 6β-hydroxycortisol, 6β-hydroxycortisone, and a combination of the two as endogenous probes for inhibition of CYP3A4 in vivo. *Clin Pharmacol Ther.* 2011;89(6):888–895.
76. Jiang X-L, Shen H-W, Yu A-M. Pinoline may be used as a probe for CYP2D6 activity. *Drug Metab Dispos.* 2009;37(3):443–446.
77. Niioka T, Uno T, Sugimoto K, Sugawara K, Hayakari M, Tateishi T. Estimation of CYP2C19 activity by the omeprazole hydroxylation index at a single point in time after intravenous and oral administration. *Eur J Clin Pharmacol.* 2007;63(11):1031–1038.

78. Yasar U, Forslund-Bergengren C, Tybring G, et al. Pharmacokinetics of losartan and its metabolite E-3174 in relation to the CYP2C9 genotype. *Clin Pharmacol Ther.* 2002;71(1):89–98.
79. Stearns RA, Chakravarty PK, Chen R, Chiu SH. Biotransformation of losartan to its active carboxylic acid metabolite in human liver microsomes. Role of cytochrome P450C and 3A subfamily members. *Drug Metab Dispos.* 1995;23(2):207–215.
80. Yun CH, Lee HS, Lee H, Rho JK, Jeong HG, Guengerich FP. Oxidation of the angiotensin II receptor antagonist losartan (DuP 753) in human liver microsomes. Role of cytochrome P4503A(4) in formation of the active metabolite EXP3174. *Drug Metab Dispos.* 1995;23(2):285–289.
81. Yang S-H, Cho Y-A, Choi J-S. Effects of ticlopidine on pharmacokinetics of losartan and its main metabolite EXP-3174 in rats. *Acta Pharmacol Sin.* 2011;32(7):967–972.
82. de Andrés F, Sosa-Macías M, Llerena A. A rapid and simple LC-MS/MS method for the simultaneous evaluation of CYP1A2, CYP2C9, CYP2C19, CYP2D6 and CYP3A4 hydroxylation capacity. *Bioanalysis.* 2014;6(5):683–696.
83. Dorado P, Gallego A, Peñas-Lledó E, Terán E, Llerena A. Relationship between the CYP2C9 IVS8-109A>T polymorphism and high losartan hydroxylation in healthy Ecuadorian volunteers. *Pharmacogenomics.* 2014;15(11):1417–1421.
84. Sekino K, Kubota T, Okada Y, et al. Effect of the single CYP2C9*3 allele on pharmacokinetics and pharmacodynamics of losartan in healthy Japanese subjects. *Eur J Clin Pharmacol.* 2003;59(8–9):589–592.
85. Gray J. A textbook of clinical pharmacology and therapeutics, 5th edition. *Clin Pharmacol Ther.* 2009;85(2):125–126.

Drug Design, Development and Therapy

Publish your work in this journal

Drug Design, Development and Therapy is an international, peer-reviewed open-access journal that spans the spectrum of drug design and development through to clinical applications. Clinical outcomes, patient safety, and programs for the development and effective, safe, and sustained use of medicines are the features of the journal, which

Submit your manuscript here: <http://www.dovepress.com/drug-design-development-and-therapy-journal>

Dovepress

has also been accepted for indexing on PubMed Central. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.