Abortion history and its association with current use of modern contraceptive methods in Luanda, Angola

Natalie Morris
Ndola Prata

Bixby Center for Population, Health and Sustainability, School of Public Health, University of California, Berkeley, CA, USA

→ Video abstract

Point your Smartphone at the code above. If you have a QR code reader the video abstract will appear. Or use: <http://youtu.be/BQR3kGhZf-M>

Background: Women in sub-Saharan Africa often use abortion as a method of limiting their fertility and spacing births. However, it is not well understood whether having an abortion influences contraceptive behavior. The goal of this study was to examine associations between abortion history and use of a modern contraceptive method among women in Luanda, Angola.

Materials and methods: We analyzed data on 1,176 women aged 15–49 years from a 2012 cross-sectional study conducted in Luanda, Angola, which aimed to obtain general information on sexual and reproductive health indicators. The outcome and exposure were based on participant reports of past induced abortions and current use of a modern method. We used a modified Poisson regression with robust standard errors to estimate the relative risks of using a modern contraceptive method, given history of induced abortion.

Results: Among all respondents, 736 (62.6%) reported using a modern contraceptive method. The clear majority of the respondents (73.21%), regardless of abortion history, were using either no method, a traditional method, or condoms. Long-acting reversible contraceptive use was very low across all respondents (2.73%). The most common family planning method reported by women with a history of abortion was condoms (32.76%). Regression analysis demonstrated that women who had a history of abortion were 1.23 times more likely to use a modern contraceptive method as compared to those who never had an abortion (relative risk: 1.23; 95% CI: 1.10–1.36), after adjusting for potential confounders. Postregression estimations of predicted probabilities demonstrated that women with a history of abortion had an 80% probability of using a modern method (95% CI: 0.76–0.84), while those who never had an abortion had a 60% probability of using a modern method (95% CI: 0.59–0.61).

Conclusion: History of induced abortion was associated with use of a modern contraceptive method in our study population. The most common contraceptive used by women with a history of induced abortion was condoms, indicating that despite adoption of a modern method, many women are still at risk for an unintended pregnancy. Further research is needed to understand the causal factors underlying women's postabortion contraceptive choices.

Keywords: Angola, abortion, contraception, reproductive health

Background

Women are often motivated to use contraception after terminating a pregnancy in order to prevent future unwanted pregnancies and subsequent abortions.^{1,2} However, various socioeconomic factors and access to family planning (FP) ultimately influence women's contraceptive behavior, regardless of their history of induced abortion.^{3,4} Understanding the relationship between induced abortion and contraceptive use is an important first step in addressing contraceptive unmet need and gaps in access to FP services.

Correspondence: Ndola Prata
Bixby Center for Population, Health and Sustainability, School of Public Health, University of California, 421 University Hall, Berkeley, CA, USA
Tel +1 510 643 4284
Email ndola@berkeley.edu

Research on FP in sub-Saharan Africa (SSA) suggests that many women use abortion as their primary method of FP, sometimes instead of other contraceptive methods.⁵ When abortion policies are extremely restrictive and access to contraceptive methods is similarly limited, terminating pregnancies is often not a safe procedure; it is estimated that unsafe abortions account for 13% of maternal deaths in SSA.^{5,6} Furthermore, studies from Europe, Asia, and Africa have shown that a key predictor of induced abortion is having previously terminated a pregnancy, indicating women are likely not getting access to modern methods after their first abortion to prevent another unwanted pregnancy.^{7,8}

Other studies have demonstrated that having an abortion may, in fact, lead to increased use of contraception.² Post-abortion care (PAC) is intended to be legally available to all women, regardless of a country's abortion policies, and research on postabortion contraceptive behavior suggests that women are more likely to use a modern contraceptive method after an abortion if PAC services are offered.^{9,10} The mixed evidence from these studies indicates a need for additional research to better understand whether women who have induced abortions in places where it is not widely available are getting the services they need to prevent future unintended pregnancies and repeat abortions.

Despite some improvements in contraceptive uptake and reduction of unintended pregnancies over the past 15 years, contraceptive prevalence remains low in SSA, particularly in the country of Angola.¹¹ It is estimated that only about 13% of married women of reproductive age nationwide and 23% of married women in the capital province of Luanda use a modern contraceptive method, as compared to the regional SSA estimate of 26%.^{11,12} The countrywide unmet need for contraception is 38%, while urban centers in Angola have an unmet need of 26%.¹² It is imperative to understand the various factors that contribute to modern contraceptive uptake in order to reduce the high levels of unmet need.

There is no up-to-date data available on abortion in Angola, in part due to the fact that induced abortion is only permitted in instances when the woman's life is in danger.¹³ As a result, there have been no recent studies focusing on the contraceptive behaviors of Angolan women who have had induced abortions. Only one study conducted in 1996 examined this relationship, but primarily explored the influence of common social indicators on the use of each type of contraceptive method.³ More recent data from Luanda suggest that women in Angola are facing limited contraceptive choices, which may affect postabortion contraceptive uptake.¹⁴ Understanding the relationship between induced

abortion history and contraceptive use is imperative to meeting the high levels of unmet contraceptive need and improving the state of FP services in the country.

The aim of the present study was to investigate the relationship between history of induced abortion and current use of a modern contraceptive method among women in Luanda, Angola.

Materials and methods

Sampling and data collection

The data used for this analysis were obtained from a cross-sectional study conducted in Luanda, Angola in 2012. The University of California, Berkeley Bixby Center for Population, Health and Sustainability collaborated with Population Services International (PSI) to implement a baseline survey of sexual and reproductive health behaviors of women in the province of Luanda. At the time of the study, this province encompassed nine municipalities including the capital city of Angola. Two of the nine municipalities were rural communities, while the other seven were urban.

We utilized a multistage random sampling design to ensure that we captured a representative sample of women of reproductive age from all municipalities in Luanda province. The size of the entire population of Luanda was estimated to be 4,901,919 in 2012, according to the National Institute of Statistics.¹⁵ Power calculations determined the sample size estimates based on municipality population size. We first distributed the target sample size proportionally to the size of each municipality and then randomly selected a number of "sampling points" (churches, hospitals, gas stations, and so on) in each municipality from a list created for that purpose. The number of sampling points chosen per municipality varied according to the total population size in each of the municipalities. Eligible study participants included women aged 15–49 years who resided in Luanda between October and November 2012. A fixed number of participants were randomly selected for recruitment from each sampling point, and one woman from each household was interviewed in order to avoid overrepresentation of women in highly similar environments and life circumstances. In total, 1,825 women of reproductive age living in Luanda were randomly selected to participate in the survey between October and November of 2012. Of these 1,825 respondents, 85% completed the survey, 8% started but did not complete the survey, 6% refused to participate, and 2% did not participate due to other reasons.

The Bixby Center and PSI Angola partnered with *Sistemas de Informação Industriais e Consultoria*, a local marketing firm, to collect the questionnaire data. A multistage

random sampling design was implemented to capture a representative sample of women of reproductive age from the Luanda province. The data collection tool was developed by researchers at the Bixby Center and drew from the Women's Questionnaire of the Demographic and Health Surveys and Angola's Malaria Indicator Survey.¹⁶ The final tool also included additional standardized questions from the PSI Tracking Results Continuously survey tools.¹⁷ The questionnaire was designed to capture information on women's knowledge, attitudes, and practices related to childbearing and FP in the region. In addition, women were asked about their experiences and preferences with FP service provision and delivery. Sociodemographic and economic indicators were also recorded. All data were collected through self-report, and interviewers were instructed not to lead or guide the respondent while she answered the questionnaire. The data collection tool was initially written in English and translated to Portuguese by the research team, with feedback from PSI Angola. The research team conducted a pilot test of the questionnaire with 30 women of reproductive age in Luanda. Feedback from this initial round was incorporated into the final Portuguese version, which was later back translated into English. The ethical boards involved in this study approved the procedure of consent, which required all women who agreed to participate to provide verbal consent prior to the interview and for the interviewer to sign to confirm that consent was given. The ethics committees also waived parental consent for adolescent participants; respondents aged 15–17 were not required to obtain parental consent for the study and instead provided verbal assent to participate. Ethical approval for this study was provided by the University of California, Berkeley Committee for Protection of Human Subjects (CPHS # 2011-08-3521), as well as the Ethical Committee at the Instituto de Saude Publica in Luanda, Angola.

Based on our predetermined exclusion criteria, respondents were dropped from the analysis if they did not complete the survey ($n=280$) or were not at risk of the outcome, meaning they would not be using a contraceptive method. This included women who were not sexually active ($n=129$), were pregnant at the time of the study ($n=123$), were actively trying to get pregnant ($n=32$), or were infecund or sterilized ($n=69$). We excluded respondents who did not answer all questions for which variables were constructed for this analysis ($n=16$). It was possible to utilize complete case analysis because these excluded respondents accounted for <10% (1.35%) of the final 1,192 respondents who were eligible for the analysis. Women who reported wanting another pregnancy at any

point in the future were included the study since they may have been using a contraceptive method at the time they were interviewed to space their births. The final sample size used in the analysis was 1,176.

Variables

The dependent variable of interest was the type of contraceptive method the respondent reported using at the time of the study. Each participant reported using one of the following methods: intrauterine device (IUD), implant, injectable, pill, condom, traditional methods (rhythm, lactational amenorrhea method, or withdrawal), or no method. In cases where multiple contraceptive methods were listed, the participant was categorized into the most effective method she reported using. The final outcome variable was then collapsed and coded as a binary variable for modern methods or traditional/no method. Modern contraceptive methods were defined as contraceptive devices or medications used to prevent pregnancy, and thereby included the condom, pill, injectable, implant, and IUD.¹⁸ The primary independent variable, abortion history, collected self-reported information on each respondent's total lifetime number of induced abortions. The questionnaire asked the respondents who had ever been pregnant, "Have you or anyone else voluntarily interrupted any of your pregnancies?" and followed up with how many and in which year these induced abortions took place. The final variable was coded as "no abortions" and "one or more abortions". The primary analysis grouped women who reported having a single abortion with those who had multiple procedures due to the rarity of the exposure in our study sample (9.94%). Additional study variables were constructed and included in the fully adjusted model based on evidence from the literature indicating their role as potential confounders in the study relationship. Sociodemographic and individual factors included age, marital status, education, number of living children, and wealth. The final wealth variable was divided into quintiles and constructed with principal components analysis (PCA) using the standard PSI toolkit for creating wealth indices.¹⁹ The questionnaire also asked the respondents, "Do you believe contraceptives are accessible to you?", which was measured as a binary "yes" or "no" variable based on the respondent's perception of her access to contraceptive methods. Variables for contraceptive self-efficacy, knowledge of contraceptive methods, and community perceptions around FP were constructed using PCA following the PSI PCA toolkit guidelines.²⁰ For contraceptive self-efficacy, data were collected for a predetermined list of indicators that measured capability of using contraceptives, ability

to use methods correctly and consistently to space births, communicate about preventing pregnancy, and negotiate contraceptive use in the face of husband or partner opposition. Knowledge of contraceptive methods was measured through a predetermined set of indicators pertaining to respondents' knowledge of different types of contraceptive methods, including proper use and side effects. Respondents were asked to list all modern contraceptive methods they knew, how they used their own prescribed method, and common side effects of their prescribed method. Finally, the variable for community perception was created using 26 questions in a five-point Likert scale format to determine community acceptability of FP use. Respondents answered questions on whether they were able to discuss FP and contraceptive use with various community members, who included partners, siblings, parents, in-laws, religious leaders, and friends. Response options ranged from "completely disagree" to "completely agree", which were later recoded as "agree/disagree" binary variables for the PCA.

Statistical analysis

We examined descriptive statistics to explore dependent and independent variables of interest. Bivariate analysis using chi-square tests assessed differences across the proportions of women who used a modern contraceptive method by induced abortion history and by each covariate included in the adjusted model.

We used a Poisson regression with robust standard errors to approximate the relative risk (RR) of using a modern method among women who had at least one abortion compared to those who had no abortions. A logistic regression would likely have resulted in inflated measures of association

because the outcome was common in the study population (21.5%). A log-binomial model was initially used to fit the data; but due to a lack of convergence, we used a Poisson model with robust standard errors in order to estimate RR.²¹

Models were constructed a priori based on findings from the background literature and variables for which we had collected data. The final, fully adjusted model included all potential confounders for which we had collected information: age, marital status, access, knowledge, wealth, education, community perception, and contraceptive self-efficacy.

Postmodel estimations of predicted probabilities for abortion history were calculated to determine the probability of use of a modern method among women who had zero abortions and among those who had one or more. Statistical significance was set at a cutoff value of $p < 0.05$. All analyses were done in Stata 14.²²

Results

Figure 1 shows the frequency and percentage of induced abortions for each year that the respondents reported having an abortion. While some induced abortions took place over 20 years before the survey, two-thirds (66%) of all induced abortions occurred within the 5 years preceding the survey.

Table 1 presents all the independent variables by the contraceptive method each respondent reported using at the time of the survey. Of the 1,176 women included in the final analytical sample, only 116 (9.86%) women reported having had one or more induced abortions. Bivariate analysis demonstrated a statistically significant difference across one or more contraceptive categories for all covariates ($p < 0.05$).

The clear majority (73.21%) of all study participants, regardless of abortion history, were using either no method,

Figure 1 Frequency and percentage of induced abortion by year.

Table 1 Distribution of contraceptive method type by study covariates

Independent variables		Contraceptive method currently using												p-value			
		No method		Traditional		Condom		Pill		Injectable		Implant			IUD		Total
		n=428	%	n=21	%	n=412	%	n=130	%	n=153	%	n=21	%	n=11	%	N=1,176	%
Abortion	0	406	94.86	20	95.24	374	90.78	106	81.54	128	83.66	16	76.19	10	90.91	1,060	90.14
	1+	22	5.14	1	4.76	38	9.22	24	18.46	25	16.34	5	23.81	1	9.09	116	9.86
	Age (years)																
	15–19	175	40.89	3	14.29	131	31.80	9	6.92	5	3.27	1	4.76	0	0.00	324	27.55
	20–24	94	21.96	4	19.05	152	36.89	35	26.92	18	11.76	6	28.57	0	0.00	309	26.28
Marital status	25–34	105	24.53	8	38.10	102	24.76	61	46.92	66	43.14	5	23.81	5	45.45	352	29.93
	35+	54	12.62	6	28.57	27	6.55	25	19.23	64	41.83	9	42.86	6	54.55	191	16.24
	Single	321	75.00	12	57.14	333	80.83	68	52.31	39	25.49	5	23.81	5	45.45	783	66.58
	Married/cohabitating	88	20.60	9	42.86	70	16.99	55	42.31	101	66.01	15	71.43	4	36.36	342	29.08
	Divorced/widowed	19	4.44	0	0.00	9	2.18	7	5.38	13	8.50	1	4.76	2	18.18	51	4.34
Access	No	211	49.30	11	52.38	83	20.15	22	16.92	22	14.38	4	19.05	2	18.18	355	30.19
	Yes	217	50.70	10	47.62	329	79.85	108	83.08	131	85.62	17	80.95	9	81.82	821	69.81
	Contraceptive self-efficacy																
	1 (lowest self-efficacy)	130	30.37	8	38.10	124	30.10	31	23.85	27	17.65	4	19.05	2	18.18	326	27.72
	2	234	54.67	9	42.86	233	85.55	74	56.92	90	58.82	9	42.86	6	54.55	655	55.70
Knowledge of contraceptives	3 (highest self-efficacy)	64	14.95	4	19.05	55	13.35	25	19.23	36	23.53	8	38.10	3	27.27	195	16.58
	1 (least knowledgeable)	46	10.75	4	19.05	25	6.07	13	10.00	10	6.54	0	0.00	2	18.18	100	8.50
	2	223	52.10	9	42.86	184	44.66	47	36.15	64	41.83	3	14.29	3	27.27	533	45.32
	3 (most knowledgeable)	159	37.15	8	38.10	203	49.27	70	53.85	79	51.63	18	85.71	6	54.55	543	46.17
	Education																
Wealth quintiles	No education/primary school	213	49.77	8	25.00	122	29.61	51	39.23	86	56.21	8	38.10	3	27.27	491	41.75
	Secondary school	163	38.08	11	62.50	217	52.67	55	42.31	44	28.76	8	38.10	5	45.45	503	42.77
	University or higher	52	12.15	2	12.50	73	17.72	24	18.46	23	15.03	5	23.81	3	27.27	182	15.48
	1 (poorest)	95	22.22	5	23.81	53	12.86	35	26.92	31	20.26	4	19.05	3	27.27	226	19.22
	2	98	22.90	4	19.05	66	16.02	32	24.65	28	18.30	4	19.05	1	9.09	233	19.81
Community perceptions of FP use	3	81	18.93	6	28.57	88	21.36	22	16.92	34	22.22	6	28.57	1	9.09	238	20.24
	4	82	19.16	5	23.81	93	22.57	18	13.85	32	20.92	5	23.81	2	18.18	237	20.15
	5 (richest)	72	16.82	1	4.76	112	27.18	23	17.69	28	18.30	2	9.52	4	36.36	242	20.58
	1 (least acceptable)	83	19.39	1	4.76	40	9.71	22	16.92	10	6.54	2	9.52	2	18.18	160	13.61
	2	301	70.33	15	71.43	316	76.70	84	64.62	103	67.32	10	47.62	7	63.64	836	71.09
Number of living children	3 (most acceptable)	44	10.28	5	23.81	56	13.59	24	18.46	40	26.14	9	42.86	2	18.18	180	15.31
	0	5	1.17	0	0.00	1	0.24	0	0.00	0	0.00	0	0.00	0	0.00	6	0.51
	1–2	122	28.50	7	33.33	108	26.21	74	56.92	80	52.29	13	61.90	5	45.45	409	34.78
	3+	301	70.33	14	66.67	303	73.54	56	43.08	73	47.71	8	38.10	6	54.55	761	64.71

Abbreviations: FP, family planning; IUD, intrauterine device.

a traditional method, or condoms. The results displayed in Table 2 demonstrate that the largest proportion (38.30%) of women who had no abortions reported use of no methods, as compared to women who had one or more induced abortions (18.97%). The method most commonly used by women who had one or more induced abortions was condoms (32.76%). Among the 116 women who had an induced abortion, only 6 (5.17%) were using an implant or IUD, which collectively represented the long-acting reversible contraceptives (LARC). In comparison, a smaller proportion of women who had no induced abortions used LARC (2.47%).

We assessed the bivariate results of history of induced abortion by study covariates. Based on the results presented in Table 3, we found that age, perceived access to contraceptives, contraceptive self-efficacy, and the number of living children all had statistically significant differences between no induced abortion and one or more induced abortions at $p < 0.05$. Chi-square analysis results in Table 4 demonstrate a statistically significant difference in use of a modern contraceptive method for abortion history ($p < 0.001$). There were also significant differences ($p < 0.05$) between use of a modern method and all covariates measured, except for contraceptive self-efficacy ($p = 0.210$).

Table 5 displays results from the Poisson regression with robust standard errors, which demonstrate that women who had one or more abortions were 1.23 times more likely to use a modern contraceptive method as compared to those who never had an abortion (95% CI: 1.10–1.36, $p < 0.001$). Women who were older, reported having access to contraceptives, and had high community acceptability of FP use were also more likely to use a modern contraceptive method. The regression results also demonstrated that all categories of older age were significantly associated with using a modern method, as was having access to contraceptives (RR: 1.80, $p < 0.001$) and high community acceptance of FP use (RR: 1.26, $p = 0.018$). Having a higher education was not significantly associated with use of a modern method, though having completed secondary school did result in a slightly higher likelihood of using a modern method (RR: 1.11, $p = 0.046$). There was no

statistically significant increase in the likelihood of using a modern method with having ever been married, having higher knowledge of contraceptives, possessing increased wealth, or having more living children ($p > 0.05$).

We then calculated the fully adjusted postregression predicted probabilities of using a modern contraceptive method for women who had no abortions and for those who had one or more abortions. Table 6 demonstrates that women who had never had an abortion had a 60% predicted probability of using a modern method (95% CI: 0.59–0.61), while those who had one or more abortions had an 80% predicted probability of using a modern method (95% CI: 0.76–0.84). The lack of overlap between the 95% CI of the two exposure groups indicates a significant difference in use of modern methods; as demonstrated in the regression results, women who had one or more abortions were more likely to use a modern method.

Discussion and conclusion

This study found that among women in Luanda, Angola, having had one or more induced abortions was significantly associated with current use of a modern contraceptive method. Various factors may explain this relationship, including greater exposure to modern methods and FP services among women who had induced abortions.

Prior studies conducted in various other settings have reported similar findings and potential reasons for this association. A systematic review of nine publications on low- and middle-income countries found that contraception uptake increased postabortion when there was access to a wide range of contraceptive methods and comprehensive sexual and reproductive health education.¹⁰ A study by Benson et al also found that offering a wide range of contraceptive methods at health clinics improved postabortion modern contraceptive uptake.²³ Though we cannot make inferences on whether the subjects of our study were seen for PAC, this particular element of FP service provision has been shown to improve uptake of modern methods.²⁴ A prospective intervention study in Zimbabwe also demonstrated an increased

Table 2 Percentage breakdown of type of contraceptive method by abortion history

Contraceptive method currently using	No method		Traditional		Condom		Pill		Injectable		Implant		IUD		Total	p-value
	n	%	n	%	n	%	n	%	n	%	n	%	n	%		
Abortion																<0.001
0	406	38.30	20	1.89	374	35.28	106	10.00	128	12.08	16	1.51	10	0.94	1,060	
1+	22	18.97	1	0.86	38	32.76	24	20.69	25	21.55	5	4.31	1	0.86	116	
Total	428	36.39	21	1.79	412	35.03	130	11.05	153	13.01	21	1.79	11	0.94	1,176	

Abbreviation: IUD, intrauterine device.

Table 3 History of induced abortion by study covariates

Covariates	Abortion						p-value
	0		1+		Total		
	n=1,060	%	n=116	%	N=1,176	%	
Age (years)							<0.001
15–19	308	29.06	16	13.79	324	27.55	
20–24	282	26.60	27	23.28	309	26.28	
25–34	298	28.11	54	46.55	352	29.93	
35+	172	16.23	19	16.38	191	16.24	
Marital status							0.064
Single	717	67.64	66	56.90	783	66.58	
Married/cohabitating	299	28.21	43	37.07	342	29.08	
Divorced/widowed	44	4.15	7	6.03	51	4.34	
Access							0.001
No	335	31.60	20	17.24	355	30.19	
Yes	725	68.40	96	82.76	801	69.81	
Contraceptive self-efficacy							0.007
1 (lowest self-efficacy)	307	28.96	19	16.38	326	27.72	
2	585	55.19	70	60.34	655	55.70	
3 (highest self-efficacy)	168	15.85	27	23.28	195	16.58	
Knowledge of contraceptives							0.721
1 (least knowledgeable)	88	8.30	12	10.34	100	8.50	
2	480	45.28	53	45.69	533	45.32	
3 (most knowledgeable)	492	46.42	51	43.97	543	45.17	
Education							0.172
No education/primary school	439	41.42	52	44.83	491	41.75	
Secondary school	462	43.58	41	35.34	503	42.77	
University or higher	159	15.00	23	19.83	182	15.48	
Wealth quintiles							0.626
1 (poorest)	201	18.96	25	21.55	226	19.22	
2	205	19.34	28	24.14	233	19.81	
3	216	20.38	22	18.97	238	20.24	
4	217	20.47	20	17.24	237	20.15	
5 (richest)	221	20.85	21	18.10	242	20.58	
Community perceptions of FP use							0.429
1 (least acceptable)	144	13.58	16	13.79	160	13.61	
2	749	70.66	87	75.00	836	71.09	
3 (most acceptable)	167	15.75	13	11.21	180	15.31	
Number of living children							0.001
0	6	0.57	0	0.00	6	0.51	
1–2	351	33.11	58	50.00	409	34.78	
3+	703	66.32	58	50.00	761	64.71	

Abbreviation: FP, family planning.

uptake of modern methods and significantly fewer unplanned pregnancies when PAC FP was offered.⁹ PAC services may have been available and accessible to the respondents in this study, which led to the higher modern method use observed among women who had an abortion.

There was a lack of significant association between wealth and modern contraceptive use, which is inconsistent with what is known about wealth inequities and fertility practices. Higher socioeconomic status has been well documented as a predictor of contraceptive use in SSA.^{25,26} However, a recent study by Creanga et al found that wealth-related

inequalities in contraceptive use have decreased in many countries in SSA, and reported that higher wealth status was associated specifically with long-acting contraceptive use.²⁷ The discrepancy between our findings and the generally accepted knowledge surrounding socioeconomic status and contraceptive use may be explained by this diminishing gap, as many modern methods, namely, condoms, are becoming more widely available in the region. Furthermore, our models included abortion as an independent variable, which may be a stronger indicator of contraceptive use in similar populations; its presence may have diminished expected associations

Table 4 Use of a modern contraceptive method by study covariates

Independent variables	Abortion						p-value
	No		Yes		Total		
	n=440	%	n=736	%	N=1,176	%	
Abortion							<0.001
0	417	94.77	643	87.36	1,060	90.14	
1+	23	5.23	93	12.64	116	9.87	
Age (years)							<0.001
15–19	178	40.45	146	19.84	324	27.55	
20–24	96	21.82	213	28.94	309	26.28	
25–34	110	25.00	242	32.88	352	29.93	
35+	56	12.73	135	18.34	191	16.24	
Marital status							<0.001
Single	328	74.55	455	61.82	783	66.58	
Married/cohabitating	93	21.14	249	33.83	342	29.08	
Divorced/widowed	19	4.32	32	4.35	51	4.34	
Access							<0.001
No	217	49.32	138	18.75	355	30.19	
Yes	223	50.68	598	81.25	821	69.81	
Contraceptive self-efficacy							0.210
1 (lowest self-efficacy)	134	30.45	192	26.09	326	27.72	
2	240	54.55	415	56.39	655	55.70	
3 (highest self-efficacy)	66	15.00	129	17.53	195	16.58	
Knowledge of contraceptives							<0.001
1 (least knowledgeable)	47	10.68	53	7.20	100	8.50	
2	229	52.05	304	41.30	533	45.32	
3 (most knowledgeable)	164	37.27	379	51.49	543	46.17	
Education							<0.001
No education/primary school	216	49.09	275	37.36	491	41.75	
Secondary school	171	38.86	332	45.11	503	42.77	
University or higher	53	12.05	129	17.53	182	15.48	
Wealth quintiles							0.010
1 (poorest)	98	22.27	128	17.39	226	19.22	
2	101	22.95	132	17.93	233	19.81	
3	85	19.32	153	20.79	238	20.24	
4	83	18.86	154	20.92	237	20.15	
5 (richest)	73	16.59	169	22.96	242	20.58	
Community perceptions of FP use							<0.001
1 (least acceptable)	83	18.86	77	10.46	160	13.61	
2	310	70.45	526	71.47	836	71.09	
3 (most acceptable)	47	10.68	133	18.07	180	15.31	
Number of living children							0.001
0	5	1.14	1	0.14	6	0.51	
1–2	128	29.09	281	38.18	409	34.78	
3+	307	69.77	454	61.68	761	64.71	

Abbreviation: FP, family planning.

between modern method use and other factors, including wealth, education, and marital status. Future studies should, therefore, consider incorporating questions about abortion history when possible in order to provide further data on its role in contraceptive uptake.

It is important to note that the most commonly selected modern method among women in our study sample who had one or more abortions was condoms. Though condoms are

classified as a modern method and can be highly efficacious when used correctly, they have high failure rates and thereby a low effectiveness (up to 18 pregnancies per 100 women per year).²⁸ It is, therefore, difficult to conclude that women in our study who were using a modern method were using one that would effectively prevent another unwanted pregnancy, particularly since condoms were the most highly used. In addition, our results demonstrated that women who had no

Table 5 Relative risk of using a modern contraceptive method by abortion history

Risk of using a modern method, given abortion history	Use of a modern method		
	RR	95% CI	p-Value
Abortion			
0	1 (ref)	–	–
1+	1.23	1.10–1.36	<0.001
Age (years)			
15–19	1 (ref)	–	–
20–24	1.38	1.20–1.59	<0.001
25–34	1.31	1.12–1.52	0.001
35+	1.32	1.12–1.57	0.001
Marital status			
Never married	1 (ref)	–	–
Ever married	1.04	0.94–1.15	0.467
Access			
No	1 (ref)	–	–
Yes	1.80	1.56–2.06	<0.001
Knowledge			
1 (least knowledgeable)	1 (ref)	–	–
2	1.01	0.82–1.23	0.956
3 (most knowledgeable)	1.15	0.94–1.40	0.178
Education			
No education/primary school	1 (ref)	–	–
Secondary school	1.11	1.00–1.23	0.046
University or higher	1.07	0.94–1.22	0.323
Wealth quintiles			
1 (poorest)	1 (ref)	–	–
2	0.94	0.81–1.09	0.428
3	1.06	0.92–1.22	0.451
4	1.02	0.88–1.18	0.776
5 (richest)	1.06	0.92–1.22	0.446
Community perceptions of FP use			
1 (least acceptable)	1 (ref)	–	–
2	1.12	0.95–1.33	0.180
3 (most acceptable)	1.26	1.04–1.52	0.018
Number of living children			
0	1 (ref)	–	–
1–2	3.52	0.59–20.92	0.167
3+	3.58	0.60–21.32	0.162

Abbreviations: FP, family planning; RR, relative risk.

abortions most commonly reported using no method, despite responding that they were sexually active and not actively trying to get pregnant. Several factors could influence women's strong preference for condoms among those who had an induced abortion and for lack of method use among those who did not. Misinformation or skepticism about certain methods can impede uptake of more effective contraceptives, namely, LARC.^{29,30} Furthermore, condoms are currently the most widely available FP method in Luanda, followed by the oral pill and injectable methods.¹⁴ FP programs should focus on improving provision of LARC methods, particularly in the context of PAC services. Angola recently implemented

Table 6 Predicted probabilities of using a modern contraceptive method by abortion history

Currently using a modern method		
Abortion	Adjusted ^a predicted probability	95% CI
0	0.60	0.59–0.61
1+	0.80	0.76–0.84

Note: ^aAge, marital status, access, knowledge, education, wealth, community perceptions, number of living children.

a PAC program that has improved postabortion contraceptive uptake, but the results of our study indicate a high need for supplies and training to offer methods postabortion that would be more effective and long term than condoms.³¹

Further research is needed to understand to what extent abortion influences contraceptive uptake in SSA, as well as how to tailor interventions to offer a wide variety of modern methods to populations that have limited access to FP care services and want to prevent unwanted pregnancies. This analysis provides insight into the relationship between past induced abortion and current contraceptive use, but prospective studies examining contraceptive use prior to and after an abortion would be more informative in understanding how abortion could directly influence women's contraceptive practices.^{32,33} These types of studies could provide information on whether women currently using modern methods changed their contraceptive method after terminating a pregnancy, and why they are selecting less effective modern contraceptives as opposed to LARC methods. Studies with larger sample sizes could potentially assess the association of past induced abortion with use of specific types of contraceptive methods or examine differences among women who had multiple abortions. This study sample did not have adequately large cell sizes (<10 observations) to be able to make statistical inferences on these particular assessments. Ultimately, studies that are designed to assess the causal factors underlying the relationship between abortion and contraceptive use are needed for increasing uptake of modern methods in Angola.

A major limitation of our study is its cross-sectional design, which does not allow us to establish causality. It is unknown when in time the respondent began using her current contraceptive method, and therefore, whether she began using it after terminating her pregnancy or used it both before the induced abortion and continued using it afterward. Though modern methods prevent pregnancies at a greater rate than traditional methods, they may still fail and lead to an unintended pregnancy. We do not know if the respondent had an

abortion as a result of method failure or because she was not using any contraceptive method. This does not allow us to establish temporality between the primary exposure and the outcome of interest and prevents us from being able to make causal inferences. However, this is likely not a major limitation due to the fact that in SSA, a vast majority of unintended pregnancies and subsequent induced abortions occur when a woman is not using any contraceptive method.³⁴ Other limitations and potential sources of bias included that the study population consisted of women in the province of the capital city of Angola, which may not be generalizable to the rural or other urban areas of the country. In addition, there may have been information bias due to misclassification of study participants who reported using multiple contraceptive methods. Respondents were grouped into “no method/traditional” and “modern” contraceptive methods based on the most effective method they were using at the time. This may not be reflective of the method they used most frequently, resulting in women who reported using both types of contraceptives being more frequently misclassified as using a modern method. This could lead to non-differential misclassification of the outcome, as it would have occurred independently of the exposure, and result in bias of the measure of association toward the null.

This study also had several strengths. There are, to date, no comprehensive reports containing information about abortion in the country of Angola. The Demographic and Health Survey for Angola includes information about FP and fertility but does not include nationwide abortion facts or statistics.¹² It is difficult to get accurate abortion information because many abortions in SSA are performed illegally and women usually underreport abortion.³⁵ The fact that our study was the first to collect data on abortion in Angola might indicate there was social desirability bias in our study, as some women may have been uncomfortable admitting they had terminated a pregnancy, either lawfully or otherwise. However, it is unlikely that social desirability bias influenced the reporting of modern method use since condoms and other modern contraceptives are not legally restricted in Angola.¹⁴ This would indicate that our observed association would not decrease if there were actually more induced abortions in our sample than reported, as the numbers would have presumably been distributed equally across the outcome variable categories.

Results from our study provide basic insight into Angolan women’s contraceptive behavior and its relationship with the history of induced abortion. While some of the respondents’ abortions occurred many years prior to the survey, the fact

that 66% of reported abortions occurred within 5 years of the survey taking place indicates that the observed association was not solely based on the known improvements to access to contraceptives over the past two decades in Angola.³⁶ These results contribute to the growing body of evidence on abortion and contraceptive use in a country with so little information on the topic. Data from this analysis can inform future studies that examine what aspects of abortion might influence use of modern contraceptives in Angola, with the intention of implementing these practices to increase uptake of more highly effective contraceptive methods.

Acknowledgments

We would like to thank Karen Weidert, MPH, for her assistance with language editing and proofreading. We would also like to thank Dr Mahasin Mujahid, PhD, for her guidance and support during prior iterations of the analyses.

Disclosure

The authors report no conflicts of interest in this work.

References

1. Padmadas SS, Lyons-Amos M, Thapa S. Contraceptive behavior among women after abortion in Nepal. *Int J Gynecol Obstet*. 2017;127:132–137.
2. Sedlecky K, Stanković Z. Contraception for adolescents after abortion. *Eur J Contracept Reprod Health Care*. 2016;21:4–14.
3. Carvalho A, Laudari C, Marini M, Faundes A. Characteristics of contraceptive acceptors in Luanda, Angola. *Afr J Fertil Sexual Reprod Heal*. 1996;1:109–114.
4. Prata N, Bell S, Weidert K, Nieto-Andrade B, Carvalho A, Neves I. Varying family planning strategies across age categories: differences in factors associated with current modern contraceptive use among youth and adult women in Luanda, Angola. *Open Access J Contracept*. 2016;7:1–9.
5. Lauro D. Abortion and contraceptive use in sub-Saharan Africa: how women plan their families. *Afr J Reprod Health*. 2011;15:13–23.
6. Cohen SA. Access to safe abortion in developing world: saving lives while advancing rights. 2012.
7. Benson J, Andersen K, Samandari G. Reductions in abortion-related mortality following policy reform: evidence from Romania, South Africa and Bangladesh. *Reprod Health* 2011;8:39.
8. Heikinheimo O, Gissler M, Suhonen S. Age, parity, history of abortion and contraceptive choices affect the risk of repeat abortion. *Contraception*. 2008;78:149–154.
9. Johnson BR, Ndhlovu S, Farr SL, Chipato T. Reducing unplanned pregnancy and abortion in Zimbabwe through postabortion contraception. *Stud Fam Plann*. 2002;33:195–202.
10. Rogers C, Dantas JAR. Access to contraception and sexual and reproductive health information post-abortion: a systematic review of literature from low- and middle-income countries. *J Fam Plann Reprod Health Care*. 2017;43(4):309–318.
11. Sharan M, Ahmed S, May J, Soucat A. Family planning trends in Sub-Saharan Africa: progress, prospects, and lessons learned. *Yes Africa Can*. 445, 2011.
12. *Key Findings of the 2015-16 Angola IIMS*. (Instituto Nacional de Estatística (INE), Ministério da Saúde (MINSa), Ministério da Planeamento e do Desenvolvimento Territorial (MPDT) and ICF, 2017).
13. Masangkay, N. *Abortion Policies: Afghanistan to France*. United Nations Publications, 2001.

14. Nieto-Andrade B, Fidel E, Simmons R, et al. Women's limited choice and availability of modern contraception at retail outlets and public-sector facilities in Luanda, Angola, 2012–2015. *Glob Health Sci Pract*. 2017;5:75–89.
15. National Statistical Office – Angola [webpage on the Internet]. Available from: <http://www.nber.org/africa/display/1095>. Accessed April 13, 2018.
16. ICF International. *Demographic and Health Surveys Methodology – Questionnaires: Household, Woman's, and Man's*. CA: ICF International; 2011.
17. Population Services International (2003). Project TRaC: Tracking Results Continuously. Retrieved September 15, 2017.
18. Hubacher D, Trussell J. A definition of modern contraceptive methods. *Contraception*. 2015;92:420–421.
19. Wealth-Quintile-Guide.pdf.
20. TRaC-Dashboard-Toolkit.pdf.
21. Zou G. A modified poisson regression approach to prospective studies with binary data. *Am J Epidemiol*. 2004;159:702–706.
22. StataCorp. Stata Statistical Software: Release 14. College Station, TX: StataCorp LP; 2015.
23. Benson J, Andersen K, Brahmi D, et al. What contraception do women use after abortion? An analysis of 319,385 cases from eight countries. *Glob Public Health*. 2016;1–16.
24. Gallagher M, Chukmuwalu K, Baunach S. Uptake of Post-abortion Care (PAC) services and acceptance of post-abortion contraception in remote areas of Puntland, Somalia. *Ann Glob Health*. 2017;83:97.
25. Dias JG, de Oliveira IT. Multilevel effects of wealth on women's contraceptive use in Mozambique. *PLoS One*. 2015;10(3):e0121758.
26. WHO | Low Use of Contraception Among Poor Women in Africa: An Equity Issue [webpage on the Internet]. *WHO* Available from: <http://www.who.int/bulletin/volumes/89/4/10-083329/en/>. Accessed April 13, 2018.
27. Creanga AA, Gillespie D, Karklins S, Tsui AO. Low use of contraception among poor women in Africa: an equity issue. *Bull World Health Organ*. 2011;89:258–266.
28. Guttmacher Institute (2016) [homepage on the Internet]. Contraceptive Use in the United States. Available from: <https://www.gutt.org>. Accessed September 15, 2017.
29. Kakaire O, Nakiggude J, Lule JC, Byamugisha JK. Post abortion women's perceptions of utilizing long acting reversible contraceptive methods in Uganda. A qualitative study. *Open J Obstet Gynecol*. 2014;04:1087.
30. Williamson LM, Parkes A, Wight D, Petticrew M, Hart GJ. Limits to modern contraceptive use among young women in developing countries: a systematic review of qualitative research. *Reprod Health*. 2009;6:3.
31. Bell S, Weidert K, Fraser A, Prata N. *Health Facility Assessment Report: Cacucaco and Viana Municipalities*. UC Berkeley Bixby Center: Angola Comprehensive Family Planning and Post-Abortion Care Project. 2013.
32. Borges ALV, OlaOlorun F, Fujimori E, Hoga LAK, Tsui AO. Contraceptive use following spontaneous and induced abortion and its association with family planning services in primary health care: results from a Brazilian longitudinal study. *Reprod Health*. 2015;12:94.
33. Prata N, Bell S, Holston M, Gerds C, Melkamu Y. Factors associated with choice of post-abortion contraception in Addis Ababa, Ethiopia. *Afr J Reprod Health*. 2011;15:51–57.
34. Collumbien M, Gerressu M, Cleland J. Non-use and use of ineffective methods of contraception. 2004.
35. Induced Abortion Worldwide [webpage on the Internet]. *Guttmacher Institute* (2016). Available from: <https://www.guttmacher.org/fact-sheet/induced-abortion-worldwide>. Accessed March 18, 2017.
36. Decker M, Constantine NA. Factors associated with contraceptive use in Angola. *Afr J Reprod Health*. 2011;15:68–77.

Open Access Journal of Contraception

Publish your work in this journal

Open Access Journal of Contraception is an international, peer-reviewed, open access, online journal, publishing original research, reports, reviews and commentaries on all areas of contraception. In addition to clinical research, demographics and health-related aspects, the journal welcomes new findings in animal and preclinical studies

Submit your manuscript here: <https://www.dovepress.com/open-access-journal-of-contraception-journal>

relating to understanding the biological mechanisms and practical development of new contraceptive agents. The manuscript management system is completely online and includes a very quick and fair peer-review system. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Dovepress