Vitiligo and social acceptance

Reza Bidaki¹
Niloofar Majidi²
Amir Moghadam Ahmadi³
Hamid Bakhshi⁴
Rezvan Sadr Mohammadi⁵
Seyed-Ali Mostafavi⁶
Mohammad Kazemi
Arababadi⁷
Maryam Hadavi⁸
Afshin Mirzaei⁹

¹Research Center of Addiction and Behavioral Sciences, Diabetes Research Center, Shahid Sadoughi University of Medical Sciences, Yazd, Iran; ²Department of Psychiatry, Rafsanjan University of Medical Sciences, Rafsanjan, Iran; ³Department of Neurology, Rafsanjan University of Medical Sciences, Rafsanjan, Iran; ⁴Continuing Medical Education Center, Rafsanjan University of Medical Sciences, Rafsanjan, Iran; ⁵Department of Psychiatry, Lorestan University of Medical Sciences, Khorramabad, Iran; ⁶Psychiatry Research Center, Roozbeh Hospital, Tehran University of Medical Sciences, Tehran, Iran; ⁷Department of Laboratory Sciences, Faculty of Paramedicine, Rafsanjan University of Medical Sciences, Rafsanjan, Iran; ⁸Department of Anesthesiology, Paramedical Faculty, Molecular Medicine Research Center, Rafsanjan University of Medical Sciences, Rafsanjan, Iran; ⁹Rafsanjan University of Medical Sciences, Rafsanjan, Iran

Correspondence: Maryam Hadavi
Department of Anesthesiology, Paramedical
Faculty, Molecular Medicine Research
Center, Rafsanjan University of Medical
Sciences, Rafsanjan, 7718796755, Iran
Tel +98 34 3425 8397
Email hadavimaryam@yahoo.com

Background and aim: Vitiligo is a chronic skin disease characterized by a total or partial loss of melanocytes from the epidermis and other tissues of the skin. It is placed in the class of secondary psychiatric disorders and can also lead to psychological problems. The main aim of this study was to assess social acceptance in vitiligo patients.

Methods: This cross-sectional study was conducted on all of the patients (n=150) with vitiligo who were referred to dermatology clinics in Rafsanjan, Iran. The patients completed a social acceptability questionnaire (Marlowe–Crowne Social Desirability Scale), and information regarding their demographic characteristics was also collected. Data were gathered and analyzed with descriptive and inferential statistics using SPSS-19 software.

Results: The mean age of the patients was 27.56 ± 10.53 years and 65.9% were female. Mean score of social acceptance among the patients was 13.51 ± 7.08 . The results showed that the mean scores of social acceptance were significantly lower in women, in those with single marital status, in those with face and neck lesions, and in those with disease duration less than 5 years.

Conclusion: The results showed that certain groups of patients with vitiligo are at greater risk of experiencing lower social acceptance.

Keywords: vitiligo, social acceptance, skin disease, secondary psychiatric disorders

Introduction

Vitiligo is a common pigmentary skin disorder, characterized by a total or partial loss of melanocytes from the epidermis of skin. The prevalence of vitiligo is 0.1%–4% and is estimated to be about 1% in white individuals.¹ The incidence of vitiligo is similar between men and women, and half of the cases occur before the age of 20 years.²

As a psycho-dermatology disease, vitiligo is one of the aspects of the new series called “secondary psychiatric disorders”.³ Vitiligo does not impose serious health problems for the patients, but causes serious beauty problems; meanwhile, most doctors do not assess mood and quality of life in these patients.

The psychological burden of this dermatological problem is relatively high.⁴ Researchers report on the burden of vitiligo that ~75% of those concerned estimate their appearance as moderately to severely intolerable.⁵ Vitiligo reduces the patients’ beauty and has deep effects on their private and social life, and causes social malfunction and decreases their quality of life.⁶ This illness lowers their self-esteem and may cause shame in social interactions.⁷ The patients may experience emotional stress, particularly if vitiligo develops on visible areas of the body like face. Some feel embarrassed, shameful (with decreased self-confidence), depressed, or worried about how others will react.^{7,8} A study showed that more than half of the vitiligo patients

declared that people stare at them, 20% said that they are labeled, and 25% said that their disease causes problems in their relationships with strangers.⁹

Social acceptance of chronic diseases such as vitiligo will be encouraged by social network, including family, friends, colleagues, and neighbors.³ Social acceptance is a vital aspect in the life of patients with chronic diseases and may be associated with an improved prognosis.¹⁰ Furthermore, some studies have mentioned the importance of social acceptance in the quality of life of the patients,¹¹ while some patients complain that society does not accept them because of their appearance.¹² It seems that these patients are not properly supported by their families or society.¹²

Based on the fact that social acceptance in the patients is dependent on social culture, the aim of this study was to determine social acceptance of vitiligo patients in Rafsanjan, a city in Iran.

Methods

This cross-sectional study was done on the vitiligo patients (n=150) referred to the dermatology clinics in Rafsanjan. Among the 150 patients referred to the dermatology clinic, 126 patients responded to the questionnaire; the response rate was 84%. The inclusion criteria were the patients should be suffering from vitiligo and referred to the dermatology clinic and have the ability to understand the concept of questionnaires. Exclusion criterion was unwillingness to participate in the research.

Data gathering tools

Marlowe–Crowne Social Desirability Scale (MC-SDS) for assessing social acceptance

MC-SDS questionnaire is one of the most widely used scales for detecting social desirability. MC-SDS consists of 33 items with 2 options for each item, yes (1) or no (0). Total score, which is the sum of all scores, was calculated and ranges from 0 to 33; scores between 0 and 8, 9 and 19, and 20 and 33 indicate low, moderate, and high levels of social desirability, respectively.

Marlowe–Crowne Social Desirability scale achieves an acceptable internal consistency coefficient of 0.88 and has reported 0.89 for reliability of retest.¹³

Inventory of demographic characteristics

This inventory included demographic characteristics such as gender, marital status, education, duration of illness, family history, and location of the lesion.

Statistical analysis

Descriptive statistics are presented as frequency and percentage or mean $\pm$ standard deviation (SD). The data were analyzed with independent *t*-test and ANOVA. *P*-values less than 0.05 were considered significant. Data analyses were performed using SPSS-19 software.

Ethics

All participants provided written informed consent. In the case of subjects under the age of 18 years, confirmation of written informed consent from the parents or legal guardians was also obtained. The study was approved by the ethics committee board of Rafsanjan University of Medical Sciences.

Results

This cross-sectional study was conducted on 126 patients with vitiligo (43 men and 83 women). The mean and SD age of the patients was 27.56 ± 10.53 years (range 13–60 years), and the average duration of the disease was 7.27 ± 7.76 years. In terms of educational status, most patients had diploma (56.3%, N=71) and primary education (21.4%, N=27). In terms of family history, the majority of patients had no positive family history (88.1%) for vitiligo. About 70.6% (89 patients) had diffuse lesions, 10.3% (13 patients) had lesions on the extremities, and 7.1% (9 patients) had lesions on their face. Other demographic characteristics of participants are presented in Table 1.

The mean score of social acceptance in the patients was 13.51 ± 7.08 . The average scores for social acceptance with regard to different subgroups of gender, marital status, educational level, location of lesion, family history, and time since the occurrence of the disease are displayed in Table 2.

Table 1 Characteristics of participants in the study of vitiligo and social acceptance

Characteristics	N	%
Gender		
Men	43	34.1
Women	83	65.9
Marital status		
Married	56	44.4
Single	60	47.6
Divorced	10	8
District		
City	93	74.6
Village	23	19
Countryside	10	6.4

Table 2 Maximum, minimum, mean, and standard deviation of social acceptance score based on demographic variables

Variables	Social acceptance (mean $\pm$ SD)	Minimum	Maximum	P-value
Age (years)				
13–22	8.65 $\pm$ 6.70	1	33	<0.001**
23–32	13.86 $\pm$ 5.05	7	29	
33–42	16.91 $\pm$ 6.37	6	30	
43–52	20.25 $\pm$ 4.57	15	25	
>53	24.60 $\pm$ 6.88	18	33	
Gender				
Men	15.27 $\pm$ 7.68	1	33	0.044*
Women	12.60 $\pm$ 6.62	1	33	
Marital status				
Single	9.86 $\pm$ 5.81	1	33	0.001**
Married	17.58 $\pm$ 6.51	1	33	
Divorced	12.60 $\pm$ 4.88	7	22	
Education level				
Primary	12.59 $\pm$ 10.63	1	33	0.850
Higher	14.75 $\pm$ 8.39	6	30	
Diploma	13.39 $\pm$ 5.99	1	30	
BSc	14.57 $\pm$ 3.77	9	23	
MSc or higher	17	17	17	
Location of lesion				
Face	7.66 $\pm$ 4.30	1	13	<0.001**
Neck	7.85 $\pm$ 5.01	1	13	
Extremities	18.38 $\pm$ 10.09	1	33	
Trunk	17.87 $\pm$ 8.80	5	33	
Diffuse	13.44 $\pm$ 6.04	7	31	
Family history				
Positive	15.46 $\pm$ 6.39	7	27	0.258
Negative	13.25 $\pm$ 7.16	1	33	
Duration of disease (years)				
<5	10.32 $\pm$ 5.94	1	33	0.001**
6–10	13.96 $\pm$ 6.12	1	27	
11–15	18.50 $\pm$ 7.62	8	30	
>15	15.94 $\pm$ 4.67	7	25	

Notes: * $P < 0.05$; ** $P \leq 0.001$.

Discussion

In this study, we have reported the results of a questionnaire distributed to all vitiligo patients referred to dermatology clinics in Rafsanjan, Iran. Other assessment scales have reported the adverse impact of vitiligo on patient quality of life; however, the MC-SDS has never been used in vitiligo patients. The current report is valuable as additional support that vitiligo negatively impacts patient quality of life. Also, this study focused on the social acceptance in patients with vitiligo. The mean score of social acceptance in vitiligo shows low social acceptability for this disease.

In a study by Karelson et al, the quality of life of patients with vitiligo and psoriasis, as case and healthy control groups, was compared between 2009 and 2011. The average score of the Dermatology Life Quality Index (DLQI) in the vitiligo group was 7.4, which was significantly higher than that of

healthy controls (DLQI = 0.6).¹⁴ The average DLQI score obtained for vitiligo patients in Karelson's study was similar to those obtained for studies performed in Belgium (DLQI = 4.9), the UK (DLQI = 4.8), and Indonesia (DLQI = 4.4).^{15–17} In contrast, the DLQI score and social acceptance of vitiligo patients was much better in our study. Similar to our study, the mean DLQI score among the vitiligo patients was higher in Japan (9.5) when compared with studies conducted in other parts of the world.¹⁸

In a study performed by Ghajarzadeh et al in Tehran between 2009 and 2010, the quality of life in 300 patients (100 patients with alopecia areata, 100 patients with psoriasis, and 100 patients with vitiligo) was assessed, which showed that the average DLQI score in the vitiligo patients was 9.6 $\pm$ 4.8.¹⁹ Another study conducted in 2008 by Dolatshahi et al reported the DLQI score of vitiligo patients to be 16.8.²⁰

In Karelson et al's study, the quality of life in patients with vitiligo and psoriasis, as case and healthy control groups, was compared between the years 2009 and 2011. The results showed that vitiligo has a greater impact on feelings and relationships in women. They also showed lower quality of life during the active phase/the first episode of the disease.¹⁴

In another study that assessed the quality of life in 119 vitiligo patients and 162 psoriasis patients, women (compared to men) had significantly higher DLQI scores (45.6 vs 13.3), and gender was associated with DLQI score only in psoriasis patients.¹⁸ However, in the study by Ghajarzadeh et al, women had a higher DLQI score compared with men.¹⁹ Wang et al showed that patients with vitiligo feel embarrassed and, in the selection of clothing, have serious concerns about covering skin imperfections.^{20, 21}

Conclusion

This study indicates that certain groups of patients with vitiligo are at greater risk of experiencing a low acceptance and quality of life. Hence, it can be inferred that these patients require more serious efforts to treat the disease. Accordingly, it seems that the psychological interventions in patients who experience a worse quality of life are very important.

Acknowledgment

We acknowledge all patients with vitiligo who agreed to participate in this survey.

Disclosure

The authors report no conflicts of interest in this work.

References

- Sahni K, Parsad D, Kanwar AJ, Mehta SD. Autologous noncultured melanocyte transplantation for stable vitiligo: can suspending autologous melanocytes in the patients' own serum improve repigmentation and patient satisfaction? *Dermatol Surg*. 2011;37(2):176–182.
- Habif TP, Campbell JL, Chapman MS, Dinulos JGH, Zug KA. *Skin Disease, Diagnosis and Treatment*. Tehran: Lofty Ideas; 2011.
- Jin H, Atkinson JH, Yu X, et al. Depression and suicidality in HIV/AIDS in China. *J Affect Disord*. 2006;94(1):269–275.
- Grimes PE. New insights and new therapies in vitiligo. *JAMA*. 2005;293(6):730–735.
- Basra M, Fenech R, Gatt R, Salek M, Finlay AY. The Dermatology Life Quality Index 1994–2007: a comprehensive review of validation data and clinical results. *Br J Dermatol*. 2008;159(5):997–1035.
- Parsad D, Dogra S, Kanwar AJ. Quality of life in patients with vitiligo. *Health Qual Life Outcomes*. 2003;1(1):1–3.
- Ahmed I, Ahmed S, Nasreen S. Frequency and pattern of psychiatric disorders in patients with vitiligo. *J Ayub Med Coll Abbottabad*. 2007;19(3):19–21.
- Maleki M, Javidi Z, Kyafr B, Saadatian V, Saremi AK. Prevalence of depression in vitiligo patients. *Q J Fundamentals Men Health*. 2005;7(25–26):5–11.
- Porter JR, Beuf AH, Lerner A, Nordlund J. Psychosocial effect of vitiligo: a comparison of vitiligo patients with “normal” control subjects, with psoriasis patients, and with patients with other pigmentary disorders. *J Am Acad Dermatol*. 1986;15(2):220–224.
- Sherr L, Lampe F, Fisher M, et al. Suicidal ideation in UK HIV clinic attenders. *AIDS*. 2008;22(13):1651–1658.
- Lee J, Strickland D. Malignant melanoma: social status and outdoor work. *Br J Cancer*. 1980;41(5):757.
- AlGhamdi KM, Moussa NA, Mandil A, et al. Public perceptions and attitudes toward vitiligo. *J Cutan Med Surg*. 2012;16(5):334–340.
- Johnson TP, Fendrich M, Hubbell A. A validation of the Crowne-Marlowe social desirability scale. Paper presented at: 57th Annual Meeting of the American Association for Public Opinion Research; 2002.
- Karelson M, Silm H, Kingo K. Quality of life and emotional state in vitiligo in an Estonian sample: comparison with psoriasis and healthy controls. *Acta Derm Venereol*. 2013;93(4):446–450.
- Kent G, al-Abadie M. Factors affecting responses on Dermatology Life Quality Index items among vitiligo sufferers. *Clin Exp Dermatol*. 1996;21(5):330–333.
- Mashayekhi V, Javidi Z, Kiafar B, et al. Quality of life in patients with vitiligo: a descriptive study on 83 patients attending a PUVA therapy unit in Imam Reza Hospital, Mashad. *Indian J Dermatol Venereol Leprol*. 2010;76(5):592.
- Ongenae K, Van Geel N, De Schepper S, Naeyaert J. Effect of vitiligo on self-reported health-related quality of life. *Br J Dermatol*. 2005;152(6):1165–1172.
- Tanioka M, Yamamoto Y, Kato M, Miyachi Y. Camouflage for patients with vitiligo vulgaris improved their quality of life. *J Cosmet Dermatol*. 2010;9(1):72–75.
- Ghajarzadeh M, Ghiasi M, Kheirkhah S. Associations between skin diseases and quality of life: a comparison of psoriasis, vitiligo, and alopecia areata. *Acta Med Iran*. 2012;50(7):511–515.
- Dolatshahi M, Ghazi P, Feizy V, Hemami MR. Life quality assessment among patients with vitiligo: comparison of married and single patients in Iran. *Indian J Dermatol Venereol Leprol*. 2008;74(6):700.
- Wang KY, Wang KH, Zhang ZP. Health-related quality of life and marital quality of vitiligo patients in China. *J Eur Acad Dermatol Venereol*. 2011;25(4):429–435.

Clinical, Cosmetic and Investigational Dermatology

Publish your work in this journal

Clinical, Cosmetic and Investigational Dermatology is an international, peer-reviewed, open access, online journal that focuses on the latest clinical and experimental research in all aspects of skin disease and cosmetic interventions. This journal is included on PubMed. The manuscript management system is completely online

Submit your manuscript here: <https://www.dovepress.com/clinical-cosmetic-and-investigational-dermatology-journal>

Dovepress

and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors