

# Hypoglycemic activity and constituents analysis of blueberry (*Vaccinium corymbosum*) fruit extracts

Weifeng Huang<sup>1</sup>  
Liangliang Yao<sup>2</sup>  
Xiao He<sup>1</sup>  
Lei Wang<sup>1</sup>  
Mingxi Li<sup>3</sup>  
Youxin Yang<sup>3</sup>  
Chunpeng Wan<sup>3-5</sup>

<sup>1</sup>Department of Microbiology and Immunology, Medical College, China Three Gorge University, Yichang, Hubei 443002, China;

<sup>2</sup>Affiliated Hospital of Jiangxi University of Traditional Chinese Medicine, Nanchang, Jiangxi 330006, China; <sup>3</sup>College of Agronomy, Jiangxi Agricultural University, Nanchang 330045, China; <sup>4</sup>Jiangxi Key Laboratory for Postharvest Technology and Nondestructive Testing of Fruits & Vegetables, Nanchang 330045, China;

<sup>5</sup>Collaborative Innovation Center of Post-Harvest Key Technology and Quality Safety of Fruits and Vegetables, Jiangxi Agricultural University, Nanchang 330045, China

**Background:** To investigate hypoglycemic activity and elucidate the active composition of the fruit blueberry (*Vaccinium corymbosum*).

**Methods:** Methanol extracts of blueberry (MEB) were separated using a D101 macroporous resin column to yield quinic acid derivative (Fr.1)- and flavonoid (Fr.2)-rich fractions. The effects of the blueberry extracts on mRNA expression of GLUT-2 (glucose transporter type 2) and PPAR $\gamma$  (peroxisome proliferator-activated receptor- $\gamma$ ), as well as on the activities of PPRE (peroxisome proliferator response element) and NF- $\kappa$ B were analyzed in LO2 normal liver cells. Real-time PCR was used to detect the expression of GLUT-2, PPAR $\gamma$ , TNF- $\alpha$ , IL-1 $\beta$ , and IL-6 mRNA. The PPRE and NF- $\kappa$ B activities were detected by a luciferase reporter assay. Western blotting was used to detect the levels of PPAR $\gamma$ , GLUT-2, and p65. The active compositions were isolated using various chromatography columns, and were analyzed by NMR.

**Results:** mRNA and protein expression of GLUT-2 and PPAR $\gamma$  were significantly increased upon treatment with 400  $\mu$ g/mL extracts of blueberry ( $P < 0.05$ ). The PPRE activity was also significantly increased in a dose-dependent manner upon administration of MEB ( $P < 0.05$ ). Furthermore, the NF- $\kappa$ B activity induced by lipopolysaccharides was inhibited by MEB ( $P < 0.05$ ). No fraction separated from MEB exhibited PPRE activation or NF- $\kappa$ B inhibition activity. Blueberry extract may execute its hypoglycemic activity by stimulating expression of GLUT-2 and PPAR $\gamma$ , and by inhibiting the inflammatory pathway. Together, quinic acid derivatives and flavonoids may result in a synergistic effect. Fourteen phenolic acids, including eight flavonoids, four quinic acid derivatives, and two other phenolic acids, were isolated and identified, and caffeoylquinic acid derivatives and quercetin glycosides were found to be the major constituents of blueberry.

**Conclusion:** Blueberry may have hypoglycemic activity that functions through synergistic effects with caffeoylquinic acid derivatives and quercetin glycosides.

**Keywords:** blueberry, hypoglycemic activity, liver cells, caffeoylquinic acid derivatives, flavonoids, synergistic effect

## Introduction

Diabetes is a type of endocrine and metabolic disease caused by an absolute deficiency (type I) or relative deficiency (type II, insulin resistance) of insulin secretion, which is characterized by hyperglycemia, as well as complications, such as chronic impairment of the eyes, kidney, heart, blood vessels, and nervous system.<sup>1</sup> Currently, diabetes is one of the most harmful diseases to human health, and the number of diabetes patients in China has reached more than 150 million.<sup>2</sup> Patients with type II diabetes account for more than 90% of the total number of diabetes patients, and the incidence of diabetes is associated with many factors, including changes in dietary patterns, unhealthy

Correspondence: Chunpeng Wan  
College of Agronomy, Jiangxi Agricultural University, 1101 Zhimin Road, Changbei District, Nanchang 330045, China  
Tel +86-791-83813185  
Email chunpengwan@jxau.edu.cn

lifestyles, reduction of physical activity, increase in mental stress, environmental pollution, smoking, lack of public awareness and self-awareness, and population aging.<sup>3</sup> The main target drugs for the treatment of type II diabetes include insulin secretagogues (sulfonylureas, glibenclamides), insulin sensitizers (thiazolidinediones, biguanides), amylin analogs (Planklin),  $\alpha$ -glucosidase and amylase inhibitors (acarbose), DPP-VI inhibitors, glucagon-like peptide-1 (GLP-1), sodium-glucose cotransporter-2 (SGLT-2) inhibitors, protein tyrosine phosphatase 1B (PTP-1B) inhibitors, and peroxisome proliferator-activated receptor- $\gamma$  (PPAR $\gamma$ ) agonists, among others.<sup>4-9</sup> Glucose transporter 2 (GLUT-2) is a membrane protein that mediates glucose transport. Hepatic cell expression of GLUT-2 not only plays an important role in glucose metabolism, but also has a close association with diabetes or its associated complications. In the diabetic rat model, glucose-impaired insulin secretion has been proven to be associated with the reduction or inhibition of GLUT-2 expression by islet b cells, which plays an important role in the development of diabetes.<sup>10</sup> Recent studies have shown that inhibition of CB1R can downregulate GLUT-2 expression and reduce glucose reabsorption, which may support the rationale for clinical testing of peripherally restricted CB1R antagonists or the development of novel renal-specific GLUT-2 inhibitors against diabetic nephropathy.<sup>11</sup> PPAR $\gamma$  is a nuclear receptor that is mainly expressed in the large intestine, adipose tissue, and liver. PPAR $\gamma$  binds to PPRE (peroxisome proliferator response element), which has a variety of biological effects and plays an important role in adipocyte differentiation, glucose and lipid metabolism, insulin resistance, and the inflammatory response. PPAR $\gamma$  is also the effective target of thiazolidinedione (TZD) hypoglycemic drugs for the treatment of type II diabetes. PPAR $\gamma$  expression is down-regulated in liver tissues in type II diabetic db/db mice, and amelioration of hyperlipidemia and hyperglycemia are related to the up-regulation of PPAR $\gamma$  expression.<sup>12</sup> The transcription factor nuclear factor (NF- $\kappa$ B) mediates inflammation and stress signals. Numerous studies have indicated that there is a close relationship between NF- $\kappa$ B signaling and diabetes. Inhibition of the NF- $\kappa$ B cell signaling pathway improves insulin sensitivity in the liver.<sup>13</sup> Suppression of the NF- $\kappa$ B signaling pathway prevents diabetic liver injury in type II diabetic rat models.<sup>14</sup>

Blueberry belongs to the genus *Vaccinium* of Ericaceae. *Vaccinium* are perennial shrub fruit trees that originated in North America. There are more than 400 species of *Vaccinium* all over the world. Blueberries are mainly grown in the US, and are also known as American blueberries. Wild blueberries

in China are mostly grown on Changbai Mountain, the Da Hinggan Mountains, and in the forest area of the Xiao Hinggan Mountains, most of which are in the Da Hinggan Mountain region. Previous research has shown that blueberries contain abundant nutritional elements, such as anthocyanins, organic acids, phenolic acids, superoxide dismutase (SOD), pectin, polysaccharides, and pterostilbene, among others.<sup>15-18</sup> Blueberry is one of the five healthy fruits recommended by the Food and Agriculture Organization of the United Nations (FAO), because dietary consumption of blueberries has been demonstrated to be very beneficial to human health and provides effective protection against diseases, including lowering blood pressure, inhibiting tumorigenesis, and potentially preventing neurodegenerative disease.<sup>19-21</sup> These effects play an important role in the treatment of diseases such as diabetes, liver disease, cancer, cardiovascular disease, and anemia, among others.<sup>22,23</sup> A previous report showed that blueberry extract has good hypoglycemic activity. The study demonstrated that anthocyanins from blueberry have the potency to alleviate symptoms of hyperglycemia using a diabetic mice model;<sup>24</sup> however, its effective mechanism is not clear. To investigate the underlying mechanism of blueberry extract in decreasing the blood glucose level, the intent of the present research was to investigate the effect of blueberry extract on GLUT-2 and PPAR $\gamma$  mRNA expression, as well as on PPRE and NF- $\kappa$ B activity in liver cells, and to identify the chemical composition of the main active components by means of separation using various chromatography columns to clarify the hypoglycemic mechanism of blueberry.

## Materials and methods

### General experimental procedures

<sup>1</sup>H and <sup>13</sup>C nuclear magnetic resonance (NMR) data were recorded on a Varian 500 MHz instrument (Varian Inc., Palo Alto, USA) with TMS as the internal standard. Electrospray ionization mass spectral (ESI-MS) data were acquired on a Q-Star Elite mass spectrometer (Applied Biosystems MDS, Waltham, MA, USA). The UV spectra were measured on a SHIMADZU UV-2450 UV-visible spectrophotometer (Shimadzu Corporation, Kyoto, Japan). High-performance liquid chromatography (HPLC) was performed on a Hitachi Elite LaChrom system (Elite Lachrom Hitachi, Japan) consisting of a L2130 pump, L-2200 autosampler, and L-2455 diode array detector, all of which were operated by EZChrom Elite software (Scientific Software, Agilent Technologies, Santa Clara, USA). All solvents were of either analytical or HPLC grade and were purchased from Wilkem Scientific (Thermo Fisher Scientific, Shanghai, China).

## Cell culture

Dulbecco's modified Eagle's medium (DMEM) and fetal bovine serum (FBS) were purchased from Gibco (Grand Island, NY, USA). Human non-tumor hepatic LO2 cells were purchased from the Chinese Academy of Sciences (Shanghai, China). Cells were maintained in DMEM supplemented with 10% FBS and incubated in a humidified incubator at 37°C in 5% CO<sub>2</sub>.

## Extraction and isolation

The fruits of the blueberry species (*Vaccinium corymbosum*) were collected locally from Xinqizhou blueberry farms (Nanchang, Jiangxi, China) in June 2015. Fresh fruits (10.0 kg, fresh weight) were extracted exhaustively with MeOH (3×20 L) at room temperature to yield MeOH extracts. A part of the extracts (50 g) was dissolved in distilled deionized water (250 mL), filtered, and then loaded onto a D101 macroporous resin column (5×60 cm).<sup>25</sup> A 5-fold column volume of water was first used to wash the polysaccharide and protein materials. Next, a 5-fold column volume of 30% ethanol was used to wash the quinic acid derivatives (Fr.1), and 70% ethanol was further used to wash the flavonoid (Fr.2) constituents.

Fr.1 (15 g) was chromatographed on a C18 MPLC column (3.5×40 cm) and eluted with a gradient system of MeOH/H<sub>2</sub>O (1:9 to 8:2, v/v) to afford five sub-fractions (A<sub>1</sub>–A<sub>5</sub>), which were combined based on analytical HPLC analyses. Fraction A<sub>2</sub> was separated using a Sephadex LH-20 column (3.5×120 cm), eluted with MeOH, further separated by semi-preparative HPLC, and eluted with a gradient system of MeOH/H<sub>2</sub>O to yield compounds 9–12. Fraction A<sub>4</sub> was separated by semi-preparative HPLC and eluted with a gradient system of MeOH/H<sub>2</sub>O to yield compounds 13 and 14.

Fr.2 (11 g) was separated using a Sephadex LH-20 column (3.5×120 cm) and eluted with MeOH to afford three sub-fractions (B<sub>1</sub>–B<sub>3</sub>). Fraction B<sub>2</sub> was separated by a C18 MPLC column (3.5×40 cm) and eluted with a gradient system of MeOH/H<sub>2</sub>O (2:8 to 8:2, v/v) to afford five sub-fractions (C<sub>1</sub>–C<sub>5</sub>), which were combined based on analytical HPLC analyses. Sub-fraction C<sub>2</sub> was separated by semi-preparative HPLC and eluted with MeOH/H<sub>2</sub>O to yield compound 6. Sub-fraction C<sub>3</sub> was separated by semi-preparative HPLC and eluted with a gradient system of MeOH/H<sub>2</sub>O to yield compounds 1–4. Sub-fraction C<sub>4</sub> was separated by semi-preparative HPLC and eluted with a gradient system of MeOH/H<sub>2</sub>O to yield compounds 5, 7, and 8.

## Luciferase reporter assays

For detecting PPRE activity, LO2 cells were co-transfected with p-PPRE-luc and pSV40-β-galactosidase.<sup>26</sup> After 6 h,

the cells were subjected to different fractions of blueberry extracts for 24 h and then harvested to measure the luciferase activity. For detection of the NF-κB activity, LO2 cells were co-transfected with p-NF-κB-luc and pSV40-β-galactosidase. After 6 h, the cells were subjected to lipopolysaccharides or different fractions of blueberry extracts for 24 h and then harvested for measurement of the luciferase activity.

## Quantitative real-time RT-PCR analyses

Total RNA was isolated from hepatocytes using TRIzol reagent according to the manufacturer's instructions (Invitrogen Life Technology, Carlsbad, CA, USA). Total RNA was reverse-transcribed to cDNA using ReverTraAce (TOYOBO, Tokyo, Japan), as instructed. Quantitative real-time PCR was performed by standard methods using species-specific primer pairs (Table 1). The 18S rRNA expression levels were amplified and used for the calibration of real-time RT-PCR.

## Western blotting

LO2 cells were washed in ice cold saline, collected, and lysed with a lysis buffer contained protease inhibitors cocktail (Roche Diagnostics, Neuilly SurSeine, France). The total protein was extracted by NE-PER (Pierce Biotechnology, Rockford, IL, USA), and the concentration of protein was quantified by bicinchoninic acid (BCA) protein assay kit (Thermo Fisher Scientific, Waltham, MA, USA) according to the protocols of manufacturer. Samples (40 μg proteins per lane) were boiled for 100°C for 5 min and loaded on 8%–12% SDS–polyacrylamide gel electrophoresis (SDS-PAGE). The separated proteins of each lane representing one mouse were transferred to a polyvinylidene difluoride (PVDF) membrane (Millipore, MA, USA) and blocked in 5% nonfat milk solution for 1 h at room temperature with gentle shaking. Membranes were incubated with primary antibodies for GAPDH, GLUT-2, PPARγ, and NF-κB p65 (Abcam, MA, USA), diluted in 5% nonfat milk/TBST at 4°C overnight. Subsequently, each membrane was washed with TBST three times following incubation with horseradish peroxidase-conjugated secondary antibodies (rabbits and mouse) for 1 h at room temperature. Finally, specific bands of the membrane were performed using chemiluminescence (ECL) reagents (Millipore) and detected by Fluorescent Image Analyzer (FUJIFILM Corp, Tokyo, Japan).

## Statistical analyses

All statistical analyses were performed with GraphPad Prism software. Values are expressed as the mean±SEM. Pairwise

**Table 1** Primers used for quantitative real-time PCR

Gene	Primer direction	Sequences
GLUT-2 (NM_000340)	Forward	5'-GGACCTGCAATTCATTGTAGCT-3'
	Reverse	5'-GAACAGGGTAAAGGCCAGGAG-3'
PPAR $\gamma$ (NM_138712)	Forward	5'-GCACTGGAATTAGATGACAGCG-3'
	Reverse	5'-GTGGTTCAGCTTCAGCTGGAG-3'
TNF- $\alpha$ (NM_000594)	Forward	5'-ACCTCTCTCTAATCAGCCCTCT-3'
	Reverse	5'-GGGTTTGCTACAACATG GGCTA-3'
IL-1 $\beta$ (NM_000576)	Forward	5'-GCAATGAGGATGACTTGTCTTTTG-3'
	Reverse	5'-CAGAGGTCCAG GTCCTGGAA-3'
IL-6 (NM_000600)	Forward	5'-AGCCACTCACCTCTT CAG AAC-3'
	Reverse	5'-ACATGTCTCCTTTCTCAGGGC-3'
18S rRNA (NR_003286)	Forward	5'-GGTCATAAGCTTGCCTGTGATTAAG-3'
	Reverse	5'-CTACGGAAACCTTGTACGACTTT-3'

**Abbreviations:** PCR, polymerase chain reaction; GLUT-2, glucose transporter type 2; PPAR $\gamma$ , peroxisome proliferator-activated receptor- $\gamma$ .

comparisons were performed with Student's t-test (two-tailed), and multiple-group comparisons were performed with one-way ANOVA with Bonferroni's post-hoc test. A  $P$ -value < 0.05 was considered to be significant.

## Results

### Effects of blueberry extracts on PPRE activity


Quinic acid derivatives and flavonoid glycosides were two major classes of the chemical constituents of blueberry. Methanol extracts of blueberry (MEB) were separated using a D101 macroporous resin column. A 5-fold column volume of water was used to wash the polysaccharide and protein materials. Next, a 5-fold column volume of 30% ethanol was used to wash the quinic acid derivatives (Fr.1), and 70% ethanol was further used to wash the flavonoids (Fr.2). The UV spectra of Fr.1 and Fr.2 were typical of quinic acid derivatives and flavonol, respectively. PPRE activity in liver cells has been linked to beneficial hypoglycemic effects. To determine the potential hypoglycemic activities of blueberry extracts, the effects of blueberry extracts on PPRE activity were determined by luciferase reporter assays. Three extracts (MEB, Fr.1, and Fr.2) from blueberry were used to treat human non-tumor hepatic LO2 cells. As shown in Figure 1A, MEB treatment (200 and 400  $\mu$ g/mL) stimulated PPRE activity by 1.3- and 1.8-fold compared with the control. As indicated, there was no significant difference in the PPRE activity upon administration of extracts Fr.1 and Fr.2 of blueberry (Figures 1B and 1C). These results indicate that MEB may affect the hypoglycemic activity of blueberry through the synergistic effects of quinic acid derivatives and flavonols.

### Effects of blueberry extracts on NF- $\kappa$ B activity

Taking into account the importance of the NF- $\kappa$ B signaling pathway in diabetic liver injury and insulin resistance,<sup>27</sup> we evaluated the effects of blueberry extracts on NF- $\kappa$ B activity using luciferase reporter assays. In consistent with the results of previous reports, LPS (10 ng/mL) activated the reporter gene by 6.9-fold compared with the control.<sup>28</sup> MEB (100, 200, and 400  $\mu$ g/mL) markedly repressed the NF- $\kappa$ B-driven luciferase gene by 15%, 27%, and 40% (Figure 2A), respectively, compared with the levels in LPS-stimulated cells. However, extracts Fr.1 and Fr.2 of blueberry had no inhibitory effect on LPS-increased NF- $\kappa$ B activity (Figures 2B and 2C). To further explore the influence of MEB on the major inflammatory signaling pathway of NF- $\kappa$ B, p65 was then examined by Western blotting (Figure 2D). LPS increased protein levels of p65 compared to the control cell, while MEB inverted these variations (Figure 2D). Moreover, to test whether MEB suppressed downstream inflammatory cytokines of the pathway in mRNA, RT-qPCR was utilized to detect the expression of TNF- $\alpha$ , IL-1 $\beta$ , and IL-6 following LPS treatment (Figure 2E). Interestingly, MEB did produce a marked change with a significant decrease in these inflammation mediators when compared to LPS treated cells. These results confirmed that MEB are effective components of blueberry with regards to its hypoglycemic activity.


### Effects of MEB on the expression of GLUT-2 and PPAR $\gamma$ mRNA and protein

To further detect the hypoglycemic activity of MEB, mRNA expression of GLUT-2 and PPAR $\gamma$  were measured in LO2



**Figure 1** Effects of blueberry extract on PPRE activity. **(A)** Effects of MEB on PPRE activity. **(B)** Effects of extract Fr.1 of blueberry on PPRE activity. **(C)** Effects of extract Fr.2 of blueberry on PPRE activity. LO2 cells were co-transfected with p-PPRE-luc and pSV40-β-galactosidase. After 6 h, the cells were subjected to different extracts of blueberry for 24 h and then harvested for measurement of the luciferase activity. All of the results are presented as the means±SD of three independent experiments (n=3). \*p<0.05; \*\*\*p<0.001.

**Abbreviations:** PPRE, peroxisome proliferator response element; MEB, methanol extracts of blueberry; RLU, relative luciferase unit; SD, standard deviation.




**Figure 2** Effects of blueberry extracts on NF-κB activity. **(A)** Effects of MEB on NF-κB activity. **(B)** Effects of extract Fr.1 of blueberry on NF-κB activity. **(C)** Effects of extract Fr.2 of blueberry on NF-κB activity. LO2 cells were co-transfected with p-NF-κB-luc and pSV40-β-galactosidase. After 6 h, the cells were subjected to LSP (10 ng/mL) and/or different extracts of blueberry for 24 h, and then harvested for measurement of the luciferase activity. **(D)** Effects of MEB on p65 expression. **(E)** Effects of MEB on mRNA expression of TNF-α, IL-1β, and IL-6. The LO2 cells were subjected to LPS (10 ng/mL) and/or different extracts of blueberry for 24 h and then harvested for Western blotting and quantitative real-time RT-PCR. All results are presented as the means±SD of three independent experiments (n=3). \*p<0.05; \*\*p<0.01; \*\*\*p<0.001.

**Abbreviations:** NF-κB, transcriptional factor nuclear factor; MEB, methanol extracts of blueberry; LPS, lipopolysaccharides; RT-PCR, real time polymerase chain reaction; SD, standard deviation.

cells under treatment with MEB. We found that MEB induced GLUT-2 and PPARγ mRNA expression. Blueberry extract at a dose of 400 μg/mL enhanced GLUT-2 and PPARγ mRNA expression by 1.5- and 1.4-fold (Figures 3A and 3B), respectively, compared with the control. The effects of MEB on the mRNA expression of GLUT-2 and PPARγ were confirmed by the Western blotting results in Figures 3C and 3D.

## Chemical compositions analysis of blueberry extracts

To elucidate the chemical compositions of the blueberry extracts, MEB were first fractionated using a D101 macroporous resin column to yield the Fr.1 and Fr.2 fractions. The two fractions were further purified using various chromatographic columns, including a Sephadex LH-20, octadecylsilane, and



**Figure 3** Effects of MEB on the mRNA expression of GLUT-2 (A), PPAR $\gamma$  (B), and protein expression of GLUT-2, PPAR $\gamma$  (C and D). LO2 cells were maintained in DMEM supplemented with 10% FBS and incubated in a humidified incubator at 37°C in 5% CO<sub>2</sub> for 24 h. Then, the cells were further incubated with MEB (100, 200, and 400 μg/mL) for another 24 h. Cells were then collected for quantitative real-time RT-PCR and Western blotting. All of the results are presented as the means $\pm$ SD of three independent experiments (n=3). \* $p$ <0.05; \*\* $p$ <0.01; \*\*\* $p$ <0.001.

**Abbreviations:** MEB, methanol extracts of blueberry; GLUT-2; glucose transporter type 2; PPAR $\gamma$ , peroxisome proliferator-activated receptor- $\gamma$ ; DMEM, Dulbecco's Modified Eagle's Medium; FBS, fetal bovine serum; RT-PCR; real time polymerase chain reaction; SD, standard deviation.

semi-preparative HPLC, to yield pure compounds. The isolated compounds were characterized by <sup>1</sup>H-NMR, <sup>13</sup>C-NMR, and ESI-MS. Fourteen phenolic acids, including eight flavonoids, four quinic acid derivatives, and two other phenolic acids, were isolated, and their structures were identified as isoquercetin (1),<sup>29</sup> hyperin (2),<sup>29</sup> guajavarin (3),<sup>29,30</sup> quercitrin (4),<sup>29</sup> astragaln (5),<sup>31</sup> isomyricitrin (6),<sup>32</sup> helichryoside (7),<sup>33</sup> tiliroside (8),<sup>33,34</sup> 5-*O*-caffeoylquinic acid (9),<sup>31,35</sup> 3-*O*-caffeoylquinic acid (10),<sup>35</sup> 5-*O*-caffeoylquinic acid methyl ester (11),<sup>33</sup> 3-*O*-caffeoylquinic acid methyl ester (12),<sup>35</sup> methyl cinnamate (13),<sup>33</sup> and methyl caffeate (14).<sup>33</sup>

Compounds 1–4 were obtained as yellowish amorphous powders. The UV spectrum showed a  $\lambda_{\max}$  at approximately 255 nm (band II) and 355 nm (band I). The <sup>1</sup>H-NMR spectrum (Table 2) showed two meta proton peaks at approximately  $\delta$  6.10 (1H, d,  $J$ =2.0 Hz) and 6.30 (1H, d,  $J$ =2.0 Hz) ppm, consistent with the H-6 and H-8 on A-ring of flavonoid, and an ABX system at approximately 7.25–7.74 (1H, d,  $J$ =2.2 Hz, H-2'), 7.20–7.48 (1H, dd,  $J$ =2.2 Hz, 8.4 Hz, H-6'), and 6.78 (1H, d,  $J$ =8.4 Hz, H-5'), corresponding to the catechol protons on the B-ring. One anomeric proton signal was found at [1:  $\delta$  5.15(1H, d,  $J$ =7.6), 2:  $\delta$  5.07(1H, d,  $J$ =7.6), 3:  $\delta$  5.07(1H, d,  $J$ =6.6), 4:  $\delta$  5.27(1H, d,  $J$ =1.1)]. These signals showed that

compounds 1–4 contained the same aglycone of quercetin. Additionally, one methyl signal at  $\delta$  0.87(3H, d,  $J$ =6.1) of compound 4 was found, which can be easily identified as quercitrin (4). Through careful comparison of the <sup>13</sup>C-NMR data of the sugar moiety carbon signals of compounds 1–3 at [1: 102.8(C-1''), 74.3(C-2''), 76.7(C-3''), 69.8(C-4''), 77.0(C-5''), 61.1(C-6''), 2: 103.9(C-1''), 71.7(C-2''), 73.7(C-3''), 68.6(C-4''), 75.7(C-5''), 60.5(C-6''), 3: 102.3(C-1''), 71.4(C-2''), 72.7(C-3''), 67.7(C-4''), 65.6(C-5'')], the compounds were identified as isoquercetin (1), hyperin (2), and guajavarin (3), respectively. Compounds 5 and 6 showed the same sugar moiety proton signals as those of compound 1 in the <sup>1</sup>H-NMR and <sup>13</sup>C-NMR data, although the B-rings were different. An AA'BB' system was found at 7.97 (2H, d,  $J$ =8.8 Hz, H-2', 6') and 6.84 (2H, d,  $J$ =7.8 Hz, H-3', 5') of compound 5 and at 7.30 (2H, s, H-2', 6') of compound 6; therefore, compounds 5 and 6 were identified as astragaln (5) and isomyricitrin (6).

Compound 7 showed the same aglycone as seen in compounds 1–4. The UV spectrum showed  $\lambda_{\max}$  values at 259 nm (band II) and 314 nm (band I), typical of the UV spectrum of the coumaroyl substituent quercetin glycoside.<sup>36</sup> <sup>1</sup>H-NMR data confirmed this speculation, showing signals of quercetin

glycoside signals [ $\delta$  7.60 (1H, d,  $J=2.0$  Hz, H-2'), 7.55 (1H, dd,  $J=8.4$ , 2.0 Hz, H-6'), 6.81 (1H, d,  $J=8.4$  Hz, H-5'), 6.28 (1H, d,  $J=2.0$  Hz, H-8), 6.11 (1H, d,  $J=2.0$  Hz, H-6), 5.28 (1H, d,  $J=7.6$  Hz, H-1''), 4.33 (1H, d,  $J=11.5$  Hz, H-6''a), 4.21 (1H, dd,  $J=11.5$ , 6.7 Hz, H-6''b), 3.55–3.40 (3H, H-2'', 3'', 5''), and 3.32 (1H, t,  $J=9.6$  Hz, H-4'')]. One additional coumaroyl signal was detected at  $\delta$  7.40 (1H, d,  $J=15.8$  Hz, H-7'''), 7.30 (2H, d,  $J=8.2$  Hz, H-2''', 6'''), 6.78 (2H, d,  $J=8.2$  Hz, H-3''', 5'''), and 6.07 (1H, d,  $J=15.8$  Hz, H-8'''). The downfield shift of H-6 ( $\delta$  4.33 and 4.21) indicated that coumaroyl was linked to C-6 of the sugar. Compound 7 was identified as helichryoside (7), which was further confirmed by HR-ESI-MS,  $m/z$  611.1097 [M+H]<sup>+</sup> (calcd. for C<sub>30</sub>H<sub>27</sub>O<sub>14</sub>, 611.1401). Compound 8 showed the same aglycone as that of 5, and the UV spectrum showed  $\lambda_{\max}$  values at 266 nm (band II) and 312 nm (band I), which is the typical UV spectrum of the coumaroyl substituent kaempferol glycoside.<sup>36</sup> Thus, compound 8 was identified as tiliroside (8). <sup>1</sup>H-NMR (500 MHz, CD<sub>3</sub>OD)  $\delta$  7.98 (2H, dd,  $J=8.8$ , 2.0 Hz, H-2', 6'), 6.82 (2H, d,  $J=8.8$ , 2.0 Hz, H-3', 5'), 7.40 (1H, d,  $J=15.9$  Hz, H-7'''), 7.31 (2H, d,  $J=8.2$  Hz, H-2''', 6'''), 6.80 (2H, d,  $J=8.2$

Hz, H-3''', 5'''), 6.30 (1H, d,  $J=2.0$  Hz, H-8), 6.12 (1H, d,  $J=2.0$  Hz, H-6), 6.00 (1H, d,  $J=15.9$  Hz, H-8'''), 5.24 (1H, d,  $J=7.6$  Hz, H-1''), 4.30 (1H, brd,  $J=11.5$  Hz, H-6''a), 4.19 (1H, dd,  $J=11.5$ , 6.7 Hz, H-6''b), 3.40–3.48 (3H, H-2'', 3'', 5''), 3.31 (1H, m, H-4'').

Compounds 9–12 showed similar <sup>1</sup>H-NMR spectra (Table 3) and UV spectra, with  $\lambda_{\max}$  values at 327, 298 (shoulder), and 242 nm, suggesting that compounds 9–12 were caffeoyl-substituted quinic acid derivatives.<sup>35</sup> The substituted position of caffeoyl was used to identify the oxygenated methine protons of the quinic acid core, as detailed in the discussion previously.<sup>35</sup> Compounds 9 and 10 were identified as 5-*O*-caffeoylquinic acid (9) and 3-*O*-caffeoylquinic acid (10). There was one additional OCH<sub>3</sub> signal found in the spectra of compounds 11 and 12, for which the other proton signals were very similar to those of compounds 9 and 10. Compounds 11 and 12 were identified as 5-*O*-caffeoylquinic acid methyl ester (11) and 3-*O*-caffeoylquinic acid methyl ester (12).  $m/z$  201.0480 [M+Na]<sup>+</sup> (calcd. for C<sub>10</sub>H<sub>10</sub>NaO<sub>3</sub>, 201.0528). <sup>1</sup>H-NMR (500 MHz, CD<sub>3</sub>OD):  $\delta$  7.60 (1H, d,  $J=16.1$  Hz, H-7), 7.44 (2H, d,  $J=8.3$  Hz, H-2, 6), 6.80 (2H,

**Table 2** <sup>1</sup>H-NMR (500 MHz, CD<sub>3</sub>OD) characteristics of flavonol glycosides 1–6 isolated from blueberry extracts

No. H proton	1	2	3	4	5	6
	$\delta$ H (J Hz)	$\delta$ H (J Hz)	$\delta$ H (J Hz)	$\delta$ H (J Hz)	$\delta$ H (J Hz)	$\delta$ H (J Hz)
6	6.10 (1H, d, 2.0)	6.10 (1H, d, 2.0)	6.10 (1H, d, 2.0)	6.09 (1H, d, 2.0)	6.10 (1H, d, 2.0)	6.10 (1H, d, 2.1)
8	6.29 (1H, d, 2.0)	6.30 (1H, d, 2.0)	6.29 (1H, d, 2.0)	6.26 (1H, d, 2.0)	6.28 (1H, d, 2.0)	6.27 (1H, d, 2.1)
2'	7.61 (1H, d, 2.2)	7.74 (1H, d, 2.2)	7.65 (1H, d, 2.1)	7.25 (1H, d, 2.0)	7.97 (1H, d, 8.8)	7.30 (1H, s)
3'					6.84 (1H, d, 7.8)	
5'	6.76 (1H, d, 8.5)	6.77 (1H, d, 8.5)	6.78 (1H, d, 8.5)	6.82 (1H, d, 8.3)	6.84 (1H, d, 7.8)	
6'	7.48 (1H, dd, 2.2, 8.5)	7.48 (1H, dd, 2.2, 8.5)	7.48 (1H, dd, 2.1, 8.5)	7.20 (1H, dd, 2.0, 8.3)	7.97 (1H, d, 8.8)	7.30 (1H, s)
1''	5.15 (1H, d, 7.6)	5.07 (1H, d, 7.6)	5.07 (1H, d, 6.6)	5.27 (1H, d, 1.1)	5.15 (1H, d, 7.6)	5.15 (1H, d, 7.6)
CH <sub>3</sub>				0.87 (3H, d, 6.1)		

**Abbreviation:** <sup>1</sup>H-NMR, <sup>1</sup>H-nuclear magnetic resonance.

**Table 3** <sup>1</sup>H-NMR (500 MHz, CD<sub>3</sub>OD) characteristics of compounds 9–12 from blueberry extracts

No. H proton	9	10	11	12
	$\delta$ H (J Hz)	$\delta$ H (J Hz)	$\delta$ H (J Hz)	$\delta$ H (J Hz)
2	1.96–2.17 (2H, m)	1.94–2.14 (2H, m)	1.90–2.12 (2H, m)	1.90–2.13 (2H, m)
3	4.08 (1H, ddd, 1.8, 4.9, 4.9)	5.23 (1H, brd, 4.1)	4.07 (1H, brs)	5.18 (1H, ddd, 1.8, 4.9, 4.9)
4	3.65 (1H, dd, 3.1, 8.8)	3.63 (1H, dd, 3.1, 8.5)	4.96 (1H, dd, 3.1, 9.4)	3.63 (1H, dd, 3.1, 7.5)
5	5.25 (1H, ddd, 4.5, 9.4, 9.4)	4.07 (1H, ddd, 3.6, 8.5, 8.5)	5.24 (1H, ddd, 4.5, 9.4, 9.4)	4.05 (1H, ddd, 4.5, 9.0, 9.0)
6	1.96–2.17 (2H, m)	1.94–2.14 (2H, m)	1.90–2.12 (2H, m)	1.90–2.13 (2H, m)
2'	6.96 (1H, d, 2.1)	6.95 (1H, d, 2.1)	7.09 (1H, d, 1.9)	6.94 (1H, d, 2.0)
5'	6.68 (1H, d, 8.2)	6.68 (1H, d, 8.2)	6.70 (1H, d, 7.9)	6.68 (1H, d, 8.2)
6'	6.86 (1H, dd, 2.1, 8.2)	6.86 (1H, dd, 2.1, 8.2)	6.96 (1H, dd, 1.9, 7.9)	6.85 (1H, dd, 2.0, 8.2)
7'	7.48 (1H, d, 15.9)	7.47 (1H, d, 15.9)	7.52 (1H, d, 15.9)	7.42 (1H, d, 15.9)
8'	6.19 (1H, d, 15.9)	6.18 (1H, d, 15.9)	6.25 (1H, d, 15.9)	6.13 (1H, d, 15.9)
OCH <sub>3</sub>			3.88 (3H, s)	3.80 (3H, s)

**Abbreviation:** <sup>1</sup>H-NMR, <sup>1</sup>H-nuclear magnetic resonance.

d,  $J=8.3$  Hz, H-3, 5), 6.30 (1H, d,  $J=16.1$  Hz, H-8), 3.75 (3H, s, OCH<sub>3</sub>). Compound **13** was identified as methyl cinnamate.

The UV spectrum of compound **14** showed  $\lambda_{\max}$  at 325, 297, 240 nm, (+) HR-ESIMS,  $m/z$  217.0453 [M+Na]<sup>+</sup> (calcd for C<sub>10</sub>H<sub>10</sub>NaO<sub>4</sub>, 217.0477). <sup>1</sup>H-NMR (500 MHz, CD<sub>3</sub>OD):  $\delta$  7.57 (1H, d,  $J=15.9$  Hz, H-7), 7.04 (1H, d,  $J=1.8$  Hz, H-2), 6.94 (1H, dd,  $J=8.2, 1.8$  Hz, H-6), 6.79 (1H, d,  $J=8.2$  Hz, H-5), 6.28 (1H,  $J=15.9$  Hz, H-8), 3.79 (3H, s, OCH<sub>3</sub>). Compound **14** was identified as methyl caffeate.

## Discussion

Most experimental and clinical studies have strongly suggested that blueberry is a source of bioactive compounds for the treatment of obesity and type II diabetes.<sup>37</sup> However, the mechanisms have not been illuminated in detail. A D101 macroporous resin column was used to separate MEB, and the effect of the resulting fractions on PPRE and NF- $\kappa$ B activity were detected. Inflammation contributes to the pathogenesis of type II diabetes, and anti-inflammation strategies for the treatment of this disease simultaneously lower blood glucose levels and potentially reduce the severity and prevalence of the associated complications.<sup>38</sup> Previous research has shown that blueberry polyphenol extract effectively inhibits the LPS-induced inflammatory response and decreases the activity of NF- $\kappa$ B,<sup>39</sup> lowering proinflammatory mediators, including nitric oxide (NO), tumor necrosis factor- $\alpha$  (TNF- $\alpha$ ), and interleukin-1 beta (IL-1 $\beta$ ), and down-regulating NO synthase (iNOS) and cyclooxygenase 2 (COX2). Inconsistent with the results of previous reporting, MEB markedly repressed the NF- $\kappa$ B-driven luciferase gene. MEB were separated into two groups, quinic acid derivatives (Fr.1) and flavonoid glycosides (Fr.2). Interestingly, neither of the separated MEB fractions demonstrated inhibitory effects on NF- $\kappa$ B activity. This phenomenon supported the notion that quinic acid derivatives and flavonoids result in a synergistic effect in the repression of NF- $\kappa$ B activity. Hyperglycemia induces inflammatory responses, producing free radicals that can lead to type II diabetes.<sup>40</sup> Polyphenols exert their protective effects against diabetes mellitus by suppressing inflammatory responses.<sup>41,42</sup> It is supposed that the hypoglycemic effects of MEB are mediated, at least in part, by the inhibition of NF- $\kappa$ B activity. Activation of PPARs in the treatment of type II diabetes mellitus (T2DM) has proven effective in improving insulin sensitivity, hyperglycemia, and lipid metabolism.<sup>43</sup>

PPARs are important in controlling the expression of genes involved in the regulation of glucose, lipid, and cholesterol metabolism; cell growth; and cell differentiation by binding to specific PPREs at the enhancer sites of

target genes.<sup>44</sup> Activation of PPARs contributed to anti-inflammatory effects in several cell types, including liver cells.<sup>12</sup> The anti-inflammatory properties of PPARs are often associated with the repression of transcriptional pathways involved in inflammatory responses, such as modulation of NF- $\kappa$ B signaling. Currently, MEB have been shown to stimulate PPRE activity and induce up-regulation of PPAR $\gamma$  mRNA expression. This suggested that the repression of NF- $\kappa$ B activity by MEB is closely related to the simultaneous PPRE activity. This result is further supported by the observation that quinic acid derivatives and flavonoids have no effect on PPRE activity. However, the existing result could not rule out the contribution of PPAR $\alpha$  and PPAR $\beta$  to PPRE activity. The potential mechanism of PPRE activity mediation by MEB requires further study.

Recently, GLUT-2 has drawn attention as a molecule that might be involved in the pathogenesis of diabetes mellitus. Improvement of insulin signaling in HepG2 is involved in enhancing the GLUT-2 expression.<sup>45</sup> GLUT-2 is a glucose transporter that is mainly present in the plasma membrane of pancreatic  $\beta$ -cells. GLUT-2 has been shown to have blood glucose regulation functions through the control of insulin secretion,<sup>46</sup> and it was reported that GLUT-2 mRNA and protein expression were reduced by hyperinsulinemia and increased by hyperglycemia in the liver of diabetic rats.<sup>47</sup> Rutin, a flavonoid isolated from *Toona sinensis* Roem, has the ability to enhance insulin-dependent receptor kinase (IRK) activity and glucose transporter 4 (GLUT4) translocation in differentiated myotubes.<sup>48</sup> Chlorogenic acid (CGA), a common dietary polyphenol with numerous biologically activities, reversed the downregulation of GLUT-2 induced by a HFD (high-fat diet).<sup>49</sup> Consistently, the present study demonstrated that MEB stimulated GLUT-2 mRNA expression in liver cells. Polyphenol-rich *Chrysanthemum morifolium* extract (CME) also showed the ability to reverse the decline of PPAR $\alpha/\gamma$  and GLUT-2 induced by alloxan. Chemical constituents analysis showed that chlorogenic acid, dicaffeoylquinic acid, and apigenin were the major polyphenols of CME, and those polyphenols might exert a synergic hypoglycemic effect via PPAR $\alpha/\gamma$ -mediated mechanisms.<sup>50</sup> *Pongamia glabra* (PBME) and *Ficus glomerata* (FBME) produced a synergistic hypoglycemic effect with combined therapy at low doses. The primary constituents in the two plants were flavonoids, furanoflavonoids, sterols, saponins, glycosides, glaucinol, tannins, and other polyphenol compounds.<sup>51</sup> Through inhibition of oxidative stress, polyphenols protect against the effects of chronic diseases mediated by inflammatory responses.<sup>52</sup>



Taken together, blueberry extract may exert hypoglycemic properties through the synergistic effects of caffeoylquinic acid derivatives and quercetin glycosides, and the hypoglycemic effect is involved in an increase of GLUT-2 and PPAR $\gamma$  expression and inhibition of the relevant inflammatory pathways. Although further efforts are needed to define the hypoglycemic mechanisms of blueberry extract in diabetic animal model, a diet rich in blueberry extract may be a potential chemopreventive tool useful for the management of diabetes.

## Acknowledgments

The project was supported by the National Natural Science Foundation of China (31500286) and the Natural Science Foundation of Jiangxi Province (20161BAB214167). We thank Nature Research Editing Service for its linguistic assistance during the preparation of this manuscript.

## Author contributions

Weifeng Huang and Chunpeng Wan conceived and designed the experiments; Liangliang Yao, Xiao He, Lei Wang, Mingxi Li, and Youxin Yang performed the experiments; Weifeng Huang and Chunpeng Wan analyzed the data; Chunpeng Wan contributed reagents/materials/analysis tools; Weifeng Huang and Chunpeng Wan wrote the paper. All authors contributed toward data analysis, drafting and revising the paper and agree to be accountable for all aspects of the work.

## Disclosure

The authors declare no conflicts of interest in this work.

## References

- American Diabetes Association. Diagnosis and classification of diabetes mellitus. *Diabetes Care*. 2014;37(S1):S81–S90.
- Ma RC, Lin X, Jia W. Causes of type 2 diabetes in China. *Lancet Diabetes Endocrinol*. 2014;2(12):980–991.
- Chan JC, Zhang Y, Ning G. Diabetes in China: a societal solution for a personal challenge. *Lancet Diabetes Endocrinol*. 2014;2(12):969–979.
- Deacon CF, Mannucci E, Ahrén B. Glycaemic efficacy of glucagon-like peptide-1 receptor agonists and dipeptidyl peptidase-4 inhibitors as add-on therapy to metformin in subjects with type 2 diabetes—a review and meta analysis. *Diabetes Obes Metab*, 2012;14(8):762–767.
- Nair S, Wilding JP. Sodium glucose cotransporter 2 inhibitors as a new treatment for diabetes mellitus. *J Clin Endocrinol Metab*. 2010;95(1):34–42.
- Bhandari MR, Jong-Anurakkun N, Hong G, Kawabata J.  $\alpha$ -Glucosidase and  $\alpha$ -amylase inhibitory activities of Nepalese medicinal herb Pak-hanbhed (*Bergenia ciliata*, Haw.). *Food Chem*. 2008;106(1):247–252.
- Green BD, Flatt PR, Bailey CJ. Dipeptidyl peptidase IV (DPP IV) inhibitors: a newly emerging drug class for the treatment of type 2 diabetes. *Diab Vasc Dis Res*. 2006;3(3):159–165.
- An T, Hong D, Hu L, Li J. *Protein tyrosine phosphatases 1B inhibitors from traditional Chinese medicine*. ACS Publications; 2006:143–156.
- Huang TH, Peng G, Kota BP, et al. Anti-diabetic action of Punica granatum flower extract: Activation of PPAR $\gamma$  and identification of an active component. *Toxicol Appl Pharmacol*. 2005;207(2):160–169.
- Abe N, Watanabe T, Ozawa S, et al. Pancreatic endocrine function and glucose transporter (GLUT)–2 expression in rat acute pancreatitis. *Pancreas*. 2002;25(2):149–153.
- Jourdan T, Szanda G, Rosenberg AZ, et al. Overactive cannabinoid 1 receptor in podocytes drives type 2 diabetic nephropathy. *Proc Natl Acad Sci USA*. 2014;111(50):E5420–E5428.
- Zhong J, Gong W, Lu L, et al. Irbesartan ameliorates hyperlipidemia and liver steatosis in type 2 diabetic db/db mice via stimulating PPAR $\gamma$ , AMPK/Akt/mTOR signaling and autophagy. *Int Immunopharmacol*. 2017;42:176–184.
- Ke B, Zhao Z, Ye X, et al. Inactivation of NF- $\kappa$ B p65 (RelA) in liver improves insulin sensitivity and inhibits cAMP/PKA pathway. *Diabetes*. 2015;64(10):3355–3362.
- Han L, Li C, Sun B, et al. Protective effects of celastrol on diabetic liver injury via TLR4/MyD88/NF- $\kappa$ B signaling pathway in type 2 diabetic rats. *J Diabetes Res*. 2016;2016:2641248
- Su X, Zhang J, Wang H, et al. Phenolic acid profiling, antioxidant, and anti-inflammatory activities, and miRNA regulation in the polyphenols of 16 blueberry samples from China. *Molecules*. 2017;22(2):pii: E312.
- Khalid S, Barfoot KL, May G, Lamport DJ, Reynolds SA, Williams CM. Effects of acute blueberry flavonoids on mood in children and young adults. *Nutrients*. 2017;9(2):pii: E158.
- Diaconeasa Z, Leopold L, Rugină D, Avvaz H, Socaciu C. Antiproliferative and antioxidant properties of anthocyanin rich extracts from blueberry and blackcurrant juice. *Int J Mol Sci*. 2015;16(2):2352–2365.
- Castrejón ADR, Eichholz I, Rohn S, Kroh LW, Huyskens-Keil S. Phenolic profile and antioxidant activity of highbush blueberry (*Vaccinium corymbosum* L.) during fruit maturation and ripening. *Food Chem*. 2008;109(3):564–572.
- Szajdek A, Borowska EJ. Bioactive compounds and health-promoting properties of berry fruits: a review. *Plant Foods Hum Nutr*. 2008;63(4):147–156.
- Neto CC. Cranberry and blueberry: evidence for protective effects against cancer and vascular diseases. *Mol Nutr Food Res*. 2007;51(6):652–664.
- Rodríguez-Mateos A, Ishisaka A, Mawatari K, Vidal-Diez A, Spencer JP, Terao J. Blueberry intervention improves vascular reactivity and lowers blood pressure in high-fat-, high-cholesterol-fed rats. *Br J Nutr*. 2013;109(10):1746–1754.
- Stull AJ. Blueberries' impact on insulin resistance and glucose intolerance. *Antioxidants*. 2016;5(4):pii: E44.
- Bingül İ, Başaran-Küçükgergin C, Tekkeşin MS, Olgaç V, Doğru-Abbasoğlu S, Uysal M. Effect of blueberry pretreatment on diethylnitrosamine-induced oxidative stress and liver injury in rats. *Environ Toxicol Pharmacol*. 2013;36(2):529–538.
- Grace MH, Ribnicky DM, Kuhn P, et al. Hypoglycemic activity of a novel anthocyanin-rich formulation from lowbush blueberry, *Vaccinium angustifolium* Aiton. *Phytomedicine*. 2009;16(5):406–415.
- Shi D, Xu M, Ren M, et al. Immunomodulatory effect of flavonoids of blueberry (*vaccinium corymbosum* L.) leaves via the NF- $\kappa$ B signal pathway in LPS-stimulated RAW 264.7 cells. *J Immunol Res*. 2017;2017:5476903.
- Smale ST. Luciferase assay. *Cold Spring Harb Protoc*. 2010;2010(5):pdb. prot5421.
- Baker RG, Hayden MS, Ghosh S. NF- $\kappa$ B, inflammation, and metabolic disease. *Cell Metab*. 2011;13(1):11–22.
- Chassin C, Hornef MW, Bens M, et al. Hormonal control of the renal immune response and antibacterial host defense by arginine vasopressin. *J Exp Med*. 2007, 204(12):2837–2852.
- Lu Y, Foo LY. Identification and quantification of major polyphenols in apple pomace. *Food Chem*. 1997;59(2):187–194.

30. Fraisse D, Heitz A, Carnat A, Carnat AP, Lamaison JL. Quercetin 3-arabinopyranoside, a major flavonoid compound from *Alchemilla xanthochlora*. *Fitoterapia*. 2000;71(4):463–464.
31. Tan C, Wang Q, Luo C, Chen S, Li Q, Li P. Yeast  $\alpha$ -glucosidase inhibitory phenolic compounds isolated from *Gynura medica* leaf. *Int J Mol Sci*. 2013;14(2):2551–2558.
32. Scharbert S, Holzmann N, Hofmann T. Identification of the astringent taste compounds in black tea infusions by combining instrumental analysis and human bioresponse. *J Agric Food Chem*. 2004;52(11):3498–3508.
33. Wan C, Yuan T, Cirello AL, Seeram NP. Antioxidant and  $\alpha$ -glucosidase inhibitory phenolics isolated from highbush blueberry flowers. *Food Chem*. 2012;135(3):1929–1937.
34. Zhang Y, Seeram NP, Lee R, Feng L, Heber D. Isolation and identification of strawberry phenolics with antioxidant and human cancer cell antiproliferative properties. *J Agric Food Chem*. 2008;56(3):670–675.
35. Wan C, Li S, Liu L, Chen C, Fan S. Caffeoylquinic acids from the aerial parts of *chrysanthemum coronarium* L. *Plants*. 2017;6(1):pii. E10.
36. Wang XM, Wan CP, Zhou SR, Qiu Y. Two new flavonol glycosides from *Sarcopyramis bodinieri* var. *delicate*. *Molecules*. 2008;13(6):1399–1405.
37. Shi M, Loftus H, McAinch AJ, Su XQ. Blueberry as a source of bioactive compounds for the treatment of obesity, type 2 diabetes and chronic inflammation. *J Funct Foods*. 2017;30:16–29.
38. Donath MY, Shoelson SE. Type 2 diabetes as an inflammatory disease. *Nat Rev Immunol*. 2011;11(2):98–107.
39. Lau FC, Joseph JA, McDonald JE, Kalt W. Attenuation of iNOS and COX2 by blueberry polyphenols is mediated through the suppression of NF- $\kappa$ B activation. *J Funct Foods*. 2009;1(3):274–283.
40. Pickup JC. Inflammation and activated innate immunity in the pathogenesis of type 2 diabetes. *Diabetes Care*. 2004;27(3):813–823.
41. Tzeng TF, Liu WY, Liou SS, Hong TY, Liu IM. Antioxidant-rich extract from plantaginis semen ameliorates diabetic retinal injury in a streptozotocin-induced diabetic rat model. *Nutrients*. 2016;8(9):pii. E572.
42. Liu HW, Wei CC, Chang SJ. Low-molecular-weight polyphenols protect kidney damage through suppressing NF- $\kappa$ B and modulating mitochondrial biogenesis in diabetic db/db mice. *Food Funct*. 2016;7(4):1941–1949.
43. Davidson MA, Mattison DR, Azoulay L, Krewski D. Thiazolidinedione drugs in the treatment of type 2 diabetes mellitus: past, present and future. *Crit Rev Toxicol*. 2018;48(1):52–108.
44. Berger J, Moller DE. The mechanisms of action of PPARs. *Annu Rev Med*. 2002;53:409–435.
45. Cordero-Herrera I, Martín MA, Bravo L, Goya L, Ramos S. Cocoa flavonoids improve insulin signalling and modulate glucose production via AKT and AMPK in HepG2 cells. *Mol Nutr Food Res*. 2013;57(6):974–985.
46. Guillam MT, Hümmler E, Schaerer E, et al. Early diabetes and abnormal postnatal pancreatic islet development in mice lacking *Glut-2*. *Nat Genet*. 1997;17(3):327–330.
47. Burcelin R, Eddouks M, Kande J, Assan R, Girard J. Evidence that GLUT-2 mRNA and protein concentrations are decreased by hyperinsulinaemia and increased by hyperglycaemia in liver of diabetic rats. *Biochem J*. 1992;288(2):675–679.
48. Hsu CY, Shih HY, Chia YC, et al. Rutin potentiates insulin receptor kinase to enhance insulin-dependent glucose transporter 4 translocation. *Mol Nutr Food Res*. 2014;58(6):1168–1176.
49. Peng BJ, Zhu Q, Zhong YL, Xu SH, Wang Z. Chlorogenic acid maintains glucose homeostasis through modulating the expression of SGLT-1, GLUT-2, and PLG in different intestinal segments of Sprague-Dawley rats fed a high-fat diet. *Biomed Environ Sci*. 2015;28(12):894–903.
50. Shang X, Zhu ZY, Wang F, Liu JC, Liu JY, Xie ML. Hypoglycemic effect of *Chrysanthemum morifolium* extract on alloxan-induced diabetic mice is associated with peroxisome proliferator-activated receptor  $\alpha/\gamma$ -mediated hepatic glycogen synthesis. *J Appl Biomed*. 2017;15(1):81–86.
51. Heroor S, Beknal A, Mahurkar N. Synergistic activity of bark extracts of *Pongamia glabra* and *Ficus glomerata* in alloxan-induced diabetic rats. *World J Pharm Pharm Sci*. 2013;2(6):6640–6652.
52. Mitjavila MT, Moreno JJ. The effects of polyphenols on oxidative stress and the arachidonic acid cascade. Implications for the prevention/treatment of high prevalence diseases. *Biochem Pharmacol*. 2012;84(9):1113–1122.

## Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy

Dovepress

### Publish your work in this journal

Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy is an international, peer-reviewed open-access journal committed to the rapid publication of the latest laboratory and clinical findings in the fields of diabetes, metabolic syndrome and obesity research. Original research, review, case reports, hypothesis formation, expert

opinion and commentaries are all considered for publication. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/diabetes-metabolic-syndrome-and-obesity-targets-and-therapy-journal>