Co-production of MCR-I and NDM-5 in *Escherichia coli* isolated from a colonization case of inpatient

Siyuan Feng^{1,2,*}Cong Shen^{1,2,*}Hongtao Chen³Xiaobin Zheng⁴Yong Xia⁵Lan-Lan Zhong^{1,2}Xi Huang⁶Xinwei Wu⁷Guo-Bao Tian^{1,2}

¹Department of Immunology, Zhongshan School of Medicine, Sun Yat-sen University, Guangzhou, China; ²Key Laboratory of Tropical Diseases Control, Sun Yat-sen University, Ministry of Education, Guangzhou, China; ³Department of Laboratory, The Fifth Affiliated Hospital of Sun Yat-sen University, Zhuhai, China; ⁴Department of Respiratory Medicine, The Fifth Affiliated Hospital of Sun Yat-sen University, Zhuhai, China; ⁵Department of Clinical Laboratory Medicine, The Third Affiliated Hospital of Guangzhou Medical University, Guangzhou, China; ⁶Program of Pathobiology and Immunology, The Fifth Affiliated Hospital of Sun Yat-sen University, Zhuhai, China; ⁷Department of Microbiology, Guangzhou Center for Disease Control and Prevention, Guangzhou, China

*These authors contributed equally to this work

Correspondence: Guo-Bao Tian
Department of Immunology, Zhongshan School of Medicine, Sun Yat-sen University, 74 Zhongshan 2nd Road, Guangzhou 510080, China
Tel +86 208 733 5387
Fax +86 208 733 5387
Email tiangb@mail.sysu.edu.cn

Xinwei Wu
Department of Microbiology, Guangzhou Center for Disease Control and Prevention, No. 1, Qide Road, Guangzhou 510440, China
Email tomwu@126.com

Abstract: Colistin is increasingly used as an antibiotic of last resort for treating carbapenem-resistant Enterobacteriaceae. Mobile colistin resistance gene *mcr-1* has been increasingly reported in Enterobacteriaceae around the world. Of particular concern is the spread of *mcr-1* into carbapenemase-producing Enterobacteriaceae, which results in highly drug-resistant strains that are potentially untreatable. Notably, such *mcr-1*-carrying isolates harboring carbapenemase genes have been reported in animals and patients with infection. Here, we report an *Escherichia coli* strain carrying co-transferable *mcr-1*-harboring IncX4 and *bla*_{NDM-5}-harboring IncX3 plasmids, which was recovered in the context of fecal colonization.

Keywords: *Escherichia coli*, MCR-1, NDM-5, co-production

Introduction

Colistin is increasingly used as an antibiotic of last resort for treating multidrug-resistant Gram-negative bacteria, especially carbapenem-resistant Enterobacteriaceae. In December 2015, transmissible colistin resistance gene *mcr-1*, encoding phosphoethanolamine transferase, was reported from China. The gene *mcr-1* has since been increasingly reported in Enterobacteriaceae around the world.^{1,2} Of particular concern is spread of *mcr-1* into carbapenemase-producing Enterobacteriaceae, which results in highly drug-resistant strains that are potentially untreatable.

Methods and results

A total of 349 non-duplicate *Escherichia coli* clinical isolates were collected from serial fecal specimens in a hospital with capacity of 3,000 beds in Guangzhou, China in 2016 for the purpose of routine drug resistance monitoring. All isolates were characterized by API 20E system (BioMerieux, Marcy l'Etoile, France) and 16S rDNA sequencing. The *mcr-1* gene was detected using polymerase chain reaction (PCR) and sequencing.² As a result, 88 (25.2%) isolates were positive for *mcr-1*. This *mcr-1* positivity was higher than found in our previous study (11.3%),² indicating that the prevalence of *mcr-1* in *E. coli* is still increasing in this area. Susceptibility testing was performed for routine resistance observation; the results showed that one of the strains, *E. coli* GB788, was resistant to both colistin and carbapenem (Table 1). PCR and sequencing confirmed that *E. coli* GB788 carried both *mcr-1* and *bla*_{NDM-5} genes. *E. coli* GB788 was then subjected to phylogenetic typing and multilocus sequence typing for molecular epidemiology.² As a result, *E. coli* GB788 was classified as phylogenetic group A and sequence type 46 which has been sporadically identified among drug-resistant *E. coli* strains.³

Table 1 Characterization of the *Escherichia coli* GB788 and its transconjugants

Characteristics	<i>E. coli</i> GB788	Transconjugants		
		<i>E. coli</i> C600 GB788-CL/IMP	<i>E. coli</i> C600 GB788-CL	<i>E. coli</i> C600 GB788-IMP
Rescore	Inpatient			
Isolation site	Feces			
MLST	ST46			
Phylogenetic group	A			
Plasmid replicon type	IncX3, IncX4	IncX3, IncX4	IncX4	IncX3
Minimum inhibitory concentration (µg/mL)				
Colistin	16	16	16	0.25
Polymyxin B	16	16	16	0.5
Tigecycline	0.5	0.25	0.25	0.5
Ampicillin	>256	>256	64	>256
Amoxicillin–clavulanic acid	>256	256	8	16
Cefotaxime	128	64	≤1	64
Ceftazidime	>256	>256	≤1	>256
Cefepime	32	8	≤0.5	8
Gentamicin	16	≤1	≤1	≤1
Amikacin	4	4	2	2
Ertapenem	16	8	≤0.125	8
Imipenem	16	4	0.25	4
Meropenem	16	8	≤0.125	8
Fosfomycin	64	16	16	32
Nitrofurantoin	32	≤8	≤8	≤8
Ciprofloxacin	1	≤0.03	≤0.03	≤0.03

Abbreviations: MLST, multilocus sequence typing; CL, colistin; IMP, imipenem.

E. coli GB788 was recovered from a 52-year-old female patient, who was admitted to the hospital after suffering from periarthritis shoulder. The patient did not have any signs of infection, was not receiving any antibiotics, and did not have any significant travel history.

Streptomycin-resistant *E. coli* C600 was used as the recipient for the conjugation experiment to identify the mobility of drug resistance genes. Three different types of transconjugants (*E. coli* C600 GB788-colistin (CL), *E. coli* C600 GB788-imipenem (IMP), and *E. coli* C600 GB788-CL/IMP) were obtained, including those harboring *mcr-1*, *bla*_{NDM-5}, or both (Table 1), with 6–7 of 15 colonies positive for both genes. For detecting the stability of these plasmids, plasmid stability experiment was performed. Three independent lineages of each plasmid-containing strain were grown overnight at 37°C in 1 mL of Luria-Bertani (LB) broth. Bacterial cells were washed by spinning down overnight cultures and resuspending the pellet in 1 mL of LB without antibiotics. An aliquot of 1 µL of washed cells was transferred into 1 mL of LB and incubated overnight at 37°C. Serial passaging of 1 µL of overnight culture to 1 mL of LB was performed daily, approximating ten generations of growth per passage. Every 20 generations, samples were diluted and plated on LB plates. Then, 50 colonies from each lineage were screened on LB plates supplemented without colistin to determine the fraction of plasmid-containing cells. The results showed

that no plasmid was lost even after 100 generations for all three types of transconjugants, which indicated that *mcr-1* and *bla*_{NDM-5} can transfer alone or co-transfer at a certain frequency and subsequent stability.⁴ PCR-based replicon typing was performed to detect the plasmid type, which was determined by a PCR-based replicon typing method. Plasmid DNA was amplified by five multiplex and three simplex PCRs using 18 pairs of primers as reported previously that are recognized as Inc (incompatibility) replicon types: FIA, FIB, FIC, HI1, HI2, I1-Ic, L/M, N, P, W, T, A/C, K, B/O, X, Y, F, and FIIA. Pulsed field gel electrophoresis analysis of S1 nuclease-digested DNA (S1-PFGE), followed by southern blotting were conducted for identifying the location of *mcr-1* and *bla*_{NDM-5} carrying plasmid. Whole-cell DNA of all *mcr-1*-producing isolates was extracted and embedded in gold agarose gel plugs (SeaKem® Gold Agarose; Lonza, Rockland, ME, USA). The plugs were digested with S1 nuclease (TaKaRa, Dalian, China), and the DNA fragments were separated by PFGE. Southern blotting hybridizations of plasmid DNA were performed with a digoxin-labeled *mcr-1* probe according to the manufacturer's instructions (Roche Diagnostics, Mannheim, Germany). The results showed that *mcr-1* was located on ~33 kb IncX4 plasmid, while *bla*_{NDM-5} was located on ~45 kb IncX3 plasmid (Figure S1). IncX4 and IncX3 are the common host-range plasmids for *mcr-1* and *bla*_{NDM-5}, respectively.

For detailed genome characteristics of *E. coli* GB788, whole genome sequencing was performed. The genome DNA of *E. coli* GB788 was extracted and DNA libraries were constructed with 350 and 6,000 bp DNA fragments. With Illumina Hiseq4000 platform, 13,531,019 50-bp paired-end reads were produced from 6,000 bp DNA fragment library, while 5,333,992 150-bp paired-end reads were produced from 350 bp library. Reads were assembled with SPAdes.⁵ The predicted open reading frames were annotated using the gene ontology, Kyoto Encyclopedia of Genes and Genomes, Swiss-Prot, non-redundant, and cluster of orthologous groups of proteins databases. Assembled contigs were aligned with the reference genome of *E. coli* strain K-12 substrain MG1655 (NC_000913) by using BLASTp and BLASTn. As a result, several genomic insertion and deletion segments were identified in *E. coli* GB788 (Figure S2A). Forty-five resistance genes including *mcr-1* and *bla*_{NDM-5} were identified according to the Comprehensive Antibiotic Resistance Database⁶ using identity matches of >60% and E values of <1e-5 (Figure S2B). This Whole Genome Shotgun project has been deposited at DDBJ/ENA/GenBank under the accession NEFV000000000. The version described in this paper is version NEFV01000000.

To obtain the complete sequence of *mcr-1* and *bla*_{NDM-5}-harboring plasmids, pGB788-CL and pGB788-IMP were extracted from transconjugants. The DNA libraries were constructed with 350 bp DNA fragments. Thus, 1,201,548 and 1,236,894 150-bp paired-end reads were produced with Illumina Hiseq4000 platform for the two plasmids, respectively. The scaffolds were assembled with SPAdes,⁵ and the scaffold with targeted genes of *mcr-1* and *bla*_{NDM-5} was selected as the candidates. Assembly results showed that the size of *mcr-1*-harboring plasmid pGB788-CL and *bla*_{NDM-5}-harboring plasmid GB788-IMP were 30,988 and 45,793 bp, respectively, which were consistent with the results of plasmid size determination by S1-PFGE and southern blotting (Figure S3).

Conclusion

Although the coexistence of *bla*_{NDM-5} and *mcr-1* has been reported in infection cases from specimens such as bloodstream, urinary tract, kidney, bile, and liver infections, this study is the first to report such an occurrence from a fecal specimen of a colonized individual. Emergence of coexistence of *mcr-1* and carbapenemase genes in Enterobacteriaceae may seriously compromise the effectiveness of antimicrobial therapy. Therefore, close monitoring and routine investigations are required to establish measures to control dissemination of such strains in both infected and colonized individuals. The clinical isolate samples used in this research were part of the routine hospital laboratory procedure. Oral consent of the patients was obtained.

Disclosure

The authors report no conflicts of interest in this work.

References

1. Wang Y, Tian GB, Zhang R, et al. Prevalence, risk factors, outcomes, and molecular epidemiology of *mcr-1*-positive Enterobacteriaceae in patients and healthy adults from China: an epidemiological and clinical study. *Lancet Infect Dis*. 2017;17(4):390–399.
2. Zhong LL, Zhang YF, Doi Y, et al. Coproduction of MCR-1 and NDM-1 by colistin-resistant *Escherichia coli* isolated from a healthy individual. *Antimicrob Agents Chemother*. 2017;61(1):e01962-16.
3. Vignaroli C, di Sante L, Leoni F, et al. Multidrug-resistant and epidemic clones of *Escherichia coli* from natural beds of Venus clam. *Food Microbiol*. 2016;59:1–6.
4. Sandegren L, Linkevicius M, Lytsy B, Melhus Å, Andersson DI. Transfer of an *Escherichia coli* ST131 multiresistance cassette has created a *Klebsiella pneumoniae*-specific plasmid associated with a major nosocomial outbreak. *J Antimicrob Chemother*. 2012;67(1):74–83.
5. Nurk S, Bankevich A, Antipov D, et al. Assembling single-cell genomes and mini-metagenomes from chimeric MDA products. *J Comput Biol*. 2013;20(10):714–737.
6. Jia B, Raphenya AR, Alcock B, et al. CARD 2017: expansion and model-centric curation of the comprehensive antibiotic resistance database. *Nucleic Acids Res*. 2017;45(D1):566–573.

Supplementary material

Figure S1 Location of *mcr-1* and *bla*_{NDM-5} in *Escherichia coli* GB788.

Notes: (A) S1-PFGE analysis. (B) Southern hybridization of with the *mcr-1* and *bla*_{NDM-5} probe, respectively. Lane M, DNA ladder (New England Bio-Labs, Beverly, MA, USA); lane 1, *E. coli* GB788; lane 2, *E. coli* C600 GB788-CL; lane 3, *E. coli* C600 GB788-IMP; lane 4, *E. coli* C600 GB788-CL/IMP.

Abbreviations: S1-PFGE, pulsed field gel electrophoresis analysis of S1 nuclease-digested DNA; CL, colistin; IMP, imipenem.

Figure S2 Genomic maps and resistance genes of *Escherichia coli* GB788.

Notes: (A) Comparison of GB788 and reference sequence NC_000913. Orange bands represent assembly result of GB788 that has similar sequence with reference sequence. Outer bands with different colors represent different COG function annotations. (B) Resistance genes distribution in GB788 genome. Different color bands represent different COG function annotations.

Abbreviation: COG, cluster of orthologous groups of proteins.

Figure S3 Structure map of (A) pGB788-IMP and (B) pGB788-CL.

Note: Plasmid analysis of pGB788-IMP and pGB788-CL using GCview plasmid sequencing.

Abbreviations: CDS, coding sequence; GC, guanine-cytosine; CL, colistin; IMP, imipenem.

Infection and Drug Resistance

Publish your work in this journal

Infection and Drug Resistance is an international, peer-reviewed open-access journal that focuses on the optimal treatment of infection (bacterial, fungal and viral) and the development and institution of preventive strategies to minimize the development and spread of resistance. The journal is specifically concerned with the epidemiology of antibiotic

Submit your manuscript here: <https://www.dovepress.com/infection-and-drug-resistance-journal>

resistance and the mechanisms of resistance development and diffusion in both hospitals and the community. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Dovepress