High Spy1 expression predicts poor prognosis in colorectal cancer

Qin Jin^{1,*}
Gang Liu^{2,*}
Luri Bao^{3,*}
Yuzhen Ma⁴
Huidong Qi⁵
Zhizhong Yun⁴
Yanfeng Dai²
Shu Zhang¹

¹Department of Pathology, Affiliated Hospital of Nantong University, Nantong, People's Republic of China; ²State Key Laboratory of Reproductive Regulation and Breeding of Grassland Livestock, College of Life Science, Inner Mongolia University, Hohhot, People's Republic of China; ³Department of Pathology, Inner Mongolia Medical University, Hohhot, People's Republic of China; ⁴Centre of Reproductive Medicine, Inner Mongolia Hospital, Hohhot, People's Republic of China; ⁵Medical School of Nantong University, Nantong, People's Republic of China

*These authors contributed equally to this work

Correspondence: Gang Liu
State Key Laboratory of Reproductive Regulation and Breeding of Grassland Livestock, College of Life Science, Inner Mongolia University, 235 West Univ. Road, Hohhot, Inner Mongolia 010021, People's Republic of China
Tel +86 471 499 2944
Email liugang@life.imu.edu.cn

Shu Zhang
Department of Pathology, Affiliated Hospital of Nantong University, 20 Xisi Road, Nantong 226001, Jiangsu, People's Republic of China
Tel +86 513 8505 2222
Email tzyfzs@163.com

Background: Spy1 (SPDYA) is a new discovered cell cycle protein capable of promoting cell proliferation dependent on cyclin-dependent kinase-2 activation. However, to the best of our knowledge, the expression of Spy1 in colorectal cancer (CRC) tissues remains virtually unknown.

Materials and methods: In this retrospective study, we investigated the mRNA and protein expression levels of Spy1 in CRC tissues and corresponding non-cancerous tissues with the analyses of quantitative real-time polymerase chain reaction, western blotting, and immunohistochemistry. In our research, the prognostic significances of Spy1 expression were further explored by univariate and multivariate survival analyses of 203 patients who were followed up.

Results: The results demonstrated that the levels of *Spy1* mRNA were significantly higher in CRC tissues compared with corresponding non-cancerous tissues ($p=0.0002$). The results of immunohistochemistry demonstrated that the expressions of Spy1 were significantly associated with clinicopathological parameters, including T stage ($\chi^2=7.126$, $p=0.028$) and TNM stage ($\chi^2=9.461$, $p=0.009$). Kaplan-Meier analysis results indicated that high Spy1 expression (HR=2.573, $p<0.001$) and TNM stage (HR=1.494, $p=0.011$) were independent factors to predict poor prognosis for patients with CRC.

Conclusion: We concluded that high Spy1 expression is significantly associated with unfavorable prognosis in CRC and could serve as a potential prognostic marker in clinical diagnosis of CRC.

Keywords: Spy1, colorectal cancer, prognostic marker, clinical diagnosis

Introduction

Colorectal cancer (CRC) is the third most common cancer worldwide, and is also associated with the second highest malignancy mortality worldwide. In 2014, there were projected to be 136,830 individuals newly diagnosed with CRC and 50,310 CRC deaths in the United States, with over 2.4 million new cases expected to be diagnosed worldwide by 2035.^{1,2} The incidence of CRC is increasing each year, with decreasing age of CRC patients. Therefore, CRC has become a major public health issue.

The fecal occult blood test (FOBT)³ and serum diagnostic biomarkers such as carcinoembryonic antigen⁴ and carbohydrate antigen 19-9⁵ are used for CRC diagnosis in clinics, but the accuracies of the FOBT and these biomarkers remain unsatisfactory and require improvement.⁶ Alternatively, colonoscopy could significantly improve the diagnosis of CRC.⁷ However, this diagnostic method has shortcomings, such as cost, risk, and inconvenience.⁸ CT colonography is highly sensitive for CRC detections in clinical, but there is still some limitations such as heterogeneous sensitivity.⁹ Therefore, it is vital to discover new CRC biomarkers with high specificity and sensitivity.

Cell cycle progression is regulated by sequential activation and inactivation of a series of cyclin-dependent kinases (CDKs).¹⁰ Spy1 (speedy/RINGO cell cycle regulator family member A; also known as SPDYA) is a member of the Speedy/Ringo families of cell cycle regulators, which control cell proliferation and survival through the atypical activations of CDKs.¹¹ The expression of Spy1 enhances CDK2-dependent p27^{kip1} degradation during late G1 phase and throughout S phase by directly promoting p27 degradation.¹² Furthermore, Spy1 has been shown to override DNA damage response,¹³ functions to inhibit DNA damage-induced apoptosis, and promote mammalian cell survival in response to a number of genotoxic agents, including hydroxyurea, cisplatin, and camptothecin.¹⁴ Spy1 expression has been associated with breast cancer,^{15,16} ovarian cancer,¹⁷ hepatocellular carcinoma,¹⁸ and multiple myeloma.¹⁹ However, to the best of our knowledge, the expression of Spy1 in CRC remains virtually unknown. Therefore, in the present study, we compared the expression of Spy1 protein in CRC tissues and adjacent tissues (as corresponding non-cancerous tissues) using quantitative real-time polymerase chain reaction (qRT-PCR), immunohistochemistry, and western blot analysis. We also investigated its associations with clinical and pathologic factors, as well as the prognostic implications.

Materials and methods

Patients

In total, 40 pairs of fresh CRC tissues and adjacent tissues were collected from the Department of Pathology, Affiliated Hospital of Nantong University. Simultaneously, a total of 203 paraffin-embedded CRC tissues and 203 matched adjacent tissues were collected from the Department of Pathology, Affiliated Hospital of Nantong University between January 2009 and June 2011. Diagnoses of CRC were validated by two independent pathologists in the department, according to the International Union Against Cancer and the TNM Classification of Malignant Tumors 7th edition. None of the included patients received radiotherapy or chemotherapy before surgery. Each patient signed written informed consent for this present study. All cases had corresponding clinical data and follow-up records. The follow-up rate was 100%. The study protocol was approved by the Ethics Committee of the Affiliated Hospital of Nantong University, and all experiments were performed in accordance with approved guidelines of the Affiliated Hospital of Nantong University.

qRT-PCR analyses

In total, 30 pairs of the collected fresh CRC tissues and adjacent tissues were used for qRT-PCR analysis.

Total RNA was extracted from tissues with TrizolTM (79306, Gibco, Shanghai, People's Republic of China), and cDNA was synthesized using commercial kits (Prime ScriptTM RT reagent Kit with gDNA Eraser, RR047A; TaKaRa, Beijing, People's Republic of China). Amplifications were performed in an ABI Verti 96 well thermal cycler or a Thermo Pikoreal using commercial kits (Premix TaqTM; TaKaRa) and specific synthesized primers. The ubiquitously expressed β -actin gene was used as an internal control. The integrity of all the DNA fragments amplified by PCRs was confirmed by sequencing. All experiments were performed in triplicate.

The primers for *Spy1* were as follows: forward primer 5'-ATT GGG AAA CCA AAA TGA GGC-3' and reverse primer 5'-TCC TGG TAT GCT CAC TTA TAG-3'. β -actin was used as an internal control, and the primers for β -actin were as follows: forward primer 5'-TAA TCT TCG CCT TAA TAC TT-3' and reverse primer 5'-AGC CTT CAT ACA TCT CAA-3'. The relative *Spy1* mRNA expression was calculated using the $2^{-\Delta\Delta C_t}$ method.

Antibodies and western blotting

Quantified protein lysates from 40 pairs of fresh CRC tissues and adjacent tissues were measured with a Protein BCA assay kit (163-2086; Bio-Rad Laboratories Inc., Hercules, CA, USA) according to manufacturer's instructions. The proteins in lysates were resolved on SDS-PAGE gels (P1200; Solarbio, Beijing, People's Republic of China), transferred onto polyvinylidene fluoride membranes (IPVH00010; EMD Millipore, Billerica, MA, USA), and immunoblotted with anti-human antibodies to GAPDH (ab8245; Abcam, Shanghai, People's Republic of China) and Spy1 (ab153965; Abcam). The blots were visualized with enhanced chemiluminescence (Amersham Biosciences, Shanghai, People's Republic of China) after exposure to X-ray films. GAPDH was stained as a loading control.

Immunohistochemistry analysis

A total of 203 CRC and adjacent tissue paraffin sections were deparaffinized with xylene and rehydrated in graded ethanol (100-95-85-75%), and then washed with phosphate-buffered saline solution (PBS, 0.01 M, pH 7.0). Antigen retrieval was achieved by boiling under pressure in citrate buffer (0.01 M, pH 6.0). Non-specific binding was blocked through incubation with 5% goat blocking serum (SL039; Solarbio) in PBS for 30 min at 37°C. All paraffin sections were incubated with rabbit anti-Spy1 antibody (1:300, ab153965; Abcam) and subsequently with goat anti-rabbit HRP secondary antibody (ZDR-5306; ZSGB-BIO, Beijing, People's Republic of China). For a color reaction, sections were incubated with the

DAB substrate chromogen solution (DA1010; Solarbio) for 8 minutes. Subsequently, sections were counterstained with hematoxylin.

Immunostained sections were scored by two independent pathologists under blinded experimental conditions according to intensity and percentage of *Spy1*-positive cells under light microscope (Axio imager A2; Zeiss, Shanghai, People's Republic of China). The intensity of *Spy1*-positive cells was scored as follows: 0 (0%, negative), 1 (1%–33%, poor), 2 (34%–66%, moderate), and 3 ($\geq 67\%$, strong).²⁰ The multiplication of the intensity and percentage scores led to the final *Spy1* staining score and was defined as follows: staining score less than 3 was considered as low expression, while staining score of 4 or more was considered as high expression.

Comparison of *Spy1* genes expression level in different cancers

The GEPIA (<http://gepia.cancer-pku.cn/index.html>) is a newly developed interactive web server for analyzing the RNA sequencing expression data of 9,736 tumors and 8,587 normal samples from the TCGA and the GTE projects using a standard processing pipeline.²¹ It provides customizable functions such as tumor and normal differential expression analysis, and we could demonstrate the *Spy1* gene expression levels in different cancers such as adrenocortical carcinoma (ACC), kidney renal clear cell carcinoma (KIRC) and colon adenocarcinoma (COAD).

Statistical analysis

The expression levels of *Spy1* mRNA in fresh CRC tissues and adjacent tissues were calculated by the Wilcoxon signed-rank nonparametric test. We used Pearson's χ^2 to examine the correlations between *Spy1* protein expression and clinicopathological parameters. Kaplan-Meier and log-rank test were performed to calculate the survival curves. Factors of prognostic significance in the univariate analysis were further analyzed by the model of multivariate Cox regression. For all tests, *p*-values less than 0.05 were considered significant. All statistical data were analyzed by the Statistical Package for the Social Sciences software (SPSS, IBM, version 19.0). All data are represented as mean $\pm$ SEM.

Results

Spy1 mRNA expression in CRC tissues by qRT-PCR analysis and western blotting

The mRNA expression levels of *Spy1* were detected by qRT-PCR analyses of 30 pairs of CRC tissues and their corresponding adjacent tissues. Compared with the transcript levels in adjacent tissues, the transcript levels of *Spy1* were

significantly higher in CRC tissues (0.8157 ± 0.1289 vs 0.6993 ± 0.1149 , 1.95-fold, $p=0.0002$, with Wilcoxon signed-rank nonparametric test) (Figure 1).

The expression of *Spy1* protein in the CRC tissues and adjacent tissues was confirmed by western blot analysis. The representative results are shown in Figure 2. There were 27 cases (90%) of CRC tissues with higher protein level of *Spy1* and two cases (6.67%) of CRC tissues with similar expression level of *Spy1* protein, compared with the adjacent tissues. There was one case (3.33%) of CRC tissue with lower protein level of *Spy1* than that of adjacent tissue. The results indicated that the expressions of *Spy1* protein in CRC tissues were significantly higher than those in adjacent tissues.

The location and expression of *Spy1* protein in CRC tissues by immunohistochemistry

The immunohistochemistry analyses were performed to determine the protein location and expression of *Spy1* in CRC

Figure 1 Quantitative real-time polymerase chain reaction (qRT-PCR) analysis was used to detect *Spy1* mRNA expression levels in colorectal cancer (CRC) tissues and the adjacent tissues.

Note: The *Spy1* mRNA levels in CRC tissues were significantly higher than those in the adjacent tissues.

Figure 2 The representative *Spy1* expression levels in colorectal cancer (CRC) tissues were detected by western blot assay and compared with adjacent tissues.

Note: The *Spy1* levels in CRC tissue were significantly overexpressed compared to those in the adjacent tissues, and GAPDH was used as a loading control.

Abbreviations: A, adjacent tissues; C, cancer tissues.

Figure 3 The representative images of the Spy1 protein location and expression in colorectal cancer (CRC).

Notes: (A) No immunohistochemistry (IHC) staining of Spy1 in the cell nucleus of normal intestinal mucosa. (B) IHC staining of Spy1 in the nuclear of normal intestinal mucosa. (C) No IHC staining of Spy1 in the nuclear of CRC cells. (D) Moderate IHC staining of Spy1 in the nuclear of CRC cells. (E) High IHC staining of Spy1 in the nuclear of CRC cells. Original magnification, $\times 400$ in A–E.

tissues and adjacent tissues. As shown in Figure 3, the location of Spy1 expression was mainly detected in the cell nucleus of CRC tissues. High Spy1 expression level was detected in 61.08% (124/203) of CRC tissues, which was significantly higher ($\chi^2=19.951$, $p<0.05$) than that in adjacent tissues (38.92% [79/203]).

Correlations between Spy1 protein expression and clinicopathological attributes of CRC patients

The correlations between Spy1 protein levels and clinicopathological attributes of CRC patients were investigated (Table 1). Higher expressions of Spy1 were significantly associated with T stage ($\chi^2=7.126$, $p=0.028$) and TNM stage ($\chi^2=9.461$, $p=0.009$). However, Spy1 protein expressions in CRC were not significantly associated with age, sex, tumor size, histological type, location, and differentiation (Table 1).

High Spy1 protein expression predicts poor prognosis of CRC patients

Subsequently, univariate analyses were conducted to evaluate Spy1 protein expressions and clinicopathological factors on prognosis of CRC patients. The result showed that the high expression level of Spy1 (HR=2.846, $p<0.001$), T stage (HR=1.335, $p=0.026$), and TNM stage (HR=1.609, $p=0.003$) were significantly associated with poor overall survival

Table 1 Association of Spy1 expression with clinical characteristics and selected biological markers of CRC

Characteristic	n	Spy1 expression (%)		χ^2	p
		High	Low or no		
Age (years)				1.293	0.255
≤60	75	42 (56.0)	33 (44.0)		
>60	128	82 (64.1)	46 (35.9)		
Sex				0.048	0.826
Male	124	75 (60.5)	49 (39.5)		
Female	79	49 (62.0)	30 (38.0)		
Tumor size (cm)				0.096	0.757
≤5	157	95 (60.5)	62 (39.5)		
>5	46	29 (63.0)	17 (37.0)		
Histological classification				0.523	0.470
Tubular+papillary	181	109 (60.2)	72 (39.8)		
Other	22	15 (68.2)	7 (31.8)		
Location				4.642	0.200
Ascending colon	66	42 (63.6)	24 (36.4)		
Transverse colon	27	18 (66.7)	9 (33.3)		
Descending colon+sigmoid colon	60	30 (50.0)	30 (50.0)		
Rectum	50	34 (68.0)	16 (32.0)		
Differentiation				0.284	0.594
Well+moderate	150	90(60.0)	60(40.0)		
Poor	53	34(64.2)	19(35.8)		
T stage				7.126	0.028*
T1	45	23 (51.1)	22 (48.9)		
T2	87	49 (56.3)	38 (43.7)		
T3+T4	71	52 (73.2)	19 (26.8)		

(Continued)

Table 1 (Continued)

Characteristic	n	Spy1 expression (%)		χ^2	p
		High	Low or no		
N stage				5.334	0.069
N0	122	78 (63.9)	44 (36.1)		
N1	56	36 (64.3)	20 (35.7)		
N2	25	10 (40.0)	15 (60.0)		
TNM stage				9.461	0.009*
I	24	10 (41.7)	14 (58.3)		
II	53	27 (50.9)	26 (49.1)		
III+IV	136	87 (69.0)	39 (31.0)		

Note: * $p < 0.05$.

Abbreviation: CRC, colorectal cancer.

of CRC patients (Table 2). These prognostic factors were further evaluated by multivariate Cox proportional hazards regression model analysis. With the Kaplan-Meier analysis, high Spy1 expression (HR=2.573, $p < 0.001$) and TNM stage (HR=1.494, $p = 0.011$) were found to be independent prognostic markers of poor 5-year overall survival (Table 2). CRC patients with high expression of Spy1 protein had significantly shorter overall survival compared with those with low or no Spy1 expression (Figure 4). CRC patients with TNM stage III+IV had significantly shorter overall survival compared with those with stage I and stage II (Figure 4).

Survival analysis of target genes

High expressions of Spy1 confer survival disadvantages to ACC patients (log rank $p = 0.0062$, HR [high]=3), COAD patients (log rank $p = 0.03$, HR [high]=1.7) and KIRC patients [(log rank $p = 2.3 \times 10^{-7}$, HR [high]=2.3) in the GEPIA dataset with TCGA samples (Figure 5).

Discussion

CRC is the most malignant disease in developed and developing countries, largely because of its propensity to metastasize. The therapeutic regimens of fluoropyrimidine chemotherapy (parenteral 5-fluorouracil/leucovorin or oral capecitabine) in combination with oxaliplatin (FOLFOX or XELOX) or irinotecan (FOLFIRI) and bevacizumab (Avastin; Genentech, Inc., South San Francisco, CA, USA) have improved overall survival of CRC.²² Furthermore, targeted treatments like anti-VEGF, anti VEGFR, or anti-EGFR depending on the all-RAS status^{23,24} and anti-PDL-1 (pembrolizumab) according to the microsatellite instability (MSI) status and PDL-1 expression^{25,26} have been used for clinical trials. However, the therapeutic responses vary individually. Additional genetic targeted therapies for CRC still need exploration.²⁷

Tumor cells accumulate alterations that result in constitutive mitogenic signaling and defective responses to anti-mitogenic signals that contribute to unscheduled

Table 2 Univariate and multivariate analysis of prognostic factors in CRC for 5-year overall survival

	Univariate analysis			Multivariate analysis		
	HR	p-value	95% CI	HR	p-value	95% CI
Expression of Spy1						
High vs low or no expression	2.846	<0.001*	1.828–4.429	2.573	<0.001*	1.647–4.019
Age (years)						
≤60 vs >60	1.031	0.879	0.696–1.527			
Sex						
Male vs female	0.874	0.501	0.592–1.292			
Tumor size (cm)						
≤5 vs >5	1.133	0.584	0.724–1.772			
Histological classification						
Tubular+papillary vs other	0.897	0.732	0.481–1.673			
Location						
Ascending colon vs transverse colon vs descending colon + sigmoid colon	0.873	0.103	0.742–1.028			
Differentiation						
Well+moderate vs poor	0.765	0.168	0.522–1.120			
T stage				1.266	0.075	0.977–1.641
T1 vs T2 vs T3+T4	1.335	0.026*	1.034–1.723			
N stage						
N0 vs N1 vs N2	1.203	0.201	0.906–1.597			
TNM stage						
I vs II vs III+IV	1.609	0.003*	1.178–2.197	1.494	0.011*	1.096–2.037

Note: * $p < 0.05$.

Abbreviation: CRC, colorectal cancer.

Figure 4 The survival analysis of colorectal cancer (CRC) patients by Kaplan-Meier analysis.

Notes: Overall survival rates in CRC patients with high expression level of Spy1 (green line) were statistically lower than those in CRC patients with low or no Spy1 expression (blue line). Overall survival rates in CRC patients with high expression levels of TNM stage III+IV (yellow line) were statistically lower than those in CRC patients with TNM stage I (blue line) and II (green line).

Figure 5 Expression of *Spy1* in cancers.

Notes: Overall survival rates in adrenocortical carcinoma (ACC), colon adenocarcinoma (COAD) and kidney renal clear cell carcinoma (KIRC) patients with high expression levels of *Spy1* (red line) were statistically lower than those in patients with low and no *Spy1* expression (blue line).

Abbreviations: SPADY, *Spy1*; TPM, transcripts per kilobase of exonmodel per million mapped reads; HR, hazard ratio.

proliferation. These mutations, taken together, result not only in proliferative advantages but also in increased susceptibility to the accumulation of additional genetic alterations that contribute to tumor progression and acquisition of more aggressive phenotypes.

In addition, most tumors acquire genomic instability (GIN) that leads to additional mutations. The acquisition of GIN is a crucial feature in tumor development, and there are at least three distinct pathways in CRC: the CIN, MSI, and CpG island methylator phenotype pathways.^{28,29} Most cases of CRC arise through the CIN pathway by widespread imbalances in chromosome number (aneuploidy) and loss of heterozygosity due to the defects in chromosomal segregation, telomere stability, and the DNA damage response.

The basic cell cycle defects such as GIN or CIN are mediated, directly or indirectly, by misregulations of CDKs. During the past two decades, a vast body of literature has illustrated the relevance of cell cycle deregulation in human cancer, and targeting the cell cycle in general and CDK in particular presents unique opportunities for drug discovery.³⁰

Firstly, isolated in *Xenopus*, *Xenopus-Spy1* could induce rapid meiotic maturation and premature activation of CDKs.^{15,31} Subsequently, *Spy1* was found to be expressed in a variety of human tissues, and its overexpression enhances G1-S phase progression, p27^{kip} degradation,¹² and cellular proliferation by activating CDK2.^{32,33} *Spy1* is also required for cell-cycle re-entry in maturing vertebrate oocytes.³⁴ There are multiple isoforms of *Spy1* that are now classified into a larger family of proteins, known as the *Spy1*/RINGO family. Furthermore, abrogation of *Spy1* expression through the use of siRNA hinders cell proliferation, thus demonstrating that *Spy1* is an essential

component of cell proliferation pathways.¹⁸ These findings open up new possibilities that this gene may be involved in human cell cycle disorders such as oncogenesis.³⁵ To date, no study has been performed for *Spy1* expression using the group of CRC specimens, and the present study is the first to analyze the *Spy1* expression in CRC, with respect to possible associations with clinicopathological data as well as patients' poor prognosis.

In this study, the expression levels of *Spy1* in CRC tissues, with adjacent tissues, were detected with the observation that both *Spy1* mRNA and protein levels were significantly higher in CRC tissues than those in the adjacent tissues. High *Spy1* protein level was associated with poor overall survival for CRC patients, which indicated that *Spy1* protein level may be used as an independent clinical marker in predicting unfavorable prognosis.

GEPIA is an interactive web application for gene expression analysis based on 9,736 tumors and 8,587 normal samples from the TCGA and the GTE databases.²¹ Using the GEPIA datasets, we have revealed that *Spy1* expression levels are positively correlated with the survival of ACC, COAD, and KIRC cancer patients. Combined with the expression model, we predict that *Spy1* plays an important role in cancer progression and may be a candidate biomarker.

Taken together, we analyzed the *Spy1* expression in CRC and demonstrated that *Spy1* was involved in CRC development. The findings indicated that *Spy1* may act as an oncogene during CRC development and it could be identified as a novel prognostic biomarker in CRC clinical treatment. Related studies concerning the potential mechanisms of *Spy1* in CRC and other kinds of human cancers are being performed by our research group.

Acknowledgment

This work was supported by the Inner Mongolia Science and Technology Research Project (2017ZD04 to Yanfeng Dai and 201502107 and 2017MS08137 to Yuzhen Ma).

Disclosure

The authors report no conflicts of interest in this work.

References

1. Siegel R, Desantis C, Jemal A. Colorectal cancer statistics, 2014. *CA Cancer J Clin*. 2014;64(2):104–117.
2. Day LW, Velayos F. Colorectal cancer screening and surveillance in the elderly: updates and controversies. *Gut Liver*. 2015;9(2):143–151.
3. Zavoral M, Suchanek S, Majek O, et al. Colorectal cancer screening: 20 years of development and recent progress. *World J Gastroenterol*. 2014;20(14):3825–3834.
4. Bajenova O, Chaika N, Tolkunova E, et al. Carcinoembryonic antigen promotes colorectal cancer progression by targeting adherens junction complexes. *Exp Cell Res*. 2014;324(2):115–123.
5. Wang YR, Yan JX, Wang LN. The diagnostic value of serum carcinoembryonic antigen, alpha fetoprotein and carbohydrate antigen 19-9 for colorectal cancer. *J Cancer Res Ther*. 2014;10 Suppl:307–309.
6. Shibutani M, Maeda K, Nagahara H, et al. Significance of CEA and CA19-9 combination as a prognostic indicator and for recurrence monitoring in patients with stage II colorectal cancer. *Anticancer Res*. 2014;34(7):3753–3758.
7. Brenner H, Stock C, Hoffmeister M. Effect of screening sigmoidoscopy and screening colonoscopy on colorectal cancer incidence and mortality: systematic review and meta-analysis of randomised controlled trials and observational studies. *BMJ*. 2014;348:g2467.
8. Byun JY, Yoon SJ, Oh IH, Kim YA, Seo HY, Lee YH. Economic burden of colorectal cancer in Korea. *J Prev Med Public Health*. 2014;47(2):84–93.
9. Pickhardt PJ, Hassan C, Halligan S, Marmo R. Colorectal cancer: CT colonography and colonoscopy for detection—systematic review and meta-analysis. *Radiology*. 2011;259(2):393–405.
10. Lim S, Kaldis P. Cdks, cyclins and CKIs: roles beyond cell cycle regulation. *Development*. 2013;140(15):3079–3093.
11. McAndrew CW, Gastwirt RF, Donoghue DJ. The atypical CDK activator Spy1 regulates the intrinsic DNA damage response and is dependent upon p53 to inhibit apoptosis. *Cell Cycle*. 2009;8(1):66–75.
12. McAndrew CW, Gastwirt RF, Meyer AN, Porter LA, Donoghue DJ. Spy1 enhances phosphorylation and degradation of the cell cycle inhibitor p27. *Cell Cycle*. 2007;6(15):1937–1945.
13. Ferraiuolo RM, Tubman J, Sinha I, Hamm C, Porter LA. The cyclin-like protein, SPY1, regulates the ER α and ERK1/2 pathways promoting tamoxifen resistance. *Oncotarget*. 2017;8(14):23337–23352.
14. Barnes EA, Porter LA, Lenormand JL, Dellinger RW, Donoghue DJ. Human Spy1 promotes survival of mammalian cells following DNA damage. *Cancer Res*. 2003;63(13):3701–3707.
15. Al Sorkhy M, Ferraiuolo RM, Jalili E, et al. The cyclin-like protein Spy1/RINGO promotes mammary transformation and is elevated in human breast cancer. *BMC Cancer*. 2012;12:45.
16. Zucchi I, Mento E, Kuznetsov VA, et al. Gene expression profiles of epithelial cells microscopically isolated from a breast-invasive ductal carcinoma and a nodal metastasis. *Proc Natl Acad Sci U S A*. 2004;101(52):18147–18152.
17. Lu S, Liu R, Su M, et al. Spy1 participates in the proliferation and apoptosis of epithelial ovarian cancer. *J Mol Histol*. 2016;47(1):47–57.
18. Ke Q, Ji J, Cheng C, et al. Expression and prognostic role of Spy1 as a novel cell cycle protein in hepatocellular carcinoma. *Exp Mol Pathol*. 2009;87(3):167–172.
19. Fei M, Hang Q, Hou S, Ruan C. Cell adhesion to fibronectin down-regulates the expression of Spy1 and contributes to drug resistance in multiple myeloma cells. *Int J Hematol*. 2013;98(4):446–455.
20. Patel SH, Kneuert PJ, Delgado M, et al. Clinically relevant biomarkers to select patients for targeted inhibitor therapy after resection of hepatocellular carcinoma. *Ann Surg Oncol*. 2011;18(12):3384–3390.
21. Tang Z, Li C, Kang B, Gao G, Li C, Zhang Z. GEPIA: a web server for cancer and normal gene expression profiling and interactive analyses. *Nucleic Acids Res*. 2017;45(W1):W98–W102.
22. Berlin J, Bendell JC, Hart LL, et al. A randomized phase II trial of vismodegib versus placebo with FOLFOX or FOLFIRI and bevacizumab in patients with previously untreated metastatic colorectal cancer. *Clin Cancer Res*. 2013;19(1):258–267.
23. Wang JX, Wu HL, Zhu M, Zhou R. Role of anti-epidermal growth factor receptor therapy compared with anti-vascular endothelial growth factor therapy for metastatic colorectal cancer: an update meta-analysis of randomized clinical trials. *Pathol Oncol Res*. Epub 2018 Jan 30.
24. Khattak MA, Martin H, Davidson A, Phillips M. Role of first-line anti-epidermal growth factor receptor therapy compared with anti-vascular endothelial growth factor therapy in advanced colorectal cancer: a meta-analysis of randomized clinical trials. *Clin Colorectal Cancer*. 2015;14(2):81–90.
25. Sehdev A, Cramer HM, Ibrahim AA, Younger AE, O'Neil BH. Pathological complete response with anti-PD-1 therapy in a patient with microsatellite instable high, BRAF mutant metastatic colon cancer: a case report and review of literature. *Discov Med*. 2016;21(117):341–347.
26. Toh JWT, de Souza P, Lim SH, et al. The potential value of immunotherapy in colorectal cancers: review of the evidence for programmed death-1 inhibitor therapy. *Clin Colorectal Cancer*. 2016;15(4):285–291.
27. Zhang J, Zhou L, Zhao S, Dicker DT, El-Deiry WS. The CDK4/6 inhibitor palbociclib synergizes with irinotecan to promote colorectal cancer cell death under hypoxia. *Cell Cycle*. 2017;16(12):1193–1200.
28. Pino MS, Chung DC. The chromosomal instability pathway in colon cancer. *Gastroenterology*. 2010;138(6):2059–2072.
29. Issa JP. Colon cancer: it's CIN or CIMP. *Clin Cancer Res*. 2008;14(19):5939–5940.
30. Vermeulen K, Van Bockstaele DR, Berneman ZN. The cell cycle: a review of regulation, deregulation and therapeutic targets in cancer. *Cell Prolif*. 2003;36(3):131–149.
31. Karaiskou A, Perez LH, Ferby I, Ozon R, Jessus C, Nebreda AR. Differential regulation of Cdc2 and Cdk2 by RINGO and cyclins. *J Biol Chem*. 2001;276(38):36028–36034.
32. Porter LA, Kong-Beltran M, Donoghue DJ. Spy1 interacts with p27Kip1 to allow G1/S progression. *Mol Biol Cell*. 2003;14(9):3664–3674.
33. Porter LA, Dellinger RW, Tynan JA, et al. Human Speedy: a novel cell cycle regulator that enhances proliferation through activation of Cdk2. *J Cell Biol*. 2002;157(3):357–366.
34. Arumugam K, MacNicol MC, Wang Y, et al. Ringo/cyclin-dependent kinase and mitogen-activated protein kinase signaling pathways regulate the activity of the cell fate determinant Musashi to promote cell cycle re-entry in *Xenopus* oocytes. *J Biol Chem*. 2012;287(13):10639–10649.
35. Nebreda AR. CDK activation by non-cyclin proteins. *Current Opinion in Cell Biology*. 2006;18(2):192–198.

Cancer Management and Research

Dovepress

Publish your work in this journal

Cancer Management and Research is an international, peer-reviewed open access journal focusing on cancer research and the optimal use of preventative and integrated treatment interventions to achieve improved outcomes, enhanced survival and quality of life for the cancer patient. The manuscript management system is completely online and includes

a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/cancer-management-and-research-journal>