A vascular access and midlines program can decrease hospital-acquired central line-associated bloodstream infections and cost to a community-based hospital

Rahul Pathak
Sumalatha Gangina
Falina Jairam
Kimberly Hinton

University of Central Florida School
of Medicine and Heart of Florida
Regional Medical Center, Davenport,
FL, USA

Background: We propose that substituting central lines with midlines can help reduce the total number of central line catheter-days as well as central line-associated bloodstream infections (CLABSIs), thereby reducing the associated costs and the resulting increased length of stay. A midline or vascular access program in a community hospital can bring about these positive changes.

Objectives: Our objective is to evaluate whether the institution of a midline program for vascular access at a community hospital can reduce the number of central line catheter-days and the associated CLABSI rate, incidence, and cost.

Materials and methods: We collected data on the number of central line catheter-days per year starting from 2012. We also collected data on the total number of patient-days during this period and the number of CLABSIs. We started Centers for Disease Control and Prevention (CDC)-based recommendations to help decrease CLABSIs in June 2014; this included the use of the central venous catheters (CVC) insertion bundles and CVC maintenance bundle. Chlorhexidine baths were also given to all patients with central lines. In June 2015, we started a midline program and tracked data till June 2017. We then compared the infection rates during these periods.

Results: We conclude that instituting CDC recommendations to decrease CLABSIs did bring down the CLABSI rate; this decrease was not statistically significant. However, the addition of the midline program to replace central lines whenever possible, combined with universal CDC recommendations, did result in a significant decrease in both the number of central line days per patient-day and the CLABSI rate.

Conclusion: We recommend hospitals to develop a midline program to help reduce the use of central line catheters when possible to reduce the total number of catheter-days and the CLABSI rate associated with them.

Keywords: central line-associated bloodstream infections, midline, venous access, hospital-acquired infections, hospital length of stay, quality of care

Introduction

Central venous catheters are the most common source of hospital-acquired bloodstream infections in the US. Each infection has an estimated additional cost of \$25,000/- per episode.¹ There are up to 250,000 episodes reported in the US per year, each episode increasing the cost and duration of the hospital stay.² The Centers for Disease Control and Prevention (CDC) has released many guidelines to help prevent and reduce catheter-related bloodstream infections, and this has helped bring down

Correspondence: Rahul Pathak
University of Central Florida, 40100
US-27, Davenport, FL 33837, USA
Email rahul.pathak@ucf.edu

the rate of infections significantly.^{3,4} However, the number of central venous catheter-days remains close to 15 million catheter-days per year.⁵ The CDC reports a 50% decrease in central line-associated bloodstream infection (CLABSI) rate from 2008 to 2014 and a 9% decrease between 2013 and 2014. The CDC introduced general guidelines to decrease and prevent CLABSI in 2009 (hospital acquired infections data reports on CDC website). Nevertheless, about 30,100 CLABSIs still occur in the intensive care units and wards of US acute care facilities each year.⁶

In addition to bloodstream infection, central lines are also associated with significant complications including bleeding, pneumothorax, and arrhythmias.⁷

Midline catheters are intravenous (IV) devices placed in deep peripheral veins (eg, basilic, brachial, cephalic) usually under ultrasound guidance. They are between 3 and 10 inches in length (8–25 cm), and their tips reside distal to the axillary vein.^{8,9} These catheters are termed “midlines” because they are longer than peripheral IV catheters, which are generally 1–3 inches long, and shorter than peripherally inserted central catheters (PICCs), which extend into the vena cava. Several studies have demonstrated that central line use can be decreased by the use of midline catheters.^{10–13} Two studies recently showed that the use of central venous catheters was decreased by 80%–85% in an emergency department setting over the span of 5 years by using midlines instead.^{11,13} Another study recently demonstrated a decrease in the rate of CLABSI by introducing a midline program in a ventilator unit.¹²

In a recent 2-year study, the rate of bacteremia in midline catheters was reported to be 0.3%.¹⁴ We wanted to study whether introducing a midline program can help bring the rate and incidence of CLABSI down in a small community hospital.

Materials and methods

This was a retrospective review of the data collected over the past 5 years from our infection control department. This study was determined to be a non-human subjects research, and a waiver for informed consent was granted by the University of Central Florida institutional review board. There were no patient identifiers used in this data for our study. We collected data on the number of central line catheter-days per year starting from 2012. We also collected data on the total number of patient days during this period and the number of CLABSI.

We started CDC-based recommendations to help decrease CLABSI in June 2014; this included the use of the central venous catheters (CVC) insertion bundles and CVC

maintenance bundle. Chlorhexidine baths were also given to all patients with a central line.

We started a midline program in June 2015. With the introduction of this program, we replaced central lines in the hospital whenever possible by midlines. We had a vascular access team available to do this. All emergency central lines and central lines placed for total parental nutrition and ionotropic medications were replaced when possible by midlines. Central lines placed for difficult IV access were replaced by midlines. We then compared the bloodstream infection rate from these periods. We excluded infections in central line ports for chemotherapy and infections in hemodialysis catheters, as these cannot be replaced by midlines. Various data including the organism cultured, and the location in the hospital where the central line was placed, were also accumulated.

The number of catheter-days for any calendar year was determined by the sum total number of central line catheters in the hospital each day for that calendar year. Thus, the annual CLABSI rate is reported as the total number of CLABSI/total number of catheter-days $\times$ 1,000 for each calendar year.

The data from 2012 and 2013 were compared to 2014 after the institution of CDC recommendations for CVC maintenance bundle. Then, the data from 2012 and 2013 were compared to 2015 onward – the period wherein we instituted the combined use of CVC maintenance bundle and the midline protocol.

For our statistical analysis, we used the data analysis firm called Instat Services Inc. (San Diego, CA, USA). They compared the rates of infections using the Fisher’s exact test to compare different interventions.

Results

When we introduced the universal CVC bundle recommended by the CDC, the CLABSI rate decreased, but not significantly ($P=0.5768$). However, the addition of midline program to replace central lines reduced the CLABSI rate significantly as compared to the use of CVC bundle alone (Table 1). Moreover, the combination of the CVC bundle plus the midline program was very significant in terms of CLABSI reduction when compared with the previous period during which there were no measures in place ($P=0.003$ and $P=0.0113$, respectively).

The total number of catheter-days also fell significantly (Table 2), and the catheter-days per patient-day had a significant drop as well ($P<0.001$).

Fisher’s exact test was used to calculate significance (analysis provided by Instat Services Inc.).

Table 1 Data broken up into periods when CDC guidelines were implemented alone and when CDC guidelines were implemented along with the midline program

	Prior to CVC guidelines	After CVC guidelines alone	After CVC guidelines and midline program
Infections/catheter-days	23/5,977 (0.385%)	9/3,117 (0.289%)	2/4,275 (0.047%)
Catheter-days/patient-days	5,977/104,353	3,117/59,189	4,275/127,401

Abbreviations: CDC, Centers for Disease Control and Prevention; CVC, central venous catheters.

Discussion

Central lines are necessary and help in improving the management of patients. When needed for the delivery of vesicant medications, inotropic medications, and management of intensive care unit patients, there are few options besides large-bore central line access. However, with the associated complications and CLABSI, we should try other alternatives as well. Midlines are an excellent mode of venous access that, like central lines, can deliver large-volume fluids and can be placed for up to 4 weeks at a time.^{9,10,12} With a bloodstream infection rate of only 0.3%, they are much safer and less complicated than central lines.¹⁴

Midline catheters are peripheral IV access devices between 3 and 10 inches in length (8–25 cm). These catheters are generally placed under ultrasound guidance in proximal veins such as the brachial or cephalic with the tip distal to the axillary vein.¹² Newer midlines made of novel co-polymers and treated for reduced biofilm formation are safer than the older ones and cause less thrombophlebitis. Midlines can easily be inserted by a trained vascular access nurse instead of a physician, unlike central lines. We do not need X-rays to confirm the position, as we do for central lines. It is an easy procedure and takes only a few months to set up a vascular access team with the experience to place midlines and PICC lines. Some hospitals already have special nurses trained to insert PICC lines; to teach them midline insertion takes very little time.

Studies have already shown that starting a midline program can decrease the overall number of central lines inserted in a hospital emergency room.^{10–13} The greatest number of

infections associated with central lines are from those inserted in emergency situations where standard precautions may not be possible. Other studies have shown that a midline program can reduce the overall number of catheter-days and CLABSI rate in a ventilator unit.

Our study shows that midlines can decrease the overall incidence of CLABSI in a community hospital after a midline program is initiated.

It saves the cost in a number of ways:

1. Inserted by a nurse rather than a physician.
2. No post-insertion X-ray needed to confirm position.
3. Decreases the incidence of CLABSI and the associated costs to the hospital.
4. Patients who require to be discharged with IV medications can go home or to the nursing home with a midline.
5. No complications such as bleeding, pneumothorax, or cardiac arrhythmias.
6. The chance of getting a deep vein thrombosis is <2% as opposed to 1%–38.5% with a PICC line.^{14,15}

Setting up a midline program will also help other nursing units when there is difficulty in IV access. Midlines will eliminate the need for central lines for conditions such as long-term venous access, difficult venous access, irritable medication infusions, and large volume infusions. More recently invented midlines boast a near-zero bloodstream infection rate with proper insertion techniques.

Conclusion

The setting up of a vascular access program in a hospital where midlines are used to replace central lines when

Table 2 Total catheter-days and infections in the divided time periods along with the total patient-days and the CLABSI rate/1,000 catheter-days

Period	Catheter-days	Infections	CLABSI rate/1,000 catheter-days	Total number of patient-days
July 2012–June 2013	3,655	13	3.55	51,135
July 2013–June 2014	2,232	10	4.48	53,218
July 2014–June 2015	3,117	9	2.88	59,189
July 2015–June 2016	2,547	2	0.78	62,835
July 2016–June 2017	1,728	0	0	64,566

Abbreviation: CLABSI, central line-associated bloodstream infection.

feasible significantly decreases the number of catheter-days within the hospital and also significantly reduces the rate of CLABSI alone or in combination with universal CDC recommendations, including CVC bundle. We recommend hospitals to develop a midline program to help reduce the use of central line catheters when possible. This will help to reduce the total number of catheter-days and the infection rate associated with them.

Disclosure

The authors report no conflicts of interest in this work.

References

1. Pittet D, Tarara D, Wenzel RP. Nosocomial bloodstream infection in critically ill patients. Excess length of stay, extra costs, and attributable mortality. *JAMA*. 1994;271(20):1598–1601.
2. Haddadin Y, Regunath H. Central line associated blood stream infections (CLABSI). *StatPearls*. Epub 2017 Mar 28.
3. O'Grady NP, Alexander M, Burns LA, et al. Guidelines for the prevention of intravascular catheter-related infections. *Clin Infect Dis*. 2011; 52(9):e162–e193.
4. O'Grady NP, Alexander M, Burns LA, et al. Summary of recommendations: guidelines for the prevention of intravascular catheter-related infections. *Clin Infect Dis*. 2011;52(9):1087–1099.
5. McGee DC, Gould MK. Preventing complications of central venous catheterization. *N Engl J Med*. 2003;348(12):1123–1133.
6. Bell T, O'Grady NP. Prevention of central line-associated bloodstream infections. *Infect Dis Clin North Am*. 2017;31(3):551–559.
7. Kornbau C, Lee KC, Hughes GD, Firstenberg MS. Central line complications. *Int J Crit Illn Inj Sci*. 2015;5(3):170–178.
8. O'Grady NP, Chertow DS. Managing bloodstream infections in patients who have short-term central venous catheters. *Cleve Clin J Med*. 2011; 78(1):10–17.
9. Deutsch GB, Sathyanarayana SA, Singh N, Nicastro J. Ultrasound-guided placement of midline catheters in the surgical intensive care unit: a cost-effective proposal for timely central line removal. *J Surg Res*. 2014;191(1):1–5.
10. Shokoohi H, Boniface K, McCarthy M, et al. Ultrasound-guided peripheral intravenous access program is associated with a marked reduction in central venous catheter use in noncritically ill emergency department patients. *Ann Emerg Med*. 2013;61(2):198–203.
11. Schoenfeld E, Shokoohi H, Boniface K. Ultrasound-guided peripheral intravenous access in the emergency department: patient-centered survey. *West J Emerg Med*. 2011;12(4):475–477.
12. Pathak R, Patel A, Enuh H, Adekunle O, Shrisantharajah V, Diaz K. The incidence of central line-associated bacteremia after the introduction of midline catheters in a ventilator unit population. *Infect Dis Clin Pract (Baltim Md)*. 2015;23(3):131–134.
13. Au AK, Rotte MJ, Grzybowski RJ, et al. Decrease in central venous catheter placement due to use of ultrasound guidance for peripheral intravenous catheters. *Am J Emerg Med*. 2012;30(9):1950–1954.
14. Goetz AM, Miller J, Wagener MM, Muder RR. Complications related to intravenous midline catheter usage. A 2-year study. *J Intraven Nurs*. 1998;21(2):76–80.
15. Cummings M, Hearse N, Mccutcheon H, Deuter K. Improving antibiotic treatment outcomes through the implementation of a midline: piloting a change in practice for cystic fibrosis patients. *J Vasc Nurs*. 2011;29(1): 11–15.

Therapeutics and Clinical Risk Management

Publish your work in this journal

Therapeutics and Clinical Risk Management is an international, peer-reviewed journal of clinical therapeutics and risk management, focusing on concise rapid reporting of clinical studies in all therapeutic areas, outcomes, safety, and programs for the effective, safe, and sustained use of medicines. This journal is indexed on PubMed Central, CAS,

Submit your manuscript here: <http://www.dovepress.com/therapeutics-and-clinical-risk-management-journal>

Dovepress

EMBASE, Scopus and the Elsevier Bibliographic databases. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.