

Undernutrition and associated factors among children aged 6–59 months living in slum areas of Gondar city, northwest Ethiopia: a cross-sectional study

Atanaw Gelu¹
Melkie Edris²
Terefe Derso²
Zegeye Abebe²

¹Comprehensive Specialized Hospital, College of Medicine and Health Sciences, University of Gondar, Gondar, Ethiopia; ²Department of Human Nutrition, Institute of Public Health, College of Medicine and Health Sciences, University of Gondar, Gondar, Ethiopia

Background: Many people, especially in developing countries such as Ethiopia, are migrating to cities. The majority of these people have settled in slum areas, which often have poor sanitation and housing conditions. Therefore, this study was conducted to gather evidence on nutritional status and associated factors among children aged 6–59 months living in slum areas of Gondar city, northwest Ethiopia.

Methods: A community-based cross-sectional study was carried out from 3 to 28 May 2017. A total of 593 children from the slum areas aged 6–59 months were included in the study. Bivariable and multivariable logistic regression analyses were carried out to identify the independent determinants of stunting and wasting.

Results: The overall prevalences of stunting and wasting were 42.3% (95% CI 38.34, 46.3%) and 7.3% (95% CI 5.3, 9.4%), respectively. Poor wealth status (adjusted odds ratio [AOR]=1.79; 95% CI 1.19, 2.70) and age of child 24–35 months (AOR=2.56; 95% CI 1.32, 4.96), 36–47 months (AOR=2.27; 95% CI 1.14, 4.54) and 48–59 months (AOR=2.69; 95% CI 1.35, 5.32) were independently associated with stunting. Similarly, presence of a fever in the previous 2 weeks (AOR=2.29; 95% CI 1.20, 4.38) and paternal control over resources (AOR=3.66; 95% CI 1.12, 11.04) were associated with wasting. Children of uneducated mothers (AOR=3.30; 95% CI 1.29, 8.46) were also more likely to be wasted.

Conclusion: This study illustrates that undernutrition is prevalent in the slum areas of Gondar city and is a critical public health problem. Therefore, attention should be targeted at economically disadvantaged children living in slum areas. In addition, there is a need to improve medical awareness of families with young children in these areas and increase the health-seeking behavior of these families, primarily by focusing on maternal education. Increasing women's decision-making autonomy over household resources is also recommended to address the problem.

Keywords: stunting, wasting, undernutrition, slum area, Ethiopia

Correspondence: Zegeye Abebe
Department of Human Nutrition,
Institute of Public Health, College of
Medicine and Health Sciences, University
of Gondar, PO Box 196, Gondar,
Ethiopia
Tel +25 196 587 6452
Email zegeye24@gmail.com

Background

Undernutrition generates economic losses associated with impaired brain development, poor school performance, and impaired productivity and earnings.^{2–4} Despite economic improvements in Ethiopia over the past few decades, children aged under 5 years (under-fives) are still suffering from undernutrition.^{4,5} As an illustration, 28% of all cases of child mortality, 12% of repeating grades in primary school and 16% of loss of gross domestic product are associated with undernutrition.⁴

The causes of undernutrition are complex and an individual's nutritional status is influenced by three broad categories of factors: food, health and care.⁶ In particular, the amount and severity of undernutrition are shown to be higher in slums, as a result of poverty and inferior living conditions compared to developed city areas.^{6–8} Poor housing conditions, overcrowding, poor quality of drinking water and inadequate sanitation are major threats to the health and nutritional status of children in slum areas.^{9,10} In addition, there are usually poor waste-disposal systems in slums, so their inhabitants often rely on shared toilets, while others use open areas or plastic bags (flying toilets) for defecation. Children may put contaminated fingers, pica or fomites into their mouth while crawling or playing around the contaminated environments associated with poor waste-disposal practices.¹¹ Thus, children living in such conditions are always at a high risk of developing undernutrition and other health problems.⁹ In urban slum areas, thousands of children die every day from preventable diseases related to the inadequate provision of water and sanitation, and lack of access to health care.^{8,9}

Studies in different slum settings have explored various factors, including advanced age of the child, male sex, illness in the preceding 2 weeks, prelacteal feeding, early or late initiation of complementary feeding, not being exclusively breastfed for 6 months and duration of breastfeeding <2 years, and found that these are independently associated with stunting. In addition, poor maternal education, non-working status of the mother, poor socio-economic status, and separated marital status of mother and father are independent factors associated with stunting.^{12–15} Similarly, suboptimal breastfeeding, poor wealth status, poor maternal education, larger family size, diarrheal and respiratory tract morbidities, and unavailability of latrines are associated with increased odds of wasting among children aged under 5 years.^{16–21}

The Government of Ethiopia has endorsed a National Nutrition Program and child feeding strategy with special focus on infant and young child nutritional practices. Currently in Ethiopia, especially in the Amhara region, a huge number of under-fives (46%) are suffering from undernutrition and its complications.^{4,5} There is a dearth of evidence on the nutritional status of under-fives in slum areas of Ethiopia, with the majority of previous local studies focused predominantly on rural areas.^{5,22–24} Many people, especially in developing countries such as Ethiopia, are migrating to cities and the majority of these settle in slum areas, neglected parts of the city with very limited basic services.²⁵ The slums normally have poor housing conditions and poor sanitation, and

the inhabitants are from lower socio-economic classes, which directly impacts on food security.²⁶ Hence, the evidence gathered from this study is useful to all stakeholders of interventions involved in control of undernutrition. The information is also appropriate in policy, planning and decision making at the local as well as the national level. Therefore, this study was conducted to gather evidence on nutritional status and associated factors among children aged 6–59 months living in slum areas of Gondar city, in northwest Ethiopia.

Key message

In the slum areas of Gondar city, the prevalence of undernutrition is high and it is a critical public health problem.

Methods

Study setting and design

A community-based cross-sectional study was carried out from 3 to 28 May 2017 in the slum areas of Gondar City. Gondar is located in North Gondar Administrative Zone, which is 750 km from Addis Ababa, the capital of Ethiopia. According to the recent administration system, the city has 12 sub-cities. The city administration identified three slum areas. The total estimated population of the city is 264,964.

Study population

All children aged 6–59 months who lived in any of the three slum areas of Gondar city were included in the survey. The sample size was computed using a single proportion formula considering the following assumptions: a prevalence of stunting in Amhara region of 46%,⁵ a 95% level of confidence and a 5% margin of error. A 10% non-response rate was also added to achieve a minimum sample size of 420. However, according to the city administration, the total number of children aged 6–59 months living in the three slum areas of the city was 600. Thus, to increase the power of the study, all children (593) fulfilling the inclusion criteria were included in the study.

Data collection instrument and procedure

A structured interviewer-administered questionnaire composed of questions on socio-demographic factors, maternal and child health care, food security, hygiene and sanitation was used to collect the data. The questionnaire was administered via house-to-house visits. The questionnaire was prepared in English and translated into the local language, Amharic, and then translated back into English to check the consistency. Six diploma nurse data collectors and three

BSc field supervisors participated in the data collection. Anthropometric measurements of both weight and length/height were carried out using the procedure stipulated by the World Health Organization (WHO) (2006).²⁷ The weight of children aged 6–23 months was measured by a hanging scale (serial number 3541317009, with graduation of 0.1 kg and measuring up to 20 kg; Seca, Hamburg, Germany) to the nearest 0.1 kg with light clothing and no shoes. Similarly, the weight of children aged 24–59 months was measured by the Seca digital balance (serial number 5755107131646, with graduation of 0.1 kg and measuring up to 160 kg). The instrument was calibrated before weighing each child and by setting it to zero. Furthermore, the weighing scale was checked daily against the standard weight for accuracy. The length of a child (aged 6–23 months) was measured using a horizontal wooden length board in a recumbent position and length was read to the nearest 0.1 cm. Height was measured using the Seca vertical height scale (model number 213) following the standard procedure. The occipital (back of the head), shoulder blades, buttocks and heels were touching the measuring board and height was recorded to the nearest 0.01cm. Two days training on the technique of interviewing and anthropometric measurements was given for data collectors and supervisors. Then the tool was pretested in 5% of the sample size in the study area. Necessary modifications were made, such as identifying and replacing ambiguous words. During data collection, close supervision, spot-checking and reviewing the completed questionnaire were undertaken by the supervisors and principal investigator on a daily basis. Data clean-up and cross-checking were carried out before analysis.

Assessment of nutritional status

Anthropometric indices were calculated using the WHO Anthro software. The Z-scores of indices, height-for-age Z-score (HAZ) and weight-for-height Z-score (WHZ) were computed and compared using the WHO Multicenter Growth Reference Standard. A child with an HAZ less than -2 SD from the reference population was defined as stunted and a child with a WHZ less than -2 SD from the reference population was classified as wasted; otherwise they were considered normal.¹

Assessment of household food security, dietary diversity and wealth status

Household food security status (HFSS) was assessed using the Household Food Insecurity Access Scale. HFSS was determined using a 1-month recall period and two types of

questions: an occurrence question followed by a question on the frequency of occurrence of the event. Finally, the HFSS was categorized as food secure for a score ≤ 1 , and food insecure for a score ≥ 2 .²⁸

The dietary diversity score (DDS) of a child was assessed by interviewing the mother and asking her to list all food and drink taken by the child in the previous 24 hours. Finally, food items were categorized into seven food groups; a DDS of 4 was considered as the minimum acceptable dietary diversity and DDS of < 4 was classified as poor dietary diversity.²⁹

Household wealth status was determined from the key household asset ownership variables (such as quantity of cereal products, house, livestock and agricultural land ownership) and categorized as low, medium and high based on the lower, middle and higher score tertiles using principal component analysis.³⁰

Data processing and analysis

Data were edited, coded and entered into Epi Info version 7, and exported to SPSS version 20 for analysis. Descriptive statistics (means, frequencies and proportions) were used to present the findings. A bivariable analysis was conducted for stunting and wasting separately. Variables with *P*-values of < 0.2 in the bivariable analysis were entered into a multivariable logistic regression analysis to identify the independent predictors of stunting and wasting. Both the crude odds ratio (COR) and adjusted odds ratio (AOR) with corresponding 95% CI were computed to show the strength of the association. A Hosmer–Lemeshow goodness-of-fit test was checked ($P=0.9$ for stunting and $P=0.6$ for wasting). Variables with a *P*-value of < 0.05 in the multivariate logistic regression analysis were taken as statistically significant.

Results

Socio-economic and demographic characteristics

A total of 593 mother/caregiver–child pairs were included in the study. The mean \pm SD age of the children was 31.06 ± 15.08 months. More than two-thirds (73.9%) of mothers were married and were Orthodox Christian (68.6%). About 61.4% of mothers were unemployed and 33.7% had no formal education (Table 1).

Child, maternal and household-related characteristics

The majority of the mothers (93.8%) had received antenatal care during pregnancy. All children were breastfed (100%) and almost all were vaccinated (97.5%). The majority had

received deworming (57.8%) and vitamin A supplementation (82.1%) in the previous 6 months. However, over a quarter (27%) had a history of diarrhea and 27.8% of children had had a fever within the 2 weeks before the date of the survey. More than three-quarters (78.6%) of children had started breastfeeding within 1 hour of birth. Most children (70.3%) received complementary feeding at the age of 6 months, but more than half (60.1%) had an unvaried diet. Regarding the availability of latrines and food security, around one-quarter (26%) of households had no latrine and 40.6% did not have a secure food supply.

Prevalence of stunting and wasting

Out of 593 children, 42.3% (95% CI 38.34, 46.3%) and 7.3% (95% CI 5.3, 9.4%) were stunted and wasted, respectively. About 15.9% of children surveyed were severely stunted (Table 2).

Associated factors with stunting

In the bivariable analysis, mother's occupation, number of under-fives, decision making over resources, the age of the children and low household wealth were associated with stunting. The age of the children and household wealth status remained significantly associated with stunting in the multivariable logistic regression analysis. Thus, higher proportions of children are stunted in households with poor socio-economic status (AOR=1.79; 95% CI 1.19, 2.70). Likewise, higher odds of stunting were observed in higher age groups, 24–35 months (AOR=2.56; 95% CI 1.32, 4.96), 36–47 months (AOR=2.27; 95% CI 1.14, 4.54) and 48–59 months (AOR=2.69; 95% CI 1.35, 5.32) (Table 3).

Factors associated with wasting

As presented in Table 4, fever in 2 weeks before the date of the survey, maternal education and paternal control over resources were significantly associated with wasting. Accordingly, a higher odds of wasting was observed among children who had fever (AOR=2.29; 95% CI 1.20, 4.38), children of uneducated mothers (AOR=3.30; 95% CI 1.29, 8.46) and children whose fathers controlled the resources at home (AOR=3.66; 95% CI 1.12, 11.04).

Discussion

Despite various interventions to alleviate childhood under-nutrition in Ethiopia, the current study found that the prevalence of stunting and wasting remains unacceptably high. Using the WHO cut-off point for declaring the public

Table 1 Socio-economic and demographic characteristics of children and their parents in slum areas of Gondar city, northwest Ethiopia, 2017 (n=593)

Characteristics	Frequency	Percentage
Age of child (months)		
6–11	60	10.1
12–23	150	25.3
24–35	152	25.6
36–47	113	19.1
48–59	118	19.9
Birth order of child		
First child	276	46.5
Second or third child	258	43.5
Fourth or above	59	9.9
Birth interval of child		
First child (no interval)	274	46.3
1–2 year interval	142	23.9
>2 year interval	177	29.8
Father's education		
No formal education	170	28.7
Primary education	162	27.3
Secondary and above	261	44
Mother's marital status		
Single	55	9.3
Married	438	73.9
Divorced	86	14.5
Widowed	14	2.4
Mother's education		
No formal education	200	33.7
Primary education	183	30.9
Secondary and above	210	35.4
Mother's occupation		
Unemployed	364	61.4
Government employee	132	22.3
Private worker	97	16.4
Father's occupation		
Unemployed	222	37.4
Government employed	182	30.7
Private worker	189	31.9
Number of children under 5		
1	460	76.7
≥2	140	23.3
Total family size		
≤4	338	57.0
≥5	255	43.0
Head of household		
Mother	336	56.7
Father	254	42.8
Sister	3	0.5
Decision maker over resources		
Husband	227	38.3
Wife	198	33.4
Both jointly	168	28.3
Religion		
Orthodox	407	68.6
Muslim	184	31.0
Protestant	2	0.3

(Continued)

Table 1 (Continued)

Characteristics	Frequency	Percentage
Ethnicity		
Amhara	579	97.6
Others ^a	14	2.4
Household wealth status		
Poor	198	33.4
Medium	197	33.2
Rich	198	33.4

Note: ^aTigray, Oromo, Kimant.

Table 2 Nutritional status of children in urban slums areas of Gondar city, northwest Ethiopia (n=593)

Indicators	Number	Percentage
Weight for height		
Normal	550	92.7
Wasted	43	7.3
Severely wasted	14	2.4
Height for age		
Normal	342	57.7
Stunted	251	42.3
Severely stunted	94	15.9
Weight for age		
Normal	456	76.9
Underweight	137	23.1
Severely underweight	34	5.7

health importance of undernutrition, the study found that stunting was a critical health problem, with a prevalence of 42.3%, and wasting was a medium public health problem, with a prevalence of 7.3% among the children inhabiting the slums of Gondar city.³¹ This indicates that a considerable number of children suffered from the consequences of undernutrition. The findings should stimulate the concerned body to develop new initiatives and investments and to strengthen existing interventions to decrease the burden of undernutrition.

The prevalence of stunting in the current report, 42.3%, is in line with a study conducted in slum areas of Bahir Dar town, also in the Amhara region of Ethiopia (42%),¹² and in Kenya (47%).¹⁴ But the prevalence seen in this study was higher than in studies conducted in other countries, such as Bangladesh (19.5%),³² Iran (9.53%)³³ and India (30.3%).³⁴ The discrepancy may be due to the difference in study settings and subjects. In this study, most of the mothers were uneducated. Uneducated mothers may not easily understand the nutritional requirements of their children, and may not follow the recommended procedures for feeding children.^{13,35}

In this study, the prevalence of wasting, 7.3%, was consistent with findings of studies conducted in slum areas of Ethiopia (6.4%),¹² Iran (8.19%)³³ and Botswana (5.5%).³⁶ However, the prevalence of wasting in the present study was higher than that reported in Kenya (2.6%).¹⁴ The high prevalence of wasting may be due to the mothers' low educational levels as well as differences in design between studies.

In the current study, a higher proportion of children were stunted in the higher age groups compared to the younger ones. This finding is supported by a report from Ludhiana, India.³⁷ This may be due to higher nutritional demands for growth and development. In addition, the child may be put on a family diet and so suffer from a lack of adequate dietary diversity and meal frequency. It was observed that a large number of children received a unvaried diet. Therefore, the risk of impaired linear growth increased as the age of the child advanced.^{36,38}

Consistent with the established facts,^{39,40} a higher odds of stunting was found among children from poorer households: the odds of stunting were 1.79 times higher among children from poorer households compared to children from richer households. This finding is supported by the findings of a study from Dabat district in Ethiopia.⁴¹ One possible explanation may be related to the negative effects of poor wealth status on the household's access to food, food purchasing power and sanitation facilities.⁴² Thus, children in poor households received poor-quality food, an inadequate amount and frequency, and a monotonous diet, which had a direct effect on nutritional status.⁴³

In this study, children who had fever in the 2 weeks before the date of the survey were more likely to be wasted. Basic services, such as housing, safe drinking water and sanitation facilities, are lacking in slum areas, making slum inhabitants more vulnerable to infectious diseases, which further negatively affects their nutritional status. Thus, infection facilitates losses of nutrients through decreased intestinal absorption, direct loss of nutrients, increased metabolic response, or loss of appetite and thus decreased food intake.^{44,45}

The educational status of mothers was an independent factor on wasting in children. The likelihood of wasting was increased by 30% among children of uneducated mothers. Educated mothers are more able to understand the consequences of nutritional deficiency, and more likely to maintain good hygiene practices and provide a high quality of care to their children. In addition, educated mothers can make independent decisions over food preference and preparation, and are more likely to have control over household resources compared to uneducated mothers.⁴⁶

Table 3 Factors associated with stunting among children in slum areas of Gondar city, northwest Ethiopia, 2017

Variables	Stunting		COR (95% CI)	AOR (95% CI)
	Yes (n)	No (n)		
Mother's occupation				
Unemployed	148	216	1.06 (0.67, 1.68)	1.09 (0.66, 1.78)
Government employee	65	67	1.51 (0.89, 2.56)	1.26 (0.71, 2.25)
Private worker	38	59	1.00	1.00
Number of children under 5				
One	185	268	1.00	1.00
Two and above	66	74	1.29 (0.88, 1.89)	1.28 (0.85, 1.95)
Decision making over resources				
Husband	99	128	1.36 (0.90, 2.04)	1.19 (0.91, 2.11)
Wife	91	107	1.49 (0.98, 2.27)	1.36 (0.88, 2.10)
Both jointly	61	107	1.00	1.00
Age of child (months)				
6–11	16	44	1	1.00
12–23	57	93	1.69 (0.87, 3.26)	1.75 (0.90, 3.41)
24–35	71	81	2.41 (1.25, 4.64)	2.56 (1.32, 4.96)
36–47	49	64	2.11 (1.06, 4.17)	2.27 (1.14, 4.54)
48–59	58	60	2.66 (1.35, 5.23)	2.69 (1.35, 5.32)
Household wealth status				
Poor	101	97	1.74 (1.17, 2.61)	1.79 (1.19, 2.70)
Medium	76	121	1.05 (0.7, 1.58)	1.06 (0.70, 1.61)
Rich	74	124	1.00	1.00

Abbreviations: COR, crude odds ratio; AOR, adjusted odds ratio.

Table 4 Factors associated with wasting among study children in slum area of Gondar city, Northwest Ethiopia, 2017

Variables	Wasting		COR (95% CI)	AOR (95% CI)
	Yes (n)	No (n)		
Number of children ever born				
1	19	236	0.64 (0.26, 1.61)	0.77 (0.30, 2.03)
2 or 3	17	258	0.53 (0.21, 1.33)	0.70 (0.27, 1.84)
≥4	7	56	1.00	1.00
ANC visit				
Yes	5	32	2.13 (0.79, 5.78)	1.33 (0.47, 3.73)
No	38	518	1.00	1.00
Fever in the previous 2 weeks				
Yes	20	145	2.43 (1.30, 4.55)	2.29 (1.20, 4.38)
No	23	405	1.00	1.00
Decision making over resources				
Husband	21	206	4.18 (1.45, 13.26)	3.66 (1.12, 11.04)
Wife	18	180	4.10 (1.36, 12.37)	3.00 (0.98, 9.23)
Both jointly	4	164	1.00	1.00
Mother's education				
No formal education	22	178	4.20 (1.67, 10.60)	3.30 (1.29, 8.46)
Primary education	15	168	3.04 (1.15, 8.00)	2.49 (0.93, 6.63)
Secondary and above	6	204	1.00	1.00

Abbreviations: COR, crude odds ratio; AOR, adjusted odds ratio; ANC, antenatal care.

Finally, household decision making was associated with wasting. Previous studies confirmed that a lack of female decision making and autonomy negatively impacts children's nutritional status. Autonomous mothers are more likely to spend time and resources in caring for their children. In addition, they are more likely to participate in various social activities outside the home, which may allow for exchange of

information about child feeding practices, caring and nutrition. In turn, this has long-term cumulative effects on the health, growth and nutritional status of children.⁴⁰

Limitations

The study was conducted by including all children in the study area. However, the study has some limitations. First,

the study did not consider the quantity and quality of food consumed by the children. Second, the study may not free from social desirability bias in responding to questions on the type and frequency of foods given to children.

Conclusions

The study confirmed the critical nature of undernutrition as a public health problem in the slum areas of Gondar city. The age of the children and poor household wealth status were associated with stunting. Similarly, lack of maternal education, fever in the 2 weeks preceding the interview and paternal control over household resources were associated with wasting. Therefore, attention should be targeted to economically disadvantaged children living in the slum areas. In addition, health education and education on health-seeking behavior regarding sick children, focusing on maternal education and increasing women's decision-making autonomy over household resources are recommended to address the problem.

Ethics approval and consent to participate

Ethical approval for this study was obtained from the Institutional Review Board of the University of Gondar, who also approved the use of verbal informed consent from the parents/caretakers. A permission letter was also obtained from Gondar city administration. Verbal consent was obtained from the parent/caretaker of each study participant after the purpose of the study had been explained.

Acknowledgments

The authors would like to thank study participants for their willingness to participate in the study, and the University of Gondar and Gondar City Health Department for material support. Finally, we would like to thank Mr Joe Kennedy for language editing. No funding was secured for this study.

Author contributions

AG, TD and ZA conceived the study, developed the tool, coordinated data collection, carried out the statistical analysis and drafted the manuscript. ME, AG, ZA and TD participated in the design of the study, developed the tool, participated in the statistical analysis and drafted the manuscript. All authors read and approved the final manuscript.

Disclosure

The authors report no conflicts of interest in this work.

References

1. World Health Organization/UNICEF. *WHO Child Growth Standards and the Identification of Severe Acute Malnutrition in Infants and Children: A Joint Statement by the World Health Organization and the United Nations Children's Fund*. Geneva: World Health Organization; 2009.
2. Pollitt E, Gorman KS, Engle PL, Martorell R, Rivera J. Early supplementary feeding and cognition: effects over two decades. *Monogr Soc Res Child Dev*. 1993;58(7):1–99; discussion 111–118.
3. Grantham-Mcgregor S, Cheung YB, Cueto S, et al. Developmental potential in the first 5 years for children in developing countries. *Lancet*. 2007;369(9555):60–70.
4. UNICEF. *The Cost of Hunger in Ethiopia. The Social and Economic Impact of Child Undernourishment in Ethiopia. Summary Report*. Addis Ababa: UNICEF; 2014.
5. Central Statistical Agency [Ethiopia] and ICF International. *Ethiopia Demographic and Health Survey 2011*. Addis Ababa/Calverton, MD: Central Statistical Agency/ICF International; 2016.
6. Ramji S. Impact of infant & young child feeding & caring practices on nutritional status & health. *Indian J Med Res*. 2009;130(5):624–626.
7. Awasthi S, Agarwal S. Determinants of childhood mortality and morbidity in urban slums in India. *Indian Pediatr*. 2003;40(12):1145–1160.
8. Goudet SM, Griffiths PL, Bogin BA, Madise NJ, Cochrane Public Health Group. Nutritional interventions for preventing stunting in children (0 to 5 years) living in urban slums. *Cochrane Database Syst Rev*. 2015;(5):CD011695.
9. Unger A. Children's health in slum settings. *Arch Dis Child*. 2013;98(10):799–805.
10. UN-HABITAT. *Defining Slums: Towards an Operational Definition for Measuring Slums. Background Paper 2, Expert Group Meeting on Slum Indicators*. Nairobi: United Nations; 2002.
11. Moya J, Bearer CF, Etzel RA. Children's behavior and physiology and how it affects exposure to environmental contaminants. *Pediatrics*. 2004;113(4 Suppl):996–1006.
12. Demilew YM, Abie DD. Undernutrition and associated factors among 24–36-month-old children in slum areas of Bahir Dar city, Ethiopia. *Int J Gen Med*. 2017;10:79–86.
13. Ahsan KZ, Arifeen SE, Al-Mamun MA, Khan SH, Chakraborty N. Effects of individual, household and community characteristics on child nutritional status in the slums of urban Bangladesh. *Arch Public Health*. 2017;75(9):9.
14. Olack B, Burke H, Cosmas L, et al. Nutritional status of under-five children living in an informal urban settlement in Nairobi, Kenya. *J Health Popul Nutr*. 2011;29(4):357–363.
15. Dhok R, Thakre S. Chronic undernutrition amongst under-five in an urban slum of Central India. *Int J Community Med Public Health*. 2016;3(3):700–704.
16. Onsa ZO, Ahmed NMK. Impact of Exclusive Breast Feeding on the Growth of Sudanese Children (0–24 Months). *Pak J Nutr*. 2014;13(2):99–106.
17. Engle PL, Menon P, Haddad L. Care and Nutrition: Concepts and Measurement. *World Dev*. 1999;27(8):1309–1337.
18. Bantamen G, Belaynew W, Dube J. Assessment of factors associated with malnutrition among under five years age children at Machakel Woreda, Northwest Ethiopia: a case control study. *J Nutr Food Sci*. 2014;4(1):256.
19. Ayana AB, Hailemariam TW, Melke AS. Determinants of acute malnutrition among children aged 6–59 months in Public Hospitals, Oromia region, West Ethiopia: a case–control study. *BMC Nutr*. 2015;1(1):34.
20. Gebretsadik A, Worku A, Berhane Y, et al. Factors associated with acute respiratory infection in children under the age of 5 years: evidence from the 2011 Ethiopia demographic and health survey. *Neuropsychiatr Dis Treat*. 2015;11:2159–2175.
21. Tariku B, Mulugeta A, Tsadik M, Azene G. Prevalence and Risk Factors of Child Malnutrition in Community Based Nutrition Program Implementing and Nonimplementing Districts from South East Amhara, Ethiopia. *OALib*. 2014;01(03):1–17.

22. Asfaw M, Wondaferash M, Taha M, Dube L. Prevalence of undernutrition and associated factors among children aged between six to fifty nine months in Bule Hora district, South Ethiopia. *BMC Public Health*. 2015;15(1):41.
23. Fikadu T, Assegid S, Dube L. Factors associated with stunting among children of age 24 to 59 months in Meskan district, Gurage Zone, South Ethiopia: a case-control study. *BMC Public Health*. 2014;14(1):800.
24. Tariku A, Woldie H, Fekadu A, Adane AA, Ferede AT, Yitayew S. Nearly half of preschool children are stunted in Dembia district, Northwest Ethiopia: a community based cross-sectional study. *Arch Public Health*. 2016;74(1):13.
25. Akter T. *Migration and Living Conditions in Urban Slums: Implications for Food Security*. Dhaka: Unnayan Onneshan; 2009.
26. Rahman MK, Muhibbullah M, Islam MS. Socio-economic status of slum dwellers: a case study of Uttara periphery, Dhaka. *Asian Review: Journal of Asian University of Bangladesh*. 2015;4(1):14–27.
27. World Health Organization. *Who Child Growth Standards: Length/Height for Age, Weight-for-Age, Weight-for-Length, Weight-for-Height And Body Mass Index-for-Age, Methods and Development*. Geneva: World Health Organization; 2006.
28. Coates J, Swindale A, Bilinsky P. *Household Food Insecurity Access Scale (HFIAS) for Measurement of Food Access: Indicator Guide*. Washington, DC: Food and Nutrition Technical Assistance Project, Academy for Educational Development; 2007.
29. Kennedy GL, Pedro MR, Seghieri C, Nantel G, Brouwer I. Dietary diversity score is a useful indicator of micronutrient intake in non-breast-feeding Filipino children. *J Nutr*. 2007;137(2):472–477.
30. Abebe Z, Gebeye E, Tariku A. Poor dietary diversity, wealth status and use of un-iodized salt are associated with goiter among school children: a cross-sectional study in Ethiopia. *BMC Public Health*. 2017;17(1):44.
31. World Health Organization. *Nutrition Landscape Information System (NLIS) Country Profile Indicators: Interpretation Guide*. Geneva: World Health Organization; 2010.
32. Rahman L, Miah ZI, Biswas H, et al. Maternal Factors Causing Under-Nutrition among the under-Five Slum Children in Dhaka City. *Med Col J*. 2013;6(1):92–96.
33. Kavosi E, Hassanzadeh Rostami Z, Nasihatkon A, et al. Prevalence and Determinants of Under-Nutrition Among Children Under Six: a Cross-Sectional Survey in Fars Province, Iran. *Int J Health Policy Manag*. 2014;3(2):71–76.
34. Mukherjee K. Malnutrition in under five children in Turbhe stores urban slums in Navi Mumbai. *Int J Med Sci Public Health*. 2014;3(9):1127–1131.
35. Brhane G, Regassa N. Nutritional status of children under five years of age in Shire Indaselassie, North Ethiopia: Examining the prevalence and risk factors. *Kontakt*. 2014;16:161–170.
36. Nnyepi M, Bandeke T, Mahgoub S. Factors affecting prevalence of malnutrition among children under three years of age in Botswana. *Afr J Food Agric Nutr Dev*. 2006;6(1):1–15.
37. Sengupta P, Philip N, Benjamin AI. Epidemiological correlates of under-nutrition in under-5 years children in an urban slum of Ludhiana. *HPPI*. 2010;33(1):1–9.
38. Ricci JA, Becker S. Risk factors for wasting and stunting among children in Metro Cebu, Philippines. *Am J Clin Nutr*. 1996;63(6):966–975.
39. Fentahun W, Wubshet M, Tariku A. Undernutrition and associated factors among children aged 6-59 months in East Belesa District, northwest Ethiopia: a community based cross-sectional study. *BMC Public Health*. 2016;16:506.
40. Rahman MM, Saima U, Goni MA. Impact of Maternal Household Decision-Making Autonomy on Child Nutritional Status in Bangladesh. *Asia Pac J Public Health*. 2015;27(5):509–520.
41. Derso T, Tariku A, Bikis GA, Wassie MM. Stunting, wasting and associated factors among children aged 6-24months in Dabat health and demographic surveillance system site: A community based cross-sectional study in Ethiopia. *BMC Pediatr*. 2017;17(1):96.
42. UNICEF. *Strategy for Improved Nutrition of Women and Children in Developing Countries. A UNICEF Policy Review*. New York: UNICEF; 1990.
43. Belew AK, Ali BM, Abebe Z, Dachew BA. Dietary diversity and meal frequency among infant and young children: a community based study. *Ital J Pediatr*. 2017;43(1):73.
44. Scrimshaw NS. Effect of infection on nutritional status. *Proc Natl Sci Counc Repub China B*. 1992;16(1):46–64.
45. Gabriela M, Marón A, Clará W, et al. Association between nutritional status and severity of dengue infection in children in El Salvador. *Am J Trop Med Hyg*. 2010;82(2):324–329.
46. Panigrahi A, Das SC. Undernutrition and its correlates among children of 3-9 years of age residing in slum areas of Bhubaneswar, India. *Sci World J*. 2014;2014:1–9.

Pediatric Health, Medicine and Therapeutics

Publish your work in this journal

Pediatric Health, Medicine and Therapeutics is an international, peer-reviewed, open access journal publishing original research, reports, editorials, reviews and commentaries. All aspects of health maintenance, preventative measures and disease treatment interventions are addressed within the journal. Practitioners from all disciplines are invited to submit

their work as well as healthcare researchers and patient support groups. The manuscript management system is completely online and includes a very quick and fair peer-review system. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <http://www.dovepress.com/pediatric-health-medicine-and-therapeutics-journal>

Dovepress