Quantitative assessment of aberrant *P16^{INK4a}* methylation in ovarian cancer: a meta-analysis based on literature and TCGA datasets

Jie Ruan^{1,*}Peipei Xu^{2,3,*}Wei Fan⁴Qiaoling Deng³Mingxia Yu³

¹Key Laboratory for Medical Molecular Diagnostics of Guangdong, Guangdong Medical University, Dongguan, Guangdong, 523808, China;

²Department of Clinical Laboratory, The Third Affiliated Hospital of Zhengzhou University, Zhengzhou, Henan, 450072, China; ³Department of Clinical Laboratory, Zhongnan Hospital of Wuhan University, Wuhan, Hubei, 430071, China; ⁴Department of Pathology, Zhongnan Hospital of Wuhan University, Wuhan, Hubei, 430071, China

*These authors contributed equally to this work

Abstract: Epigenetic alteration of *P16^{INK4a}* is conventionally thought to induce the initiation of carcinoma. However, the role of *P16^{INK4a}* methylation in ovarian cancer still remains controversial. Therefore, we performed a meta-analysis to further elucidate the relationship between *P16^{INK4a}* promoter methylation and ovarian cancer. A total of 24 studies, including 20 on risk, 10 on clinicopathological features, and 3 on prognosis, were included in our meta-analysis. Our results indicated that the frequency of *P16^{INK4a}* methylation in cancer tissues was significantly higher than normal tissues and low malignant potential tumor tissues (odds ratio [OR] =5.01, 95% CI=1.55–16.14; OR=1.88, 95% CI=1.10–3.19, respectively), but similar to benign tissues (OR=1.18, 95% CI=0.52–2.65). Furthermore, *P16^{INK4a}* promoter methylation was not strongly correlated with age, clinical stage, tumor differentiation, or histological subtype in patients with ovarian cancer. Additionally, survival analysis showed that patients with *P16^{INK4a}* promoter methylation had a shorter progression-free survival in univariate and multivariate Cox regression models (hazard ratio =1.68, 95% CI=1.26–2.24; hazard ratio =1.55, 95% CI=1.15–2.08; respectively). In The Cancer Genome Atlas datasets, the methylation levels of seven out of nine CpG sites were significantly increased in the ovarian tumor tissues compared with the normal tissues. In conclusion, the present meta-analysis suggests that *P16^{INK4a}* promoter methylation may be useful in distinguishing malignant cancer from healthy ovarian tissues, and it may be a potential predictive marker for prognosis in patients with ovarian cancer.

Keywords: ovarian cancer, *P16^{INK4a}* promoter methylation, TCGA datasets, meta-analysis

Introduction

Ovarian cancer is the fifth leading cause of cancer-related deaths in women. According to the GLOBOCAN 2012 database, the incidences of ovarian cancer are 9.1 per 1,00,100 in developed countries and 5.0 per 1,00,000 in developing countries.¹ Therefore, approximately 70% is high-grade serous carcinomas.² Up to now, despite the effective treatments including radical resection, systemic chemotherapy, and targeted drugs for patients, the average 5-year survival is still only at 46%.³ Ovarian cancer is a multifactorial disease caused by the interaction of genetic and epigenetic factors.^{4,5} DNA methylation, as the most common epigenetic alteration, could occur at CpG island in the promoter region, 5' or 3' untranslated regions, and even in gene body of tumor suppressor genes (TSGs). Hypermethylation in the proximal promoter region often contributes to the transcriptional downregulation but methylation in exons is associated with active transcription.^{6,7} Recently, mounting evidences demonstrated that DNA methylation was involved in ovarian cancer.^{8–10} Therefore, identifying the role of TSG methylation in patients with ovarian cancer is of value.

Correspondence: Mingxia Yu
Department of Clinical Laboratory,
Zhongnan Hospital of Wuhan University,
169 Donghu Road, Wuhan, Hubei,
430071, China
Tel +86 027 6781 1772
Email dewrosy520@163.com

P16^{INK4a} (also known as *CDKN2A*), a classical TSG, is located on chromosome 9p21 and plays an important role in cell cycle regulation by decelerating cells progression from G1 to S phase.^{11,12} It has become clear that the expression of *P16* is reduced by DNA methylation.^{13–15} Also, *P16^{INK4a}* inactivation upregulates retinoblastoma (RB) protein by stimulating the cyclin-dependent kinases (CDKs) and RB pathway, which leads to dysfunction of cell proliferation and apoptosis, thereby further facilitating carcinogenesis.¹⁶ Indeed, several types of cancer, including ovarian cancer, exhibit a methylation phenotype of *P16^{INK4a}*.^{17–19}

To date, even though abundant studies have been conducted to explore the role of *P16^{INK4a}* promoter methylation in ovarian cancer, the results are still inconclusive. Several studies reported that *P16^{INK4a}* promoter methylation was associated with an increasing trend in ovarian cancer,^{20–23} while, other studies suggested that *P16^{INK4a}* promoter methylation was not related to the occurrence of ovarian cancer.^{24–30} Interestingly, even the conclusions in two published meta-analyses were inconsistent. Xiao et al reported that aberrant methylation of *P16^{INK4a}* was significantly associated with ovarian carcinogenesis,³¹ while Jiang et al suggested no association between *P16^{INK4a}* methylation and epithelial ovarian cancer.³²

Considering these conflicting conclusions on the role of methylated *P16^{INK4a}* in ovarian cancer, we performed an adaptive synthesized analysis to quantitatively evaluate the occurrence frequency, clinicopathological features, and potential prognostic significance of *P16^{INK4a}* promoter methylation in ovarian cancer. Moreover, we searched The Cancer Genome Atlas (TCGA) database, collecting hundreds of ovarian cancer samples with whole genome DNA methylation datasets to validate our meta-analysis.

Materials and methods

Search strategy and selection criteria

PubMed, Embase, Web of Science, and China National Knowledge Infrastructure were searched up to April 12, 2017, by the following keywords and search items: (*P16* OR *P16^{INK4a}* OR *CDKN2A*) AND (methylation OR hypermethylation OR demethylation) AND (ovarian OR ovary) AND (cancer OR carcinoma OR neoplasm). The search was limited to human studies, without language restriction. Moreover, a manual search of the relevant references was implemented to identify the potentially additional articles.

The following criteria were used for screening eligible studies: 1) case-control studies evaluating the association between *P16^{INK4a}* promoter methylation and ovarian cancer

risk, or case only studies evaluating the association of *P16^{INK4a}* promoter methylation with clinicopathological features or prognosis in ovarian cancer; 2) articles providing sufficient information for calculating an odds ratio (OR) and corresponding 95% CI, or study offering hazard ratio (HR) and 95% CI directly; 3) sample types limited to tissues; and 4) studies with full-text articles. It is worth noting that when multiple reports were published from a same study population, only the most recent or complete information was included in this meta-analysis. Meanwhile, studies with Newcastle Ottawa Scale (NOS) scores greater than or equal to five were enrolled.

Data extraction and quality assessment

With a preformed unified form, data were extracted independently by two investigators, and disagreements were resolved by discussion till consensus was achieved. The following information was extracted from studies: the first author's name, publication year, country, geographical location, sample size, age of patients in the case group, the frequencies of methylation in the case and control groups, methods for detecting methylation, methylation site, disease stage, tumor grade, histological subtype, and effects on survival outcomes.

The quality of eligible case-control studies was assessed according to the NOS criteria.³³ The NOS criteria are based on three aspects: 1) subject selection: 0–4; 2) comparability of subject: 0–2; 3) clinical outcome: 0–3.

Statistical analysis

Statistical analysis was conducted with Review Manager 5.2 (Cochrane Collaboration, Oxford, UK) and the Stata 12.0 (Stata Corporation, College Station, TX, USA). ORs with corresponding 95% CIs were calculated to estimate the association between *P16^{INK4a}* promoter methylation and ovarian cancer risk or clinicopathological features. Meanwhile, HRs and 95% CIs were used to assess the prognosis of *P16^{INK4a}* promoter methylation on ovarian cancer. Inter-study heterogeneity was estimated with the Cochran's *Q* statistic and *I*² tests. *P* < 0.05 or *I*² > 50% indicated substantial heterogeneity, and then the random-effects model was applied. Otherwise, the fixed-effects model was selected. We also explored sources of heterogeneity using meta-regression and subgroup analyses by publication year, geographical location, method, and case sample size. Additionally, sensitivity analysis was performed to investigate the influence of individual study. Publication bias was evaluated by funnel plots and Begg's test, and *P* < 0.05 was considered statistically significant. It is

worth mentioning that, for some trials containing no events in both case and control arms, as no information supplied about the likely magnitude of the effect, we excluded such trials when synthesizing data.³⁴

TCGA datasets extraction and analysis

We collected DNA methylation datasets of 582 ovarian cancer cases and 12 ovarian normal tissues from TCGA (“TCGA-ovary [OV]” project) program.³⁵ The methylation measurement was performed using Illumina HumanMethylation27 BeadChip. Beta value of each CpG site was extracted to assess the methylation level of *CDKN2A* gene. Beta value was calculated based on the intensities of the methylated (M) and unmethylated (U) bead types: beta value = M/(M+U).³⁶ The difference of DNA methylation level of CpG sites between ovarian tumor tissues and normal ovarian tissues in TCGA

database was analyzed by Student’s *t*-test on the means. *P16^{INK4a}* gene expression value (fragments per kilobase of transcript per million mapped reads) in ovarian tumor tissues (TCGA, “TCGA-OV” project) was also extracted. Pearson’s product-moment correlation between *P16^{INK4a}* gene expression levels and methylation of its CpG islands was evaluated. Data analysis was performed using R software (R i386 3.4.0). *P*-values were adjusted via Bonferroni correction.

Results

Identification of relevant studies

The procedure of study selection is outlined in Figure 1. We identified 233 articles in the initial literature search. A total of 153 references remained after removing duplicates. After reading titles and abstracts, 84 records were identified for further full-text assessment, which further excluded 60 more

Figure 1 Flow diagram of study selection.

articles. Finally, 24 studies from 1997 to 2015 were included in this meta-analysis.^{17,20,22–30,37–49}

Baseline characteristics of included studies

Out of the 24 studies, 11 studies were conducted in Asia, 7 in Europe, 4 in America, 1 in Africa, and 1 in Oceania. The detection methods of methylation in 20 studies were methylation-specific PCR (MSP) and real-time quantitative MSP, while methylation-specific multiplex ligation-dependent

probe amplification was used in two studies, MethyLight was used in one study, and Southern analysis was used in one study. Among the 24 articles, 20 studies^{17,20,22–30,37–40,42,45–47,49} addressed the risk of *P16^{INK4a}* promoter methylation in ovarian cancer, 10 studies^{20,25,28,29,38,41,43,44,47,48} covered clinicopathological features, and 3 studies^{20,42,43} discussed prognosis. To explore the relationship between *P16^{INK4a}* promoter methylation and ovarian cancer risk, three groups, that is, normal tissues, benign tissues, and low malignant potential or borderline tumor tissues (LMP), were compared. The NOS scores

Table I Characteristics of studies included for the association between *P16^{INK4a}* methylation and ovarian cancer risk

Authors	Year	Country	Geographical location	Sample size ^a	Case number				Age (years) ^d	Sample type	Method	Methylation site	NOS score
					C (M/n)	LMP (M/n)	B (M/n)	NT (M/n)					
Moselhy et al ¹⁷	2015	Saudi Arabia	Asia	Small	14/18	NA	12/32	NA	52.3±12.1	FFPET	MSP	Promoter	7
Bhagat et al ²⁰	2014	India	Asia	Large	58/134	5/23	11/26	0/15	49.55±9.72	FFT	qMSP	Promoter	7
Ozdemir et al ²⁴	2012	Turkey ^b	Asia	Large	1/75	NA	NA	0/75	NA	Tissue	MS-MLPA	Promoter	7
Ho et al ³⁷	2012	Taiwan	Asia	Small	1/47	NA	6/29	NA	50(32–66)	FFT	MS-MLPA	Promoter	7
Cul'bová et al ³⁸	2011	Slovakia	Europe	Small	5/13	0/2	5/19	NA	54.8 (34–74)	FFT	MSP	Promoter	6
Abou-Zeid et al ²²	2011	Egypt ^c	Africa	Large	21/52	NA	9/43	4/40	60(49–74)	FFT	qMSP	Promoter	7
Gu et al ³⁹	2009	China	Asia	Large	8/87	NA	13/42	NA	51(21–69)	Tissue	MethyLight	Promoter	7
Shen et al ²⁵	2008	China	Asia	Large	13/63	NA	NA	0/30	52.8 (33–76)	FFT	MSP	Promoter	6
Wu et al ⁴⁰	2007	Norway	Europe	Large	0/52	0/2	0/2	NA	NA	FFT	MSP	Promoter	6
Tam et al ²⁶	2007	Hong Kong	Asia	Large	17/89	1/16	1/19	4/16	53.1±1.4	FFT	MSP	Promoter	7
Wiley et al ⁴²	2006	Italy	Europe	Large	89/215	4/19	NA	NA	57.7±11.4	FFT	MSP	Promoter	7
Li et al ²⁷	2006	China	Asia	Small	6/18	NA	NA	0/10	NA	Tissue	MSP	Promoter	5
Makarla et al ²⁹	2005	USA	America	Small	7/23	5/23	3/23	0/16	51.5 (20–86)	FFT	MSP	Promoter	7
Liu et al ²⁸	2005	USA	America	Large	13/52	NA	NA	15/40	61.5±9.4	FFT	MSP	Promoter	5
Dhillon et al ²³	2004	India	Asia	Small	10/25	NA	NA	1/75	NA	Tissue	MSP	Promoter	7
Rathi et al ³⁰	2002	USA	America	Small	5/49	NA	NA	0/16	56(40–79)	FFT	MSP	Promoter	7
Strathdee et al ⁴⁶	2001	UK	Europe	Large	0/93	NA	NA	0/18	NA	FFT	MSP	Promoter	6
Brown et al ⁴⁵	2001	UK	Europe	Small	0/30	0/13	0/14	NA	NA	FFT	MSP	Promoter	5
McCluskey et al ⁴⁷	1999	USA	America	Small	21/37	11/15	14/20	NA	NA	FFT	MSP	Promoter	6
Shih et al ⁴⁹	1997	Australia	Oceania	Small	0/45	0/3	0/2	NA	NA	Tissue	Southern analysis	Promoter	5

Notes: ^aWe defined n<50 as small size and ≥50 as large size. ^bTurkey is a transcontinental Eurasian country and is usually assigned to Asia internationally. ^cEgypt is a transcontinental country spanning the northeast corner of Africa and southwest corner of Asia, usually assigned to Africa internationally. ^dAge data are presented as mean ± SD or median (IQR).

Abbreviations: B, benign tissues; BL, borderline; C, cancer tissues; FFPET, formalin fixed and paraffin embedded tissues; FFT, fast frozen tissues; LMP, low malignant potential or borderline tumor tissues; M, methylated; MS-MLPA, methylation-specific multiplex ligation-dependent probe amplification; MSP, methylation-specific PCR; n, number of patients in the group; NA, not available; NT, normal tissues; NOS, Newcastle Ottawa Scale; qMSP, real-time quantitative MSP.

of all case-control studies were ≥ 5 . The basic characteristics of all included studies are summarized in Tables 1 and 2.

Quantitative data synthesis

Association between P16^{INK4a} promoter methylation and ovarian cancer risk

A total of 1,217 ovarian cancers, 116 LMP cancers, 271 benign patients, and 351 normal controls were quantitatively synthesized in this analysis. Results indicated that the frequency of P16^{INK4a} promoter methylation in cancer tissues was significantly elevated than that in normal tissues (OR=5.01, 95% CI=1.55–16.14) and LMP tissues (OR=1.88, 95% CI=1.10–3.19), but similar to benign tissues (OR=1.18, 95% CI=0.52–2.65; Figure 2). Further analyses showed that the frequencies of P16^{INK4a} promoter methylation in benign tissues and LMP tissues were not higher than those in normal tissues (OR =2.28, 95% CI=0.37–14.09; OR =2.28, 95% CI=0.15–34.73, respectively; Figure 3).

With large heterogeneity, meta-regression and subgroup analyses were conducted by the publication year, geographical location, method, and case sample size in the comparison of cancer tissues vs normal tissues. Meta-regression found that case sample size was significantly correlated with the inter-study heterogeneity ($P=0.041$) while other covariates were not (Table 3). Furthermore, as shown in Table 3, subgroup analyses revealed that the OR was 5.69 (95% CI=0.42–76.14) for the publication year ≤ 2005 and 4.71 (95% CI=1.30–17.07) for >2005 under the random-effects model. For geographical location, the OR was 7.85 (95% CI=1.33–46.32) in Asia, 2.31 (95% CI=0.24–22.01) in America, and 6.10 (95% CI=1.89–19.69) in Africa under random-effects model. For

test method, the OR for MSP was 4.49 (95% CI=0.97–20.64) under random-effects model and 8.11 (95% CI=2.93–22.40) for other methods under fixed-effects model. In addition, the OR was 15.75 (95% CI=4.05–61.34) for sample size <50 in fixed-effects model and 2.21 (95% CI=1.33–3.67) for that ≥ 50 in random-effects model.

Association between P16^{INK4a} promoter methylation and clinicopathological features in patients with ovarian cancer

Ten studies comprising 680 samples were enrolled to assess whether or not the abnormal P16^{INK4a} promoter methylation was associated with ovarian cancer clinicopathological characteristics. As displayed in Figure 4, no statistically significant correlation was found between P16^{INK4a} promoter methylation and age of patients (≥ 60 vs <60 : OR =1.39, 95% CI=0.66–2.92), clinical stage (III–IV vs I–II: OR =1.21, 95% CI=0.81–1.82), grade (3 vs 1–2: OR=1.20, 95% CI=0.82–1.1.75) as well as histological subtype (serous vs non-serous: OR=1.09, 95% CI=0.76–1.55).

Prognostic value of P16^{INK4a} promoter methylation in patients with ovarian cancer

Only two studies^{42,43} containing 464 patients evaluated the P16^{INK4a} promoter methylation on progression-free survival (PFS) and three studies^{20,42,43} containing 600 patients on overall survival (OS). The combined results revealed P16^{INK4a} promoter methylation was significantly associated with a poor PFS by univariate Cox proportional hazards regression model (HR=1.68, 95% CI=1.26–2.24; Figure 5A). After considering potential confounders by adjusting for age at diagnosis

Table 2 Characteristics of studies included for the association between P16^{INK4a} methylation and clinicopathological features of ovarian cancer

Authors	Year	Country	Geographical location	Number of patients	Age (years) ^a	Tumor stage		Tumor grade		Histological subtype	
						I–II (M/n)	III–IV (M/n)	1–2 (M/n)	3 (M/n)	Serous (M/n)	Non-serous (M/n)
Bhagat et al ²⁰	2014	India	Asia	134	49.55 $\pm$ 9.72	19/41	39/93	18/45	40/89	32/76	26/58
Cul'bová et al ³⁸	2011	Slovakia	Europe	13	54.8 (34–74)	NA	NA	NA	NA	2/6	3/7
Shen et al ²⁵	2008	China	Asia	63	52.8 (33–76)	4/22	9/41	6/36	7/27	7/34	6/29
Yang et al ⁴¹	2006	Hong Kong	Asia	49	48.8 (26–79)	4/24	5/25	6/22	3/25	3/17	6/32
Makarla et al ²⁹	2005	USA	America	23	51.5 (20–86)	NA	NA	NA	NA	2/9	5/14
Liu et al ²⁸	2005	USA	America	52	61.5 $\pm$ 9.4	NA	NA	10/41	3/11	NA	NA
Katsaros et al ⁴³	2004	Italy	Europe	249	57(19–82)	22/68	68/152	26/75	64/141	40/86	50/141
Hashiguchi et al ⁴⁴	2001	Japan	Asia	46	NA	4/21	2/20	7/33	0/13	2/14	5/32
McCluskey et al ⁴⁷	1999	USA	America	29	NA	NA	NA	NA	NA	1/14	1/15
Milde-Langosch et al ⁴⁸	1998	Germany	Europe	44	NA	NA	NA	NA	NA	3/11	13/33

Note: ^aAge data are presented as mean $\pm$ SD or median (IQR).

Abbreviations: M, methylated; n, number of patients in the group; NA, not available.

Figure 2 Forest plots for the association between *P16^{INK4a}* methylation and ovarian cancer risk.

Notes: (A) Cancer tissues vs normal tissues; (B) cancer tissues vs benign tissues; (C) cancer tissues vs LMP tissues.

Abbreviations: LMP, low malignant potential or borderline tumor tissues; M-H, Mantel-Haenszel.

or surgery, disease stage, histological grade, and residual tumor size, the pooled HR was 1.55 (1.15–2.08; Figure 5B). Survival analysis also showed that *P16^{INK4a}* promoter methylation reduced OS in univariate and multivariate Cox regression models (HR=1.28, 95% CI=0.97–1.68; HR=1.16, 95% CI=0.87–1.55, respectively; Figure 5C and D), but the differences were not statistically significant.

Sensitivity analysis and publication bias

As presented in Figure 6A–C, no single study significantly affected the pooled ORs in the sensitivity analysis, indicating our analysis was relatively stable and credible. Funnel plots and Begg's test were used to evaluate the publication bias. The funnel plots were largely symmetric suggesting there were no publication biases in the meta-analysis of *P16^{INK4a}*

Figure 3 Forest plots for the association between *P16^{INK4a}* methylation and ovarian diseases.

Notes: (A) Benign tissues vs normal tissues; (B) LMP tissues vs normal tissues.

Abbreviations: LMP, low malignant potential or borderline tumor tissues; M-H, Mantel-Haenszel.

Table 3 Meta-regression and subgroup analyses of *P16^{INK4a}* methylation in comparison of cancer tissues vs normal tissues

Stratified analysis	Number of studies	Pooled OR (95% CI)		Meta-regression P-value	Heterogeneity	
		Random	Fixed		I ² (%)	P-value
Publication year				0.376		
≤2005	4	5.69 (0.42–76.14)	2.17 (1.17–4.05)		84	0.0003
>2005	6	4.71 (1.30–17.07)	4.65 (2.32–9.30)		56	0.05
Geographical location				0.161		
Asia	6	7.85 (1.33–46.32)	5.87 (2.70–12.78)		70	0.005
America	3	2.31 (0.24–22.01)	1.15 (0.56–2.37)		68	0.05
Africa	1	6.10 (1.89–19.69)	6.10 (1.89–19.69)		–	–
Method				0.651		
MSP	7	4.49 (0.97–20.64)	2.33 (1.38–3.92)		77	0.0003
Others	3	6.79 (2.43–18.94)	8.11 (2.93–22.40)		0	0.57
Case sample size				0.041		
<50	4	17.21 (4.54–65.28)	15.75 (4.05–61.34)		0	0.58
≥50	6	2.21 (1.33–3.67)	2.74 (0.71–10.53)		75	0.001

Abbreviations: MSP, methylation-specific PCR; OR, odds ratio.

promoter methylation and ovarian cancer risk, which was confirmed by the Begg's test (Figure 6D–F).

Methylation level of *P16^{INK4a}* measured by TCGA program

To further explore the methylation level of *P16^{INK4a}* in ovarian tumor tissues, we extracted DNA methylation data of *P16^{INK4a}* CpG sites measured with Illumina HumanMethylation27 BeadChip from TCGA program. As shown in Table 4, the

beta values of 582 ovarian tumor tissues and 12 normal ovarian tissues were extracted for analysis. Obviously, the methylation levels of seven out of nine CpG sites were significantly increased in the ovarian tumor tissues compared with the normal tissues (cg03079681, cg07752420, cg09099744, cg10895543, cg11653709, cg12840719, and cg26673943). Among these regions, methylation level of probe cg26673943 region (located at the promoter region of *P16^{INK4a}*) was negatively associated with *P16^{INK4a}* expression

Figure 4 Forest plots for the association between *P16^{INK4a}* methylation and clinicopathological features in ovarian cancer.

Notes: (A) Age; (B) clinical stage; (C) tumor grade; (D) histological subtype.

Abbreviation: M-H, Mantel-Haenszel.

in ovarian cancer patients (adjusted P -value <0.000001). However, methylation levels of the rest six probes, which are located at non-promoter region tended to be positively associated with *P16^{INK4a}* gene expression. Additionally, we found that methylation level of probe cg13479669 region

was lower in tumor tissues compared with normal tissues, and negatively associated with *P16^{INK4a}* gene expression in tumor tissues. These results suggest that hypermethylation of *P16^{INK4a}* might be correlated with ovarian carcinogenesis and development. Nevertheless, it seems that the methylation at

Figure 5 Forest plots for P16^{INK4a} methylation on survival analysis in univariate and multivariate Cox regression model.

Notes: (A) PFS in univariate Cox regression model; (B) PFS in multivariate Cox regression model; (C) OS in univariate Cox regression model; (D) OS in multivariate Cox regression model.

Abbreviations: OS, overall survival; PFS, progression-free survival; SE, standard error.

promoter region or non-promoter region has contrary effects on P16^{INK4a} gene expression.

Discussion

Ovarian cancer is one of the leading causes of cancer-related deaths in women.⁵⁰ Identification of early disease indicators for diagnosis and prognosis is of clinical value. P16^{INK4a}, which resembles classic TSGs such as P53, is an important negative regulator of cell growth and proliferation.¹⁶ It has been synthetically evaluated for aberrant P16^{INK4a} methylation in numerous cancers,^{51–54} including ovarian cancer.^{31,32} Considering the conflicting conclusions in two meta-analyses and the lack of comprehensive assessment on the role of methylated P16^{INK4a} in ovarian cancer, we performed an

adaptive synthesized analysis to investigate the relationships between P16^{INK4a} promoter methylation and ovarian cancer risk, as well as clinicopathological features and prognostic value in ovarian cancer. Meanwhile, we searched TCGA data to validate our meta-analysis.

Our meta-analysis demonstrated that P16^{INK4a} promoter methylation in cancer tissues was significantly higher than that in normal tissues ($P<0.05$), but not much increased than that in benign tissues. Compared with normal tissues, the frequency of P16^{INK4a} promoter methylation was 2.28-fold higher in both benign tissues and LMP tissues ($P>0.05$), but the differences were not statistically significant. The reason for this phenomenon may be that the transformation of normal cells to cancer cells is a long-term, gradual, and

Figure 6 Sensitivity analyses and Begg's test for publication bias of *P16^{INK4a}* methylation during the carcinogenesis of ovarian cancer.

Notes: (A and D) Cancer tissues vs normal tissues; (B and E) sensitivity analysis for the comparison of cancer tissues vs benign tissues; (C and F) sensitivity analysis for the comparison of cancer tissues vs LMP tissues.

Abbreviations: LMP, low malignant potential or borderline tumor tissues; SE, standard error.

multiphase process.⁵⁵ Although not establishing a strong correlation between *P16^{INK4a}* promoter methylation and cancer progression, the above results do suggest a possibility that epigenetic alteration of *P16^{INK4a}* promoter methylation might play a certain role in ovarian carcinogenesis and might be useful in distinguishing malignant tumor from healthy ovarian tissues. Considering the evident heterogeneity, we conducted subgroup analyses based on probable

covariates in the comparison of cancer tissues vs normal tissues. For geographical location, *P16^{INK4a}* promoter methylation is a risk factor in Asia and Africa, but not in America. The divergence may be underscored in a large part to a combination of differences in allele frequencies and complex epistasis or gene–environment interactions.⁵⁶ A review also outlined that some factors such as distinct physical appearance, behavior, and response to environ-

Table 4 Methylation of *P16^{INK4a}* CpG sites on Illumina HumanMethylation 27 BeadChip from TCGA datasets

Probe (Illumina HumanMethylation 27)	CpG island location (chromosome: DNA range)	Normal tissue beta value (mean, n=12)	Tumor tissue beta value (mean, n=582)	Adjusted P-value ^a	Pearson correlation coefficient	Adjusted P-value ^b
cg00718440	9: 21983444–21986348	0.016	0.016	0.960249	0.194104	0.001719
cg03079681	9: 21983444–21986348	0.015	0.026	<0.000001	0.012972	1.0
cg07752420	9: 21958106–21958899	0.149	0.653	<0.000001	0.569887	<0.000001
cg09099744	9: 21958106–21958899	0.099	0.642	<0.000001	0.630768	<0.000001
cg10895543	9: 21958106–21958899	0.120	0.651	<0.000001	0.624147	<0.000001
cg11653709	9: 21958106–21958899	0.144	0.610	<0.000001	0.555400	<0.000001
cg12840719	9: 21958106–21958899	0.092	0.594	<0.000001	0.627484	<0.000001
cg13479669	9: 21983444–21986348	0.045	0.027	0.004226	−0.150891	0.0333435
cg26673943	9: 21983444–21986348	0.047	0.056	0.042428	−0.269361	<0.000001

Notes: ^aP-value of t-test of the difference between normal tissue beta value and tumor tissue beta value. ^bP-value of Pearson's correlation between the tumor tissues beta value and CDKN2A expression (n=368).

Abbreviation: TCGA, The Cancer Genome Atlas.

mental agents and drugs between human populations could have contributed to the epigenetic variations.⁵⁷ Similar findings appeared in the subgroup analyses of different methods and publication year. Kurdyukov and Bullock⁵⁸ suggested that it was essential to choose an appropriate method in a suitable region to answer a particular biological question in studies of DNA methylation. Additionally, the 95% CI was large in the group of small sample size while relatively small in the group of large sample size, implying the conclusion may not be reliable unless studies should be conducted using a sufficient number of samples. Previous studies also demonstrated that the methylation status in blood samples or fluids might be different from that in tissues.^{59,60} Thus, our results should be interpreted with caution because sample types were limited to tissues in studies included in this meta-analysis.

Previous studies indicated that *P16^{INK4a}* promoter methylation was associated with poorly differentiated tumors and was different in histological subtype in ovarian cancer.^{22,43} However, we could not establish any significant correlations between *P16^{INK4a}* promoter methylation and clinicopathological features, including age, clinical stage, tumor differentiation or histological subtype in this study. Therefore, it might not be essential to predict the invasion and metastasis of ovarian cancer.

Katsaros et al⁴³ and Wiley et al⁴² reported association of *P16^{INK4a}* promoter methylation with PFS and OS in ovarian cancer, while Bhagat et al²⁰ found no significant value in predicting prognosis. In the present study, we discovered that *P16^{INK4a}* promoter methylation represented a risk factor for PFS. For OS, patients with *P16^{INK4a}* promoter methylation also had a slightly elevated risk, though the differences are not statistically significant. This trend

was also observed in other types of cancer.^{51,54} However, its statistical confirmation requires large studies. The data from TCGA also indicated that methylation level of probe cg26673943 region (located at the promoter region of *P16^{INK4a}*) in the ovarian tumor tissues was higher than normal ovarian tissues. Increased methylation of CpG island at the promoter region was negatively associated with *P16^{INK4a}* gene expression, while methylation of CpG islands at non-promoter regions was positively associated with *P16^{INK4a}* expression.

Compared with previous meta-analyses,^{31,32} our meta-analysis had several improvements. First, the development of ovarian cancer is a multistep procedure involving normal tissues, benign disease, LMP or borderline tumor, and malignant tumor.²⁰ We compared malignant ovarian cancer with LMP tumors, benign disease, and normal samples to give more rigorously to the analysis. Second, with 1,217 malignant ovarian cancer patients, 116 LMP, 271 benign patients, and 351 normal samples, the sample size in our study is much larger than that of all previous meta-analyses. Finally, we included the clinicopathological features and prognostic significance of *P16^{INK4a}* promoter methylation in ovarian cancer for more comprehensive understanding of the underlying pathogenesis of ovarian cancer. These strengths make our study a useful effort in seeking better understanding of the *P16^{INK4a}* promoter methylation in ovarian cancer.

Limitations

Several potential limitations in our current study should be noted. First, the heterogeneity was still large after subgroup analyses in the assessment of the association between *P16^{INK4a}* promoter methylation and ovarian cancer risk, which may

affect the statistical power. Second, as a retrospective study, a potential unidentified confounding information and selection bias may exist in our meta-analysis. We could not eliminate the possibility of publication bias, where positive results are likely published than negative results. Third, the total sample size was still relatively small for reliably assessing the prognostic value of *P16^{INK4a}* promoter methylation in ovarian cancer. Fourth, none of the studies included in our meta-analysis defined the region considered as promoter or provided specific methylation sites. Therefore, we are unable to establish whether or not they focused on the same sequence of *P16^{INK4a}* gene. However, the impact of methylation on transcriptional potential depends on the density of the methylated CpG islands and their location relative to the transcription start site. This highlights the importance of a uniform and full-scale reporting of study designs and outcomes. Additionally, previous researches showed that the occurrence of *P16^{INK4a}* promoter methylation may depend on the histological subtype.^{41,48,61} However, we are unable to extract sufficient data to analyze the association between *P16^{INK4a}* promoter methylation and high-grade serous carcinomas because no detailed information of *P16^{INK4a}* promoter methylation in high-grade serous carcinomas was provided in the eligible articles.

Although with certain limitations, our study is a comprehensive meta-analysis focusing on the correlation of aberrant *P16^{INK4a}* promoter methylation with the initiation, development, and prognosis of ovarian cancer to provide a new insight into the pathogenesis of ovarian cancer.

Conclusion

In conclusion, our meta-analysis suggests that aberrant methylation of *P16^{INK4a}* promoter may be essential to the initiation of ovarian cancer and in distinguishing malignant from healthy ovarian tissues. Besides, *P16^{INK4a}* promoter methylation is a potential predictive factor for poor prognosis in ovarian cancer. This study indicates the need for multicenter large-scale studies to confirm the role of *P16^{INK4a}* promoter methylation in ovarian cancer.

Acknowledgments

This study was supported by National Natural Science Funds (numbers 81472033 and 30901308), the National Science Foundation of Hubei Province (numbers 2013CFB233 and 2013CFB235), the Scientific and Technological Project of Wuhan City (number 2014060101010045), Hubei Province Health and Family Planning Scientific Research Project (WJ2015Q021), and Training Program of the Science and

Technology Innovation from Zhongnan Hospital of Wuhan University (cxpy20160054). We thank the Beijing Circle & Dot Technology Co., Ltd. for assisting in analyzing the TCGA database.

Disclosure

The authors report no conflicts of interest in this work.

References

1. Torre LA, Bray F, Siegel RL, Ferlay J, Lortet-Tieulent J, Jemal A. Global cancer statistics, 2012. *CA Cancer J Clin*. 2015;65(2):87–108.
2. Prat J. New insights into ovarian cancer pathology. *Ann Oncol*. 2012;23(Suppl 10):x111–x117.
3. Siegel RL, Miller KD, Jemal A. Cancer statistics, 2016. *CA Cancer J Clin*. 2016;66(1):7–30.
4. Bai H, Cao D, Yang J, Li M, Zhang Z, Shen K. Genetic and epigenetic heterogeneity of epithelial ovarian cancer and the clinical implications for molecular targeted therapy. *J Cell Mol Med*. 2016;20(4):581–593.
5. You JS, Jones PA. Cancer genetics and epigenetics: two sides of the same coin? *Cancer Cell*. 2012;22(1):9–20.
6. Kazanets A, Shorstova T, Hilmi K, Marques M, Witcher M. Epigenetic silencing of tumor suppressor genes: paradigms, puzzles, and potential. *Biochim Biophys Acta*. 2016;1865(2):275–288.
7. Maunakea AK, Chepelev I, Cui K, Zhao K. Intragenic DNA methylation modulates alternative splicing by recruiting MeCP2 to promote exon recognition. *Cell Res*. 2013;23(11):1256–1269.
8. Dong A, Lu Y, Lu B. Genomic/Epigenomic alterations in ovarian carcinoma: translational insight into clinical practice. *J Cancer*. 2016;7(11):1441–1451.
9. Koukoura O, Spandidos DA, Daponte A, Sifakis S. DNA methylation profiles in ovarian cancer: implication in diagnosis and therapy (Review). *Mol Med Rep*. 2014;10(1):3–9.
10. Gloss BS, Samimi G. Epigenetic biomarkers in epithelial ovarian cancer. *Cancer Lett*. 2014;342(2):257–263.
11. Sharpless NE, Depinho RA. The INK4A/ARF locus and its two gene products. *Curr Opin Genet Dev*. 1999;9(1):22–30.
12. Serrano M, Hannon GJ, Beach D. A new regulatory motif in cell-cycle control causing specific inhibition of cyclin D/CDK4. *Nature*. 1993;366(6456):704–707.
13. Qin Y, Liu JY, Li B, Sun ZL, Sun ZF. Association of low p16INK4a and p15INK4b mRNAs expression with their CpG islands methylation with human hepatocellular carcinogenesis. *World J Gastroenterol*. 2004;10(9):1276–1280.
14. Kim BN, Yamamoto H, Ikeda K, et al. Methylation and expression of p16INK4 tumor suppressor gene in primary colorectal cancer tissues. *Int J Oncol*. 2005;26(5):1217–1226.
15. Gao SJ, Zhang GF, Zhang RP. High CpG island methylation of p16 gene and loss of p16 protein expression associate with the development and progression of tetralogy of Fallot. *J Genet*. 2016;95(4):831–837.
16. Sharpless NE. INK4a/ARF: a multifunctional tumor suppressor locus. *Mutat Res*. 2005;576(1–2):22–38.
17. Moselhy SS, Kumosani TA, Kamal IH, Jalal JA, Jabaar HS, Dalol A. Hypermethylation of P15, P16, and E-cadherin genes in ovarian cancer. *Toxicol Ind Health*. 2015;31(10):924–930.
18. di Vinci A, Perdelli L, Banelli B, et al. p16(INK4a) promoter methylation and protein expression in breast fibroadenoma and carcinoma. *Int J Cancer*. 2005;114(3):414–421.
19. Shima K, Noshio K, Baba Y, et al. Prognostic significance of CDKN2A (p16) promoter methylation and loss of expression in 902 colorectal cancers: cohort study and literature review. *Int J Cancer*. 2011;128(5):1080–1094.
20. Bhagat R, Kumar SS, Vaderhobli S, et al. Epigenetic alteration of p16 and retinoic acid receptor beta genes in the development of epithelial ovarian carcinoma. *Tumour Biol*. 2014;35(9):9069–9078.

21. Bammidi LS, Neerukonda GN, Murthy S, Kanapuram RD. p16 gene alterations in human ovarian cancers: comparison between tissue and blood samples. *Int J Gynecol Cancer*. 2012;22(4):553–560.
22. Abou-Zeid AA, Azzam AZ, Kamel NA. Methylation status of the gene promoter of cyclin-dependent kinase inhibitor 2A (CDKN2A) in ovarian cancer. *Scand J Clin Lab Invest*. 2011;71(7):542–547.
23. Dhillon VS, Aslam M, Husain SA. The contribution of genetic and epigenetic changes in granulosa cell tumors of ovarian origin. *Clin Cancer Res*. 2004;10(16):5537–5545.
24. Ozdemir F, Altinisik J, Karateke A, Coksuer H, Buyru N. Methylation of tumor suppressor genes in ovarian cancer. *Exp Ther Med*. 2012;4(6):1092–1096.
25. Shen WJ, Dai DQ, Guo KJ, Xm L. Promoter hypermethylation of RASSF1A, BRCA1 and p16 gene in epithelial ovarian cancer and its clinical significance. *Chin J Cancer Prev Treat*. 2008;15(7):530–533.
26. Tam KF, Liu VW, Liu SS, et al. Methylation profile in benign, borderline and malignant ovarian tumors. *J Cancer Res Clin Oncol*. 2007;133(5):331–341.
27. Li M, Huang ZJ, Dong WH, et al. Disfigurement of p16INK4A gene expression in development of ovarian cancer and the mechanism. *Zhonghua Fu Chan Ke Za Zhi*. 2006;41(6):408–412.
28. Liu Z, Wang LE, Wang L, et al. Methylation and messenger RNA expression of p15INK4b but not p16INK4a are independent risk factors for ovarian cancer. *Clin Cancer Res*. 2005;11(13):4968–4976.
29. Makarla PB, Saboorian MH, Ashfaq R, et al. Promoter hypermethylation profile of ovarian epithelial neoplasms. *Clin Cancer Res*. 2005;11(15):5365–5369.
30. Rath A, Virmani AK, Schorge JO, et al. Methylation profiles of sporadic ovarian tumors and nonmalignant ovaries from high-risk women. *Clin Cancer Res*. 2002;8(11):3324–3331.
31. Xiao X, Cai F, Niu X, Shi H, Zhong Y. Association between P16INK4a promoter methylation and ovarian cancer: a meta-analysis of 12 published studies. *PLoS One*. 2016;11(9):e0163257.
32. Jiang Y, Yan F, Liang L, Wan Y, Liu J, Cheng W. Meta-analysis demonstrates no association between P16^{INK4A} promoter methylation and epithelial ovarian cancer. *Arch Gynecol Obstet*. 2017;295(3):697–704.
33. Stang A. Critical evaluation of the Newcastle-Ottawa scale for the assessment of the quality of nonrandomized studies in meta-analyses. *Eur J Epidemiol*. 2010;25(9):603–605.
34. Bradburn MJ, Deeks JJ, Berlin JA, Russell Localio A. Much ado about nothing: a comparison of the performance of meta-analytical methods with rare events. *Stat Med*. 2007;26(1):53–77.
35. National Institutes of Health; National Cancer Institute; National Human Genome Research Institute. *The Cancer Genome Atlas*. Available from: <https://cancergenome.nih.gov/>. Accessed August 19, 2018.
36. Cancer Genome Atlas Research Network. Integrated genomic analyses of ovarian carcinoma. *Nature*. 2011;474(7353):609–615.
37. Ho CM, Huang CJ, Huang CY, Wu YY, Chang SF, Cheng WF. Promoter methylation status of HIN-1 associated with outcomes of ovarian clear cell adenocarcinoma. *Mol Cancer*. 2012;11:53.
38. Cul'bová M, Lasabova Z, Stanclová A, et al. Metylácia vybraných tumorsupresorických génov v benigných a maligných ovariálnych nádoroch [Methylation of selected tumor-suppressor genes in benign and malignant ovarian tumors]. *Ceska /Gynekol*. 2011;76(4):274–279. Czech.
39. Gu XH, Lu Y, Ma D, Liu XS, Guo SW. Model of aberrant DNA methylation patterns and its applications in epithelial ovarian cancer. *Zhonghua Fu Chan Ke Za Zhi*. 2009;44(10):754–759.
40. Wu Q, Lothe RA, Ahlquist T, et al. DNA methylation profiling of ovarian carcinomas and their in vitro models identifies HOXA9, HOXB5, SCGB3A1, and CRABP1 as novel targets. *Mol Cancer*. 2007;6:45.
41. Yang HJ, Liu VW, Wang Y, Tsang PC, Ngan HY. Differential DNA methylation profiles in gynecological cancers and correlation with clinico-pathological data. *BMC Cancer*. 2006;6:212.
42. Wiley A, Katsaros D, Chen H, et al. Aberrant promoter methylation of multiple genes in malignant ovarian tumors and in ovarian tumors with low malignant potential. *Cancer*. 2006;107(2):299–308.
43. Katsaros D, Cho W, Singal R, et al. Methylation of tumor suppressor gene p16 and prognosis of epithelial ovarian cancer. *Gynecol Oncol*. 2004;94(3):685–692.
44. Hashiguchi Y, Tsuda H, Yamamoto K, Inoue T, Ishiko O, Ogita S. Combined analysis of p53 and RB pathways in epithelial ovarian cancer. *Hum Pathol*. 2001;32(9):988–996.
45. Brown I, Milner B, Rooney P, Haites N. Inactivation of the p16INK4A gene by methylation is not a frequent event in sporadic ovarian carcinoma. *Oncol Rep*. 2001;8(6):1359–1362.
46. Strathdee G, Appleton K, Illand M, et al. Primary ovarian carcinomas display multiple methylator phenotypes involving known tumor suppressor genes. *Am J Pathol*. 2001;158(3):1121–1127.
47. McCluskey LL, Chen C, Delgadillo E, Felix JC, Mudderspach LI, Dubeau L. Differences in p16 gene methylation and expression in benign and malignant ovarian tumors. *Gynecol Oncol*. 1999;72(1):87–92.
48. Milde-Langosch K, Ocon E, Becker G, Löning T. p16/MTS1 inactivation in ovarian carcinomas: high frequency of reduced protein expression associated with hyper-methylation or mutation in endometrioid and mucinous tumors. *Int J Cancer*. 1998;79(1):61–65.
49. Shih YC, Kerr J, Liu J, et al. Rare mutations and no hypermethylation at the CDKN2A locus in epithelial ovarian tumours. *Int J Cancer*. 1997;70(5):508–511.
50. Ferlay J, Soerjomataram I, Dikshit R, et al. Cancer incidence and mortality worldwide: sources, methods and major patterns in GLOBOCAN 2012. *Int J Cancer*. 2015;136(5):E359–E386.
51. Xing X, Cai W, Shi H, et al. The prognostic value of CDKN2A hypermethylation in colorectal cancer: a meta-analysis. *Br J Cancer*. 2013;108(12):2542–2548.
52. Li J, Zhou C, Zhou H, et al. The association between methylated CDKN2A and cervical carcinogenesis, and its diagnostic value in cervical cancer: a meta-analysis. *Ther Clin Risk Manag*. 2016;12:1249–1260.
53. Wang X, Zhu YB, Cui HP, Yu TT, Ti Y. Aberrant promoter methylation of p15 (INK4b) and p16 (INK4a) genes may contribute to the pathogenesis of multiple myeloma: a meta-analysis. *Tumour Biol*. 2014;35(9):9035–9043.
54. Tang B, Li Y, Qi G, et al. Clinicopathological significance of CDKN2A promoter hypermethylation frequency with pancreatic cancer. *Sci Rep*. 2015;5:13563.
55. Hogan C. Impact of interactions between normal and transformed epithelial cells and the relevance to cancer. *Cell Mol Life Sci*. 2012;69(2):203–213.
56. Fraser HB, Lam LL, Neumann SM, Kobor MS. Population-specificity of human DNA methylation. *Genome Biol*. 2012;13(2):R8.
57. Kader F, Ghai M. DNA methylation-based variation between human populations. *Mol Genet Genomics*. 2017;292(1):5–35.
58. Kurdyukov S, Bullock M. DNA methylation analysis: choosing the right method. *Biology*. 2016;5(1):3.
59. Chang H, Yi B, Li L, et al. Methylation of tumor associated genes in tissue and plasma samples from liver disease patients. *Exp Mol Pathol*. 2008;85(2):96–100.
60. Zhu W, Qin W, Hewett JE, Sauter ER. Quantitative evaluation of DNA hypermethylation in malignant and benign breast tissue and fluids. *Int J Cancer*. 2010;126(2):474–482.
61. Niederacher D, Yan HY, An HX, Bender HG, Beckmann MW. CDKN2A gene inactivation in epithelial sporadic ovarian cancer. *Br J Cancer*. 1999;80(12):1920–1926.

Cancer Management and Research**Dovepress****Publish your work in this journal**

Cancer Management and Research is an international, peer-reviewed open access journal focusing on cancer research and the optimal use of preventative and integrated treatment interventions to achieve improved outcomes, enhanced survival and quality of life for the cancer patient. The manuscript management system is completely online and includes

a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/cancer-management-and-research-journal>