Natural borneol is a novel chemosensitizer that enhances temozolomide-induced anticancer efficiency against human glioma by triggering mitochondrial dysfunction and reactive oxide species-mediated oxidative damage

Wen-Jian Liu,^{1,*} Yi-Bo Yin,^{1,*}
Jing-Yi Sun,^{2,*} Sai Feng,³
Jin-Kui Ma,⁴ Xiao-Yan Fu,⁵
Ya-Jun Hou,⁵ Ming-Feng
Yang,⁵ Bao-Liang Sun,^{1,5}
Cun-Dong Fan⁵

¹Department of Neurology, Affiliated Hospital of Taishan Medical University, Taian, Shandong, People's Republic of China; ²Department of Orthopaedics, Wonju Severance Christian Hospital, Yonsei University Wonju College of Medicine, Wonju, Gangwon, Korea; ³Guangzhou New BenFu Technology Co., Ltd, Guangzhou, Guangdong, People's Republic of China; ⁴Faculty of Bioresource Sciences, Akita Prefectural University, Shimoshinjo-Nakano, Akita-shi, Akita, Japan; ⁵Key Lab of Cerebral Microcirculation in Universities of Shandong, Taishan Medical University, Taian, Shandong, People's Republic of China

*These authors contributed equally to this work

Correspondence: Bao-Liang Sun;
Cun-Dong Fan
Department of Neurology, Taishan Medical University, Yingsheng East Road 2, Taian, Shandong 271000, People's Republic of China
Tel/fax +86 538 623 0030;
+86 538 623 0027
Email tbsun66@163.com;
tcdfan66@163.com

Background: Temozolomide (TMZ)-based chemotherapy represents an effective way for treating human glioma. However, its clinical application is limited because of its side effects and resistance to standard chemotherapy. Hence, the search for novel chemosensitizers to augment their anticancer efficiency has attracted much attention. Natural borneol (NB) has been identified as a potential chemosensitizer in treating human cancers. However, the synergistic effect and mechanism of NB and TMZ in human glioma have not been investigated yet.

Materials and methods: U251 human glioma cells were cultured, and the cytotoxicity and apoptosis of NB and/or TMZ were examined by MTT assay, flow cytometric analysis and Western blot. Nude mice tumor model was also employed to evaluate the in vivo anticancer effect and mechanism.

Results: The results showed that the combined treatment of NB and TMZ more effectively inhibited human glioma growth via triggering mitochondria-mediated apoptosis in vitro, accompanied by the caspase activation. Combined treatment of NB and TMZ also caused mitochondrial dysfunction through disturbing Bcl-2 family expression. Further investigation revealed that NB enhanced TMZ-induced DNA damage through inducing reactive oxide species (ROS) overproduction. Moreover, glioma tumor xenograft growth in vivo was more effectively inhibited by the combined treatment with NB and TMZ through triggering apoptosis and anti-angiogenesis.

Conclusion: Taken together, our findings validated that the strategy of using NB and TMZ could be a highly efficient way to achieve anticancer synergism.

Keywords: glioma, borneol, temozolomide, DNA damage, apoptosis

Introduction

Glioma is the most common primary malignant tumor in the adult central nervous system.¹ The 5-year survival rate is less than 5%.²⁻⁴ Because of its high recurrence rate, single surgery, chemotherapy or radiotherapy was not appropriate for the treatment of human glioma.⁵ Combined therapies have emerged as excellent strategies. Studies have shown that postoperative chemotherapy had achieved great outcome for human glioma.⁶ However, the existence of blood-brain barrier (BBB) confined the clinical application of chemotherapy.⁷⁻⁹ Hence, the search for novel chemosensitizers

to promote the BBB permeability and improve the chemotherapy of human glioma is urgently needed.¹⁰

Natural borneol (NB), a simple bicyclic monoterpene, was isolated from the resin of *Dryobalanops aromatica* Gaertn F, which showed multiple pharmacological properties. It is reported that NB could improve the oral bioavailability of anti-tumor drugs by regulating the permeability of the BBB.¹⁰ Increased evidence have confirmed that NB can increase cell membrane mobility and drug solubility, decrease mucus stickiness and elasticity and inhibit drug efflux.^{11,12} Recently, studies showed that NB can act as a chemosensitizer to enhance anticancer drug-induced apoptosis in human cancer cells through triggering DNA damage and regulation of MAPKs and AKT pathways.^{13,14} However, the potential of NB as a chemosensitizer in human glioma has not been elucidated yet.

In this study, U251 human glioma cells and nude mice-bearing glioma xenograft were employed to evaluate the combined effect and mechanism of NB and TMZ against human glioma cell growth in vitro and in vivo. The results indicated that NB had the potential to enhance TMZ-mediated anticancer efficiency against human glioma through triggering mitochondrial dysfunction and reactive oxide species (ROS)-induced DNA damage.

Materials and methods

Chemicals

NB was obtained from BenFu Technology Co., Ltd. (Guangzhou, China) DMEM, PBS, FBS, 2',7'-dichlorofluorescein diacetate (DCFH-DA) probe and mitochondria-targeted MitoSOX probe (M36008) were all purchased from Thermo Fisher Scientific (Waltham, MA, USA). Temozolomide (TMZ), MTT and propidium iodide (PI) were obtained from Sigma-Aldrich Co. (St Louis, MO, USA). TUNEL-DAPI kit, MitoSOX probe and BCA assay kit were purchased from Beyotime Institute of Biotechnology (Shanghai, China). All antibodies used in this study were obtained from Cell Signaling Technology (Beverly, MA, USA). All solvents used were of high performance liquid chromatography (HPLC) grade.

Cell culture, drug treatment and determination of cell viability

U251 human glioma cells were acquired from American Type Culture Collection (ATCC, Manassas, VA, USA) and were cultured in DMEM with 10% FBS, penicillin (100 U/mL) and streptomycin (100 U/mL) at 37°C in a 5% CO₂ incubator. Cell was seeded in 96-well plates and treated with 10–160 µM TMZ or 5–80 µg/mL of NB for 48 hours. For combined treatment, the cells were treated with TMZ (20 and 40 µM)

and NB (40 and 80 µg/mL) for 48 hours. Cell viability of U251 cells was detected by the MTT assay. After treatment, 20 µL/well of MTT solution (5 mg/mL) was added to the cells and incubated for 5 hours at 37°C. The supernatants were aspirated and 150 µL of DMSO was added. Then, the absorbance was measured on a scanning multi-well spectrophotometer. Cell viability was calculated as a percentage of control (as 100%).

Flow cytometric analysis

Cell cycle distribution and apoptosis were assessed by flow cytometric analysis as previously used.¹⁵ The cells treated with NB and/or TMZ were harvested by centrifugation and washed with PBS, and then the cells were fixed with 70% alcohol and stained with PI solution for 2 hours. Labeled cells were washed with PBS and then analyzed by flow cytometry (FCM). The proportions of cells in G0/G1, S and G2/M phases were represented as DNA histograms. Apoptotic cells with hypodiploid DNA contents were measured by quantifying the sub-G1 peak. For each experiment 10,000 events per sample were recorded.

Detection of early and late apoptoses

Early and late apoptoses of U251 cells were detected by Annexin V–PI staining and TUNEL-DAPI staining, respectively. Cells cultured in a chamber slide were treated with 3.7% formaldehyde for 10 minutes and permeabilized in PBS with 0.1% Triton X-100. Then, the cells were administered with the Annexin V–PI staining and TUNEL-DAPI staining kits for measuring the early and late apoptoses, respectively. The experiments were conducted according to the manufacturer's instructions. The images were obtained from a fluorescence microscope.

Evaluation of mitochondrial dysfunction

Mitochondrial function was evaluated by the mitochondrial membrane potential ($\Delta\psi_m$) and mitochondrial morphology, which were examined by JC-1 and Mito-Tracker probes, respectively. Briefly, the cells seeded in a 6 cm plate were exposed to NB and/or TMZ. After treatment, the cells were washed and incubated with 10 µM JC-1 or Mito-Tracker for 15 minutes in the dark. Then, the cells were washed and imaged under an inverted fluorescence microscope.

Measurement of ROS generation and superoxide anion

The intracellular ROS and superoxide anion were measured by DCFH-DA and MitoSOX probes in live cells, respectively. Briefly, after treatment with NB and/or TMZ, the cells

were incubated with 10 μ M DCFH-DA or 0.5 μ M MitoSOX. After reaction, the cells were washed and observed by an inverted fluorescence microscope for the detection of ROS (green fluorescence) and superoxide anion (red fluorescence). The images shown here were obtained from three independent experiments.

Western blot analysis

Total protein was prepared from the cells treated with NB and/or TMZ and quantified by the BCA kit. SDS-PAGE was performed in 10% tricine gel, and 40 μ g/lane total protein was loaded. After electrophoresis, proteins were transferred from the gel onto a nitrocellulose membrane at 110 V for 1 hour. Then, the membrane was blocked with 5% nonfat milk for 1 hour, and the membranes were hatched with primary antibodies at 1:1,000 dilution in 5% nonfat milk over night at 4°C. Finally, the membrane was treated with horseradish peroxidase-coupled secondary antibody at 1:2,000 for 1 hour at room temperature. Protein was imaged under an Image-System (Bio-Rad Laboratories Inc., Hercules, CA, USA). β -Actin was used as the equal loading.

In vivo study

U251 cell suspension (10^9 cells) was inoculated in the armpit of 6-week-old Balb/c nude mice. After 1-week growth, the tumor-bearing mice were randomly divided into four groups (10 mice/group). The control group was given the corresponding solvent, and the other groups were given 20 mg/kg/day NB and/or 10 mg/kg/day TMZ every day for 2 weeks. At the end of the experiment, tumor was separated, and the tumor volume and weight were measured, using the formula: $\text{volume} = l \times w^2 / 2$, with l being the maximal length and w being the width. Sections of tumors were used for Western blotting and immunohistochemical (IHC) assay. All animal experiments were approved by the Animal Experimentation Ethics Committee of Shandong in China.

Statistical analysis

All data and images were done from three independent experiments at least, and all data were indicated as mean $\pm$ standard error. Statistical analysis was performed by SPSS statistical package (SPSS 13.0 for Windows; SPSS Inc., Chicago, IL, USA). The data of two groups were analyzed by Student's t -test, and the data of three or more groups were analyzed by one-way analysis of variance multiple comparisons to assess significant differences, and differences with $P < 0.05$ or $P < 0.01$ were considered statistically significant. Bars with different letters indicate statistical difference at the $P < 0.05$ level.

Ethical statement

This study was performed in Key Lab of Cerebral Micro-circulation in Universities of Shandong, Taishan Medical University. All the in vitro and in vivo experiments were performed in accordance with the relevant guidelines and regulations of Taishan Medical University. The in vivo experiments were approved by the Taishan Medical University Ethics Committee. All surgeries were performed under 10% chloral hydrate, and all efforts were made to minimize suffering.

Results

NB enhances TMZ-induced cytotoxicity in human glioma cells

In order to optimize the dosages for combined treatment, the cytotoxicity of NB or TMZ alone toward human glioma cells was screened by the MTT assay. As shown in Figure 1A and B, treatment of U251 cells with TMZ (0–40 μ M) for 48 hours caused no significant cytotoxicity. Only treatment with 80 and 160 μ M TMZ significantly inhibited U251 cell growth. U251 cells exposed to NB (0–80 μ M) for 48 hours showed no cytotoxicity. However, combined treatment of NB and TMZ significantly inhibited U251 cell growth (Figure 1C). For instance, the treatment of cells with 80 μ M NB or 40 μ M TMZ alone showed no significant changes in the U251 cell viability. However, combined treatment of NB (80 μ M) and TMZ (40 μ M) significantly inhibited the cell viability to 52.8%. Similar cytotoxicity effect of NB and/or TMZ was also detected in U87 human glioma cells and BV-2 mice microglial cells (Figure S1). Combined treatment-induced growth inhibitory effect was further confirmed by the morphological changes in U251 cells. As shown in Figure 1D, the treatment of U251 cells with NB or TMZ alone caused no significant change in cell morphology. However, combined treatment of NB and TMZ dramatically resulted in the cell morphological changes, such as the cell shrinkage, decrease in the cell number and loss of cell-to-cell contact. Taken together, these results indicated that NB had the potential to enhance TMZ-induced cytotoxicity in human glioma cells.

NB enhances TMZ-induced apoptosis in human glioma cells

To verify the cell death mechanism, U251 cells after treatment with NB and/or TMZ were analyzed by FCM. As shown in Figure 2A, the treatment of U251 cells with NB or TMZ alone caused no significant apoptosis and no effects on cell cycle distribution. However, enhanced cell apoptosis was dramatically detected after combined treatment of NB and TMZ.

Figure 1 NB enhances TMZ-induced cytotoxicity in human glioma cells.

Notes: Cytotoxicity of TMZ (**A**) and NB (**B**) toward U251 cells. (**C**) NB enhances TMZ-induced cytotoxicity in U251 cells. U251 cells were seeded in 96-well plate and treated with NB and/or TMZ for 48 hours. The cell viability was detected by the MTT assay. (**D**) Cell morphological changes. Cells were treated with 80 $\mu\text{g/mL}$ NB and/or 40 μM TMZ for 48 hours, and cell morphology was observed by a microscope (magnification, 200 $\times$). Scale bar=50 μm . All data and images were obtained from three independent trials. Bars with "*" or "**" indicate the statistical difference at the $P<0.05$ and $P<0.01$ levels, respectively. Bars with different characters indicate the statistical difference at the $P<0.05$ level.

Abbreviations: NB, natural borneol; TMZ, temozolomide.

For instance, the combined treatment of NB (40 and 80 μM) and TMZ (20 μM) triggered U251 cell apoptosis to 28.1% and 49.2%, respectively. Annexin V-PI and TUNEL-DAPI staining further confirmed combined treatment-induced early and late apoptoses, respectively (Figure 2B and C). The statistical analysis of early apoptosis further confirmed this conclusion (Figure S2). Moreover, the enhanced cell apoptosis was further confirmed in the protein level. As shown in Figure 2D, enhanced activation of caspase-3, -7 and -9 was detected after combined treatment of NB and TMZ. The activation of caspase-7 and caspase-9 indicated the contribution of mitochondria-mediated apoptosis. Taken together, these

results demonstrated that NB had the potential to enhance TMZ-induced apoptosis in human glioma cells.

NB enhances TMZ-induced mitochondria dysfunction by regulating Bcl-2 family

Mitochondrion integrates the internal and external apoptotic signals and plays a significant role in deciding cells' fate. The depletion of mitochondrial membrane potential ($\Delta\psi_m$) will lead to the release of multiple apoptosis factors from the mitochondria into the cytoplasm. Hence, both the $\Delta\psi_m$ and the mitochondrial morphological change were evaluated using JC-1 and Mito-Tracker probes, respectively. As shown

Figure 2 NB enhances TMZ-induced apoptosis in human glioma cells.

Notes: (A) FCM analysis of cell apoptosis. Cells seeded in six-well plate were treated with NB and/or TMZ for 48 hours. Then, the cells were stained with PI and analyzed by FCM. Detection of cell early apoptosis (B) and late apoptosis (C). Scale bar in (B) is 20 μm. Cells after treatment were stained by Annexin V-PI and TUNEL-DAPI kits for the detection of early apoptosis and late apoptosis, respectively. (D) Caspase activation. Total protein was prepared, and the protein expression was examined by the Western blotting method. Scale bar in (C) is 50 μm. All data and images were obtained from three independent experiments.

Abbreviations: FCM, flow cytometry; NB, natural borneol; PI, propidium iodide; TMZ, temozolomide.

in Figure 3A, U251 cells exposed to treatment with NB or TMZ alone showed no significant change in the $\Delta\psi_m$ and the mitochondrial morphology. However, the combined treatment of NB and TMZ significantly caused the loss of $\Delta\psi_m$, as reflected by the fluorescence shift from red to green. The statistical analysis of $\Delta\psi_m$ further confirmed this conclusion (Figure S3). Moreover, combined treatment also resulted in the mitochondrial fragmentation. As shown in Figure 3B, the mitochondria in healthy cells and in cells treated with NB or TMZ alone all showed filamentous network with extensive interconnection throughout the cytoplasm. However, combined treatment markedly caused the mitochondrial fragmentation from protonema to punctiform. Bcl-2 family proteins have been depicted as essential regulators of the

mitochondrial apoptosis pathway. Therefore, the Bcl-2 family was also identified. As shown in Figure 3C, exposure of cells to NB or TMZ alone caused no obvious effects on the Bcl-2 family expression. Combined treatment of NB and TMZ effectively increased the expression of Bax and Bad, but decreased the expression of Bcl-2 and Bcl-XL. Taken together, these abovementioned results revealed that NB had the potential to enhance TMZ-induced mitochondrial dysfunction through affecting Bcl-2 family balance.

NB enhances TMZ-induced DNA damage through ROS accumulation

ROS could enter nucleus to cause DNA damage by which many anticancer agents induce cancer cell apoptosis.¹⁶

Figure 3 NB enhances TMZ-induced mitochondria dysfunction by regulating Bcl-2 family.

Notes: NB enhances TMZ-induced depletion of $\Delta\psi_m$ (A) and mitochondrial fragmentation (B). Scale bar in (A) is 20 μm. Scale bar in (B) is 10 μm. Cells after treatment were stained by JC-1 and Mito-Tracker probes, respectively. (C) Bcl-2 family expression. All images were obtained from three independent trials.

Abbreviations: NB, natural borneol; TMZ, temozolomide.

Therefore, the oxidative status in U251 cells treated with NB and/or TMZ was examined. As shown in Figure 4A, treatment with NB or TMZ alone only caused slight ROS generation. However, the combined treatment of NB and TMZ significantly caused ROS accumulation, as indicated by the enhanced green fluorescence. The statistical analysis of ROS generation further confirmed combined treatment-induced ROS accumulation (Figure S4). Furthermore, the mitochondrial superoxide overproduction was also detected by MitoSOX, a specific mitochondria-targeted probe, and the results suggested that enhanced production of superoxide was vividly observed in U251 cells after combined treatment with NB and TMZ (Figure 4B). ROS generation caused DNA damage, which is effective to combat human cancers. Hence, combined treatment-induced DNA damage was investigated. As shown in Figure 4C, no significant DNA damage was observed after treatment with NB or TMZ alone. As expected, combined treatment of NB and TMZ significantly caused DNA damage, as demonstrated by the enhanced phosphorylation level of histone (Ser139), p53 (Ser15), ATR (Ser428) and ATM (Ser1981), all of which were DNA damage markers.¹⁷ ROS inhibition by glutathione (GSH) effectively inhibited

combined treatment-induced cell death (Figure S5), which further confirmed the significant role of ROS in combined treatment-induced cell growth inhibition against human glioma cells. Taken together, these results suggested that NB had the potential to enhance TMZ-induced DNA damage through ROS overproduction.

NB enhances TMZ-induced anticancer efficiency in vivo

To explore the combined treatment-induced anticancer effect in vivo, the immunodeficient nude mice were employed. The results showed that treatment with NB or TMZ alone only slightly inhibited the glioma growth, as indicated by the decreased tumor volume and tumor weight. However, combined treatment of NB and TMZ significantly inhibited tumor volume and tumor weight compared to that in treatment with NB or TMZ alone (Figure 5A and B). Additionally, combined treatment-induced anticancer mechanism was similar to that of the in vitro mechanism. As shown in Figure 5C, after combined treatment of NB and TMZ, enhanced cell apoptosis and DNA damage were significantly observed, as indicated by the increased expression of active caspase-3 and the phosphorylation level

Figure 4 NB enhances TMZ-induced DNA damage through ROS accumulation.

Notes: NB enhances TMZ-induced ROS (**A**) and superoxide anion (**B**) generation. The intracellular superoxide anion and ROS were measured by MitoSOX and DCFH-DA probes, respectively. The experiments were conducted according to the "Materials and methods" section. (**C**) Detection of DNA damage. Total protein was prepared, and the protein expression was examined by the Western blotting method. Scale bar=20 μm. All data and images were obtained from three independent experiments.

Abbreviations: NB, natural borneol; ROS, reactive oxide species; TMZ, temozolomide.

Figure 5 NB enhances TMZ-induced anticancer efficiency in vivo.

Notes: NB enhances TMZ-induced inhibition against tumor volume (**A**) and tumor weight (**B**). (**C**) NB enhances TMZ-induced apoptosis and DNA damage. The protein expression was examined by Western blotting methods. (**D**) NB enhances TMZ-induced anti-proliferation and anti-angiogenesis. The protein expression was examined by IHC methods. Scale bar=50 μm. Bars with different characters indicate the statistical difference at the $P < 0.05$ level. All data and images were obtained from three independent experiments.

Abbreviations: IHC, immunohistochemical; NB, natural borneol; TMZ, temozolomide.

of histone (Ser139) and p53 (Ser15) (Figure 5C). The staining of Ki-67 and active caspase-3 in tumor tissue further confirmed combined treatment-induced anti-proliferation and apoptosis, respectively (Figure 5D). Moreover, combined treatment also abolished the tumor angiogenesis. As shown in Figure 5D, HE staining revealed that combined treatment significantly decreased the number of erythrocyte-containing blood vessels. The staining of CD-31, a marker of vascular endothelial cells, further confirmed the combined treatment-induced anti-angiogenesis. Taken together, these results suggested that NB had the potential to enhance TMZ-induced anticancer efficiency against human glioma *in vivo*.

Discussion

Single agent-based chemotherapy was not appropriate for the treatment of human glioma. Recently, combined treatment of chemosensitizer and chemotherapeutic drugs has been applied in clinic to improve the anti-tumor effect. Novel chemosensitizers can improve the efficiency of existing anti-cancer drugs and overcome their resistance and side effects, and hence has been accepted as a new strategy in the field of chemistry and pharmacology.¹⁸ Induction of tumor cell apoptosis and/or cell cycle arrest all contributed to the cytotoxic mechanisms for anti-cancer effect of medicines to kill cancer cells.¹⁹ Apoptosis plays an important role in the development, homeostasis and prevention of cancer. Mitochondria-mediated (intrinsic) pathway and death receptor-mediated (extrinsic) pathway can be activated in response to apoptotic stimuli by induction of a series of caspase cascades.^{20,21} In this study, combined treatment of NB and TMZ significantly induced U251 cell apoptosis accompanied by caspase activation, indicating that NB had the potential to enhance TMZ-induced apoptosis against human glioma.

Mitochondria as the center for apoptotic cascades converges apoptosis signals from extrinsic and intrinsic apoptotic pathways, which play key role in regulating mitochondria-mediated apoptosis.²⁰ Depletion of mitochondrial membrane potential will result in the opening of the mitochondrial permeability transition pore (MPTP), and the release of cytochrome c, apoptosis-inducing factors (AIFs), SMAC/Diablo and endonuclease G, which eventually leads to the caspase activation and apoptotic cascades.²² In this study, combined treatment of NB and TMZ significantly caused the mitochondrial dysfunction, as convinced by the loss of mitochondrial membrane potential and mitochondrial fragmentation, indicating that NB had the potential to enhance TMZ-induced mitochondrial dysfunction. Bcl-2 family protein, including pro-apoptotic members (such as Bax, Bid, Bad and Bim) and anti-apoptotic members (such as

Bcl-2, Bcl-XL, Mcl-1 and Bcl-w), plays an important role in triggering mitochondrial apoptosis.²³ Bcl-2 and Bcl-XL can bind the outer membrane of mitochondria and block efflux of cytochrome c.²⁴ Overexpression of Bcl-2 and Bcl-XL in many cancer cells is one of the main causes of the resistance toward chemotherapeutic agents and radiotherapy.²⁵ Hence, the balance of pro-apoptotic and anti-apoptotic members decides the stability and the integrity of the mitochondrial membrane.²⁶ In this study, combined treatment of NB and TMZ significantly caused the imbalance of Bcl-2 family expression, which eventually resulted in mitochondria-mediated apoptosis in human glioma cells.

ROS, such as superoxide anion, hydrogen peroxide and hydroxyl radical, have been determined as potential regulators of apoptosis by modulating the extrinsic and intrinsic pathways of apoptosis.²⁷ ROS may assault cell membrane lipids, proteins and DNA, cause oxidative damage and lead to cell apoptosis. Oxidative stress can irritate metabolic activity and cause mitochondrial dysfunction. Excessive production of ROS will lead to the accumulation of oxidized products of DNA, such as DNA strand breaks (DSB).²⁸ p53 as a tumor suppressor can be activated in response to DNA damage and plays a key role in regulating cell cycle arrest and/or apoptosis.²⁹ Activation of p53 can regulate the p21 expression that inhibits cyclin-dependent kinase activity.³⁰ In this study, combined treatment of NB and TMZ significantly caused ROS accumulation, which eventually leads to the DNA damage, as convinced by the activation of DNA damage makers, such as ATM, ATR, p53 and histone. These results come to a conclusion that NB has the potential to enhance TMZ-induced DNA damage through ROS overproduction.

Conclusion

Our findings demonstrated that NB has the potential to enhance TMZ-mediated anticancer efficiency against human glioma through triggering mitochondrial dysfunction and ROS-induced DNA damage, which validated that the strategy of using NB and TMZ could be an effective way for therapy of human glioma.

Acknowledgment

This study was supported by the National Natural Science Foundation of China No 81501106 to C-DF and No 81471212 to B-LS.

Author contributions

All authors contributed toward data analysis, drafting and critically revising the paper and agree to be accountable for all aspects of the work.

Disclosure

The authors report no conflicts of interest in this work.

References

- Ostrom QT, Gittleman H, Farah P, et al. CBTRUS statistical report: Primary brain and central nervous system tumors diagnosed in the United States in 2006–2010. *Neuro Oncol*. 2013;15 Suppl 2:ii1–ii56.
- Carlsson SK, Brothers SP, Wahlestedt C. Emerging treatment strategies for glioblastoma multiforme. *EMBO Mol Med*. 2014;6(11):1359–1370.
- Chen W, Zheng R, Baade PD, et al. Cancer statistics in China, 2015. *CA Cancer J Clin*. 2016;66(2):115–132.
- Chen J, McKay RM, Parada LF. Malignant glioma: lessons from genomics, mouse models, and stem cells. *Cell*. 2012;149(1):36–47.
- Maule F, Bresolin S, Rampazzo E, et al. Annexin 2A sustains glioblastoma cell dissemination and proliferation. *Oncotarget*. 2016;7(34):54632–54649.
- Hart MG, Garside R, Rogers G, Stein K, Grant R. Temozolomide for High Grade Glioma. *Cochrane Database Syst Rev*. 2013;(4):CD007415.
- Abbott NJ, Patabendige AA, Dolman DE, Yusof SR, Begley DJ. Structure and function of the blood-brain barrier. *Neurobiol Dis*. 2010;37(1):13–25.
- Trost A, Lange S, Schroedl F, et al. Brain and Retinal Pericytes: Origin, Function and Role. *Front Cell Neurosci*. 2016;10:20.
- Attwell D, Mishra A, Hall CN, O'Farrell FM, Dalkara T. What is a pericyte? *J Cereb Blood Flow Metab*. 2016;36(2):451–455.
- Berghiche H, Houamria M, Smaghe G, Soltani N. Activity of kk-42 in combined treatment with RH-0345 or 20-hydroxyecdysone on morphometric measurements and free ecdysteroid in eggs of mealworms. *Commun Agric Appl Biol Sci*. 2005;70(4):837–841.
- Zhou Y, Li W, Chen L, Ma S, Ping L, Yang Z. Enhancement of intestinal absorption of akebia saponin D by borneol and probenecid in situ and in vitro. *Environ Toxicol Pharmacol*. 2010;29(3):229–234.
- Shen Q, Li X, Li W, Zhao X. Enhanced intestinal absorption of daidzein by borneol/menthol eutectic mixture and microemulsion. *AAPS Pharm Sci Tech*. 2011;12(4):1044–1049.
- Su J, Lai H, Chen J, et al. Natural borneol, a monoterpenoid compound, potentiates selenocystine-induced apoptosis in human hepatocellular carcinoma cells by enhancement of cellular uptake and activation of ROS-mediated DNA damage. *PLoS One*. 2013;8(5):e63502.
- Chen J, Li L, Su J, Li B, Chen T, Wong YS. Synergistic apoptosis-inducing effects on A375 human melanoma cells of natural borneol and curcumin. *PLoS One*. 2014;9(6):e101277.
- Chen T, Zheng W, Wong YS, Yang F. Mitochondria-mediated apoptosis in human breast carcinoma MCF-7 cells induced by a novel selenadi-azole derivative. *Biomed Pharmacother*. 2008;62(2):77–84.
- Mizutani H. Mechanism of DNA damage and apoptosis induced by anticancer drugs through generation of reactive oxygen species. *Yakugaku Zasshi*. 2007;127(11):1837–1842.
- Sancar A, Lindsey-Boltz LA, Unsal-Kaçmaz K, Linn S. Molecular mechanisms of mammalian DNA repair and the DNA damage checkpoints. *Annu Rev Biochem*. 2004;73:39–85.
- Kruh GD, Goldstein LJ. Doxorubicin and multidrug resistance. *Curr Opin Oncol*. 1993;5(6):1029–1034.
- Chen J, Li L, Su J, Chen T. Natural borneol enhances bisdemethoxy-curcumin-induced cell cycle arrest in the G2/M phase through up-regulation of intracellular ROS in HepG2 cells. *Food Funct*. 2015;6(3):740–748.
- Kaufmann SH, Earnshaw WC. Induction of apoptosis by cancer chemotherapy. *Exp Cell Res*. 2000;256(1):42–49.
- Riedl SJ, Shi Y. Molecular mechanisms of caspase regulation during apoptosis. *Nat Rev Mol Cell Biol*. 2004;5(11):897–907.
- van Gurp M, Festjens N, van Loo G, Saelens X, Vandenabeele P. Mitochondrial intermembrane proteins in cell death. *Biochem Biophys Res Commun*. 2003;304(3):487–497.
- Krantic S, Mechawar N, Reix S, Quirion R. Apoptosis-inducing factor: a matter of neuron life and death. *Prog Neurobiol*. 2007;81(3):179–196.
- Hikita H, Takehara T, Shimizu S, et al. Mcl-1 and Bcl-xL cooperatively maintain integrity of hepatocytes in developing and adult murine liver. *Hepatology*. 2009;50(4):1217–1226.
- Edlich F, Banerjee S, Suzuki M, et al. Bcl-x(L) retrotranslocates Bax from the mitochondria into the cytosol. *Cell*. 2011;145(1):104–116.
- Martinou JC, Youle RJ. Mitochondria in apoptosis: Bcl-2 family members and mitochondrial dynamics. *Dev Cell*. 2011;21(1):92–101.
- Pelicano H, Carney D, Huang P. ROS stress in cancer cells and therapeutic implications. *Drug Resist Updat*. 2004;7(2):97–110.
- Circu ML, Ty A. Reactive oxygen species, cellular redox systems, and apoptosis. *Free Radical Bio Med*. 2010;48:749–762.
- Cui Q, Yu JH, Wu JN, et al. P53-mediated cell cycle arrest and apoptosis through a caspase-3-independent, but caspase-9-dependent pathway in oridonin-treated MCF-7 human breast cancer cells. *Acta Pharmacol Sin*. 2007;28(7):1057–1066.
- Boulaire J, Fotedar A, Fotedar R. The functions of the cdk-cyclin kinase inhibitor p21WAF1. *Pathol Biol*. 2000;48(3):190–202.

Supplementary materials

Figure S1 Combined treatment-induced cytotoxicity in (A) U87 and (B) BV-2 cells.

Notes: U87 or BV-2 cells were treated with 80 μg/mL NB and/or 40 μM TMZ for 48 hours. Cell viability was detected by the MTT assay. All data were obtained from three independent trials. Bars with different letters indicate statistical difference at the $P < 0.05$ level.

Abbreviations: NB, natural borneol; TMZ, temozolomide.

Figure S2 Statistic analysis of early apoptosis.

Notes: U251 cells after treatment with NB and/or TMZ were stained by Annexin V-PI, and the early apoptotic cells (green fluorescence) were quantified by statistical analysis. All data were expressed as mean $\pm$ SD. Bars with different letters indicate statistical difference at the $P < 0.05$ level.

Abbreviations: NB, natural borneol; PI, propidium iodide; TMZ, temozolomide.

Figure S3 Mitochondrial membrane potential ($\Delta\Psi_m$).

Notes: U251 cells after treatment with NB and/or TMZ were stained by JC-1 probe, and the $\Delta\Psi_m$ was quantified by indicating the red fluorescence. All data were expressed as mean $\pm$ SD. Bars with different letters indicate statistical difference at the $P < 0.05$ level.

Abbreviations: NB, natural borneol; TMZ, temozolomide.

Figure S4 Statistic analysis of ROS generation.

Notes: U251 cells after treatment with NB and/or TMZ were stained by DCFH-DA probe, and the ROS generation was quantified by indicating the green fluorescence. All data were expressed as mean±SD. Bars with different letters indicate statistical difference at the $P<0.05$ level.

Abbreviations: NB, natural borneol; TMZ, temozolomide.

Figure S5 Effect of ROS inhibition on combined treatment-induced cell death in U251 cells.

Notes: U251 cells were pre-treated with 5 mM GSH for 2 hours before combined treatment. Cell viability was detected by the MTT assay. All data were obtained from three independent trials. Bars with different letters indicate statistical difference at the $P<0.05$ level.

Abbreviations: GSH, glutathione; NB, natural borneol; TMZ, temozolomide.

OncoTargets and Therapy

Publish your work in this journal

OncoTargets and Therapy is an international, peer-reviewed, open access journal focusing on the pathological basis of all cancers, potential targets for therapy and treatment protocols employed to improve the management of cancer patients. The journal also focuses on the impact of management programs and new therapeutic agents and protocols on

Submit your manuscript here: <http://www.dovepress.com/oncotargets-and-therapy-journal>

patient perspectives such as quality of life, adherence and satisfaction. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Dovepress