

# Using knowledge translation for quality improvement: an interprofessional education intervention to improve thromboprophylaxis among medical inpatients

Melissa K Myers<sup>1</sup>  
 Claire L Jansson-Knodell<sup>1</sup>  
 Darrell R Schroeder<sup>2</sup>  
 John G O'Meara<sup>3</sup>  
 Sara L Bonnes<sup>4</sup>  
 John T Ratelle<sup>5</sup>

<sup>1</sup>Department of Medicine, Mayo Clinic, Rochester, MN, USA;

<sup>2</sup>Department of Health Sciences Research, Division of Biomedical Statistics and Informatics, Mayo Clinic, Rochester, MN, USA; <sup>3</sup>Department of Pharmacy Services, Mayo Clinic, Rochester, MN, USA; <sup>4</sup>Department of Medicine, Division of General Internal Medicine, Mayo Clinic, Rochester, MN, USA; <sup>5</sup>Department of Medicine, Division of Hospital Internal Medicine, Mayo Clinic, Rochester, MN, USA

**Background:** Low-molecular-weight heparin (LMWH) is an effective means of preventing venous thromboembolism (VTE) among medical inpatients. Compared with unfractionated heparin, LMWH is equivalent or superior in efficacy and risk of bleeding. Despite its advantages, LMWH is underused in VTE prophylaxis for general-medicine patients hospitalized at our institution. Thus, a quality improvement (QI) initiative was undertaken to increase LMWH use for VTE prophylaxis among medical patients hospitalized on resident teaching services.

**Methods:** A QI team was formed, consisting of resident and attending physicians with pharmacy leaders. A systems analysis was performed, which showed gaps in resident knowledge as the greatest barrier to LMWH use. A knowledge translation framework was used to improve prescribing practices. Several Plan-Do-Study-Act cycles were executed, including resident-of-resident and pharmacist-of-resident education with performance audit and feedback.

**Results:** Pharmacist-of-resident education elicited the largest improvement and was sustained through a recurring pharmacist-led, interprofessional educational session as part of the monthly hospital orientation for incoming residents. Data analysis showed a statistically significant increase in LMWH use among treatment-eligible hospitalized medical patients, from 12.1% to 69.2%, following intervention ( $P < 0.001$ ). Extrapolated over 1 year, this improvement conserved 9,490 injections and nearly 791 hours of nurse time.

**Conclusions:** This QI project indicates that an interprofessional education intervention can lead to sustainable improvement in resident prescribing practices. This project also highlights the value of knowledge translation for the design of tailored interventions in QI initiatives.

**Keywords:** implementation science, interprofessional education, interprofessional teamwork, medical education, quality improvement

## Introduction

Venous thromboembolism (VTE) is a major cause of morbidity and death among hospitalized medical patients.<sup>1</sup> Current guidelines recommend pharmacologic prophylaxis for these patients at high risk of VTE.<sup>2,3</sup> Options include unfractionated heparin (UFH) and low-molecular-weight heparins (LMWHs). Compared with UFH, LMWH offers a similar efficacy with lower risks of both bleeding and heparin-induced thrombocytopenia.<sup>4,5</sup> In addition, LMWH is dosed once daily rather than the two or three times daily regimen for UFH, thereby increasing patient adherence and saving time for administration.<sup>6,7</sup> Thus, LMWH should be considered the high-value choice for pharmacologic VTE prophylaxis among therapy-eligible medical inpatients.<sup>8,9</sup>

Correspondence: John T Ratelle  
 Division of Hospital Internal Medicine,  
 Mayo Clinic, 200 First St SW, Rochester,  
 MN 55905, USA  
 Tel +1 507 293 3589  
 Email Ratelle.John@mayo.edu

Despite the benefits of LMWH, many US hospitals continue to use UFH for VTE prophylaxis.<sup>10</sup> Baseline data from January to April 2015 on inpatient general medicine teaching services at our institution have shown that <20% of patients eligible for VTE prophylaxis received LMWH. Further investigation detected a lack of awareness regarding the research evidence, benefits, and cost of LMWH as the primary reason behind prescriber choice of UFH at our institution.

Changing physician behavior is challenging, and many education interventions lead to only modest or inconsistent improvement in prescribing practices.<sup>11</sup> Knowledge translation (KT) is a relatively new term to describe the process of creating awareness and increasing the use of research evidence when making clinical decisions.<sup>12</sup> Lavis et al<sup>13</sup> offer the following five-step framework for KT, which may be useful for changing physician practice patterns:

1. What knowledge should be transferred (ie, what is the desired message or behavior)?
2. To whom should the knowledge be transferred (ie, who is the target audience)?
3. By whom should the knowledge be transferred (ie, who is the messenger)?
4. How should the knowledge be transferred (ie, what is the intervention or KT process)?
5. With what effect should the knowledge be transferred (ie, what is the desired outcome)?

Applying this KT framework to quality improvement (QI) initiatives offers an attractive approach to improving physician prescribing practices. Indeed, a recent Cochrane systematic review found that tailored interventions prospectively designed to overcome contextual barriers to implementation are an effective method to change physician performance.<sup>14</sup> QI initiatives have successfully increased overall rates of VTE prophylaxis in multiple settings.<sup>15,16</sup> However, we were unable to find a report that specifically aimed to increase the use of LMWH over UFH for medical inpatients.

Given the gap between existing research evidence and physician practice at our institution, a QI project was initiated to increase LMWH use for VTE prophylaxis among patients on wards that receive general medicine teaching services.

## Methods

This project is reported in accordance with the Standards for Quality Improvement Reporting Excellence 2.0 guidelines.<sup>17</sup> According to the policy activities that constitute research at Mayo Clinic, this work met criteria for operational

improvement activities and did not require ethics review by the Institutional Review Board.

## Context

This QI project involved collaboration of internal medicine residents and attending physicians with nurses and pharmacists from four general medicine teaching services at Mayo Clinic Hospital, Saint Mary's Campus. Each of these services is composed of three interns (first-year residents), one senior resident (third-year resident), and one attending physician. Residents rotate on the service in 4- or 5-week blocks, and attending physicians rotate in 2-week blocks. Clinical pharmacists and pharmacy residents are part of the team and attend rounds with the general medical services on weekdays.

In May 2015, a team was formed to increase LMWH use for VTE prophylaxis among patients who received care from the four resident services. A systems analysis showed that VTE prophylaxis was most frequently ordered by interns as part of a computerized admission order set for newly hospitalized patients. The order set identified LMWH as the preferred option for VTE prophylaxis (for patients without renal dysfunction), but most interns continued to order UFH. Further investigation observed a lack of awareness regarding the research evidence and benefits of LMWH as the primary reason for most interns who chose UFH. For this reason, resident education was the primary focus of subsequent interventions.

## Interventions

Using the Model for Improvement,<sup>18</sup> several Plan–Do–Study–Act (PDSA) cycles were performed between May and August 2015. Initial interventions consisted of resident-of-resident education in the form of didactics and posters placed in resident work areas. In addition, intern prescribing practices were audited, and resident-to-resident feedback was provided through email. Analysis of these interventions observed a small but measurable improvement in LMWH prescribing practices. Qualitative feedback from interns on the medical services indicated that some skepticism continued regarding the benefits of LMWH.

The focus of the next intervention was pharmacist-of-resident education. The decision to implement an interprofessional education intervention was based on qualitative feedback from residents during the systems analysis. Interns verbalized a strong sense of trust and respect toward the clinical expertise, judgment, and recommendations made by pharmacists on the medical unit.

At the start of each new resident rotation, a supervising clinical pharmacist (JGO) presented a 10-minute lecture about the QI project to residents, with specific data regarding safety, efficacy, costs, dosing, and periprocedural management of LMWH for VTE prophylaxis. Prescribing practices reflected a prompt and substantial improvement in LMWH use after the presentation. Subsequent PDSA cycles focused on sustainability by incorporating a standard pharmacist-led educational lecture into new-resident orientation on the first day of each clinical rotation. The benefits of this intervention appeared to reach a maximum effect by September 2015, at which time the project entered the sustainability phase, which consisted of a monthly, recurring pharmacist-led interprofessional education lecture during the new-resident rotation orientation.

From November 2015 to March 2016, the intervention was interrupted because of several competing pharmacy initiatives and relocation of the primary ward for the resident teaching services. In April 2016, the pharmacist-of-resident monthly didactic was restarted and was augmented by informal audit and feedback of interns' prescribing practices by clinical pharmacists during rounds.

## Study of the intervention and outcome measures

To measure this QI project's effect on LMWH use, data were obtained from the hospital pharmacy database regarding the use of heparin (LMWH or UFH) for patients hospitalized on the wards of general medicine teaching services between January 1, 2015, and June 30, 2016. Given stakeholder concerns about the potential increased risk of bleeding among patients receiving LMWH compared with UFH, we also evaluated the balancing measure of major bleeding among patients hospitalized during the project timeline. Our definition of major bleeding, which is based on existing literature,<sup>19</sup> is a decrease in hemoglobin of  $\geq 2.0$  g/dL in a 48-hour period.

## Analysis

For analysis purposes, LMWH use was quantified weekly as a proportion of all patients receiving thromboprophylaxis. To calculate this proportion, the sum of the number of patients receiving LMWH for the days of the given calendar week was the numerator, and the sum of the number of patients receiving LMWH or UFH for the days of the same week was the denominator. The process improvement efforts were implemented through multiple iterations of PDSA cycles undertaken from May through August 2015. Therefore, when evaluating the effectiveness of the overall improvement


effort, we divided the study period into three phases: January through April 2015 (preintervention), May through August 2015 (intervention), and September 2015 through June 2016 (postintervention). Patients with renal failure (defined as creatinine  $>2$  g/dL) were excluded from the analysis.

To account for potential underlying trends in LMWH use, we performed segmented regression.<sup>20</sup> The segmented regression model included terms to assess a linear trend in LMWH use before the initiation of the process improvement efforts, a step-change when the intervention was completed, and a potential change following intervention. As we were not interested specifically in modeling the changes in LMWH use during the intervention phase, the LMWH use data for this transition period were excluded rather than modeled as a separate segment.

The segmented regression analysis was performed with ordinary least-squares regression. The Durbin–Watson test was used to assess the assumption that the error terms associated with the weekly data points were uncorrelated. To supplement the segmented regression analysis, we compared demographic characteristics, heparin use, and bleeding complications among hospitalized patients who received LMWH vs UFH during the preintervention vs postintervention periods. In all cases, analyses were performed using SAS software (version 9.3; SAS Institute Inc).

## Results

LMWH use over the study period is summarized in Figure 1. The segmented regression analysis detected a significant increase ( $P<0.001$ ) in the use of LMWH following the intervention, with no evidence of a trend over time during the preintervention period ( $P=0.34$ ) or following intervention



**Figure 1** Weekly LMWH use before and after the knowledge translation intervention.

**Abbreviations:** LMWH, low-molecular-weight heparin; post, postintervention; pre, preintervention.

( $P=0.26$ ). The median (interquartile range) weekly LMWH use increased from 11% (5%–15%) during preintervention to 64% (58%–70%) during the postintervention phase.

Demographic characteristics of patients who received heparin were similar for preintervention and postintervention periods (Table 1). Among patients receiving some form of heparin during the preintervention, 90.5% received UFH on at least 1 day and 43.5% received UFH for 4 or more days. These percentages were significantly reduced ( $P<0.001$ ) in the postintervention period, with 34.5% of patients receiving UFH on at least 1 day and 14.5% on 4 or more days (Table 1). The percentage of patients who had major bleeding did not differ significantly between the periods (3.6% preintervention vs 3.4% postintervention,  $P=0.82$ ).

To assess the effect of the intervention on institutional costs and resource use, we performed a post hoc analysis comparing number of injections and nursing medication administration time pre and postintervention. Assuming an average census of 24 patients per day who require VTE prophylaxis injections, we estimate that our intervention,

sustained over 1 year, would directly save 9,490 injections. Assuming an average medication administration time of 5 minutes per UFH or LMWH injection, the direct saving of 9,490 injections would lead to an indirect saving of 790.8 hours of nurse time (equivalent to 0.38 full-time employment).

## Discussion

A brief interprofessional educational intervention correlated with a significant change in resident prescribing practices. A monthly pharmacist-led didactic for each new group of residents resulted in a sustained increase in LMWH use for VTE prophylaxis in treatment-eligible patients, with a corresponding decrease in UFH prescribing and no change in the rate of major bleeding. In particular, this study showed that the use of KT principles to design a relatively simple low-cost QI intervention can lead to sustained change in resident prescribing in a clinical setting where frequent turnover occurs among physician learners.

## Interpretation

Selecting the target audience is an essential step in the process of KT.<sup>12,13</sup> Health care providers often work in large teams with member rotation in the hospital, which may make this process challenging. Using stakeholder interviews, we were able to identify the important role that the intern has in the selection of VTE prophylaxis on our general medicine teaching services. This finding not only enabled us to identify the target audience for KT, but also showed the specific knowledge gaps and barriers to LMWH use among these providers. The educational intervention was then tailored to address these gaps and facilitate effective KT.

Observations from the initial PDSA cycles suggested that pharmacist-based education had a greater effect on residents' prescribing practices than peer-based education. Again, this determination fits into the framework of KT, which identifies the importance of the person from whom knowledge should be transferred. Although the content of the education was similar, informal feedback showed that residents tended to view information about the benefits of LMWH as more credible when it came from a content expert (ie, a pharmacist) rather than a peer. This finding is also consistent with existing research that has shown a positive correlation between teacher credibility and learner outcomes.<sup>21</sup>

Clinical pharmacists have an important role in the care of hospitalized patients in medical wards. Pharmacist-led interventions can substantially alter physician prescribing habits and improve patient outcomes.<sup>22,23</sup> However, the quality of the physician–pharmacist relationship appears to be a major

**Table 1** Patient demographic characteristics, heparin use, and bleeding complications

Characteristic <sup>a</sup>	Preintervention (n=497)	Postintervention (n=1,762)	P-value <sup>b</sup>
Age, mean (SD), years	63.3 (19.5)	63.0 (18.5)	0.73
<b>Sex</b>			0.40
Male	252 (50.7)	855 (48.5)	
Female	245 (49.3)	906 (51.4)	
<b>Any LMWH use</b>	60 (12.1)	1,220 (69.2)	<0.001
<b>LMWH use, days</b>			<0.001
0	437 (88.0)	542 (30.8)	
1	7 (1.4)	117 (6.6)	
2–3	26 (5.2)	639 (36.3)	
4–7	20 (4.0)	371 (21.1)	
≥8	7 (1.4)	93 (5.3)	
Median (IQR)	0 (0–0)	2 (0–4)	
<b>Any UFH use</b>	450 (90.5)	607 (34.4)	<0.001
<b>UFH use, days</b>			<0.001
0	47 (9.5)	1,155 (65.6)	
1	6 (1.2)	56 (3.2)	
2–3	228 (45.9)	296 (16.8)	
4–7	160 (32.2)	209 (11.9)	
≥8	56 (11.3)	46 (2.6)	
Median (IQR)	3 (2–5)	0 (0–2)	
<b>Bleeding<sup>c</sup></b>	18 (3.6)	60 (3.4)	0.82

**Notes:** <sup>a</sup>Values are presented as number and percentage of patients unless specified otherwise. <sup>b</sup>Categorical variables were compared between groups using chi-squared test. Age was compared between groups using two-sample t-test. Total days of LMWH use and UFH use were compared between groups using Wilcoxon rank sum test. <sup>c</sup>Defined as a decline in hemoglobin of ≥2.0 g/dL over 48 hours.

**Abbreviations:** IQR, interquartile range; LMWH, low-molecular-weight heparin; UFH, unfractionated heparin.



determinant to the adoption of pharmacist recommendations by physicians.<sup>24</sup> Interprofessional education has been proposed as a method to improve the working relationships between doctors and pharmacists.<sup>25</sup> The present project establishes the positive effect that pharmacist educators can have on the behavior of physician learners. Additionally, although we did not formally measure the effect of the intervention on interprofessional teamwork, informal feedback from project participants indicated improved collaboration between pharmacists and residents on the medicine services. Institutions aiming to improve physicians' prescribing practices should consider mechanisms to incorporate pharmacists into the education of medical trainees.

## Limitations

Although we have demonstrated that a QI project designed within the KT framework can succeed to improve inpatient LMWH use, our project has limitations. First, this was a single-institution study involving residents at a large academic center, which limits the generalizability to nonresident prescribers and other settings such as nonacademic hospitals. Yet, application of a similar KT approach that focuses on identifying the relevant prescribers, reasons for their prescribing habits, and whom they view as a credible source of information should be effective in other settings. Second, this was not a randomized study, which opens the possibility of confounding. However, we used segmented regression to attempt to identify any hospital-wide phenomenon that may have influenced LMWH use. Third, the intervention phase involved multiple PDSA cycles, making it difficult to know with certainty that the final pharmacist-of-resident education directly caused the observed changes in prescribing practice. However, when the sustainability phase of the intervention was temporarily interrupted because of relocation of the hospital unit, a corresponding decrease in LMWH prescriptions was observed. Use of LMWH subsequently increased when the intervention was reinstated, strongly suggesting that the educational intervention had a causal relationship with the increased LMWH use. This finding also emphasizes that continuity of the education intervention was important for achieving sustainable results. Finally, although research evidence suggests clinical benefits of LMWH over UFH, we did not measure the effect of the intervention on patient outcomes.

## Conclusion

This report demonstrates that a relatively low-effort, interprofessional education intervention can produce sustainable improvement in resident physicians' prescribing practices. This

initiative also highlights the value of KT framework application to QI methods. With accurate identification of knowledge gaps and misconceptions underlying current practice, similar QI endeavors may lead to educational interventions that apply across a wide range of clinical settings and specialties. Future work is needed to clarify the effect of pharmacist-of-resident education on pharmacist–doctor teamwork and collaboration.

## Acknowledgment

This study was supported in part by the Mayo Clinic Hospital Internal Medicine Small Grants Program. The funder had no involvement in study design, data acquisition, data analysis, manuscript preparation, or submission of this manuscript.

## Disclosure

The authors report no conflicts of interest in this work.

## References

1. Barbar S, Noventa F, Rossetto V, et al. A risk assessment model for the identification of hospitalized medical patients at risk for venous thromboembolism: the Padua Prediction Score. *J Thromb Haemost.* 2010;8(11):2450–2457.
2. Kahn SR, Lim W, Dunn AS, et al. Prevention of VTE in nonsurgical patients: Antithrombotic Therapy and Prevention of Thrombosis, 9th ed: American College of Chest Physicians Evidence-Based Clinical Practice Guidelines. *Chest.* 2012;141(2 Suppl):e195S–e226S.
3. Qaseem A, Chou R, Humphrey LL, et al. Venous thromboembolism prophylaxis in hospitalized patients: a clinical practice guideline from the American College of Physicians. *Ann Intern Med.* 2011;155(9):625–632.
4. Alikhan R, Cohen AT. Heparin for the prevention of venous thromboembolism in general medical patients (excluding stroke and myocardial infarction). *Cochrane Database Syst Rev.* 2009;3:CD003747.
5. Laporte S, Liotier J, Bertoletti L, et al. Individual patient data meta-analysis of enoxaparin vs. unfractionated heparin for venous thromboembolism prevention in medical patients. *J Thromb Haemost.* 2011;9(3):464–472.
6. Fanikos J, Stevens LA, Labreche M, et al. Adherence to pharmacological thromboprophylaxis orders in hospitalized patients. *Am J Med.* 2010;123(6):536–541.
7. Lloyd AC, Anderson PM, Quinlan DJ, Bearne A. Economic evaluation of the use of enoxaparin for thromboprophylaxis in acutely ill medical patients. *J Med Econ.* 2001;4(1-4):99–113.
8. Amin AN, Lin J, Lenhart G, Schulman KL. Clinical and economic outcomes in patients at risk of venous thromboembolism receiving appropriate enoxaparin or unfractionated heparin prophylaxis. *Thromb Haemost.* 2009;102(2):321–326.
9. Burleigh E, Wang C, Foster D, et al. Thromboprophylaxis in medically ill patients at risk for venous thromboembolism. *Am J Health Syst Pharm.* 2006;63(20 Suppl 6):S23–S29.
10. Tiryaki F, Nutescu EA, Hennenfent JA, et al. Anticoagulation therapy for hospitalized patients: patterns of use, compliance with national guidelines, and performance on quality measures. *Am J Health Syst Pharm.* 2011;68(13):1239–1244.
11. Cervero RM, Gaines JK. The impact of CME on physician performance and patient health outcomes: an updated synthesis of systematic reviews. *J Contin Educ Health Prof.* 2015;35(2):131–138.
12. Grimshaw JM, Eccles MP, Lavis JN, Hill SJ, Squires JE. Knowledge translation of research findings. *Implement Sci.* 2012;7:50.
13. Lavis JN, Robertson D, Woodside JM, et al. How can research organizations more effectively transfer research knowledge to decision makers? *Milbank Q.* 2003;81(2):221–248, 171–172.

14. Baker R, Camosso-Stefinovic J, Gillies C, et al. Tailored interventions to address determinants of practice. *Cochrane Database Syst Rev*. 2015;(4):CD005470.
15. Streiff MB, Lau BD, Hobson DB, et al. The Johns Hopkins Venous Thromboembolism Collaborative: multidisciplinary team approach to achieve perfect prophylaxis. *J Hosp Med*. 2016;11(Suppl 2):S8–S14.
16. Jenkins IH, White RH, Amin AN, et al. Reducing the incidence of hospital-associated venous thromboembolism within a network of academic hospitals: findings from five University of California medical centers. *J Hosp Med*. 2016;11(Suppl 2):S22–S28.
17. Ogrinc G, Davies L, Goodman D, et al. SQUIRE 2.0 (Standards for Quality Improvement Reporting Excellence): revised publication guidelines from a detailed consensus process. *Contin Educ Nurs*. 2015;46(11):501–507.
18. Langley GJ, Moen R, Nolan KM, et al. *The Improvement Guide: A Practical Approach to Enhancing Organizational Performance*. California: Jossey-Bass; 2009.
19. Schulman S, Kearon C; Subcommittee on Control of Anticoagulation of the Scientific and Standardization Committee of the International Society on Thrombosis and Haemostasis. Definition of major bleeding in clinical investigations of antihemostatic medicinal products in non-surgical patients. *J Thromb Haemost*. 2005;3(4):692–694.
20. Wagner AK, Soumerai SB, Zhang F, Ross-Degnan D. Segmented regression analysis of interrupted time series studies in medication use research. *J Clin Pharm Ther*. 2002;27(4):299–309.
21. Finn AN, Schrodt P, Witt PL, et al. A meta-analytical review of teacher credibility and its associations with teacher behaviors and student outcomes. *Commun Educ*. 2009;58(4):516–537.
22. McMullin ST, Hennenfent JA, Ritchie DJ, et al. A prospective, randomized trial to assess the cost impact of pharmacist-initiated interventions. *Arch Intern Med*. 1999;159(19):2306–2309.
23. Mekonnen AB, McLachlan AJ, BrienJ-Annee, Brien JA. Effectiveness of pharmacist-led medication reconciliation programmes on clinical outcomes at hospital transitions: a systematic review and meta-analysis. *BMJ Open*. 2016;6(2):e010003.
24. Béchet C, Pichon R, Giordan A, Bonnabry P, et al. Hospital pharmacists seen through the eyes of physicians: qualitative semi-structured interviews. *Int J Clin Pharm*. 2016;38(6):1483–1496.
25. Gallagher RM, Gallagher HC. Improving the working relationship between doctors and pharmacists: is inter-professional education the answer? *Adv Health Sci Educ Theory Pract*. 2012;17(2):247–257.

## Journal of Multidisciplinary Healthcare

### Publish your work in this journal

The Journal of Multidisciplinary Healthcare is an international, peer-reviewed open-access journal that aims to represent and publish research in healthcare areas delivered by practitioners of different disciplines. This includes studies and reviews conducted by multidisciplinary teams as well as research which evaluates the results or conduct of such teams or health

care processes in general. The journal covers a very wide range of areas and welcomes submissions from practitioners at all levels, from all over the world. The manuscript management system is completely online and includes a very quick and fair peer-review system. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/journal-of-multidisciplinary-healthcare-journal>

Dovepress