Downregulation of miR-638 promotes progression of breast cancer and is associated with prognosis of breast cancer patients

Minghong Li*

Jian Wang*

Haibo Liu

Department of Laboratory Medicine,
Yidu Central Hospital of Weifang,
Shandong 262500, People's Republic
of China

*These authors contributed equally
to this work

Background: Breast cancer is the most common tumor among women. miR-638 has been demonstrated to play an important role in various cancers.

Purpose: In this study, we aimed to investigate the function and prognostic value of miR-638 in breast cancer.

Methods: Quantitative real-time polymerase chain reaction analysis was used to evaluate the expression of miR-638 in breast cancer tissues and cell lines. The correlation of miR-638 with clinicopathological features was analyzed using the chi-squared test. Kaplan–Meier survival analysis and Cox regression assay were performed to investigate the prognostic value of miR-638 in breast cancer patients. The effects of miR-638 on the biological behavior of breast cancer cells were evaluated using functional assays.

Results: The expression of miR-638 was downregulated in breast cancer tissues and cell lines (all $P < 0.05$). Decreased expression of miR-638 was significantly correlated with lymph node metastasis ($P = 0.015$) and TNM stage ($P = 0.021$). Patients with low miR-638 expression had shorter overall survival compared with those with high levels (Log-rank $P = 0.025$). The miR-638 could be considered as an independent prognostic factor for the patients (HR = 0.321, 95% CI = 0.117–0.882, $P = 0.027$). Downregulation of miR-638 was capable of promoting cell proliferation, migration, and invasion in MDA-MB-231 and MCF-7 cells.

Conclusion: All the results indicate that miR-638 is a tumor suppressor in breast cancer and is involved in the progression of breast cancer. Thus, it may serve as a prognostic biomarker for breast cancer.

Keywords: miR-638, cell proliferation, migration, invasion, prognosis, breast cancer

Introduction

Breast cancer is a common malignancy and a leading cause of cancer-related mortality in women.¹ It is a heterogeneous disease that can be categorized into three major subtypes, of which triple-negative breast cancer (TNBC), accounting for 10%–20% of all breast cancer cases, has the worst prognosis because of high recurrence rates and early metastasis.^{2–4} Although advances in both detection and therapy have led to the improvement of prognosis, recurrence, and metastasis remain to be major challenges in breast cancer management that are related to the poor prognosis.^{5,6} Current biological markers used in the management of breast cancer are mainly estrogen receptor (ER), progesterone receptor (PR), and HER2.^{7,8} Rising evidence demonstrated that the prognosis could be improved by using some molecular biomarkers. Therefore, identification of novel precise biomarkers is important and needed for the development of prognosis in breast cancer patients.

Correspondence: Jian Wang
Department of Laboratory Medicine,
Yidu Central Hospital of Weifang,
No 4138, South Linglongshan Road,
Shandong 262500, People's Republic
of China
Tel +86 536 327 5619
Fax +86 536 327 5619
Email zou8402pidao@163.com

MicroRNAs (miRNAs) are small endogenous noncoding RNAs, which function in the transcriptional and posttranscriptional regulation of gene expression.^{9–12} Evidence has shown that miRNAs play a crucial role in different biological processes, such as inflammation, cell proliferation, migration, invasion, and differentiation.^{13–16} Furthermore, a large number of miRNAs have been identified to function as oncogenes or tumor suppressor genes in the tumorigenesis process.^{17–19} Recently, dysregulation of miR-638 has been confirmed in several types of cancer, such as gastric cancer, basal cell carcinoma, and non-small-cell lung cancer (NSCLC).^{20–22} A recent study by Tan et al²³ showed the expression of miR-638 was decreased in most non-TNBC and played an important role in TNBC progression via BRCA1 deregulation. However, the role of miR-638 in breast cancer is still limited.

In this study, we investigated the expression levels of miR-638 in breast cancer tissues and cell lines. The effects of miR-638 on biological behaviors of cancer cells were also assessed. This study aimed to identify the prognostic significance of miR-638 in breast cancer.

Methods and materials

Patients and tissue samples

Breast cancer tissues and adjacent normal tissues (located >3 cm away from the tumors) were collected from 125 breast cancer patients between January 2007 and December 2012 in Yidu Central Hospital of Weifang (Weifang, People's Republic of China). None of the patients had received radiotherapy or chemotherapy or any other treatment before surgery. The tissues were verified by experienced pathologists. All the surgical specimens of the tumor resection were snap-frozen in liquid nitrogen after surgery until use. Written informed consent was obtained from all patients that agreed to participate in our study. This study was approved by the Ethics Committee of Yidu Central Hospital of Weifang (Weifang, People's Republic of China). Detailed clinical characteristics of the patients are summarized in Table 1. A 5-year follow-up information was collected for the subsequent analysis.

Cell lines and transfection

The breast cancer cell lines (MDA-MB-231, MCF-7) and normal breast cell line MCF10A were obtained from the American Type Culture Collection (Manassas, VA, USA). These breast cell lines were maintained in Roswell Park Memorial Institute 1640 (RPMI 1640) with 10% fetal bovine serum and 1% antibiotics (Invitrogen, Carlsbad, CA, USA). MCF-10A cells were cultured in mammary epithelial cell growth medium (MEGM) (CC-3150, Lonza, Basel, Switzerland) containing 100 ng/mL of cholera toxin to make

Table 1 Relationship between miR-638 expression and clinical characteristics of breast cancer patients

Characteristics	Total number (N=125)	miR-638 expression		P-values
		High (n=55)	Low (n=70)	
Age (years)				0.363
<50	58	23	35	
≥50	67	32	35	
Tumor size (cm)				0.315
<2	54	21	33	
≥2	71	34	37	
ER status				0.676
Negative	61	28	33	
Positive	64	27	37	
PR status				0.245
Negative	55	21	34	
Positive	70	34	36	
HER2 status				0.696
Negative	57	24	33	
Positive	68	31	37	
Lymph node metastasis				0.015
Negative	53	30	23	
Positive	72	25	47	
TNM stage				0.021
I–II	65	35	30	
III–IV	60	20	40	

Abbreviations: ER, estrogen receptor; PR, progesterone receptor.

a complete growth culture medium. All cells were cultured in a humidified incubator at 37°C with 5% CO₂. A total of 1×10⁵ cells were seeded into a 6-well plate. Then, the cells were transfected with the miR-638 mimic or mimic negative control (mimic NC), miR-638 inhibitor or negative inhibitor control (inhibitor NC) using the Lipofectamine 2000 reagent (Invitrogen) according to the manufacturer's instructions. Cells transfected with only transfection reagent were used as mock. The transfection efficiency was detected by quantitative real-time PCR (qRT-PCR) analysis.

RNA extraction and qRT-PCR analysis

Total RNA was isolated from tissues or cell lines using TRIzol reagent (Invitrogen) in accordance with the manufacturer's instructions. Subsequently, purified total RNA was reverse transcribed to cDNA by using M-MLV reverse transcriptase (Promega, Madison, WI, USA) following the manufacturer's instructions. The qRT-PCR experiment was performed using SYBR Green (Takara, Kusatsu, Japan) on an ABI 7500 system (Applied Biosystems, Foster City, CA, USA). The relative miR-638 levels were detected from the threshold cycle with the 2^{−ΔΔCt} method using the U6 snRNA as an endogenous control. All the experiments were performed in triplicate.

MTT assay

The colorimetric MTT analysis was performed to examine the effects of miR-638 on cell proliferation of breast

cancer cells. In brief, the cells were grown in 96-well plates and transfected. After transfection and incubation for 5 days, to each well was added 10 μ L MTT (5 mg/mL, Sigma-Aldrich, St Louis, MO, USA) and incubated at 37°C. After 4 hours incubation, the supernatant was removed and 150 μ L DMSO (Sigma-Aldrich) was added to each well to dissolve formazan crystals. The absorbance was measured at 490 nm using a spectrophotometer (Multiskan MK3, Thermo Fisher Scientific, Waltham, MA, USA). The experiment was repeated at least three times.

Transwell migration and invasion assay

To measure the effects of miR-638 on breast cancer cell migration and invasion ability, a transwell kit (Corning, Tewksbury, MA, USA, 8.0 μ m pores) was used with a 24-well transwell chamber according to the manufacturer's instruction. A total of 5×10^4 cells transfected with miR-638 mimic or NC, miR-638 inhibitor or NC, or mock were placed in the upper chamber in RPMI 1640 with no serum. The bottom chamber contained RPMI 1640 with 10% fetal bovine serum, using as the chemotactic factor. For the invasion assay, the upper chamber was first coated with Matrigel (BD Company, Bedford, MA, USA). After incubating for 24 hours, the cells that had migrated or invaded to the bottom chamber were fixed in 3.7% formaldehyde for 5 minutes and stained with 0.1% crystal violet for 15 minutes. The number of cells on the bottom of the chamber was counted under a microscope (Olympus, Tokyo, Japan). The process of the transwell migration assay was similar to the transwell invasion assay except for the Matrigel-coated membrane.

Statistical analysis

Statistical analysis of data was performed using the SPSS 21.0 software (IBM Corporation, Armonk, NY, USA), and

the data were expressed as the mean $\pm$ SD at least three times. Student's *t*-test was performed to analyze the significance of differences between the two groups. The association between miR-638 expression and clinicopathological parameters of patients was analyzed by chi-squared test. Kaplan–Meier method was used to analyze the association between overall survival and miR-638 expression in breast cancer patients according to the log-rank test. Cox regression analysis was used to assess the prognostic factors. Differences were considered statistically significant when $P < 0.05$.

Results

Relative expression of miR-638 in breast cancer tissues and cell lines

We used qRT-PCR analysis to measure the miR-638 expression levels in breast cancer tissues. As shown in Figure 1A, the expression level of miR-638 was significantly lower in breast cancer tissues than in normal breast tissues ($P < 0.001$).

The expression of miR-638 was also evaluated in two breast cancer cell lines, MDA-MB-231 and MCF-7, compared to a normal breast cell line MCF10A. The result showed that the expression level of miR-638 was decreased significantly in breast cancer cell lines than in normal cell line (all $P < 0.01$, Figure 1B).

Correlation of miR-638 expression and clinical characteristics

According to the mean value of miR-638 expression levels in breast cancer tissues, which was used as the cutoff value, we classified the patients into low miR-638 expression group ($n=70$) and high expression group ($n=55$). As shown in Table 1, the expression of miR-638 was significantly associated with lymph node metastasis ($P=0.015$) and TNM stage

Figure 1 Expression of miR-638 in breast cancer tissues and cell lines was measured by qRT-PCR.

Notes: (A) The expression levels of miR-638 were lower in breast cancer tissues than in normal tissues. (B) The level of miR-638 in breast cancer cell lines (MDA-MB-231 and MCF-7) was downregulated compared with normal breast cell line MCF10A. ** $P < 0.01$, *** $P < 0.001$.

Abbreviation: qRT-PCR, quantitative real-time polymerase chain reaction.

Figure 2 The prognosis of miR-638 in breast cancer.

Note: Patients with low miR-638 expression showed shorter overall survival than those with high expression levels.

($P=0.021$). There was no dramatical association between miR-638 expression and other clinical parameters, such as age, tumor size, ER status, PR status, and HER2 status (all $P>0.05$, Table 1).

Prognostic significance of miR-638 in breast cancer

We used the Kaplan–Meier analysis to evaluate the association of miR-638 expression with overall survival of breast cancer patients. As shown in Figure 2, the breast cancer patients with lower miR-638 expression in tumor tissues suffered from unfavorable overall survival compared with those with higher miR-638 expression (Log-rank $P=0.025$,

Figure 2). Furthermore, multivariate Cox regression analysis results indicated that miR-638 expression (HR =0.321, 95% CI =0.117–0.882, $P=0.027$, Table 2) was an independent prognostic factor for breast cancer patients.

Overexpression of miR-638 can suppress breast cancer cell proliferation, migration, and invasion

To investigate the function of miR-638 in breast cancer cell lines, two breast cancer cell lines, MDA-MB-231 and MCF-7, were transfected with miR-638 mimic or mimic NC, miR-638 inhibitor or inhibitor NC. The qRT-PCR analysis results indicated that miR-638 mimic could increase the expression of miR-638 in MDA-MB-231 and MCF-7 cells, and miR-638 inhibitor can decrease the expression of miR-638 in MDA-MB-231 and MCF-7 cells, respectively ($P<0.05$, Figure 3A and B). MTT assays were then performed to investigate the effects of miR-638 on cell growth in vitro. We observed that the proliferation of cells was inhibited in miR-638 mimic-transfected cells and promoted in miR-638 inhibitor-transfected cells ($P<0.05$, Figure 3C and D).

To further investigate whether miR-638 affects cell metastasis, transwell assay was used in this study. The transwell migration assay results showed that cell migration was inhibited in cells transfected with miR-638 mimic, and promoted in the cells transfected with miR-638 inhibitor ($P<0.05$, Figure 4A and B). The transwell invasion assay with Matrigel results demonstrated that miR-638 mimic could inhibit the cell invasion, and miR-638 inhibitor could promote the cell invasion ($P<0.05$, Figure 4C and D).

Discussion

The incidence of breast cancer has increased rapidly with economic development in People's Republic of China in recent years.²⁴ Breast cancer patients who are diagnosed and received treatment at an early stage usually have longer overall survival, while those at advanced stage usually relapse or recur,

Table 2 Univariate and multivariate Cox analyses for miR-638 in breast cancer patients

Characteristics	Univariate analyses			Multivariate analysis		
	HR	95% CI	P-value	HR	95% CI	P-value
miR-638	0.325	0.119–0.886	0.028	0.325	0.119–0.886	0.028
Age	1.828	0.857–3.900	0.119	–	–	–
Tumor size	1.067	0.464–2.454	0.879	–	–	–
ER status	1.221	0.547–2.728	0.626	–	–	–
PR status	0.798	0.323–1.970	0.625	–	–	–
HER-2 status	1.634	0.743–3.592	0.222	–	–	–
Lymph node metastasis	0.799	0.337–1.894	0.611	–	–	–
TNM stage	1.521	0.721–3.210	0.271	–	–	–

Note: –, indicated no related data.

Abbreviations: ER, estrogen receptor; HR, hazard ratio; PR, progesterone receptor.

Figure 3 Expression of miR-638 expression on proliferation of breast cancer cells.

Notes: (A, B) qRT-PCR was used to detect the transfection efficiency of miR-638 mimic or inhibitor in MDA-MB-231 and MCF-7 cells. (C, D) MTT assay was used to determine the cell proliferation. * $P < 0.05$, *** $P < 0.001$.

Abbreviation: qRT-PCR, quantitative real-time polymerase chain reaction.

leading to poor prognosis.²⁵ Nowadays, the incidence of breast cancer in young women is more aggressive and invasive than that in older patients, leading to an adverse prognosis.^{26,27} Despite the advances of both detection and treatment, the rates of breast cancer remain high in most countries. Therefore, more accurate cancer-related biomarkers are needed to help predict patient prognosis and treatment strategies.

An increasing number of molecular markers, including miRNAs, have been identified play crucial roles in tumor pathogenesis and development, acting as biomarkers for diagnosis, prognosis, or treatment of cancers.^{28,29} Increasing miRNAs have been reported as new biological markers in cancers. For instance, Shidfar et al³⁰ found that miR-18a and miR-210 were significantly highly expressed in breast cancer, and could be used as a candidate for breast cancer risk biomarkers. Marques et al³¹ evaluated miR-195 and let-7a expression in invasive breast cancer and showed miR-195 and let-7a could be used as a noninvasive biomarker for breast cancer detection. The study by He et al³² indicated that

miR-638 might act as a sensitizer in cancer chemotherapy and accompany chemotherapy drugs to enhance chemotherapeutic efficacy. To better understand the role of miR-638 in breast cancer, its expression patterns, and prognostic significance, this study was carried out in breast cancer patients.

The study by Zavala et al³³ investigated the expression of miR-638 in TNBC and found that miR-638 was overexpressed in most TNBC tumors, consistent with the result in the TNBC samples of Tan et al.²³ In our study, we mainly measured the expression of miR-638 in the breast cancer tissues and cell lines. Similar to the non-TNBC result of Tan et al,²³ expression of miR-638 was significantly downregulated in breast cancer tissues, as well as in cell lines, compared to normal control. Moreover, the decreased expression of miR-638 was correlated with lymph node metastasis and TNM stage. Taken together, we considered that miR-638 might be a suppressor gene in breast cancer and be involved in the progression of this malignancy. Furthermore, we investigated the clinical significance of miR-638 in breast

Figure 4 Expression of miR-638 expression on migration and invasion of breast cancer cells.

Notes: (A, B) Transwell migration assay were performed in breast cancer cell lines (MDA-MB-231 and MCF-7) transfected with miR-638 mimic or mimic NC, inhibitor or inhibitor NC. (C, D) Transwell invasion assay were performed in breast cancer cell lines (MDA-MB-231 and MCF-7). * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$.

Abbreviation: NC, negative control.

cancer prognosis using the Kaplan–Meier analysis and Cox regression analysis. The Kaplan–Meier result with the data from follow-up showed that patients with low expression of miR-638 had a shorter overall survival rate than those with high miR-638 expression, which suggested that miR-638 expression was related to the prognostic value for patients with breast cancer. Multivariate Cox regression analysis results showed that miR-638 was an independent biomarker for indicating the overall survival of breast cancer patients.

To investigate the function of miR-638 in breast cancer progression, the effects of miR-638 on biological behaviors of breast cancer were also analyzed. The analysis results demonstrated that cell proliferation, migration, and invasion were suppressed by the overexpression of miR-638, which showed the potential functional role of miR-638 in breast cancer progression. Many of the studies have indicated the effects of miR-638 on biological behaviors during cancer progression.^{22,34} For instance, the study by Xia et al²² showed

that downregulated miR-638 could promote cell invasion and proliferation in NSCLC, indicating that miR-638 may play a pivotal role in the development of NSCLC. Zhang et al³⁴ revealed that miR-638 inhibited colorectal carcinoma cell growth, invasion, and functioned as a tumor suppressor in human colorectal cancer by inhibiting TSPAN1. The study by Zhao et al³⁵ found that miR-638 was downregulated in gastric cancer, and molecular mechanistic investigation results demonstrated that MeCP2, a target of miR-638, facilitated gastric cancer cell proliferation through activation of the MEK1/2-ERK1/2 signaling pathway by upregulating GIT1. Besides the cell experiments, the precise molecular mechanistic investigations underlying role of miR-638 in breast cancer need to be assessed in the further studies. Also, due to the limitation of sample size in the present study, further studies are necessary with a larger research cohort.

In conclusion, we identified that expression of miR-638 was downregulated in breast cancer tissue and was correlated

with the progression of breast cancer. This study results showed that miR-638 might be an independent prognostic biomarker for breast cancer.

Disclosure

The authors report no conflicts of interest in this work.

References

- Siegel RL, Miller KD, Jemal A. Cancer statistics, 2016. *CA Cancer J Clin*. 2016;66(1):7–30.
- di Cosimo S, Baselga J. Management of breast cancer with targeted agents: importance of heterogeneity. [corrected]. *Nat Rev Clin Oncol*. 2010;7(3):139–147.
- Metzger-Filho O, Tutt A, de Azambuja E, et al. Dissecting the heterogeneity of triple-negative breast cancer. *J Clin Oncol*. 2012;30(15):1879–1887.
- Yang Q, Liu HY, Liu D, Song YQ. Ultrasonographic features of triple-negative breast cancer: a comparison with other breast cancer subtypes. *Asian Pac J Cancer Prev*. 2015;16(8):3229–3232.
- Chaffer CL, Weinberg RA. A perspective on cancer cell metastasis. *Science*. 2011;331(6024):1559–1564.
- Scully OJ, Bay BH, Yip G, Yu Y. Breast cancer metastasis. *Cancer Genomics Proteomics*. 2012;9(5):311–320.
- Sorlie T, Perou CM, Tibshirani R, et al. Gene expression patterns of breast carcinomas distinguish tumor subclasses with clinical implications. *Proc Natl Acad Sci U S A*. 2001;98(19):10869–10874.
- Liu RZ, Garcia E, Glubrecht DD, Poon HY, Mackey JR, Godbout R. CRABP1 is associated with a poor prognosis in breast cancer: adding to the complexity of breast cancer cell response to retinoic acid. *Mol Cancer*. 2015;14:129.
- Bartel DP. MicroRNAs: genomics, biogenesis, mechanism, and function. *Cell*. 2004;116(2):281–297.
- Chen K, Rajewsky N. The evolution of gene regulation by transcription factors and microRNAs. *Nat Rev Genet*. 2007;8(2):93–103.
- Li P, Dong J, Zhou X, et al. Expression patterns of microRNA-329 and its clinical performance in diagnosis and prognosis of breast cancer. *Onco Targets Ther*. 2017;10:5711–5718.
- Huang Y, Jian W, Zhao J, Wang G. Overexpression of *HDAC9* is associated with poor prognosis and tumor progression of breast cancer in Chinese females. *Onco Targets Ther*. 2018;11:2177–2184.
- Singh RP, Massachi I, Manickavel S, et al. The role of miRNA in inflammation and autoimmunity. *Autoimmun Rev*. 2013;12(12):1160–1165.
- Chen L, Song J, Cui J, et al. microRNAs regulate adipocyte differentiation. *Cell Biol Int*. 2013;37(6):533–546.
- Lu GJ, Dong YQ, Zhang QM, et al. miRNA-221 promotes proliferation, migration and invasion by targeting TIMP2 in renal cell carcinoma. *Int J Clin Exp Pathol*. 2015;8(5):5224–5229.
- Viñas JL, Ventayol M, Brüne B, et al. miRNA let-7e modulates the Wnt pathway and early nephrogenic markers in mouse embryonic stem cell differentiation. *PLoS ONE*. 2013;8(4):e60937.
- Farazi TA, Hoell JI, Morozov P, Tuschl T. MicroRNAs in human cancer. *Adv Exp Med Biol*. 2013;774:1–20.
- Fang YX, Gao WQ. Roles of microRNAs during prostatic tumorigenesis and tumor progression. *Oncogene*. 2014;33(2):135–147.
- Jiang C, Chen X, Alattar M, Wei J, Liu H. MicroRNAs in tumorigenesis, metastasis, diagnosis and prognosis of gastric cancer. *Cancer Gene Ther*. 2015;22(6):291–301.
- Yao Y, Suo AL, Li ZF, et al. MicroRNA profiling of human gastric cancer. *Mol Med Rep*. 2009;2(6):963–970.
- Sand M, Skrygan M, Sand D, et al. Expression of microRNAs in basal cell carcinoma. *Br J Dermatol*. 2012;167(4):847–855.
- Xia Y, Wu Y, Liu B, Wang P, Chen Y. Downregulation of miR-638 promotes invasion and proliferation by regulating SOX2 and induces EMT in NSCLC. *FEBS Lett*. 2014;588(14):2238–2245.
- Tan X, Peng J, Fu Y, et al. miR-638 mediated regulation of BRCA1 affects DNA repair and sensitivity to UV and cisplatin in triple-negative breast cancer. *Breast Cancer Res*. 2014;16(5):435.
- Hong W, Dong E. The past, present and future of breast cancer research in China. *Cancer Lett*. 2014;351(1):1–5.
- Naumov GN, Townson JL, Macdonald IC, et al. Ineffectiveness of doxorubicin treatment on solitary dormant mammary carcinoma cells or late-developing metastases. *Breast Cancer Res Treat*. 2003;82(3):199–206.
- Darwish AD, Helal AM, Aly El-Din NH, Solaiman LL, Amin A. Breast cancer in women aging 35 years old and younger: the Egyptian National Cancer Institute (NCI) experience. *Breast*. 2017;31:1–8.
- Slaoui M, Mouh FZ, Ghanname I, Razine R, El Mzibri M, Amrani M. Outcome of breast ZC cancer in moroccan young women correlated to clinic-pathological features, risk factors and treatment: a comparative study of 716 cases in a single institution. *PLoS ONE*. 2016;11(10):e0164841.
- Jagarlamudi KK, Shaw M. Thymidine kinase 1 as a tumor biomarker: technical advances offer new potential to an old biomarker. *Biomark Med*. 2018;12(9):1035–1048.
- Givechian KB, Garner C, Garban H, Rabizadeh S, Soon-Shiong P. CAD/POLD2 gene expression is associated with poor overall survival and chemoresistance in bladder urothelial carcinoma. *Oncotarget*. 2018;9(51):29743–29752.
- Shidfar A, Costa FF, Scholtens D, et al. Expression of miR-18a and miR-210 in normal breast tissue as candidate biomarkers of breast cancer risk. *Cancer Prev Res*. 2017;10(1):89–97.
- Marques MM, Evangelista AF, Macedo T, et al. Expression of tumor suppressors miR-195 and let-7a as potential biomarkers of invasive breast cancer. *Clinics*. 2018;73:e184.
- He M, Lin Y, Tang Y, et al. miR-638 suppresses DNA damage repair by targeting SMC1A expression in terminally differentiated cells. *Aging*. 2016;8(7):1442–1456.
- Zavala V, Pérez-Moreno E, Tapia T, Camus M, Carvallo P. miR-146a and miR-638 in BRCA1-deficient triple negative breast cancer tumors, as potential biomarkers for improved overall survival. *Cancer Biomark*. 2016;16(1):99–107.
- Zhang J, Fei B, Wang Q, et al. MicroRNA-638 inhibits cell proliferation, invasion and regulates cell cycle by targeting tetraspanin 1 in human colorectal carcinoma. *Oncotarget*. 2014;5(23):12083–12096.
- Zhao LY, Tong DD, Xue M, et al. MeCP2, a target of miR-638, facilitates gastric cancer cell proliferation through activation of the MEK1/2-ERK1/2 signaling pathway by upregulating GIT1. *Oncogenesis*. 2017;6(7):e368.

OncoTargets and Therapy

Publish your work in this journal

OncoTargets and Therapy is an international, peer-reviewed, open access journal focusing on the pathological basis of all cancers, potential targets for therapy and treatment protocols employed to improve the management of cancer patients. The journal also focuses on the impact of management programs and new therapeutic agents and protocols on

Submit your manuscript here: <http://www.dovepress.com/oncotargets-and-therapy-journal>

patient perspectives such as quality of life, adherence and satisfaction. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Dovepress