The threat of carbapenem-resistant gram-negative bacteria in a Middle East region

Effat Davoudi-Monfared
Hossein Khalili

Department of Clinical Pharmacy,
Faculty of Pharmacy, Tehran University
of Medical Sciences, Tehran, Iran

→ Video abstract

Point your Smartphone at the code above. If you have a
QR code reader the video abstract will appear. Or use:
<http://youtu.be/MnaI9PzhAk>

Abstract: Data on the status of carbapenem-resistant microorganisms in the Middle East countries are scarce. The aim of this review was to collect available data regarding resistance to carbapenems in a Middle East region. Available data regarding carbapenem-resistant isolates were considered for evaluation in this review. Biomedical electronic databases were systematically searched to find related articles. The key terms used were “carbapenem-resistant, resistant gram-negative bacilli, *Enterobacteriaceae*, fermenting and non-fermenting gram-negative bacilli, *Pseudomonas*, *Acinetobacter*, *Klebsiella* and Iran”. After primary screening, 275 relevant articles were selected to be assessed thoroughly. Resistance rate to carbapenems was reported between 1% and 86% during years 2006–2018. Most of the carbapenem-resistant microorganisms were isolated from burn patients. Modified Hodge test was a commonly used phenotypic test. Only in few studies, genotypic assays were considered. Pattern of antibiotic use can affect emergence of resistant microorganisms. Rational use of drugs, and specifically, antibiotics is a challenging issue in developing countries. Mean number of drugs per prescription in these countries was higher than the World Health Organization standards. Overuse of antibiotics, especially injectable ones, and easy access to antibiotics without prescription is a warning alarm for future antibiotic resistance in developing countries. Establishing antimicrobial stewardship’s programs is new in the hospitals. Unfortunately, rules and regulatory issues to restrict antibiotic access in community pharmacies and prescription by general physicians are limited.

Keywords: carbapenem, resistant, gram-negative bacteria, antibiotics

Introduction

Carbapenem-resistant gram-negative bacteria are now a global concern around the world. Most of these strains are also resistant to other antimicrobial agents including aminoglycosides and fluoroquinolones.¹ Few therapeutic options without the desired efficacy are available to treat infections caused by carbapenem-resistant microorganisms.² This real threat necessitates applying reliable methods to determine prevalence of these isolates.

The methods to detect carbapenemase-producing microorganisms are divided into two classes: phenotypic and genotypic methods. Phenotypic methods are nonmolecular assays that detect structure of the carbapenemase enzyme (mostly active site) through a chemical or microbiological process.³ The carbapenemase enzymes are structurally classified according to their active sites. Ambler class A (known as *Klebsiella*-producing carbapenemase [KPC]) has serine amino acid in its active site and can be inhibited by β -lactamase inhibitors. Ambler class B has zinc in its active site and is called metallo- β -lactamase (MBL), so it can react with EDTA and dipicolinic acid (DPA). The last

Correspondence: Hossein Khalili
Department of Clinical Pharmacy, Faculty
of Pharmacy, P.O. Box 14155/6451,
Tehran University of Medical Sciences,
Tehran 1417614411, Iran
Tel/Fax +98 216 695 4709
Email Khaliliah@sina.tums.ac.ir

class, Ambler class D, has serine in its active site, but it may or may not be inhibited by β -lactamase inhibitors.⁴ According to these classes, various phenotypic methods are proposed to detect carbapenemase enzyme in gram-negative isolates.

Genotypic methods are targeted to detect resistance genes (mostly *bla* type genes) and polymerase chain reaction (PCR) is a common method. Other genotypic method is clonal typing.⁵ Although detection of resistant-encoding genes is more accurate, however, these methods are usually expensive and not available anywhere.⁶ Some other mechanisms of resistance to carbapenems including loss of porins, efflux pump mutation, and target site inactivation have been identified.⁷ Figure 1 shows different genotypic and phenotypic methods for detection of resistance to carbapenems.

Data on the status of carbapenem-resistant microorganisms in the Middle East countries are scarce. The aims of this review were to collect available data regarding methods of detection and prevalence of resistance to carbapenems in a Middle East region.

Methods

Databases

Biomedical electronic databases (Scopus, Medline, Google Scholar, and Science Direct) were systematically searched to find related articles. The key terms used were “carbapenem-resistant, resistant gram-negative bacilli, *Enterobacteriaceae*,

fermenting and non-fermenting gram-negative bacilli, *Pseudomonas*, *Acinetobacter*, *Klebsiella* and Iran”. A total of 586 articles were found. After primary screening, 275 relevant articles were selected to be assessed thoroughly.

Study settings

Of the 275 articles, case reports, systematic reviews, meta-analysis, and articles in Persian language were excluded, and finally 97 English-language articles were included in the present review. These studies were published between 2006 and 2018 and were reported from different regions of Iran. Forty-eight studies were from Tehran and most of them were done at Motahari hospital (Level I Iranian burn hospital). The number of publications from other areas were as follows: ten in Isfahan, eight in northwest of Iran, five in south of Iran, four in Shiraz, three each in Hamadan, Arak and Mazandaran, two each in Shahrekord, Kermanshah, Mashhad, and Zanjan, and one each in Kerman, Khorramabad, and Qom. All the studies considered the standards and guidelines of the clinical and laboratory standards institute (CLSI) for the preparation and interpretation of the tests. Figure 2 illustrates data processing in this review.

Clinical and microbiological setting

All the studies included inpatients, with eight also considering outpatients. If the source of the samples was specific, they

Figure 1 Screening methods for carbapenem-resistant microorganisms.

Notes: ¹MHT was the most used phenotypic method in Iranian studies. MHT originally detects KPC, but if zinc sulfate is added to MHT culture media, it can detect MBL. ²It is based on carbapenemase inhibition by betalactamase inhibitors. CDT consists of two disks: a carbapenem and the other combination of a carbapenem and a betalactamase inhibitor. If the disk with inhibitor shows a bigger inhibition zone, the result is considered positive. DDST consists of carbapenem discs at a variable distance to inhibitor discs. The observation of synergy between disks is noted as a positive result. ³These phenotypic methods can detect different Ambler classes. ⁴CIM is a newer phenotypic method that was introduced in 2015 by van der Zwaluw. A carbapenem disk is inserted in culture media of suspected strain. Then it is transferred to another culture media with known control strain. If the suspected strain contains carbapenemase enzyme, the carbapenem disk has been degraded and the control strain in second culture will grow. ⁵Some other phenotypic methods like spectrometric assays are also used and have higher sensitivity and specificity, but they are costly and time consuming.

Abbreviations: CDT, combination disk test; CIM, carbapenem inactivation method; DDST, double disk synergy test; ESBL, extended spectrum β -lactamase; KPC, *Klebsiella*-producing carbapenemase; MBL, metallo- β -lactamase; MHT, modified Hodge test; MIC, minimum inhibitory concentration; PCR, polymerase chain reaction.

Figure 2 Consort flowchart of study.

Abbreviation: ESBL, extended spectrum β -lactamase.

are mentioned and if they were collected from different sites, they are named as “different sites”.

All studies have been categorized into four distinct groups:

1. Studies that reported carbapenem-resistant microorganisms according to the disk diffusion method
2. Studies that considered phenotypic methods
3. Studies that used genotypic methods
4. Studies that applied both genotypic and phenotypic assays

Results

Studies that reported carbapenem-resistant microorganisms according to the disk diffusion method (early detection of resistance to a carbapenem)

In this category, detection of resistance to a carbapenem was described only by disk diffusion method. However, in some of them, minimum inhibitory concentration (MIC) values were determined. No phenotypic or genotypic assay was considered for carbapenemase detection. A total of 3,396 isolates were reported in this group (Table 1).

Screening of resistance to a carbapenem according to the disk diffusion method was common among the studies.^{8–23} Although this method may be applicable for many antimicrobials, however, for carbapenems more confirmatory tests beyond the disk diffusion assay are needed. In this setting, MIC would help for detection of resistant isolates.

Five studies were multicenter,^{8–12} but the sample sizes for three of them were small.^{10,13,14} The trend of resistance to a carbapenem did not follow a specific pattern. However, in two studies,^{12,15} 40% of the isolates were resistant to a carbapenem.

Studies that considered phenotypic methods

A total of 4,264 isolates were included in this category and *Acinetobacter* spp. and *K. pneumonia* were the common isolates. Modified Hodge test (MHT) was used as a common phenotypic test in most studies. MHT can detect Ambler class A (KPC), although it may be positive with the presence of other carbapenemase enzymes. Other applied tests were combination disk test (CDT) for detection of MBL or extended spectrum β -lactamase and the double disk synergy test for detection of MBL. In the study by Raategar Lari et al,²⁴ 92% of *A. baumannii* isolates were resistant to imipenem and only 24% of them were MBL positive in the CDT test, but in the next study,²⁵ 54% of isolates were resistant to imipenem and all of them were KPC producers according to the phenotypic tests.

The rates of resistance to aminoglycosides and fluoroquinolones were similar to those of carbapenems, although no special trend can be followed through different years. The results are provided in Table 2.^{26–41}

Studies that included genotypic methods

A total of 2,195 strains were included in this group and except for three studies, all assessed *Acinetobacter* spp. mostly *A. baumannii*. PCR was performed in order to determine types of *bla* genes. In some studies, MIC values were correlated with the presence of *bla* genes. In the study by Taherikalani et al the MIC values of 256 mcg/mL were detected in strains with more than two *bla*_{OXA} genes.⁴² In the study by Azizi et al, higher MIC value was associated with presence of *bla*_{OXA-24/40 like}.⁴³ Bahador et al detected *IS*_{Aba} genes.⁴⁴ When these elements were placed upstream of OXA-type genes, they enhanced the expression of OXA-type genes. Presence of

Table 1 Screening of carbapenem-resistant isolates (disk diffusion method)

Author (year)	Setting	Population	Samples	Microorganism (No.)	MIC ^a	Resistance rate to FQ and AG	Results ^b	Weak points	Strength points
Mobaraki et al (2018) ¹⁵	Hospitals in Tabriz	Inpatients and outpatients	Different sites	<i>Pseudomonas aeruginosa</i> (200)	–	TOB 57% GEN 62% CIP 62% c	<ul style="list-style-type: none"> 46.5% of isolates were resistant to IPM 	–	<ul style="list-style-type: none"> Multicenter study
Ghanbari et al (2017) ¹⁶	One hospital in Fuldshahr	General inpatients	Urine	Different strains (317)	–	c	<ul style="list-style-type: none"> 9.5% of <i>Escherichia coli</i>, 11% of <i>Klebsiella</i> and 25% of <i>Proteus spp.</i> isolates were resistant to IPM 	–	<ul style="list-style-type: none"> Large sample size
Chahoufard et al (2017) ¹⁷	One hospital in Bandar Abbas	General inpatients	Urine	Different strains (296)	–	c	<ul style="list-style-type: none"> Resistance rate to IPM: <i>Enterobacter spp.</i> 7.1% <i>Citrobacter spp.</i> 0% <i>E. coli</i> 4.9% <i>Klebsiella pneumoniae</i> 19.2% <i>P. aeruginosa</i> 21.4% <i>Acinetobacter baumannii</i> 53.2% 	–	<ul style="list-style-type: none"> Adequate sample size Clinical status of patients including duration of hospital stay and comorbidities were evaluated
Ansari et al (2017) ¹⁸	Two hospitals in Shahrekord	General inpatients	Urine, blood, sputum, wound	<i>A. baumannii</i> (30)	–	NAL 80%, OFX 86% LVX 66% –	<ul style="list-style-type: none"> About 55% of strains were resistant to IPM and DOR Intermediate resistance rate was 10%–33.3% 97% were resistant to carbapenems MIC values of all resistant isolates were above 32 mcg/mL 	<ul style="list-style-type: none"> Small sample size AGs were not tested 	<ul style="list-style-type: none"> Cloning smpA gen
Douraghi et al (2016) ⁸	Four hospitals in Tehran	General inpatients and outpatients	Different sites	<i>A. baumannii</i> (400)	✓	–	<ul style="list-style-type: none"> 97% were resistant to carbapenems MIC values of all resistant isolates were above 32 mcg/mL 	–	<ul style="list-style-type: none"> Multicenter study Large sample size
Ghasemian et al (2016) ⁹	Four hospitals in Mazandaran	General ICU patients	Different sites	<i>Acinetobacter spp.</i> (50)	✓	CIP 96% AMK 100%	<ul style="list-style-type: none"> Resistance to IPM and MEM according to E-test was 100%, and according to disk diffusion was 76% and 96%, respectively Most common site for sampling was endotracheal tube 	<ul style="list-style-type: none"> Small sample size MIC values did not report 	<ul style="list-style-type: none"> Multicenter study
Babamahmoodiet al (2015) ¹³	Three hospitals in Mazandaran	ICU patients	Wound, respiratory secretions, urine, blood	Different strains (114)	–	c	<ul style="list-style-type: none"> 14% of <i>P. aeruginosa</i>, 60% of <i>Acinetobacter spp.</i>, 33% of <i>E. coli</i>, 16% of <i>Enterobacter</i> and none of <i>K. pneumoniae</i> isolates were resistant to IMP 	<ul style="list-style-type: none"> Small sample size for each strain 	<ul style="list-style-type: none"> Multicenter study Demographic features and risk factors for hospital-acquired infection were described

(Continued)

Table 1 (Continued)

Author (year)	Setting	Population	Samples	Microorganism (No.)	MICa	Resistance rate to FQ and AG	Resultsb	Weak points	Strength points
Mohajeri et al (2014) ¹⁴	Three hospitals in Kermanshah	General inpatients	Blood, urine, sputum	<i>Acinetobacter</i> spp. (104)	–	Not defined	<ul style="list-style-type: none"> 48% of isolates were resistant to carbapenems (imipenem and meropenem). 76% of carbapenem-resistant strains were isolated from sputum samples 	<ul style="list-style-type: none"> Few antibiotics were tested 	<ul style="list-style-type: none"> Multicenter study
Babakhani et al (2014) ¹⁹	One hospital in Khorramabad	General inpatients (mostly ICU)	Different sites	<i>Klebsiella</i> spp. (80)	–	GEN 52 % AMK 7.5% CIP 82 % OFX 75% NAL 60% CIP 97%	<ul style="list-style-type: none"> 67% and 1% of isolates were resistant to IMP and MEM, respectively 	–	–
Kamalbeik et al (2013) ²⁰	One Poisoning center in Tehran	ICU patients	Different sites	<i>Acinetobacter</i> spp. (40)	–	–	<ul style="list-style-type: none"> Resistance rates to MEM and IPM were 100% and 62% (27.5% intermediate resistant to IPM) 	<ul style="list-style-type: none"> Small sample size 	–
Hashemi et al (2013) ²¹	Two hospitals in Hamadan	General inpatients and outpatients	Different sites	<i>Enterobacteriaceae</i> spp. (574)	–	–	<ul style="list-style-type: none"> Resistance rate to IPM was 32% in inpatients and 4% in outpatients. Resistance rates to IPM in inpatients: <i>Enterobacter</i> 56%, <i>Serratia</i> 54%, <i>Klebsiella</i> 35%, <i>E. coli</i> 32%, and <i>Proteus</i> 11% 	–	<ul style="list-style-type: none"> Large sample size
Shakibaie et al (2012) ¹⁰	One hospital in Kashan	ICU patients	Different sites	<i>Acinetobacter</i> spp. (15)	✓	CIP 60% GEN 60% AMK 26%	<ul style="list-style-type: none"> 73% of isolates were resistant to IPM MIC range for resistant strains was 128–240 and MIC of IPM for seven isolates was 240 mcg/mL 	<ul style="list-style-type: none"> Very small sample size 	–
Rahbar et al (2010) ²²	One hospital in Tehran	General inpatients	Different sites	<i>P. aeruginosa</i> (109) <i>Acinetobacter</i> spp. (88) <i>Stenotrophomonas maltophilia</i> (48) <i>Burkholderia cepacia</i> (12)	–	–	<ul style="list-style-type: none"> 1 %, 16%, and 98% of <i>A. baumannii</i>, <i>P. aeruginosa</i>, and <i>S. maltophilia</i> isolates were resistant to IPM, respectively 	–	<ul style="list-style-type: none"> Demographic and medical data of patients were provided

(Continued)

Table 1 (Continued)

Author (year)	Setting	Population	Samples	Microorganism (No.)	MIC ^a	Resistance rate to FQ and AG	Results ^b	Weak points	Strength points
Soroush et al (2010) ²³	One pediatric hospital in Tehran	General pediatric patients	Different sites	<i>Acinetobacter</i> spp. (145)	–	In 2007: CIP 79% AMK 58% KAN 71% GEN 65% TOB 61%	<ul style="list-style-type: none"> Resistance rate to IPM and MEM was analyzed in years 2006 and 2007 that was about 50% 	<ul style="list-style-type: none"> Carbapenem resistance was evaluated only in the last 2 years (2006–2007) 	<ul style="list-style-type: none"> Evaluated carbapenem resistance status in pediatrics Change in the antibiotics resistance patterns was considered Site of sampling with highest count of MDR strains in each year considered
Yousefi et al (2010) ¹¹	One hospital in Orumie	General inpatients and outpatients	Different sites	<i>P. aeruginosa</i> (160)	✓	–	<ul style="list-style-type: none"> Approximately 40% of isolates were resistant to IPM, which means MIC for them was 31 mcg/mL Presence of Class I integron and being in ICU and burn unit led to significantly higher IPM-resistant strains 40% and 7% of <i>P. aeruginosa</i> and 38% and 27% of <i>A. baumannii</i> isolates were resistant to MEM and IMP, respectively MIC range value for MEM was 0.5–32 mcg/mL The majority isolates of <i>P. aeruginosa</i> and <i>A. baumannii</i> had an MIC >32 µg/mL for MEM 	<ul style="list-style-type: none"> Just IPM was tested 	–
Rahbar et al (2008) ¹²	One hospital in Tehran	General inpatients and ICU patients	Not defined	Different strains (202)	✓	–	–	–	–

Notes: ^aMIC values of carbapenems (and not other antimicrobial agents) were determined. ^bResistant rates were reported according to the disk diffusion test. The unit for all reported MIC values is mcg/mL. ^cFor complete information about the resistance to AG and FQ, refer to the original reference.

Abbreviations: AG, aminoglycoside; AMK, amikacin; CDT, combination disk test; CIP, ciprofloxacin; DD, disk diffusion; DDST, double disk synergy test; DOR, doripenem; ERT, erlapenem; ESBL, extended spectrum β -lactamase; FQ, fluoroquinolones; GEN, gentamicin; ICU, intensive care unit; IPM, imipenem; KAN, kanamycin; LYX, levofloxacin; MBL, metallo- β -lactamase; MEM, meropenem; MHT, modified Hodge test; MIC, minimum inhibitory concentration; NAL, nalidixic acid; NOR, norfloxacin; OFX, ofloxacin; TOB, tobramycin.

Table 2 Studies that considered phenotypic methods for detecting carbapenemase enzymes

Author (year)	Setting	Population	Samples	Microorganism (no.)	MIC ^a	Method of carbapenemase detection	Resistance to FQ and AG	Results ^b	Weak points	Strength points
Ghotaslou et al (2018) ²⁶	Five hospitals in East and West Azerbaijan	General inpatients	Different sites	<i>Enterobacteriaceae</i> (307)	–	<ul style="list-style-type: none"> • DD • BA and DPA for MBL detection 	ε	<ul style="list-style-type: none"> • Resistance rate to carbapenem was observed only in <i>K. pneumonia</i> (57 isolates; 25%–45%) • Of these 57 isolates, phenotypic methods revealed that 15%, 7%, and 3% isolates of <i>Klebsiella</i> had MBL, KPC, and OXA-48 genes, respectively • Most of the carbapenemase strains were isolated from urine and blood 	<ul style="list-style-type: none"> • Small sample size for each species 	–
Saadatian Farivar et al (2018) ²⁷	Three hospitals in Tehran	Inpatients and outpatients	Different sites	<i>Klebsiella pneumonia</i> (81)	–	<ul style="list-style-type: none"> • DD • MHT for MBL detection 	OFX 65%, CIP 68%, NOR 66%, GEN 66%, AMK 51%, TOB 56%, KAN 79%, GEN 96%, AMK 96%, CIP 98%	<ul style="list-style-type: none"> • Resistance rate to IPM was 45% • 6 (7.4% of all) isolates were positive for MHT test 	–	–
Moosavian et al (2014) ²⁸	Two hospitals in Ahvaz	General inpatients	Different sites	<i>Acinetobacter</i> spp. (100)	–	<ul style="list-style-type: none"> • DD • MHT 	GEN 96%, AMK 96%, CIP 98%	<ul style="list-style-type: none"> • Resistance rates to IPM, MEM, and ERT were 95%, 96%, and 53%, respectively • 53% of all strains were MHT positive 	–	–

(Continued)

Table 2 (Continued)

Author (year)	Setting	Population	Samples	Microorganism (no.)	MIC ^a	Method of carbenemase detection	Resistance to FQ and AG	Results ^b	Weak points	Strength points
Jonaiddi Jafari et al (2014) ²⁹	One hospital in Tehran	General inpatients	Different sites	Different strains (350)	✓	<ul style="list-style-type: none"> • DD • CDT for ESBL detection 	–	<ul style="list-style-type: none"> • MIC was determined by E-test • 95 strains were ESBLs • Resistance rates to IPM and MEM according to MIC, respectively, were: <i>Acinetobacter baumannii</i> 42%, 37% <i>Pseudomonas aeruginosa</i> 42%, 60% <i>K. pneumonia</i> 15%, 0% 	<ul style="list-style-type: none"> • Small sample size for each microorganism 	–
Ghadiri et al (2014) ³⁰	Three laboratories in Tehran	General outpatients	Urine	<i>Escherichia coli</i> (300)		<ul style="list-style-type: none"> • DD • CDT for ESBL detection • DDST for MBL detection 	AMK 11% GEN 16%	<ul style="list-style-type: none"> • 67 isolates were ESBL producers and resistance to carbenem in this group was about 10% • 21 isolates were MBL producers and resistance to carbenems among them was almost 100% • 5 isolates were both MBL and ESBL positive 	–	<ul style="list-style-type: none"> • Large sample size

(Continued)

Table 2 (Continued)

Author (year)	Setting	Population	Samples	Microorganism (no.)	MIC ^a	Method of carbapenemase detection	Resistance to FQ and AG	Results ^b	Weak points	Strength points
Fazeli et al (2014) ³¹	Teaching hospitals in Isfahan	General inpatients and ICU patients	Different sites	<i>K. pneumonia</i> (142)	–	<ul style="list-style-type: none"> • DD • CDT for ESBL 	Resistance rate to GEN and AMK and CIP: 70%–80% LYX: 50%–60%	<ul style="list-style-type: none"> • Resistance rates to IPM, MEM, and ERT among all strains were 57%, 52%, and 47%, respectively • 101 strains were ESBL producers • Most of the ESBL producers were isolated from ICU and from urine samples 	<ul style="list-style-type: none"> • It is not clear that the study was multicenter or not 	–
Mirsalehian et al (2014) ³²	Morahari hospital	Burn patients	Probably burn wounds	<i>P. aeruginosa</i> (100)	✓	<ul style="list-style-type: none"> • DD • AmpC disk and CDT for AmpC production 	–	<ul style="list-style-type: none"> • All IPM-resistant <i>Pseudomonas</i> were chosen and mean MIC for IPM was 64 mcg/mL • AmpC disk test and CDT were positive for 54 and 17 isolates, respectively showing a probable activation of one of the efflux pump and impermeability systems, or both 	–	–

(Continued)

Table 2 (Continued)

Author (year)	Setting	Population	Samples	Microorganism (no.)	MIC ^a	Method of carbapenemase detection	Resistance to FQ and AG	Results ^b	Weak points	Strength points
Moayednia et al (2014) ³³	One hospital in Isfahan	General inpatients and outpatients	Urine	<i>K. pneumonia</i> (484) and <i>E. coli</i> (1,080)		<ul style="list-style-type: none"> • DD • Double Disk for ESBL • MHT for KPC detection • CDT for MBL detection 	Just reported for ESBL producer	<ul style="list-style-type: none"> • Resistance rate to carbapenem for <i>E. coli</i> was 1%–2% and for <i>Klebsiella</i> spp., it was 41%–46% • 390 isolates of <i>E. coli</i> and 238 <i>Klebsiella</i> were ESBL producers • From all of strains: <i>E. coli</i>: 2 MBL and 10 KPC producers <i>Klebsiella</i> spp.: 10 MBL and 186 KPC producers 	<ul style="list-style-type: none"> • MIC was done by E-test, but cutoffs were not reported 	<ul style="list-style-type: none"> • Large sample size
Erfani et al (2013) ³⁴	Three hospitals in Tehran	General inpatients	Different sites	<i>Acinetobacter</i> sp. (107)	–	<ul style="list-style-type: none"> • DD • CDT for MBL detection 	CIP 95% LVX 93% GEN 51% TOB 64%	<ul style="list-style-type: none"> • Resistance rate to IPM and MEM was about 95% and 92% of these resistant isolates were MBL positive 	–	–
Lari et al (2013) ²⁴	One hospital in Tehran (Motahhari)	Burn patients	Mostly burn wound	<i>Acinetobacter</i> spp. (69)	–	<ul style="list-style-type: none"> • SDDT for ESBL detection • CDT for MBL detection 	GEN 64.7% AMK 95.7% KAN 97.1% TOB 2.9% CIP 97.1%	<ul style="list-style-type: none"> • Resistance rate to IPM was 92% • ESBL was positive in 10% of isolates • 24% of all baumannii isolates were MBL positive • The mortality rate in this study was 20% 	–	<ul style="list-style-type: none"> • Mortality rate was reported • Detection of <i>Integron</i> genes

(Continued)

Table 2 (Continued)

Author (year)	Setting	Population	Samples	Microorganism (no.)	MIC ^a	Method of carbapenemase detection	Resistance to FQ and AG	Results ^b	Weak points	Strength points
Rastegar Lari et al (2013) ²⁵	One hospital in Tehran (Motahari)	Burn patients	Burn wounds	<i>Klebsiella spp.</i> (35)	–	<ul style="list-style-type: none"> DD MHT for KPC production 	AMK 71% GEN 72% TOB 86%	<ul style="list-style-type: none"> Resistance rate to IPM was 54% and all were KPC producers 7 out of 28 patients had two <i>Klebsiella</i> isolates with two different antibiotypes Rate of mortality in patients infected with resistant strain was 33% 	<ul style="list-style-type: none"> FQ and colistin were not tested MIC values were not reported Small sample size 	<ul style="list-style-type: none"> Mortality rate was reported
Safari et al (2013) ³⁵	Three educational hospitals in Hamadan	ICU patients	Trachea, blood, urine, sputum, wounds	<i>A. baumannii</i> (100)	✓	<ul style="list-style-type: none"> DD E-test for MBL 	Resistant: CIP 97% LYX 91% AMK 84% GEN 88%	<ul style="list-style-type: none"> Resistance rate to carbapenem was 85%–94% but according to MIC, it was 87%–99% 99% of all isolates were MBL producers 		<ul style="list-style-type: none"> Multicenter study MIC was reported
Massaeli et al (2012) ³⁶	Two hospitals in Sanandaj	General inpatients	Different sites	Different strains (423)	–	<ul style="list-style-type: none"> DD DDST for MBL detection 	GEN 70% AMK 80% CIP 37% NOR 42%	<ul style="list-style-type: none"> The result of disk diffusion for carbapenem was not reported 126 isolates (30% of all) were MBL positive Risk factors like immunosuppression and use of antibiotics during the past 2 weeks were correlated with infection by MBL producers 	<ul style="list-style-type: none"> The resistance pattern of each group of microorganisms was not fully defined 	<ul style="list-style-type: none"> Correlation between being MBL and different clinical risk factors was assessed

(Continued)

Table 2 (Continued)

Author (year)	Setting	Population	Samples	Microorganism (no.)	MIC ^a	Method of carbapenemase detection	Resistance to FQ and AG	Results ^b	Weak points	Strength points
Japoni-Nejad et al (2013) ³⁷	One hospital in Arak	General inpatients and environmental isolates	Different strains	<i>A. baumannii</i> (63)	✓	<ul style="list-style-type: none"> MHT for KPC detection E-test 	Amikacin 80% Netilmicin 54%	<ul style="list-style-type: none"> 84% were resistant to carbapenem 47 isolates were MBL producers according to E-test and three of them were MHT positive 60 strains had MIC more than 256 	<ul style="list-style-type: none"> Isolates from environment of hospital were also included 	<ul style="list-style-type: none"> Small sample size
Haji Hashemi et al (2012) ³⁹	Different hospitals in Tehran and Isfahan	General inpatients and outpatients	Different sites	<i>Klebsiella pneumoniae</i> (244)	–	<ul style="list-style-type: none"> DD MHT for KPC detection CHROMagar for KPC detection 	–	<ul style="list-style-type: none"> MHT and CHROMagar were compared From all, 30 and 36 isolates were KPC producers according to MHT and CHROMagar, respectively. The infection of burn wounds at the highest percentage: 27 cases (75 %) 	<ul style="list-style-type: none"> MIC was not reported 	<ul style="list-style-type: none"> Multicenter study

(Continued)

Table 2 (Continued)

Author (year)	Setting	Population	Samples	Microorganism (no.)	MIC ^a	Method of carbapenemase detection	Resistance to FQ and AG	Results ^b	Weak points	Strength points
Azimi et al (2012) ⁴⁰	Two burn hospitals in Tehran	Infant and pediatric patients	Not defined (probably burn wounds)	<i>Klebsiella</i> , <i>Acinetobacter</i> , and <i>Pseudomonas</i> (64 strains from 20 patients)	–	<ul style="list-style-type: none"> • DD • MHT for KPC production • CDT for ESBL detection 	Not defined	<ul style="list-style-type: none"> • 36 isolates were resistant to all tested antibiotic (not mentioned which antibiotics) except Colistin • Among these 36 isolates, 15 isolates were resistant to IPM • From these 15 isolates, 13 were KPC and 6 were ESBL producers • Resistance rate to carbapenems was 33% • MIC value for IPM was 2 and for MEM was 1 • MBL was not detected 	<ul style="list-style-type: none"> • Types of samples were not defined • The sample size is very small (7 <i>A. baumannii</i>, 16 <i>Enterobacteriaceae</i>, 12 <i>Pseudomonas</i>) • There is no precise data of resistance to each antibiotic 	
Japoni et al (2006) ⁴¹	One hospital in Shiraz	General inpatients	Different sites (mostly wound)	<i>P. aeruginosa</i> (70)	✓	<ul style="list-style-type: none"> • DD • DDST for ESBL • MHT for MBL detection 	CIP 73% AMK 93% TOB 97%		<ul style="list-style-type: none"> • Small sample size 	

Notes: ^aMIC values for carbapenems (but not other antimicrobial agents) were determined. ^bResistance rates were according to the result of disk diffusion test. ^cFor complete information about the resistance to AG and FQ, please refer to the original reference.

Abbreviations: AG, aminoglycoside; AMK, amikacin; BA, boronic acid; CDT, combination disk test; CIP, ciprofloxacin; DD, disk diffusion; DDST, double disk synergy test; DOR, doripenem; DPA, dipicolinic acid; ERT, ertapenem; ESBL, extended spectrum β -lactamase; FQ, fluoroquinolones; GEN, gentamicin; ICU, intensive care unit; IPM, imipenem; KAN, kanamycin; KPC, *Klebsiella*-producing carbapenemase; LVX, levofloxacin; MBL, metallo- β -lactamase; MEM, meropenem; MHT, modified Hodge test; MIC, minimum inhibitory concentration; NAL, nalidixic acid; NOR, norfloxacin; OFX, ofloxacin; TOB, tobramycin.

these elements in genetic contents of gram-negative isolates would lead to higher rate of resistance to carbapenems. Table 3 describes the details of each study in this category.^{42,45–61}

Studies that considered both phenotypic and genotypic methods

A total of 6,759 isolates were identified in this category. In most studies more than one phenotypic method was applied. In the study by Hosseinzadeh et al, 29 strains were carbapenem resistant, in which 27 and 23 of them were MHT positive and carried *bla*_{NDM-1} gene, respectively.⁶² However, in another study by Bina et al, although MHT tests were positive, no resistance gene was found.⁶³

The detection of resistance genes was much lower for strains other than *A. baumannii*. In the study by Lari et al in 2015, only 9 strains of *P. aeruginosa* (out of 255) contained resistance genes,⁶⁴ but in the study by Bagheri Josheghani et al, at the same time, more than 90 strains (out of 124) of *A. baumannii* harbored resistance genes.⁶⁵ Data are summarized in Table 4.^{66–105}

Discussion

Increase in the rate of resistance to carbapenem antibiotics among gram-negative isolates is a worldwide concern, especially in developing countries. The results of this review also showed this threat in the recent decade in a country of Middle East area, Iran. Although the prevalence of carbapenem-resistant gram-negative rods was reported according to the disk diffusion method in most studies, phenotypic and genotypic assays confirmed this pattern in some available surveys.

Resistance to carbapenem antibiotics in the same years in a country of neighborhood, Saudi Arabia, shows lower resistance rate in comparison with Iran.¹⁰⁶ In 2010, resistance to carbapenems was reported in 10%–66% of gram-negative isolates in Saudi Arabia, which was lower than that in Iran (86%).¹¹ The results were different in Europe. In the year 2006–2007, 4% and 85% of gram-negative isolates were carbapenem resistant in North and South of Europe, respectively. Although data were limited to year 2006, Iran had a better condition compared with the South of Europe, but worse than North of Europe.¹⁰⁷ The results of another study in year 2012 also confirmed this pattern in North and South of Europe.¹⁰⁸ In case of *P. aeruginosa*, MIC values that were reported for carbapenem-resistant strains were dramatically different from European countries. In year 2011, Castanheira et al detected a mean MIC value >2 mcg/mL for carbapenem-resistant *P. aeruginosa* in 14 European countries.¹⁰⁹ Yousefi et al in Iran reported an MIC >32 mcg/mL for most of carbapenem-

resistant *P. aeruginosa* isolates. At these times, the CLSI and European Committee on Antimicrobial Susceptibility Testing cutoff values for carbapenem resistance were 2 and 8 mcg/mL, respectively.¹¹

Incremental trend of resistance to carbapenem antibiotics was evident through 2006–2018. The least resistance rate was reported by Rahbar et al in year 2010 at Milad Hospital. In this year, 1.1% of *A. baumannii* isolates were resistant to imipenem.²² This conclusion may be misleading, because 2 years earlier (in 2008), in this hospital, 27% and 40% of the same isolates were resistant to imipenem and meropenem, respectively.¹² Tarashi et al assessed the trend of antibiotic resistance in *A. baumannii* through years 2012–2015.⁷¹ The results showed that resistance in *P. aeruginosa* increased from 83% in year 2012 to 96% in year 2015. Also 100% of *A. baumannii* were resistant to carbapenem among these years.

The resistance patterns for other microorganisms did not follow specific trends. For example, 60% of *P. aeruginosa* isolates were imipenem resistant in the year 2010,¹⁰¹ but this rate was 25% in another hospital in the year 2016.⁶⁹ Outpatients-isolated *P. aeruginosa* strains showed less resistance rates (13% in Ahangarzadeh Rezaee et al³⁸ and 4.4% in Hashemi et al²¹). However, carbapenem-resistant rates were dramatically high in studies that included ICU samples.^{9,20} In an earlier study,⁹ 100% and 62% of *P. aeruginosa* isolates were resistant to imipenem and meropenem, respectively, and in a latter study,²⁰ about 100% of them were resistant to both carbapenems.

Most of the carbapenem-resistant microorganisms were isolated from burned patients, and many studies placed in this group were from Motahari Hospital. Motahari is a referral teaching burn hospital in center of Iran and most complicated burned patients are referred from all areas of the country. Increasing resistance rates can be observed through different years in burn patients; 50%–60% in year 2008 to 90% in year 2017.^{42,46} Also, the presence of *bla*_{oxA} type genes has increased from half of the resistant strains to almost 80% of them. This can be a warning alarm regarding administration of antibiotics in burned patients.

Disk diffusion is the initial method for detection of resistant strains; however, it does not have enough accuracy for antimicrobial agents like carbapenem. Most evaluated studies considered disk diffusion method. Beside disk diffusion, MIC breakpoints can be used to confirm the results of disk diffusion (Figure 3).

Although MHT has been widely used as the preferred phenotypic method for detection of MBL, it is not recommended in latest version of CLSI, due to its low sensitiv-

Table 3 Studies that performed genotypic methods to detect genes encoding carbapenemase enzymes

Author (year)	Setting	Population	Samples	Microorganism (no.)	MIC	Method of carbapenemase detection	Resistance to FQ and AG	Results ^b	Weak points	Strength points	Found carbapenemase genes ^a
Sharikhani et al (2017) ⁴⁵	Four hospitals in Qom	General inpatients (mostly ICU)	Different sites	<i>Acinetobacter baumannii</i> (108)	–	<ul style="list-style-type: none"> • DD • PCR 	CIP and AMK 93% GEN 81% LVX 91% TOB 47%	<ul style="list-style-type: none"> • 89% isolates were non-susceptible to IPM and MEM • Among carbapenem non-susceptible isolates, 82%, 55%, 22%, and 14% isolates had <i>bla</i>_{OXA-23}, <i>bla</i>_{OXA-58}, <i>bla</i>_{OXA-40} and <i>bla</i>_{OXA-143} genes, respectively • About 90% of isolates were resistant to carbapenem • <i>bla</i>_{OXA-23} • <i>bla</i>_{OXA-24} • <i>bla</i>_{OXA-58} and <i>bla</i>_{PER} genes were detected in 90%, 38%, 1%, 60%, and 18.5% of all isolates, respectively • Significant relationship between the presence of <i>bla</i>_{OXA-24} and resistance to IPM (not MEM) • All strains had biofilm producing ability 	–	<ul style="list-style-type: none"> • First report of <i>bla</i>_{OXA-143} in Iran • Multicenter study 	<i>bla</i> _{OXA-51} <i>bla</i> _{OXA-40} <i>bla</i> _{OXA-143}
Mohammadi et al (2017) ⁴⁶	Two hospitals in Tehran	General inpatients	Burn wound and trachea	<i>A. baumannii</i> (103)	–	<ul style="list-style-type: none"> • DD • PCR 	–	<ul style="list-style-type: none"> • About 90% of isolates were resistant to carbapenem • <i>bla</i>_{OXA-23} • <i>bla</i>_{OXA-24} • <i>bla</i>_{OXA-58} and <i>bla</i>_{PER} genes were detected in 90%, 38%, 1%, 60%, and 18.5% of all isolates, respectively • Significant relationship between the presence of <i>bla</i>_{OXA-24} and resistance to IPM (not MEM) • All strains had biofilm producing ability 	–	<ul style="list-style-type: none"> • Ability of biofilm formation was assessed • Correlation between presence of genes and resistance to antibiotics was defined 	<i>bla</i> _{OXA-23} <i>bla</i> _{OXA-58} <i>bla</i> _{PER}

(Continued)

Table 3 (Continued)

Author (year)	Setting	Population	Samples	Microorganism (no.)	MIC	Method of carbapenemase detection	Resistance to FQ and AG	Results ^b	Weak points	Strength points	Founded carbapenemase genes ^a
Bahador et al (2015) ⁴⁴	One hospital in Tehran	Burn patients	Not defined (probably burn wounds)	<i>A. baumannii</i> (62)	✓	<ul style="list-style-type: none"> • DD • PCR • Clonal typing 	Not reported specifically	<ul style="list-style-type: none"> • According to MIC values, 61% were resistant to IPM • From all strains, 39 had <i>bla</i>_{OXA-23} (33 were susceptible to IPM), 9 had <i>bla</i>_{OXA-58} genes • Existence of two genes (in 1.6%–19.3% of strains) was correlated with MIC more than 32 mcg/mL for IPM 	<ul style="list-style-type: none"> • Resistance to each antibiotic was not reported specifically • Small sample size 	<ul style="list-style-type: none"> • MIC values of IPM, tigecycline, and colistin were reported • ISAb gene was assessed 	<i>bla</i> _{OXA-51} <i>bla</i> _{OXA-23} <i>bla</i> _{OXA-24} <i>bla</i> _{OXA-40} <i>bla</i> _{OXA-143}
Kooti et al (2015) ⁴⁷	Four hospitals in Shiraz	General inpatients	Different sites	<i>A. baumannii</i> (200)	–	<ul style="list-style-type: none"> • DD • PCR 	GEN 84.5% AMK 86.5% CIP and LVX 99.5%	<ul style="list-style-type: none"> • About 99% of isolates were resistant to IPM and MEM • 40% had <i>bla</i>_{OXA-23}, 7% had <i>bla</i>_{OXA-24}, 0.5% had <i>bla</i>_{OXA-58} • Two <i>bla</i> genes were detected in 4.5% of strains • The correlation between presence of genes and source of samples was not significant 	–	Multicenter study	<i>bla</i> _{OXA-23} <i>bla</i> _{OXA-24} <i>bla</i> _{OXA-58}

(Continued)

Table 3 (Continued)

Author (year)	Setting	Population	Samples	Microorganism (no.)	MIC	Method of carbapenemase detection	Resistance to FQ and AG	Results ^b	Weak points	Strength points	Founded carbapenemase genes ^a
Azizi et al (2015) ⁴³	Two hospitals in Kerman	ICU patients	Different sites	<i>A. baumannii</i> (65 MDR strains from 266)	✓	<ul style="list-style-type: none"> • DD • PCR 	CIP 100% AMK 78.5% TOB 93%	<ul style="list-style-type: none"> • MIC values for IPM and MEM for 76% of strains were >256 mcg/mL • <i>bla_{oxa-23}</i> was found in all isolates including sensitive and resistant isolates • Presence of <i>bla_{oxa-24/40}</i> (29 strains) associated with higher MIC 	–	–	<i>bla_{oxa-51}</i> , <i>bla_{oxa-23}</i> , <i>bla_{oxa-24/40}</i>
Mahdian et al (2015) ⁴⁶	One hospital in Tehran (Motahari)	Burn patients	Burn wounds, blood, urine	<i>A. baumannii</i> (37)	–	<ul style="list-style-type: none"> • DD • PCR • Clonal typing 	CIP 100% GEN 94.6%	<ul style="list-style-type: none"> • 81% of strains were resistant to carbapenem <ul style="list-style-type: none"> o All resistant isolates had at least one <i>bla_{oxa-23}</i> and/or <i>bla_{oxa-24}</i> and about half of them had both genes 	<ul style="list-style-type: none"> • Small sample size • Few antibiotics were tested 	<ul style="list-style-type: none"> • MIC values for colistin and polymyxin B were reported • <i>ISAb_a</i> genes were assessed 	<i>bla_{oxa-23}</i> , <i>bla_{oxa-24}</i> , <i>bla_{TEM}</i> , <i>bla_{PER}</i>

(Continued)

Table 3 (Continued)

Author (year)	Setting	Population	Samples	Microorganism (no.)	MIC	Method of carbapenemase detection	Resistance to FQ and AG	Results ^b	Weak points	Strength points	Founded carbapenemase genes ^a
Farsiani et al (2015) ⁴⁹	One hospital in Mashhad	General inpatients and ICU patients	Different sites	<i>A. baumannii</i> (36)	✓	<ul style="list-style-type: none"> • DD • PCR • Clonal typing 	CIP 97%	<ul style="list-style-type: none"> • 97% of strains were resistant to carbapenems • MIC values of resistant isolates were >32 • <i>bla</i>_{OXA-51} and <i>bla</i>_{OXA-23} were detected in all isolates • From all isolates, <i>bla</i>_{OXA-24}, <i>bla</i>_{TEM}, <i>bla</i>_{ADC} and <i>bla</i>_{VIM} genes were found in 23, 34, 22, and 23 strains, respectively • ISAbal was detected in 97% of isolates of <i>A. baumannii</i> 	<ul style="list-style-type: none"> • Small sample size 	<ul style="list-style-type: none"> • ISAbal genes and tet genes (related to efflux pump) were analyzed 	<i>bla</i> _{OXA-51} , <i>bla</i> _{OXA-23} , <i>bla</i> _{OXA-24} , <i>bla</i> _{TEM} , <i>bla</i> _{ADC} , <i>bla</i> _{VIM} , <i>adeB</i> , <i>ISAbal</i> , <i>tetA</i> , <i>tetB</i>
Nasrolahei et al (2014) ⁵⁰	Two hospitals in Tehran and Sari	ICU and burn patients	Trachea and burn wounds	<i>A. baumannii</i> (100)	–	<ul style="list-style-type: none"> • DD • PCR 	STP 90% GEN 83% TOB 83% KAN 80%	<ul style="list-style-type: none"> • Resistance rate to IPM and MEM was about 70% • 67% of strain carried <i>bla</i>_{OXA-23} gene 	<ul style="list-style-type: none"> • FQ are not tested 	<ul style="list-style-type: none"> • Multicenter study 	<i>bla</i> _{OXA-23}
Safari et al (2014) ⁵¹	Three hospitals in Hamadan	ICU patients	Different sites	<i>Pseudomonas aeruginosa</i> (100)	✓	<ul style="list-style-type: none"> • DD • PCR 	AMK 19% GEN 28% TOB 27% LVX 28% CIP 38%	<ul style="list-style-type: none"> • Resistance rate to carbapenems was about 20% • According to MIC values, 24% of isolates were carbapenem-resistant • Four isolates had <i>bla</i>_{IMP} 	–	–	<i>bla</i> _{IMP}

(Continued)

Table 3 (Continued)

Author (year)	Setting	Population	Samples	Microorganism (no.)	MIC	Method of carbapenemase detection	Resistance to FQ and AG	Results ^b	Weak points	Strength points	Founded carbapenemase genes ^a
Bahador et al (2014) ⁵²	Two hospitals in Tehran	ICU patients	Different sites	<i>A. baumannii</i> (100)	✓	<ul style="list-style-type: none"> DD Clonal typing 	c	<ul style="list-style-type: none"> Resistance rates were compared among years 2006 and 2011, according to MIC cutoffs In year 2006, resistance rate to IPM was 30% with MIC ≤4–64 mcg/mL. In year 2011 it was 48% and MIC values were ≤4 to ≥256 mcg/mL Genotypes I and F were the most frequent genotypes in years 2006 and 2011, respectively 		<ul style="list-style-type: none"> Range of MIC values for each antibiotic was reported 	–
Karmostaji et al (2013) ⁵³	Two hospitals in Tehran	General inpatients (mostly ICU)	Different sites	<i>Acinetobacter</i> spp. (131)	–	<ul style="list-style-type: none"> DD PCR 	<p>CIP 95% GEN 77% AMK 54%</p>	<ul style="list-style-type: none"> Resistance rate to IPM and MEM were 67% and 84%, respectively 107, 17, and 1 isolates carried of the resistant <i>bla</i>_{OXA-23}[*] and <i>bla</i>_{OXA-24}[*] and <i>bla</i>_{OXA-58}[*] respectively. From these strains, seven strains had both <i>bla</i>_{OXA-23} and <i>bla</i>_{OXA-24} 	–	–	<ul style="list-style-type: none"> <i>bla</i>_{OXA-51} <i>bla</i>_{OXA-23} <i>bla</i>_{OXA-24} <i>bla</i>_{OXA-58}

(Continued)

Table 3 (Continued)

Author (year)	Setting	Population	Samples	Microorganism (no.)	MIC	Method of carbapenemase detection	Resistance to FQ and AG	Results ^b	Weak points	Strength points	Founded carbapenemase genes ^a
Shoja et al (2013) ⁵⁴	Two hospitals in Ahvaz	ICU patients	Trachea	<i>A. baumannii</i> (206)	–	<ul style="list-style-type: none"> • DD • PCR 	GEN 83%, TOB 78%, AMK 88%, CIP 96%	<ul style="list-style-type: none"> • Resistance rate to IPM and MEM was 96% • <i>bla</i>_{OXA-23} and <i>bla</i>_{OXA-24} was found in 85% and 8% of all strains, respectively. • 62% of isolates were resistant to carbapenems • Out of 62 resistant isolates, 55 carried <i>bla</i>_{OXA-23}, 1 carried <i>bla</i>_{OXA-40} and 2 isolates had <i>bla</i>_{OXA-38} genes • 56% of isolates were resistant to IPM, although 13% intermediate resistance should be considered • Among resistant isolates, 61 and 33 isolates had <i>bla</i>_{VIM} and <i>bla</i>_{IPM} genes, 23 strains had 2 resistant genes 	–	–	<i>bla</i> _{OXA-23} <i>bla</i> _{OXA-24}
Sohrabi et al (2012) ⁵⁵	One hospital in Tabriz	General inpatients	Different sites	<i>A. baumannii</i> (100)		<ul style="list-style-type: none"> • DD • PCR 	CIP and GEN 86% LVX 84% AMK 81%	<ul style="list-style-type: none"> • 62% of isolates were resistant to carbapenems • Out of 62 resistant isolates, 55 carried <i>bla</i>_{OXA-23}, 1 carried <i>bla</i>_{OXA-40} and 2 isolates had <i>bla</i>_{OXA-38} genes • 56% of isolates were resistant to IPM, although 13% intermediate resistance should be considered • Among resistant isolates, 61 and 33 isolates had <i>bla</i>_{VIM} and <i>bla</i>_{IPM} genes, 23 strains had 2 resistant genes 	–	–	<i>bla</i> _{OXA-23} <i>bla</i> _{OXA-38} <i>bla</i> _{OXA-40}
Sepehriseresht et al (2012) ⁵⁶	One hospital in Tehran (Motahari)	Burn patients	Burn wounds	<i>P. aeruginosa</i> (483)	–	<ul style="list-style-type: none"> • DD • PCR 	–	<ul style="list-style-type: none"> • 56% of isolates were resistant to IPM, although 13% intermediate resistance should be considered • Among resistant isolates, 61 and 33 isolates had <i>bla</i>_{VIM} and <i>bla</i>_{IPM} genes, 23 strains had 2 resistant genes 	<ul style="list-style-type: none"> • Origin of <i>bla</i>_{IPM} is not clear 	–	<i>bla</i> _{VIM} <i>bla</i> _{IPM}

(Continued)

Table 3 (Continued)

Author (year)	Setting	Population	Samples	Microorganism (no.)	MIC	Method of carbapenemase detection	Resistance to FQ and AG	Results ^b	Weak points	Strength points	Founded carbapenemase genes ^a
Forozesh Fard et al (2012) ⁵⁷	One hospital in Isfahan	CF patients	Sputum	<i>P. aeruginosa</i> (11)	–	<ul style="list-style-type: none"> • DD • PCR 	CIP 0% TOB 45.4%	<ul style="list-style-type: none"> • None of the isolates was resistant to IPM. None was positive for <i>bla_{VIM}</i> gene 	<ul style="list-style-type: none"> • Very small sample size • Few antibiotics were tested 	–	–
Peymani et al (2012) ⁵⁸	One hospital in Tabriz	General inpatients	Different sites	<i>A. baumannii</i> (68)	–	<ul style="list-style-type: none"> • PCR • Clonal typing 	–	<ul style="list-style-type: none"> • All IPM-resistant isolates were chosen for study • 44% of isolates were from tracheal samples • All were positive for <i>bla_{OXA-23}</i> and <i>IS_{Aba1}</i> was present upstream of all <i>bla_{OXA-23}</i> genes 	–	–	<i>bla_{OXA-23}</i>
Asadollahi et al (2012) ⁵⁹	One teaching hospital in Tehran	ICU burn patients	Burn wounds	<i>A. baumannii</i> (23)	✓	<ul style="list-style-type: none"> • DD • PCR 	CIP 100% GEN 30% AMK 47%	<ul style="list-style-type: none"> • Resistance rate to IPM was 48% • Range of MIC values for IPM was 0.004–32 mcg/mL • From all strains, <i>bla_{PER}</i> and <i>bla_{TEM}</i> genes were present in 52% and 43% of isolates, respectively. 21% and 17% of isolates had <i>bla_{OXA}</i> and <i>bla_{OXA24}</i> genes, respectively 	Small sample size	<ul style="list-style-type: none"> • MIC values were reported for each antibiotic • <i>CarO</i> and <i>tet</i> genes were assessed 	<i>bla_{TEM}</i> <i>bla_{PER}</i> <i>bla_{OXA-23}</i> <i>bla_{OXA-24}</i> <i>bla_{SHV}</i>

(Continued)

Table 3 (Continued)

Author (year)	Setting	Population	Samples	Microorganism (no.)	MIC	Method of carbapenemase detection	Resistance to FQ and AG	Results ^b	Weak points	Strength points	Founded carbapenemase genes ^a
Taherikalani et al (2009) ⁴²	Six hospitals in Tehran	General inpatients	Different sites	<i>A. baumannii</i> (80)	✓	<ul style="list-style-type: none"> • DD • PCR • Clonal typing 	–	<ul style="list-style-type: none"> • 52% of isolates were carbapenem resistant • MIC₅₀ range for carbapenem was 0.12–256 mcg/mL • Seven isolates that had MIC =256 mcg/mL had more than two <i>bla</i>_{OXA} genes • From all strains <i>bla</i>_{OXA-23} in 25%, <i>bla</i>_{OXA-58} in 21%, and <i>bla</i>_{OXA-24} in 15% genes were detected • Coexistence of multiple genes was observed • About 50% of strains were resistant to carbapenems • MIC values for resistant species were ≥64 to ≥256 mcg/mL • For details of isolated genes, refer to original article 	<ul style="list-style-type: none"> • Number of tested antibiotics were few 	<ul style="list-style-type: none"> • Multicenter study 	<i>bla</i> _{OXA-23} <i>bla</i> _{OXA-24} <i>bla</i> _{OXA-58}
Feizabadi et al (2008) ⁶⁰	One hospital in Tehran	General inpatients	Different sites	<i>Acinetobacter</i> spp. (<i>A. baumannii</i> =108 and other <i>Acinetobacter</i> =20, total =128)	✓	<ul style="list-style-type: none"> • DD • PCR 	CIP 81% LVX 74% AMK 61% NTL 86% TOB 79% GEN 81%				<i>bla</i> _{OXA-23} <i>bla</i> _{OXA-24} <i>bla</i> _{OXA-51} <i>bla</i> _{OXA-58}

(Continued)

Table 3 (Continued)

Author (year)	Setting	Population	Samples	Microorganism (no.)	MIC	Method of carbapenemase detection	Resistance to FQ and AG	Results ^b	Weak points	Strength points	Founded carbapenemase genes ^a
Taherikalani et al (2008) ⁶¹	Two hospitals in Tehran	Burn patients	Burn wounds	<i>A. baumannii</i> (38)	–	<ul style="list-style-type: none"> DD PCR 	Resistance AMK 71% TOB 90% NTL 90% LVX 81% CIP 85%	<ul style="list-style-type: none"> Resistance rate to carbapenem was about 60% Half of the carbapenem-resistant strains had at least two <i>bla</i>_{OXA} genes 	<ul style="list-style-type: none"> Small sample size MIC values were not determined 		<i>bla</i> _{OXA-23} <i>bla</i> _{OXA-24} <i>bla</i> _{OXA-51} <i>bla</i> _{OXA-58}

Notes: ^aResults about *bla*_{OXA-51} were not provided because this gene has no correlation with occurrence of resistance, unless it has *IS*_{Ab} genes in its upstream. ^bResistance rates were according to the results of disk diffusion test. For complete information about the resistance to AG and FQ, please refer to the original reference.

Abbreviations: AG, aminoglycoside; AMK, amikacin; CIP, ciprofloxacin; DD, disk diffusion; FQ, fluoroquinolones; GEN, gentamicin; ICU, intensive care unit; IMP, imipenem; KAN, kanamycin; LVX, levofloxacin; MEM, meropenem; MIC, minimum inhibitory concentration; NAL, nalidixic acid; PCR, polymerase chain reaction; STP, streptomycin; TOB, tobramycin; NTL, netilmicin.

ity.¹¹⁰ In CLSI 2018, CarbaNP method is recommended instead. Kuchibiro et al compared some phenotypic methods and MHT and showed an acceptable specificity (100%), but the sensitivity was very low (50%). Another test, modified carbapenems inactivation method, has acceptable sensitivity and specificity values (both above 99%) and does not require specific equipment. This method, beside CarbaNP, can be recommended for phenotypic detection of carbapenemase.¹¹¹

Findings according to the phenotypic and genotypic methods were inconsistent. For example, in the study by Bina et al,⁶³ although all strains were MHT positive, none of them carry *bla*_{KPC} gene. This may be due to low sensitivity of MHT in detecting carbapenemase enzyme.

The genotypic assays were mostly applied on central regions of Iran. However, there are no data regarding carbapenemase-encoding genes in most provinces. This may be due to technical and economic restrictions in these areas (Figure 4).

The presence of some *bla*_{OXA-type} genes were not necessarily associated to carbapenem-resistant. For example, *bla*_{OXA-51} in *A. baumannii* did not contribute to carbapenem resistance at all. In the study by Azizi et al, it was shown that *bla*_{OXA-23} may be found in both imipenem-sensitive and imipenem-resistant isolates.⁴³ Also, in the study by Bahador et al, *bla*_{OXA-23} was found in both susceptible and nonsusceptible strains.⁴⁴ However, higher carbapenem resistance in isolates harboring *bla*_{OXA-23} was reported (Shoja et al⁵⁴). Table 5 shows the number of genes that were found in each city and related microorganisms.

Emergence of some genes including *bla*_{NDM-1} is a serious global warning. This gene is highly transmissible and was first discovered in a Swedish patient who had traveled to Pakistan. Although almost all identified cases were originally from Pakistan and India, the first case of *bla*_{NDM-1} that was reported in Iran in the year 2012, did not have a history of traveling to these countries. In two studies from Shiraz and Isfahan, 27 and 6 strains of *K. pneumoniae* harbored *bla*_{NDM} in years 2015 and 2017, respectively.^{62,78}

Genotypic methods to identify other mechanisms of resistance to carbapenems have recently been considered by researchers. *OmpK* and *carO* genes that regulate expression of mutated efflux pump were detected in Hashemi et al and Pajand et al studies, respectively.^{83,93} In another study, efflux pump's activity and mutations in porins were investigated.¹¹² These results can expand the view about true mechanisms of resistance to carbapenems in developing countries.

Table 4 Details of studies that used both phenotypic and genotypic methods for detection of carbapenemase-producing strains

Author (year)	Setting	Population (no.)	Samples	Microorganism	MIC ^a	Method of carbapenemase enzyme detection	Resistance to FQ and AG	Outcome	Weak points	Strength points	bla-type genes
Solgi et al (2017) ⁶⁶	Two hospitals (city was not defined)	General inpatients (mostly ICU)	Rectal swab	Enterobacteriaceae spp. (95)	✓	<ul style="list-style-type: none"> MHT for MBL detection Clonal typing 	AMK 24% GEN 35% CIP 74%	<ul style="list-style-type: none"> 36 resistant strains were detected according to the phenotypic methods that among them carbapenem-resistance rates were between 88% and 100% 47 isolates were MHT positive 7 isolates carried bla_{NDM-1}, 23 carried bla_{OXA-48} 17 isolates harbored two genes 	<ul style="list-style-type: none"> MIC values were determined, but not analyzed Small sample size 	<ul style="list-style-type: none"> Risk factors for infection with resistant strains were investigated 	bla _{NDM-1} bla _{NDM-7} bla _{OXA-48}
Khorvash et al (2017) ⁶⁷	One hospital in Isfahan	ICU patients	Different sites	<i>Pseudomonas aeruginosa</i> (48)		<ul style="list-style-type: none"> MHT PCR 	AMK 79% CIP 85%	<ul style="list-style-type: none"> Resistance rate to carbapenem was 97% bla_{IMP} and bla_{VIM} were detected in 15 and 7 isolates (all were MHT positive) In one isolate of <i>P. aeruginosa</i> with the source of urine, both bla_{IMP} and bla_{VIM} genes were detected 52% of urine samples contained bla_{IMP} 	–	<ul style="list-style-type: none"> Ratio of MDR pathogens for each sample was defined 	bla _{IMP} bla _{VIM}

(Continued)

Table 4 (Continued)

Author (year)	Setting	Population (no.)	Samples	Microorganism	MIC ^a	Method of carbapenemase enzyme detection	Resistance to FQ and AG	Outcome	Weak points	Strength points	bla-type genes ^c
Hosseinzadeh et al (2017) ⁶²	Two hospitals in Shiraz	General inpatients and ICU patients	Different sites	<i>Klebsiella pneumoniae</i> (211)	✓	<ul style="list-style-type: none"> • MHT for MBL detection • DDST for MBL detection • PCR 	AMK 78% CIP 85% GEN 89%	<ul style="list-style-type: none"> • 13% of isolates (29 ones) were resistant to IPM and MEM and had an MIC value of 1.5–32 mcg/mL (25 isolates had MIC $\geq$ 4 mcg/mL) • Among carbapenem-resistant isolates, DDST and MHT were positive for 27 strains • 2 and 27 isolates had <i>bla</i>_{OXA-48} and <i>bla</i>_{NDM-1} respectively 		<ul style="list-style-type: none"> • Large sample size 	<i>bla</i> _{OXA-48} <i>bla</i> _{NDM-1}
Akhi et al (2017) ⁶⁸	Seven hospitals in Tabriz and Orumie	General inpatients	Different sites	<i>P. aeruginosa</i> (245)	–	<ul style="list-style-type: none"> • DD • MHT • MCNP test • CIM • PCR 	CIP 66% LVX 66% AMK 25%	<ul style="list-style-type: none"> • 49% (121) of isolates were carbapenem resistant • From 121 resistant strains, <i>bla</i>_{IMP} and <i>bla</i>_{VIM} genes were positive in 29 and 6 strains, respectively, and 40, 39, and 35 isolates showed positive results for MHT, MCNP, and CIM tests, respectively 	<ul style="list-style-type: none"> • MIC was not tested 	<ul style="list-style-type: none"> • Large sample size • Sensitivity and specificity of three phenotypic methods were estimated according to PCR • Multicenter study 	<i>bla</i> _{VIM} <i>bla</i> _{IMP}

(Continued)

Table 4 (Continued)

Author (year)	Setting	Population (no.)	Samples	Microorganism	MIC ^a	Method of carbapenemase enzyme detection	Resistance to FQ and AG	Outcome	Weak points	Strength points	bla-type genes
Falahat et al (2016) ⁶⁹	Three hospitals in Arak	General inpatients	Different sites	<i>P. aeruginosa</i> (108)	–	<ul style="list-style-type: none"> MHT and CDT for KPC detection PCR 	CIP 20% AMK 21% GEN 23%	<ul style="list-style-type: none"> Resistance rate to carbapenem was about 25% 38 and 26 of all isolates were MHT and CDT positive respectively 13 isolates had <i>bla</i>_{KPC} gene 	–	<ul style="list-style-type: none"> Multicenter study 	<i>bla</i> _{KPC}
Mohammadzadeh et al (2016) ⁷⁰	Two hospitals in Tehran	General inpatients	Not defined (probably different sites)	Different strains (864)	–	<ul style="list-style-type: none"> CDT, MHT, and DDST for MBL detection PCR 	Not defined	<ul style="list-style-type: none"> 7% of isolates were IPM resistant 35, 24, and 3 isolates were <i>Acinetobacter spp.</i>, <i>Pseudomonas spp.</i>, and <i>Enterobacteriaceae</i>, respectively Carbapenem resistance was confirmed phenotypically (17% and 9%) and genotypically (15% and 9%) among <i>Acinetobacter</i> and <i>Pseudomonas</i>, respectively 	–	<ul style="list-style-type: none"> Large sample size 	<i>bla</i> _{MHT} <i>bla</i> _{IMP}

(Continued)

Table 4 (Continued)

Author (year)	Setting	Population (no.)	Samples	Microorganism	MIC ^a	Method of carbapenemase enzyme detection	Resistance to FQ and AG	Outcome	Weak points	Strength points	bla-type genes
Tarashi et al (2016) ⁷¹	One hospital in Tehran	Burn patients	Burn wounds	<i>Acinetobacter baumannii</i> (189) <i>P. aeruginosa</i> (309)	–	<ul style="list-style-type: none"> • CDT for MBL • PCR 	<p>In 2015 for <i>P. aeruginosa</i>: AMK and GEN 95% CIP 97%</p> <p>For <i>A. baumannii</i>: AMK and CIP 100% GEN 95%</p>	<ul style="list-style-type: none"> • Carbapenem resistance rate of <i>P. aeruginosa</i> was 83% in 2012 and 96% in 2015 • Carbapenem resistance rate of <i>A. baumannii</i> in all years was 100% • Among 278 resistant <i>P. aeruginosa</i> isolates, 178 strains were MBL producers • Of the 187 resistant <i>A. baumannii</i>, 85 isolates were MBL producers • <i>bla</i>_{IMP-1} and <i>bla</i>_{IMP-1} genes were isolated in 30 and 52 of resistant strains of <i>P. aeruginosa</i>, respectively • <i>bla</i>_{IMP-1} and <i>bla</i>_{IMP-1} genes were detected in 10 and 34 of resistant <i>A. baumannii</i>, respectively 	–	<ul style="list-style-type: none"> • Large sample size • Providing trend of resistance through different years 	<i>bla</i> _{OXA-51} <i>bla</i> _{IMP} <i>bla</i> _{IMP}

(Continued)

Table 4 (Continued)

Author (year)	Setting	Population (no.)	Samples	Microorganism	MIC ^a	Method of carbapenemase enzyme detection	Resistance to FQ and AG	Outcome	Weak points	Strength points	bla-type genes
Maspi et al (2016) ⁷²	One hospital in Tehran	General inpatients and ICU patients	Different sites (mostly BAL and wounds)	<i>A. baumannii</i> (86)	–	<ul style="list-style-type: none"> CDT for MBL detection PCR 	TOB 75%	<ul style="list-style-type: none"> Resistance rates to MEM and IMP were 90% and 73%, respectively 44 isolates were MBL producers 2, 13, 2, 4, and 2 isolates carried <i>bla_{VIM}</i>, <i>bla_{IMP}</i>, <i>bla_{SPM}</i>, <i>bla_{GHM}</i> and <i>bla_{SHM}</i> genes, respectively 	<ul style="list-style-type: none"> MIC was not reported 	<ul style="list-style-type: none"> First report of <i>bla_{SHM}</i>, <i>bla_{GHM}</i> and <i>bla_{SPM}</i> genes 	<i>bla_{VIM}</i> <i>bla_{IMP}</i> <i>bla_{SPM}</i> <i>bla_{GHM}</i> <i>bla_{SHM}</i>
Moghadam et al (2016) ⁷³	Two hospitals in Shiraz	General inpatients and ICU patients	Different sites	<i>A. baumannii</i> (98)	✓	<ul style="list-style-type: none"> E-test for MBL detection PCR 	AMK 69% GEN 75% CIP and LVX 100%	<ul style="list-style-type: none"> Resistance to carbapenem was about 95% and according to MIC, it was 98% Except two, all resistant isolates according to MIC had breakpoints >16 mcg/mL Out of resistant isolates, 43 strains were MBL producers Out of MBL producer strains, 14 had <i>bla_{VIM}</i> and 23 had <i>bla_{IMP}</i> genes 	–	<ul style="list-style-type: none"> Results of general and ICU patients were discussed separately 	<i>bla_{VIM}</i> <i>bla_{IMP}</i>

(Continued)

Table 4 (Continued)

Author (year)	Setting	Population (no.)	Samples	Microorganism	MIC ^a	Method of carbapenemase enzyme detection	Resistance to FQ and AG	Outcome	Weak points	Strength points	bla-type genes
Pakbaten Toupkanlou et al (2015) ⁷⁴	One hospital in Tehran	Burn patients	Not defined	<i>P. aeruginosa</i> (50)		<ul style="list-style-type: none"> • CDT for ESBL detection • PCR 	CIP 98%, AMK, GEN, and TOB 96%	<ul style="list-style-type: none"> • All resistant strains to IPM were chosen for study • 23 and 17 strains were positive in two different phenotypic methods for ESBL detection • 7, 18, 18, and 18 strains had <i>bla</i>_{PER}[*], <i>bla</i>_{OXA-10}[*], <i>bla</i>_{TEM}[*] and <i>bla</i>_{SHV} genes, respectively • 10 (20%) isolates carried bla-TEM, blaOXA-10, and blaSHV genes, simultaneously 	<ul style="list-style-type: none"> • Small sample size • Origins of samples were not defined 	–	<i>bla</i> _{OXA-10} <i>bla</i> _{SHV} <i>bla</i> _{TEM} <i>bla</i> _{PER}
Bina et al (2015) ⁶³	Three hospitals in Tehran	General inpatients	Different sites	<i>K. pneumonia</i> (270)	–	<ul style="list-style-type: none"> • MHT test for KPC detection • PCR 	GEN 41.3%	<ul style="list-style-type: none"> • Resistance rate to carbapenem was 14% (41 strains) • –14.5% (33 out of 41) strains were MHT positive and all 41 strains did not carry bla-KPC gene 	<ul style="list-style-type: none"> • Few antibiotics were tested • Large sample size • Multicenter study 	–	–

(Continued)

Table 4 (Continued)

Author (year)	Setting	Population (no.)	Samples	Microorganism	MIC ^a	Method of carbapenemase enzyme detection	Resistance to FQ and AG	Outcome	Weak points	Strength points	bla-type genes
Eftekhari and Naseh (2015) ⁷⁵	Two hospitals in Tehran	Thirty nonburn and 25 burn patients	Different sites	<i>K. pneumonia</i> (30 nonburn and 25 burn, total 55)	–	<ul style="list-style-type: none"> • CDT for ESBL detection • DDST for MBL screening • MHT for MBL detection • PCR 	Burn and nonburn patients respectively: CIP 84%, 60% AMK 52%, 33.3% GEN 84%, 56.7%	Resistance rates to carbapenem were 20% in burn and 0% in nonburn patients <ul style="list-style-type: none"> • ESBL production was positive in 83% of nonburn and 72% of burn isolates • None was an MBL producer and positive for KPC genes • MHT was positive in four strains • Carbapenem-resistant isolates (n=5) from burn wounds were resistant to all tested antibiotics 	<ul style="list-style-type: none"> • Small sample size 	<ul style="list-style-type: none"> • Comparison of resistance pattern in burn and nonburn patients 	–
Lari et al (2015) ⁶⁴	One hospital in Tehran	Burn patients	Burn wounds	<i>P. aeruginosa</i> (255)	–	<ul style="list-style-type: none"> • CDT for ESBL production • DDST for MBL production • PCR 	Were tested but not reported specifically	<ul style="list-style-type: none"> • All IPM-1-resistant strains were chosen for study • 63% were positive in CDT • None were positive for DDST • 5 and 4 strains had bla_{IMP} and bla_{IMP} genes, respectively 	Resistance to other antibiotics was not defined clearly	<ul style="list-style-type: none"> • Large sample size 	bla _{IMP} bla _{IMP}

(Continued)

Table 4 (Continued)

Author (year)	Setting	Population (no.)	Samples	Microorganism	MIC ^a	Method of carbapenemase enzyme detection	Resistance to FQ and AG	Outcome	Weak points	Strength points	bla-type genes
Bagheri Josheghani et al (2015) ⁶⁵	One hospital in Kashan	General inpatients	Different sites	<i>A. baumannii</i> (124)	–	<ul style="list-style-type: none"> • CDT for ESBL detection • DDST for MBL detection • PCR 	AMK 78% GEN 83% LVX and CIP 99%	Resistance rates to carbapenems was about 90% <ul style="list-style-type: none"> • 5% were ESBL positive and 54% isolates were MBL • 79% , 25%, and 3% carried <i>bla</i>_{OXA-23}, <i>bla</i>_{OXA-24} and <i>bla</i>_{OXA-58} genes, respectively 	–	–	<i>bla</i> _{OXA-51} <i>bla</i> _{OXA-23} <i>bla</i> _{OXA-24} <i>bla</i> _{OXA-58}
Azimi et al (2016) ⁷⁶	One hospital in Tehran	Burn patients	Burn wounds	Different strains (161)	–	<ul style="list-style-type: none"> • MHT and DDST for MBL detection • CDT for KPC detection • PCR 	–	All IPM-resistant strains were included <ul style="list-style-type: none"> • 85, 51, and 112 of them were MHT, boronic acid, and dipicolinic positive, respectively 	<ul style="list-style-type: none"> • Exact number of resistance genes was not reported 	<ul style="list-style-type: none"> • The sensitivity and specificity of each phenotypic test were reported 	<i>bla</i> _{IMP} <i>bla</i> _{OXA-23} <i>bla</i> _{OXA-48}
Azimi et al (2015) ⁷⁷	One hospital in Tehran	Burn patients	Burn wounds	<i>A. baumannii</i> (65)	–	<ul style="list-style-type: none"> • CDT for ESBL detection • PCR 	80% of strains were resistant to all AG	All IPM-resistant strains were selected for study <ul style="list-style-type: none"> • 1 strain was CDT positive • 83%, 12%, and 9% of all strains harbored <i>bla</i>_{OXA-23}, <i>bla</i>_{IMP} and <i>bla</i>_{KPC} genes, respectively. 	–	–	<i>bla</i> _{OXA-23} <i>bla</i> _{OXA-51} <i>bla</i> _{IMP} <i>bla</i> _{KPC}

(Continued)

Table 4 (Continued)

Author (year)	Setting	Population (no.)	Samples	Microorganism	MIC ^a	Method of carbapenemase enzyme detection	Resistance to FQ and AG	Outcome	Weak points	Strength points	bla-type genes
Fazeli et al (2015) ⁷⁸	One hospital in Isfahan	General inpatients	Different sites	<i>K. pneumonia</i> (112)	✓	<ul style="list-style-type: none"> MHT and E-test for MBL detection PCR 	CIP 63.4% AMK 51% GEN 64%	42% of strains (49 isolates) were resistant to IPM and MEM <ul style="list-style-type: none"> 32 out of 49 resistant isolates were positive for MHT 5 isolates were positive in E-test and 6 strains harbored NDM-1 MIC values for these six strains was > 256 mcg/mL 			<i>bla</i> _{NDM}
Khoshvagt et al (2014) ⁷⁹	Four hospitals in Zanjan	General pediatrics	Stool	<i>Escherichia coli</i> (230)	–	<ul style="list-style-type: none"> DDST for ESBL and MBL detection PCR 	CIP 37% AMK 21% GEN 29%	IPM resistance rate was 2.1% <ul style="list-style-type: none"> Out of the 36 Enterocoaggregative <i>E. coli</i> isolates, 19 (52%) were ESBL positive and none were MBL positive 15 and 12 isolates had <i>bla</i>_{TEM} and <i>bla</i>_{CTX-M} genes 	<ul style="list-style-type: none"> MIC was not tested 	<ul style="list-style-type: none"> Large sample size Comparing data of healthy children with those with diarrhea 	<i>bla</i> _{TEM} and <i>bla</i> _{CTX-M}

(Continued)

Table 4 (Continued)

Author (year)	Setting	Population (no.)	Samples	Microorganism	MIC ^a	Method of carbapenemase enzyme detection	Resistance to FQ and AG	Outcome	Weak points	Strength points	bla-type genes
Noori et al (2014) ⁸⁰	Two hospitals in Tehran	General inpatients	Different sites	<i>Acinetobacter</i> spp. (108)	✓	<ul style="list-style-type: none"> • DD • CDT for MBL 	CIP 92% AMK 80% GEN 40%	<ul style="list-style-type: none"> • 91% (99 isolates) were resistant to IPM and MEM • 86 isolates were MBL producers • Out of these 86 MBL producers, three isolates harbored <i>bla_{IMP}</i> • MIC50 values for MEM and IPM were 32 and 128 mcg/mL, respectively 	–	–	<i>bla_{IMP}</i>
Vali et al (2014) ⁸¹	Two pediatric hospitals in Tehran	Cystic fibrosis inpatients	Sputum	Different gram negative (52)	✓	<ul style="list-style-type: none"> • CDT for ESBL detection • DDST for MBL detection • PCR 	GEN 24.5% CIP 18%	<ul style="list-style-type: none"> • Resistance rate to MEM was 14% (eight strains) that MHT was positive in three of them • The EDTA disk synergy was also positive in four isolates • <i>bla_{VIM}</i> and <i>bla_{IMP}</i> were detected in five and two strains of <i>Achromobacter xylosoxidans</i> 	–	<ul style="list-style-type: none"> • Data provided from cystic fibrosis patients 	<i>bla_{CTX-M}</i> <i>bla_{IMP}</i> <i>bla_{VIM}</i>

(Continued)

Table 4 (Continued)

Author (year)	Setting	Population (no.)	Samples	Microorganism	MIC ^a	Method of carbapenemase enzyme detection	Resistance to FQ and AG	Outcome	Weak points	Strength points	bla-type genes
Azimi et al (2014) ⁸²	One hospital in Tehran	Burn patients	Burn wounds	<i>K. pneumonia</i> (28)	✓	<ul style="list-style-type: none"> MHT CarbaNP test BA PCR 	GEN 90% AMK 79%	All isolates were resistant to IPM and MEM <ul style="list-style-type: none"> 9 and 11 strains had an MIC >64 mcg/mL for IPM and MEM, respectively All phenotypic tests except BA were positive 27 isolates had <i>bla</i>_{OXA-48} gene and one had <i>bla</i>_{IMP-4} Resistance rates to IPM, MEM, and DOR were between 68% and 74% MIC range for IPM and MEM was 0.25–256 mcg/mL 48 (57%) isolates were positive for ESBL 5 isolates were MHT positive 23 isolates were AmpC positive 30 strains had <i>OmpK35</i> genes 	<ul style="list-style-type: none"> Small sample size Few antibiotics were tested 	<ul style="list-style-type: none"> CarbaNP test was done 	<i>bla</i> _{OXA-48} <i>bla</i> _{IMP-4}
Hashemi et al (2014) ⁸³	Two hospitals in Tehran	General inpatients and infants	Different sites	<i>K. pneumonia</i> (83)	✓	<ul style="list-style-type: none"> CDT AmpC screening MHT PCR 	Resistance and intermediate: CIP 39.7%, 4.8% GEN 61.5%, 3.6% AMK 80.7%, 4.8%	Resistance rates to IPM, MEM, and DOR were between 68% and 74% MIC range for IPM and MEM was 0.25–256 mcg/mL 48 (57%) isolates were positive for ESBL 5 isolates were MHT positive 23 isolates were AmpC positive 30 strains had <i>OmpK35</i> genes	–	<ul style="list-style-type: none"> Detection of porin-related genes 	<i>bla</i> _{OXA-48} <i>bla</i> _{CTX-M-15}

(Continued)

Table 4 (Continued)

Author (year)	Setting	Population (no.)	Samples	Microorganism	MIC ^a	Method of carbapenemase enzyme detection	Resistance to FQ and AG	Outcome	Weak points	Strength points	bla-type genes
Japoni-Nejad et al (2014) ⁸⁴	One hospital in Arak	ICU patients	Surgical wounds, urine, blood, respiratory secretions	<i>K. pneumonia</i> (100)	–	<ul style="list-style-type: none"> • MHT • E-test for MBL detection • DDST • CDT for KPC detection • PCR 	^d	<ul style="list-style-type: none"> • Resistance rate to carbapenems in Amp C producers and carbapenemase producers was 5% and 66%, respectively • <i>bla_{VIM}</i> and <i>bla_{GES}</i> were detected in ten and two isolates • One isolate (1%) contained both <i>bla_{VIM}</i> and <i>bla_{AmpC}</i> genes 	–	–	<i>bla_{VIM}</i> , <i>bla_{GES}</i>
Farajzadeh Sheikh et al (2014) ⁸⁵	Hospitals in Ahvaz	General inpatients	Different sites	<i>P. aeruginosa</i> (223)	✓	<ul style="list-style-type: none"> • CDT and E-test for MBL detection • PCR 	GEN 66% AMK 55% TOB 67% CIP 66%	<ul style="list-style-type: none"> • 58%, 31%, 13%, and 74% of isolates were resistant to IMP, MEM, DOR and ERT, respectively. Thirty isolates were resistant to all carbapenems • All imipenem-resistant isolates were MBL producers, except one • blaIMP and blaVIM genes were detected in 26 and 1 of the all strains, respectively • 13%–74% of resistant isolates had an MIC >16 mcg/mL for different carbapenems 	–	<ul style="list-style-type: none"> • Large sample size 	<i>bla_{VIM}</i> , <i>bla_{IMP}</i>

(Continued)

Table 4 (Continued)

Author (year)	Setting	Population (no.)	Samples	Microorganism	MIC ^a	Method of carbapenemase enzyme detection	Resistance to FQ and AG	Outcome	Weak points	Strength points	bla-type genes
Aghamiri et al (2014) ⁸⁶	Several hospitals in Tehran	General inpatients	Different sites	<i>P. aeruginosa</i> (212)	✓	<ul style="list-style-type: none"> • DD • DDST for MBL detection • PCR 		<p>47% of isolates were resistant to IPM and all had an MIC > 16 mcg/mL</p> <ul style="list-style-type: none"> • 70 isolates were MBL positive • 20 and 70 strains had <i>bla_{IMP}</i> and <i>bla_{NDH-1}</i> genes, respectively <p>Resistance rate to carbapenems was between 7% and 23% (66% of urine isolates were resistant to carbapenems)</p> <ul style="list-style-type: none"> • 25 isolates were MHT positive • <i>bla_{CTX-M}</i>, <i>bla_{SHV}</i>, <i>bla_{TEM}</i> and <i>bla_{PER}</i> genes were detected in 69%, 59.5%, 35.7%, and 16.6% of carbapenem-resistant isolates respectively • <i>bla_{NDH-1}</i>, <i>bla_{NDH-1}</i> and <i>bla_{KPC}</i> were detected in 3, 5, and 1 isolates 			<i>bla_{IMP}</i> <i>bla_{NDH-1}</i>
Nobari et al (2014) ⁸⁷	Eight hospitals in Tehran	General inpatients	Different sites	<i>K. pneumonia</i> (180)		<ul style="list-style-type: none"> • DD • MHT for MBL detection • PCR 					<i>bla_{CTX-M}</i> <i>bla_{SHV}</i> <i>bla_{TEM}</i> <i>bla_{PER}</i> <i>bla_{NDH-1}</i> <i>bla_{NDH-1}</i> <i>bla_{KPC}</i>

(Continued)

Table 4 (Continued)

Author (year)	Setting	Population (no.)	Samples	Microorganism	MIC ^a	Method of carbapenemase enzyme detection	Resistance to FQ and AG	Outcome	Weak points	Strength points	bla-type genes
Hakemi Vala et al (2014) ⁸⁸	One hospital in Tehran	Burn patients	Burn wounds	<i>P. aeruginosa</i> (47) and <i>Acinetobacter spp.</i> (28)	✓	<ul style="list-style-type: none"> • CDT for MBL and ESBL detection • PCR 	<i>P. aeruginosa</i> GEN 72% CIP 69% <i>A. baumannii</i> GEN 96% CIP 100%	<ul style="list-style-type: none"> • 100% of <i>A. baumannii</i> and about 75% of <i>P. aeruginosa</i> isolates were resistant to carbapenems • MIC range of IPM for <i>P. aeruginosa</i> was 2–128 and for <i>A. baumannii</i> was 4–128 mcg/mL • 17% of the <i>P. aeruginosa</i> isolates and 16% of the <i>A. baumannii</i> isolates were MBL producers • Mortality rate of MBL producer was 20% 	–	<ul style="list-style-type: none"> • Mortality rate was reported 	<i>bla</i> _{SPM} <i>bla</i> _{IMP} <i>bla</i> _{CTX-M.15}
Tavajjohi et al (2013) ⁸⁹	One hospital in Kashan	General inpatients and environmental isolates	Different strains and wet environment	<i>P. aeruginosa</i> (100)	–	<ul style="list-style-type: none"> • DDST for ESBL detection • PCR 	GEN 50% CIP 12%	<ul style="list-style-type: none"> • 8% of isolates were ESBL producers that <i>bla</i>_{GES} was detected in all ESBL producers and 50% (4 isolates) of them were resistance to IPM 	–	<ul style="list-style-type: none"> • Small sample size (only eight strains were evaluated for carbapenem resistance and genes) 	<i>bla</i> _{GES}

(Continued)

Table 4 (Continued)

Author (year)	Setting	Population (no.)	Samples	Microorganism	MIC ^a	Method of carbenemase enzyme detection	Resistance to FQ and AG	Outcome	Weak points	Strength points	bla-type genes
Azimi et al (2013) ⁹⁰	One hospital in Tehran (Motahari)	Burn patients	Burn wounds	<i>K. pneumonia</i> (44)		<ul style="list-style-type: none"> • DD • MHT for KPC detection • BA 	–	<ul style="list-style-type: none"> • 64% of isolates were resistant to carbenemase • MHT was positive in all of them, but none showed synergism between MEM and boronic acid • All were negative for <i>bla_{KPC}</i> • All isolates were resistant to IPM and MEM • 50 out of 70 strains were MBL producers, but only 3 strains had <i>bla_{IMP}</i> 	<ul style="list-style-type: none"> • Other antibiotics were not evaluated • Small sample size 	–	
Noori et al (2013) ⁹¹	Two hospitals in Mashhad	General inpatients	Wounds and respiratory secretions	<i>Acinetobacter</i> spp. (70)		<ul style="list-style-type: none"> • DDST for MBL production • PCR 	CIP 100% GEN 88%	<ul style="list-style-type: none"> • 85% of isolates were resistant to IPM that MIC range for them was 16–128 mcg/mL • 34% of all isolates were MBL positive (31 strains) • 25 out of MBL producers had <i>bla_{OXA-23}</i> • None was positive in DDST 			<i>bla_{IMP}</i>
Azimi et al (2013) ⁹²	One hospital in Tehran	Burn patients	Not defined (probably burn wounds)	<i>A. baumannii</i> (93) and <i>Acinetobacter lwoffii</i> (1)	✓	<ul style="list-style-type: none"> • CDT for MBL detection • PCR 	–				<i>Bla_{OXA-23}</i>

(Continued)

Table 4 (Continued)

Author (year)	Setting	Population (no.)	Samples	Microorganism	MIC ^a	Method of carbapenemase enzyme detection	Resistance to FQ and AG	Outcome	Weak points	Strength points	bla-type genes
Pajand et al (2013) ⁹³	Two hospitals in Tehran	Inpatients and burn patients	Burn wounds and other sites	<i>A. baumannii</i> (43 burn and 32 nonburn patients, total=75)	✓	<ul style="list-style-type: none"> • DDST for MBL detection • PCR 	Nonburn and burn patients: CIP 87%, 97% AMK 56%, 95% TOB 47%, 84%	Resistance rate to carbapenem in burn and nonburn patients was 98% and 68%, respectively <ul style="list-style-type: none"> • 66% of nonburn and 70% of burn isolates had <i>bla</i>_{OXA-23} • <i>bla</i>_{OXA-40}-like gene was detected in 12% and 74% of non-burn and burn isolates, respectively • MIC values for all carbapenems were reported • All isolates were positive for AmpC and <i>ISAb</i> genes and 71 of them were positive for <i>carO</i> gene 	<ul style="list-style-type: none"> • Results of DDST were not reported 	<ul style="list-style-type: none"> • Detection of OMP gene (<i>carO</i>) for evaluation of carbapenem resistance 	<i>Bla</i> _{OXA-23} <i>bla</i> _{OXA-40} <i>AmpC</i> , <i>ISAb</i> , <i>carO</i>
Azami et al (2013) ⁹⁴	Three hospitals in Tehran	General inpatients	Not defined	<i>P. aeruginosa</i> (130)	✓	<ul style="list-style-type: none"> • CDT for MBL detection • PCR 	GEN 63% CIP 59%	<ul style="list-style-type: none"> • According to MIC, 53% of isolates were resistant to IPM and MIC values were ≥16 mcg/mL for IPM • 56% had class I integrons • <i>bla</i>_{VIM} was found in 10 strains 	<ul style="list-style-type: none"> • Few antibiotics were tested • Source of sample was not clear 	<ul style="list-style-type: none"> • Assessment of integrin genes (genes for mobility of betalactamase) • Multicenter study 	<i>bla</i> _{VIM}

(Continued)

Table 4 (Continued)

Author (year)	Setting	Population (no.)	Samples	Microorganism	MIC ^a	Method of carbapenemase enzyme detection	Resistance to FQ and AG	Outcome	Weak points	Strength points	bla-type genes
Mohajeri et al (2013) ⁹⁵	Three hospitals in Kermanshah	General inpatients	Sputum, blood, urine	<i>Acinetobacter</i> spp. (104)	–	<ul style="list-style-type: none"> E-test for MBL detection DDST PCR 	CIP 69%, GAT 43% LVX 62% TOB 39% GEN 68% AMK 53%	Resistance rate to carbapenems was 75%–80% <ul style="list-style-type: none"> 84 of isolates were MBL producers The <i>bla</i>_{OXA-23} and <i>bla</i>_{OXA-24} were found among 77% and 19% of the isolates, respectively Correlation between rate of resistance to carbapenem and presence of <i>bla</i>_{OXA-23}-like gene was statistically significant 	<ul style="list-style-type: none"> MIC was not reported 	<ul style="list-style-type: none"> Multicenter study 	<i>bla</i> _{OXA-23} <i>bla</i> _{OXA-24}
Doosti et al (2013) ⁹⁶	One hospital in Zanjan	General inpatients	Different sites	<i>P. aeruginosa</i> (70)	✓	<ul style="list-style-type: none"> DD DDST for MBL detection PCR 	CIP 40% GEN 55%	63% of isolates were resistant to IPM that 41 of them had an MIC >4 mcg/mL <ul style="list-style-type: none"> DDST was positive for all of them 36 isolates were MBL producers From 41 of isolates with high MIC breakpoints, 10 isolates had <i>bla</i>_{IMP} and 23 had <i>bla</i>_{IMP} 	–	–	<i>bla</i> _{IMP} <i>bla</i> _{IMP}

(Continued)

Table 4 (Continued)

Author (year)	Setting	Population (no.)	Samples	Microorganism	MIC ^a	Method of carbapenemase enzyme detection	Resistance to FQ and AG	Outcome	Weak points	Strength points	bla-type genesc
Shahcheraghi et al (2013) ⁹⁷	Five hospitals in Tehran	General inpatients	Mostly urine and feces	Different strains (360)	–	<ul style="list-style-type: none">• MHT• MBL E-test• PCR	CIP 61.3% AMK 14.4% KAN 38.6%	<ul style="list-style-type: none">• 6%, 3%, and 1% were resistant to MEM, ERT, and IPM, respectively• MHT was positive in 11 resistant isolates	<ul style="list-style-type: none">• MIC was not reported	<ul style="list-style-type: none">• Multicenter study• Large sample size• Recognition of new <i>K. pneumonia</i> containing bla NDM	<i>bla</i> _{TEM} <i>bla</i> _{SHV} <i>bla</i> _{CTX-M} <i>bla</i> _{NDM} <i>bla</i> _{PER}
Kalantar et al (2012) ⁹⁸	One hospital in Sanandaj	General inpatients	Different sites	<i>P. aeruginosa</i> (100)	✓	<ul style="list-style-type: none">• DD• DDST for MBL detection• PCR		<ul style="list-style-type: none">• 22 isolates were MBL positive and among them 8 isolates were resistant to IPM• 8 isolates had <i>bla</i>_{VIM}			<i>bla</i> _{VIM}
Shahcheraghi et al (2011) ⁹⁹	Seven hospitals in Tehran	General inpatients	Different sites	<i>Acinetobacter</i> spp. (203)	✓	<ul style="list-style-type: none">• DD• CDT for MBL detection• PCR	Of the 100 strains that were IPM resistant: CIP 84% AMK 84%	<ul style="list-style-type: none">• 49% and 100% of isolates were resistant to IPM and MEM, respectively• All isolates had an MIC ≥8% and 47% had an MIC =64 mcg/mL• From IPM-resistant isolates, 9% were MBL producers• The <i>bla</i>_{SHV}, <i>bla</i>_{GES}[†], <i>bla</i>_{OXA-51}[†] and <i>bla</i>_{OXA-23} genes were detected by PCR among 6, 2, 94, and 84 isolates of <i>A. baumannii</i>, respectively	<ul style="list-style-type: none">• Multicenter study• Large sample size• First report of <i>bla</i>_{SHV} and <i>bla</i>_{GES}	<ul style="list-style-type: none">• Multicenter study• Large sample size• First report of <i>bla</i>_{SHV} and <i>bla</i>_{GES}	<i>bla</i> _{OXA-23} <i>bla</i> _{SHV} <i>bla</i> _{GES}

(Continued)

Table 4 (Continued)

Author (year)	Setting	Population (no.)	Samples	Microorganism	MIC ^a	Method of carbenemase enzyme detection	Resistance to FQ and AG	Outcome	Weak points	Strength points	bla-type genes
Peymani et al (2011) ¹⁰⁰	One hospital in Tabriz	General inpatients	Different sites	<i>A. baumannii</i> (100)	–	<ul style="list-style-type: none"> E-test for MBL detection PCR 	AMK 83% GEN 91% CIP 90% LVX 86%	<p>About 55% of isolates were resistant to carbenem</p> <ul style="list-style-type: none"> 31 of carbenem-resistant isolates were MBL producers and 28 of them were positive for bla genes 	–	–	<i>bla</i> _{IMP} <i>bla</i> _{IMP}
Saderi et al (2010) ¹⁰¹	One hospital in Tehran	General inpatients	Burn wounds	<i>P. aeruginosa</i> (100)	–	<ul style="list-style-type: none"> CDT for MBL detection PCR for MBL genes 	GEN 86% AMK 73% CIP 55%	<ul style="list-style-type: none"> 69% of isolates were resistant to IPM 65 isolates were CDT positive 13 resistant isolates contained <i>bla</i>_{IMP} 	<ul style="list-style-type: none"> MIC was not determined 	–	<i>bla</i> _{IMP}
Yousefi et al (2010) ¹⁰²	Two hospitals in Orumieh and Tabriz	General inpatients	Different sites	<i>P. aeruginosa</i> (324)	✓	<ul style="list-style-type: none"> DDST for detection of MBL PCR 	CIP 83% GEN 86% AMK 85%	<ul style="list-style-type: none"> Resistance rate to MEM was 86% Most of the isolates had an MIC ≥32 mcg/mL for IPM Among nonsusceptible strains to IPM, 39 isolates were MBL positive 18 out of resistant isolates carried <i>bla</i>_{IMP} and 6 isolates carried <i>bla</i>_{IMP} 	<ul style="list-style-type: none"> Large sample size 	–	<i>bla</i> _{IMP} <i>bla</i> _{IMP}

(Continued)

Table 4 (Continued)

Author (year)	Setting	Population (no.)	Samples	Microorganism	MIC ^a	Method of carbapenemase enzyme detection	Resistance to FQ and AG	Outcome	Weak points	Strength points	bla-type genes
Bahar et al (2010) ¹⁰³	One hospital in Tehran (Motahari)	Burn patients	Burn wounds	<i>P. aeruginosa</i> (186)	–	<ul style="list-style-type: none"> Disk inhibitor synergy test PCR 	It was not reported for all isolates (only reported in MBL producers)	<ul style="list-style-type: none"> EDTA disk showed that 23 strains produced MBL and all had <i>bla_{VIM}</i> gene Mortality of MBL producer strains was 82% and non-MBL producer was 22% 		<ul style="list-style-type: none"> Large sample size Mortality rates of MBL and non-MBL producers were reported 	<i>bla_{VIM}</i>
Khosravi et al (2008) ¹⁰⁴	One hospital in Ahvaz	General inpatients	Different sites	<i>P. aeruginosa</i> (100)	–	<ul style="list-style-type: none"> E-test for MBL detection PCR 	–	<ul style="list-style-type: none"> Resistance rate to carbapenems was 41% Of the resistant isolates, eight isolates were MBL producers and all contained <i>bla_{VIM}</i> 	<ul style="list-style-type: none"> FQ and AG were not tested 		<i>bla_{VIM}</i>
Yazdi et al (2007) ¹⁰⁵	Two hospitals in Tehran	Not defined	Not defined	<i>P. aeruginosa</i> (126)	✓	<ul style="list-style-type: none"> DD E-test for MBL detection PCR 	–	<ul style="list-style-type: none"> According to MIC, resistance to IPM was 29% 70 isolates were MBL producers 8 isolates had <i>bla_{VIM}</i> MIC of these eight strains was 32–64 mcg/mL for IPM 	<ul style="list-style-type: none"> Few antibiotics were tested 	–	<i>bla_{VIM}</i>

Notes: ^aMIC values for carbapenems (but not other antimicrobial agents) were considered. ^bResistance rates were according to the result of disk diffusion test. ^cResults about *bla_{OXA-51}* were not provided because this gene has no correlation with occurrence of resistance, unless it has *ISAba* genes in its upstream. ^dFor complete information about the resistance to AG and FQ, please refer to the original reference.

Abbreviations: AG, aminoglycoside; AMK, amikacin; BA, boronic acid; CDT, combination disk test; CIM, carbapenem inactivation method; CIP, ciprofloxacin; DD, disk diffusion; DDST, double disk synergy test; DOR, doripenem; ERT, eripipenem; ESBL, extended spectrum β -lactamase; FQ, fluoroquinolones; GAT, gentamicin; ICU, intensive care unit; IPM, imipenem; KAN, kanamycin; KPC, *Klebsiella*-producing carbapenemase; LVX, levofloxacin; MBL, metallo- β -lactamase; MCNP, modified CarbaNP; MEM, meropenem; MHT, modified Hodge test; MIC, minimum inhibitory concentration; NAL, nalidixic acid; NOR, norfloxacin; OFX, ofloxacin; PCR, polymerase chain reaction; TOB, tobramycin; BAL, Broncho Alveolar Lavage.

Figure 3 Carbapenem-resistance rates in different areas of Iran (according to the disk diffusion method).

Notes: The data were extracted from the latest available studies. Multicenter studies from different cities were not considered for mapping due to pooled data. The studies in the special populations (ie, pediatrics or cystic fibrosis) and outpatients were excluded. *Reported resistance rates from Fars province were conflicting (13.7% in 2017 and 96% in 2016). †Data for Isfahan province were obtained from both Kashan (a city around Isfahan) and Isfahan itself, although the sample size for Isfahan was very small. ‡Data for Khorasan area were extracted from a study in 2015 with 36 strains. §Intermediate and absolute resistance rates were 25% and 2%, respectively. ¶Data in these provinces were for 2010–2014 studies.

Most studies included biological samples from different sites like urine, trachea, and wounds. However, sites with highest or lowest resistant rates were not defined, except in a few studies. In study by Nobari et al, urine was the source with more resistant strains.⁸⁷ In the study by Khorvash et al, one *P. aeruginosa* strain that was isolated from urine samples harbored both *bla*_{IMP} and *bla*_{VIM} genes.⁶⁷

Lack of clinical insight was the main limitation of almost all studies. Only in three studies, mortality rates in patients infected with resistant strains were addressed. Mortality rate of patients infected with carbapenem-resistant *K. pneumonia* isolates was 33% in 2013 (Rastegar Lari et al study).²⁵ In the same year, infections with carbapenem-resistant *A. baumannii* strains caused 20% mortality.²⁴ In the study by Bahar et al, the mortality rate of MBL-producer *P. aeruginosa*

isolates vs non-MBL producers was 82% vs 22%.¹⁰³ In some studies, patients' conditions including requiring mechanical ventilation and days of hospital stay were considered, but no correlation between these data and acquiring carbapenem-resistant strains was found. Also in most studies, difference between infection and colonization was not clear.

Restrictions in technical facilities and standard laboratories in small cities are important issues. Access to a standard microbiological laboratory is not feasible in most regions. Disk diffusion was the main method for detecting carbapenem resistance until recently. Also, MIC breakpoints were adapted from CLSI and national data regarding these cutoffs are lacking. National antimicrobial resistance surveillance studies have not been well organized. Interpreting data from different regions of a country, considering the local standards, is essential.

Figure 4 Carbapenem-resistant rates in different areas of Iran (according to the phenotypic and genotypic methods).

Notes: The data were extracted from the latest available studies. Multicenter studies from different cities were not considered for mapping due to pooled data. The studies in the special populations (ie, pediatrics or cystic fibrosis) and outpatients were excluded.

Some genes like blaSPM and blaSHV are mostly known as genes that encode ESBL enzymes, but they were included in this map, because these genes were assessed along with carbapenemase genes, and also overexpression of these genes concomitant with harboring efflux pump may be responsible for resistance to carbapenems. *Data for these five provinces were extracted from 2011 to 2014 articles, so new data are needed.

[†]Data were extracted from a 2015 study with 36 samples; another study in 2013 confirmed the presence of the genes of resistance in 4% of strains. [‡]Data from Tehran province were conflicting; genes encoding resistance were detected in 15%–50% of isolates through 2015–2016. [§]Data from Kashan (a city in Isfahan province) showed presence of resistance genes in 80% of isolates in 2015.

[¶]In Mazandaran and West Azerbaijan, these genes were reported, but rates of resistance were not included because the studies included different cities from different provinces. ①blaOXA-23 ②blaOXA-24 ③blaOXA-58 ④blaOXA-40 ⑤blaOXA-48 ⑥blaOXA-10 ⑦blaOXA-143 ⑧blaIMP ⑨blaVIM ⑩blaSPM ⑪blaGES ⑫blaPER ⑬blaNDM ⑭blaKPC ⑮blaTEM ⑯blaSHV ⑰blaADC ⑱blaSIM ⑲blaGIM.

Abbreviations: ESBL, extended spectrum β-lactamase; KPC, *Klebsiella*-producing carbapenemase; MBL, metallo-β-lactamase.

Pattern of antibiotic use can affect emergence of resistant microorganisms. Rational use of drugs, and specifically antibiotics, is a challenging issue in developing countries. Mean number of drugs per prescription in Iran was higher than the World Health Organization standards.^{113,114} Approximately, 50% of inpatients, prescriptions contained at least one antibiotic, and this percentage was even higher in outpatient settings.¹¹⁵ Overuse of antibiotics, especially injectable ones, and easy access to antibiotics without prescription is a warning alarm for future antibiotic resistance in developing countries.

Establishing antimicrobial stewardship's programs is new in our hospitals. Unfortunately, governmental rules and supports to restrict antibiotic access in community pharmacies and prescription by general physicians are limited.

Following issues may be considered in future studies. Considering combination of antimicrobials for assessing the resistance is an important finding. Evaluating effects of combination disks (ie, a carbapenem with an aminoglycoside or a fluoroquinolone) on resistant gram-negative bacilli may be helpful.¹¹⁶ Although combination therapy may increase the

Table 5 Types of bla genes, first location of isolation, and the relevant microorganisms in Iran

Ambler classes	Types of pf bla genes	First location of isolation	Year ^a	Total genes ^b	Microorganism ^c
A	GES	Tehran, by Shahcheraghi F.	2011	12	Mostly <i>P. aeruginosa</i>
	KPC	Tehran, by Nobari S.	2014	61	Mostly <i>K. pneumonia</i>
B	NDM	Tehran, by Shahcheraghi F.	2012	44	Mostly <i>K. pneumonia</i>
	VIM	Tehran, by Rezaei Yazdi H.	2007	437	Mostly <i>P. aeruginosa</i>
	IMP	Tabriz and Orumieh, by Yousefi S.	2010	271	Mostly <i>P. aeruginosa</i>
	SIM	Tehran, By Maspi H.	2016	2	Only in <i>A. baumannii</i>
	GIM	Tehran, By Maspi H.	2016	4	Only in <i>A. baumannii</i>
D	OXA-23	Tehran, by Taherikalani M.	2008	1,287	Only in <i>A. baumannii</i>
	OXA-24	Tehran, by Taherikalani M.	2008	213	Only in <i>A. baumannii</i>
	OXA-58	Tehran, by Taherikalani M.	2008	96	Only in <i>A. baumannii</i>
	OXA-40	Tabriz, by Sohrabi N.	2012	59	Only in <i>A. baumannii</i>
	OXA-48	Tehran, by Azimi L.	2014	52	In <i>K. pneumonia</i> and others
	OXA-10	Tehran, by Pakbaten S.	2015	12	Only in <i>P. aeruginosa</i>
	OXA-143	Qom, by Sharikhani Z.	2017	14	Only in <i>A. baumannii</i>

Notes: ^aYear that the genes were isolated for the first time. ^bTotal number of genes that were reported in Iranian studies. ^cThe microorganisms in which the genes were isolated.

risk of side effects, it remains an initial therapeutic option for MDR isolates. Another issue is studying of antibiotics' resistance pattern in health-care facilities. Remarkable increase in resistance rates among isolates from health-care facilities were reported. These strains may act as KPC-producing reservoirs.¹¹⁷ A higher resistance rate in long-term care facilities in comparison with intensive care units has been identified.¹¹⁸ Although some studies in Iran have included outpatients to evaluate carbapenem resistance, patients in long-term health-care facilities have not been included.

Authors' contributions

All authors contributed to data analysis, drafting and revising the article, gave final approval of the version to be published, and agree to be accountable for all aspects of the work.

Disclosure

The authors report no conflicts of interest in this work.

References

- Walsh TR, Toleman MA, Poirel L, Nordmann P. Metallo-beta-lactamases: the quiet before the storm? *Clin Microbiol Rev.* 2005;18(2):306–325.
- Chan JD, Graves JA, Dellit TH. Antimicrobial treatment and clinical outcomes of carbapenem-resistant *Acinetobacter baumannii* ventilator-associated pneumonia. *J Intensive Care Med.* 2010;25(6):343–348.
- Sun K, Xu X, Yan J, Zhang L. Evaluation of six phenotypic methods for the detection of carbapenemases in gram-negative bacteria with characterized resistance mechanisms. *Ann Lab Med.* 2017;37(4):305–312.
- Ambler RP. The structure of b-lactamases. *Philos Trans R Soc Lond B Biol Sci.* 1980;289:321–331.
- Mcmullen AR, Yarbrough ML, Wallace MA, Shupe A, Burnham CD. Evaluation of genotypic and phenotypic methods to detect carbapenemase production in gram-negative *Bacilli*. *Clin Chem.* 2017;63(3):723–730.
- Pletz MW, Wellinghausen N, Welte T. Will polymerase chain reaction (PCR)-based diagnostics improve outcome in septic patients? A clinical view. *Intensive Care Med.* 2011;37(7):1069–1076.
- Meletis G, Exindari M, Vavatsi N, Sofianou D, Diza E. Mechanisms responsible for the emergence of carbapenem resistance in *Pseudomonas aeruginosa*. *Hippokratia.* 2012;16(4):303–307.
- Douraghi M, Ghalavand Z, Nateghi Rostami M, et al. Comparative *in vitro* activity of carbapenems against clinical isolates of *Acinetobacter baumannii*. *J Appl Microbiol.* 2016;121(2):401–407.
- Ghasemian R, Ahanjan M, Fatehi E, Shokri M. Prevalence and antibiotic resistance pattern of *Acinetobacter* isolated from patients admitted in ICUs in Mazandaran, Northern Iran. *Glob J Health Sci.* 2016;8(11):112–119.
- Shakibaie MR, Adeli S, Salehi MH. Antibiotic resistance patterns and extended-spectrum β -lactamase production among *Acinetobacter* spp. isolated from an intensive care unit of a hospital in Kerman, Iran. *Antimicrob Resist Infect Control.* 2012;1:1.
- Yousefi S, Farajnia S, Nahaei MR, et al. Class 1 integron and Imipenem Resistance in Clinical Isolates of *Pseudomonas aeruginosa*: Prevalence and Antibiotic Susceptibility. *Iran J Microbiol.* 2010; 2(3): 115–121.
- Rahbar M, Kabeh-Monnavar M, Khadem Vatan K, Fadaei-Haqi A, Shakerian F. Carbapenem resistance in gram-negative *Bacilli* isolates in an Iranian 1000-bed tertiary hospital. *Pak J Med Sci.* 2008;24(4):537–540.
- Babamahmoodi F, Ahangarkani F, Davoudi A. Hospital-acquired infections, bacterial causative agents and antibiotic resistance pattern in intensive care units at teaching hospitals in north of Iran. *Int J Med Invest.* 2015;4(1):152–160.
- Mohajeri P, Rezaei Z, Sharbati S, et al. Frequency of adhesive virulence factors in carbapenemase-producing *Acinetobacter baumannii* isolated from clinical samples in West of Iran. *Asian J Biol Sci.* 2014;7(4):158–164.
- Mobaraki S, Aghazadeh M, Soroush Barhaghi MH, et al. Prevalence of integrons 1, 2, 3 associated with antibiotic resistance in *Pseudomonas aeruginosa* isolates from Northwest of Iran. *Biomedicine.* 2018; 8(1):2–7.

16. Ghanbari F, Khademi F, Saberianpour S, et al. An epidemiological study on the prevalence and antibiotic resistance patterns of bacteria isolated from urinary tract infections in central Iran. *Avicenna J Clin Microbiol Infect*. 2017;4(3):e42214.
17. Chahoofard A, Dehghan F, Karmostaji A, Zolghadri N. Hospital-acquired urinary tract infection, microbial causative agents and antibiotic resistance pattern in southern Iran: a prospective study. *J Glob Pharma Tech*. 2017;1(9):52–58.
18. Ansari H, Doosti A, Kargar M, Bijanzadeh M, Jafariny M. Antimicrobial resistant determination and prokaryotic expression of smpA Gene of *Acinetobacter baumannii* isolated from admitted patients. *Jundishapur J Microbiol*. 2017;10(11):e59370.
19. Babakhani S, Shokri Derikvand S, Nazer MR, Kazemi MJ. Comparison frequency and determination antibiotic resistance pattern of *Klebsiella* spp. isolated from Nosocomial infection in Khorramabad Shohadaye Ashayer hospital. *Bull Env Pharmacol Life Sci*. 2014;3(12): 149–154.
20. Kamalbeik S, Koucheh M, Baseri Salehi M, Fallah F, Malekan MA, Talaie H. Prevalence of Class 2 integrons in multidrug-resistant *Acinetobacter Baumannii* in toxicological ICU patients in Tehran. *Iran J Toxicol*. 2013;7(22):900–906.
21. Hashemi SH, Esna-Ashari F, Tavakoli S, Mamani M. The prevalence of antibiotic resistance of *Enterobacteriaceae* strains isolated in community- and hospital-acquired infections in teaching hospitals of Hamadan, West of Iran. *J Res Health Sci*. 2013;29(13):75–80.
22. Rahbar M, Mehragan H, Haji Ali Akbari N. Prevalence of drug resistance in nonfermenter gram-negative *Bacilli*. *Iran J Pathol*. 2010;5(2):90–96.
23. Soroush S, Haghi-Ashtiani MT, Taheri-Kalani M, et al. Antimicrobial resistance of nosocomial strain of *Acinetobacter baumannii* in Children's Medical Center of Tehran: a 6-year prospective study. *Acta Med Iran*. 2010;48(3):178–184.
24. Lari AR, Mohammadi Barzelighi H, Arjomandzadegan M, Nosrati R, Owlia P. Distribution of Class I integron among isolates of *Acinetobacter baumannii* recovered from burn patients. *J Med Bacteriol*. 2013;2(1):1–11.
25. Rastegar Lari A, Azimi L, Rahbar M, Fallah F, Alaghebandan R. Phenotypic detection of *Klebsiella pneumoniae* carbapenemase among burns patients: first report from Iran. *Burns*. 2013;39(1):174–176.
26. Ghotaslou R, Sadeghi M, Akhi M, Hasani A, Asgharzadeh M. Prevalence and antimicrobial susceptibility patterns of ESBL, AmpC and carbapenemase-producing *Enterobacteriaceae* isolated from hospitalized patients in Azerbaijan, Iran. *Iran J Pharm Res*. 2018;17(Special Issue):79–88.
27. Saadatani Farivar A, Nowroozi J, Eslami G, Sabokbar A. RAPD PCR profile, antibiotic resistance, prevalence of *armA* Gene, and detection of KPC enzyme in *Klebsiella pneumoniae* isolates. *Can J Infect Dis Med Microbiol*. 2018;Article ID 6183162:7.
28. Moosavian M, Shams N, Siros M. Detection of carbapenemases emerging in *Acinetobacter baumannii* clinical isolates by modified Hodge test. *J Med Microbiol Infect Dis*. 2014;2(4):163–166.
29. Jonaidi Jafari N, Izadi M, Hajia M, Qorbanalizadgan M, Saburi A. The susceptibility evaluation of multiresistant Gram-negative *Bacilli* to meropenem and imipenem. *Int J Travel Med Global Health*. 2014;2(1):1–3.
30. Ghadiri H, Vaez H, Razavi-Azarkhiavi K, et al. Prevalence and antibiotic susceptibility patterns of extended-spectrum β -lactamase and metallo- β -lactamase-producing uropathogenic *Escherichia coli* isolates. *Lab Med Fall*. 2014;45:290–294.
31. Fazeli H, Kamali Dolatabadi R, Taraghian A, Nasr Isfahani B, Moghim S, Norouzi M. Carbapenem resistance pattern of multiple drug-resistant and extended-spectrum beta-lactamase-positive *Klebsiella pneumoniae* in Isfahan. *Int J Enteric Pathogens*. 2014;2(4):e21495.
32. Mirsalehian A, Kalantar-Neyestanaki D, Nourijelyani K, et al. Detection of AmpC- β -lactamases producing isolates among carbapenem resistant *P. aeruginosa* isolated from burn patient. *Iran J Microbiol*. 2014;6(5):306–310.
33. Moayednia R, Shokri D, Mobasherizadeh S, Baradaran A, Fatemi SM, Merrikh A. Frequency assessment of β -lactamase enzymes in *Escherichia coli* and *Klebsiella* isolates in patients with urinary tract infection. *J Res Med Sci*. 2014;19(1):S41–S45.
34. Erfani Y, Farahbakhsh M, Godarzi H, Eslami G, Hashemi A. Detection of metallo- β -lactamase (MBL) producing *Acinetobacter* at 3 hospitals in Iran, Tehran. *Res J Biol Sci*. 2013;8(4):88–93.
35. Safari M, Saidijam M, Bahador A, Jafari R, Alikhani MY. High prevalence of multidrug resistance and metallo-beta-lactamase (M β L) producing *Acinetobacter baumannii* isolated from patients in ICU Wards, Hamadan, Iran. *J Res Health Sci*. 2013;13(2):162–167.
36. Masaeli M, Faraji T, Ramazanadeh R. Risk factors associated with resistance in metallo beta-lactamase producing *Enterobacteriaceae* isolated from patients in Sanandaj Hospitals. *Curr Drug Ther*. 2012;7(3): 179–183.
37. Japoni-Nejad A, Sofian M, van Belkum A, Ghaznavi-Rad E. Nosocomial outbreak of extensively and pan drug-resistant *Acinetobacter baumannii* in Tertiary Hospital in Central Part of Iran. *Jundishapur J Microbiol*. 2013;6(8):e9892.
38. Ahangarzadeh Rezaee M, Langarizadeh N, Aghazadeh M. First Report of Class 1 and Class 2 integrons in multidrug-resistant *Klebsiella pneumoniae* isolates from Northwest Iran. *J Infect Dis*. 2012;65:256–259.
39. Haji Hashemi B, Farzanehkhah M, Dolatyari A, et al. Study on prevalence of KPC producing from *Klebsiella pneumoniae* using modified Hodge test and CHROMagar in Iran. *Ann Biol Res*. 2012;3(12):5659–5664.
40. Azimi L, Lari AR, Alaghebandan R, Alinejad F, Mohammadpoor M, Rahbar M. KPC producer gram negative bacteria among burned infants in Motahari hospital, Tehran: first report from Iran. *Ann Burns Fire Disasters*. 2012;25(2):74–77.
41. Japoni A, Alborzi A, Kalani M, Nasiri J, Hayati M, Farshad S. Susceptibility patterns and cross-resistance of antibiotics against *Pseudomonas aeruginosa* isolated from burn patients in the South of Iran. *Burns*. 2006;32(3):343–347.
42. Taherikalani M, Fatolahzadeh B, Emameini M, Soroush S, Feizabadi MM. Distribution of different carbapenem resistant clones of *Acinetobacter baumannii* in Tehran hospitals. *New Microbiol*. 2009;32(3):265–271.
43. Azizi O, Shakibaie MR, Modarresi F, Shahcheraghi F. Molecular detection of class-D OXA carbapenemase genes in biofilm and non-biofilm forming clinical isolates of *Acinetobacter baumannii*. *Jundishapur J Microbiol*. 2015;8(1):e21042.
44. Bahador A, Rao An R, Farshadzadeh Z, et al. The prevalence of IS Aba 1 and IS Aba 4 in *Acinetobacter baumannii* species of different international clone lineages among patients with burning in Tehran, Iran. *Jundishapur J Microbiol*. 2015;8(7):e17167.
45. Sarikhani Z, Nazari R, Nateghi Rostami M. First report of OXA-143-lactamase producing *Acinetobacter baumannii* in Qom, Iran. *Iran J Basic Med Sci*. 2017;20(11):1282–1286.
46. Mohammadi M, Soroush S, Delfani S, et al. Distribution of class D carbapenemase and extended-spectrum β -lactamase genes among *Acinetobacter baumannii* isolated from burn wound and ventilator associated pneumonia infections. *J Clin Diagn Res*. 2017;11(7):DC19–DC23.
47. Kooti S, Motamedifar M, Sarvari J. Antibiotic resistance profile and distribution of oxacillinase genes among clinical isolates of *Acinetobacter baumannii* in Shiraz teaching hospitals, 2012–2013. *Jundishapur J Microbiol*. 2015;8(8):e20215.
48. Mahdian S, Sadeghifard N, Pakzad I, et al. *Acinetobacter baumannii* clonal lineages I and II harboring different carbapenem-hydrolyzing- β -lactamase genes are widespread among hospitalized burn patients in Tehran. *J Infect Public Health*. 2015;8(6):533–542.
49. Farsiani H, Mosavat A, Soleimanpour S, et al. Limited genetic diversity and extensive antimicrobial resistance in clinical isolates of *Acinetobacter baumannii* in north-east Iran. *J Med Microbiol*. 2015;64(7):767–773.

50. Nasrolahei M, Zahedi B, Bahador A, et al. Distribution of bla OXA-23, IS Aba, aminoglycosides resistant genes among burned & ICU patients in Tehran and Sari, Iran. *Ann Clin Microbiol Antimicrob.* 2014;13(1):38.
51. Safari M, Alikhani MY, Arabestani MR, Kamali Kakhki R, Jafari R. Prevalence of metallo- β -lactamases encoding genes among *Pseudomonas aeruginosa* strains isolated from the bedridden patients in the intensive care units. *Avicenna J Clin Microbiol Infect.* 2014;1(1):e19216.
52. Bahador A, Raoofian R, Taheri M, Pourakbari B, Hashemizadeh Z, Hashemi FB. Multidrug resistance among *Acinetobacter baumannii* isolates from Iran: changes in antimicrobial susceptibility patterns and genotypic profile. *Microb Drug Resist.* 2014;20(6):632–640.
53. Karmostaji A, Najar Peerayeh S, Hatef Salmanian A. Distribution of OXA-type class D β -Lactamase genes among nosocomial multi drug resistant *Acinetobacter baumannii* isolated in Tehran hospitals. *Jundishapur J Microbiol.* 2013;6(5):e8219.
54. Shoja S, Moosavian M, Peymani A, Tabatabaiefar MA, Rostami S, Ebrahimi N. Genotyping of carbapenem resistant *Acinetobacter baumannii* isolated from tracheal tube discharge of hospitalized patients in intensive care units, Ahvaz, Iran. *Iran J Microbiol.* 2013;5(4):315–322.
55. Sohrabi N, Farajnia S, Akhi MT, et al. Prevalence of OXA-type β -lactamases among *Acinetobacter baumannii* isolates from Northwest of Iran. *Microb Drug Resist.* 2012;18(4):385–389.
56. Sepehriseresh S, Boroumand MA, Pourgholi L, Sotoudeh Anvari M, Habibi E, Sattarzadeh Tabrizi M. Detection of vim- and ipm-type metallo-beta-lactamases in *Pseudomonas aeruginosa* clinical isolates. *Arch Iran Med.* 2012;15(11):670–673.
57. Forozsh Fard M, Irajian G, Takantape Moslehi Z, Fazeli H, Salehi M, Rezaia S. Drug resistance pattern of *Pseudomonas aeruginosa* strains isolated from cystic fibrosis patients at Isfahan AL Zahra hospital, Iran (2009–2010). *Iran J Microbiol.* 2012;4(2):94–97.
58. Peymani A, Higgins PG, Nahaei M-R, Farajnia S, Seifert H. Characterisation and clonal dissemination of OXA-23-producing *Acinetobacter baumannii* in Tabriz, northwest Iran. *Int J Antimicrob Agents.* 2012;39(6):526–528.
59. Asadollahi P, Akbari M, Soroush S, et al. Antimicrobial resistance patterns and their encoding genes among *Acinetobacter baumannii* strains isolated from burned patients. *Burns.* 2012;38(8):1198–1203.
60. Feizabadi MM, Fathollahzadeh B, Taherikalani M, et al. Antimicrobial susceptibility patterns and distribution of blaOXA genes among *Acinetobacter* spp. isolated from patients at Tehran hospitals. *Jpn J Infect Dis.* 2008;61(4):274–278.
61. Taherikalani M, Etemadi G, Geliani KN, Fatollahzadeh B, Soroush S, Feizabadi MM. Emergence of multi and pan-drug resistance *Acinetobacter baumannii* carrying blaOXA-type-carbapenemase genes among burn patients in Tehran, Iran. *Saudi Med J.* 2008;29(4):623–624.
62. Hosseinzadeh Z, Sedigh Ebrahim-Saraie H, Sarvari J, et al. Emergence of blaNDM-1 and blaOXA-48-like harboring carbapenem-resistant *Klebsiella pneumoniae* isolates from hospitalized patients in south-western Iran. *J Chin Med Assoc.* 2018;81(6):536–540.
63. Bina M, Pournajaf A, Mirkalantari S, Talebi M, Irajian G. Detection of the *Klebsiella pneumoniae* carbapenemase (KPC) in *K. pneumoniae* isolated from the clinical samples by the phenotypic and genotypic methods. *Iran J Pathol.* 2015;10(3):199–205.
64. Lari AR, Azimi L, Soroush S, Taherikalani M. Low prevalence of metallo-beta-lactamase in *Pseudomonas aeruginosa* isolated from a tertiary burn care center in Tehran. *Int J Immunopathol Pharmacol.* 2015;28(3):384–389.
65. Bagheri Josheghani S, Moniri R, Firoozeh F, Sehat M, Dasteh Goli Y. Susceptibility pattern and distribution of oxacillinases and blaPER-1 genes among multidrug resistant *Acinetobacter baumannii* in a teaching hospital in Iran. *J Pathog.* 2015;957259:7.
66. Solgi H, Badmasti F, Aminzadeh Z, et al. Molecular characterization of intestinal carriage of carbapenem-resistant *Enterobacteriaceae* among inpatients at two Iranian university hospitals: first report of co-production of bla_{NDM-7} and bla_{OXA-48}. *Eur J Clin Microbiol Infect Dis.* 2017;36(11):2127–2135.
67. Khorvash F, Yazdani M, Shabani S, Soudi A. *Pseudomonas aeruginosa*-producing metallo- β -lactamases (VIM, IMP, SME, and AIM) in the clinical isolates of intensive care units, a university hospital in Isfahan, Iran. *Adv Biomed Res.* 2017;30(6):147.
68. Akhi MT, Khalili Y, Ghotaslou R, et al. Carbapenem inactivation: a very affordable and highly specific method for phenotypic detection of carbapenemase-producing *Pseudomonas aeruginosa* isolates compared with other methods. *J Chemother.* 2017;29(3):144–149.
69. Falahat S, Shojapour M, Sadeghi A. Detection of KPC carbapenemase in *Pseudomonas aeruginosa* isolated from clinical samples using Modified Hodge Test and boronic acid phenotypic methods and their comparison with the polymerase chain reaction. *Jundishapur J Microbiol.* 2016;9(9):e27249.
70. Mohammadzadeh M, Tavakoli M, Mohebi A, Aghayi S. Phenotypic and genotypic detection of metallo-beta-lactamases among imipenem resistant gram negative isolates. *J Med Bacteriol.* 2016;5(1–2):36–42.
71. Tarashi S, Goudarzi H, Erfanimanesh S, Pormohammad A, Hashemi A. Phenotypic and molecular detection of metallo-beta-lactamase genes among imipenem resistant *Pseudomonas aeruginosa* and *Acinetobacter baumannii* strains isolated from patients with burn injuries. *Arch Clin Infect Dis.* 2016;11(4):e39036.
72. Maspi H, Mahmoodzadeh Hosseini H, Amin M, Imani Fooladi AA. High prevalence of extensively drug-resistant and metallo beta-lactamase-producing clinical *Acinetobacter baumannii* in Iran. *Microb Pathog.* 2016;98:155–159.
73. Moghadam M, Motamedifar M, Sarvari J, Sedigh E-SH, Mousavi SM, Moghadam F. Emergence of multidrug resistance and metallo-beta-lactamase producing *Acinetobacter baumannii* isolated from patients in Shiraz, Iran. *Ann Med Health Sci Res.* 2016;6(3):162–167.
74. Pakbaten Toupanlou S, Najar Peerayeh S, Pirhajati Mahabadi R. Class A and D extended-spectrum β -lactamases in imipenem resistant *Pseudomonas aeruginosa* isolated from burn patients in Iran. *Jundishapur J Microbiol.* 2015;8(8):e18352.
75. Eftekhar F, Naseh Z. Extended-spectrum β -lactamase and carbapenemase production among burn and non-burn clinical isolates of *Klebsiella pneumoniae*. *Iran J Microbiol.* 2015;7(3):144–149.
76. Azimi L, Talebi M, Owlia P, et al. Tracing of false negative results in phenotypic methods for identification of carbapenemase by real-time PCR. *Gene.* 2016;576(1 Pt 1):166–170.
77. Azimi L, Talebi M, Pourshafie M-R, Owlia P, Rastegar Lari A. Characterization of carbapenemases in extensively drug resistance *Acinetobacter baumannii* in a burn care center in Iran. *Int J Mol Cell Med.* 2015;4(1):46–53.
78. Fazeli H, Norouzi-Barough M, Ahadi AM, Shokri D, Solgi H. Detection of New Delhi Metallo-Beta-Lactamase-1 (NDM-1) in carbapenem-resistant *Klebsiella pneumoniae* isolated from a university hospital in Iran. *Hippokratia.* 2015;19(3):205–209.
79. Khoshvaght H, Haghi F, Zeighami H. Extended spectrum betalactamase producing enteroaggregative *Escherichia coli* from young children in Iran. *Gastroenterol Hepatol Bed Bench.* 2014;7(2):131–136.
80. Noori M, Karimi A, Fallah F, et al. High prevalence of metallo-beta-lactamase producing *Acinetobacter baumannii* isolated from two hospitals of Tehran, Iran. *Arch Pediatr Infect Dis.* 2014;2(1):e15439.
81. Vali P, Shahcheraghi F, Seyfipour M, Zamani MA, Allahyar MR, Feizabadi MM. Phenotypic and genetic characterization of carbapenemase and ESBLs producing gram-negative bacteria (GNB) isolated from patients with cystic fibrosis (CF) in Tehran hospitals. *J Clin Diag Res.* 2014;8(1):26–30.
82. Azimi L, Nordmann P, Lari AR, Bonnin RA. First report of OXA-48-producing *Klebsiella pneumoniae* strains in Iran. *GMS Hyg Infect Control.* 2014;7:9.
83. Hashemi A, Fallah F, Erfanimanesh S, Hamedani P, Alimehr S, Goudarzi H. Detection of β -lactamases and outer membrane porins among *Klebsiella pneumoniae* strains isolated in Iran. *Scientifica.* 2014;726179:6.
84. Japoni-Nejad A, Ghaznavi-Rad E, van Belkum A. Characterization of plasmid-mediated AmpC and carbapenemases among Iranian nosocomial isolates of *Klebsiella pneumoniae* using phenotyping and genotyping methods. *Osong Public Health Res Perspect.* 2014;5(6):333–338.

85. Farajzadeh Sheikh A, Rostami S, Jolodar A, et al. Detection of metallo-beta lactamases among carbapenem-resistant *Pseudomonas aeruginosa*. *Jundishapur J Microbiol*. 2014;7(8):e12289.
86. Aghamiri S, Amirmozafari N, Fallah Mehrabadi J, Fouladatan B, Samadi Kafil H. Antibiotic resistance pattern and evaluation of metallo-beta lactamase genes including *bla*_{IMP} and *bla*_{VIM} types in *Pseudomonas aeruginosa* isolated from patients in Tehran Hospitals. *ISRN Microbiol*. 2014;2014(1):1–6.
87. Nobari S, Shahcheraghi F, Rahmati Ghezlegh F, Valizadeh B. Molecular characterization of carbapenem-resistant strains of *Klebsiella pneumoniae* isolated from Iranian patients: first identification of *bla*_{KPC} gene in Iran. *Microb Drug Resist*. 2014;20(4):285–293.
88. Hakemi Vala M, Hallajzadeh M, Hashemi A, et al. Detection of Ambler class A, B and D β -lactamases among *Pseudomonas aeruginosa* and *Acinetobacter baumannii* clinical isolates from burn patients. *Ann Burn Fire Disast*. 2014;27(1):8–13.
89. Tavajjohi Z, Moniri R, Zarrabi M. Detection of GES-2, a class A β -lactamase produced by *Pseudomonas aeruginosa* in a teaching hospital in Iran. *Jundishapur J Microbiol*. 2013;6(10):e8166.
90. Azimi L, Lari AR, Talebi M, Ebrahimzadeh Namvar A, Soleymanzadeh-Moghadam S. Evaluation of phenotypic methods for detection of *Klebsiella pneumoniae* carbapenemase-producing K. *Pneumoniae in Tehran. J Med Bacteriol*. 2013;2(3–4):26–31.
91. Noori N, Vandyosefi J, Sabet F, Ashrafi S, Ghazvini K. Frequency of IMP-1 and VIM genes among metallo-beta-lactamase producing *Acinetobacter* spp. isolated from health care associated infections in Northeast of Iran. *J Med Bacteriol*. 2013;2(3–4):11–16.
92. Azimi L, Lari AR, Talebi M, Namvar AE, Jabbari M. Comparison between phenotypic and PCR for detection of OXA-23 type and metallo-beta-lactamases producer *Acinetobacter* spp. *GMS Hyg Infect Control*. 2013;8(2):Doc16.
93. Pajand O, Rezaee MA, Nahaei MR, et al. Study of the carbapenem resistance mechanisms in clinical isolates of *Acinetobacter baumannii*: comparison of burn and non-burn strains. *Burns*. 2013;39(7):1414–1419.
94. Azami S, Abdi Ali A, Asgarani E. Association between metallo- β -lactamases and integrons with multi-drug resistance in *Pseudomonas aeruginosa* isolates. *J Med Microbiol Infect Dis*. 2013;1(1):46–51.
95. Mohajeri P, Farahani A, Feizabadi MM, Ketabi H, Abiri R, Najafi F. Antimicrobial susceptibility profiling and genomic diversity of *Acinetobacter baumannii* isolates: a study in western Iran. *Iran J Microbiol*. 2013;5(3):195–202.
96. Doosti M, Ramazani A, Garshasbi M. Identification and characterization of metallo- β -lactamases producing *Pseudomonas aeruginosa* clinical isolates in University Hospital from Zanjan Province, Iran. *Iran Biomed J*. 2013;17(3):129–133.
97. Shahcheraghi F, Nobari S, Rahmati Ghezlegh F, et al. First report of New Delhi metallo-beta-lactamase-1-producing *Klebsiella pneumoniae* in Iran. *Microb Drug Resist*. 2013;19(1):30–36.
98. Kalantar E, Torabi V, Salimizand H, Soheili F, Beiranvand S, Soltan Dallal MM. First survey of metallo- β -lactamase producers in clinical isolates of *Pseudomonas aeruginosa* from a Referral Burn Center in Kurdistan Province. *Jundishapur J Nat Pharm Prod*. 2012;7(1):23–26.
99. Shahcheraghi F, Abbasalipour M, Feizabadi M, Ebrahimipour G, Akbari N. Isolation and genetic characterization of metallo- β -lactamase and carbapenemase producing strains of *Acinetobacter baumannii* from patients at Tehran hospitals. *Iran J Microbiol*. 2011;3(2):68–74.
100. Peymani A, Nahaei MR, Farajnia S, et al. High prevalence of metallo-beta-lactamase-producing *Acinetobacter baumannii* in a teaching hospital in Tabriz, Iran. *Jpn J Infect Dis*. 2011;64(1):69–71.
101. Sadari H, Lotfalipour H, Owlia P, Salimi H. Detection of metallo- β -lactamase producing *Pseudomonas aeruginosa* isolated from burn patients in Tehran, Iran. *Lab Med*. 2010;41(10):609–612.
102. Yousefi S, Farajnia S, Nahaei MR, et al. Detection of metallo- β -lactamase-encoding genes among clinical isolates of *Pseudomonas aeruginosa* in northwest of Iran. *Diagn Microbiol Infect Dis*. 2010;68(3):322–325.
103. Bahar MA, Jamali S, Samadikuchaksaraei A. Imipenem-resistant *Pseudomonas aeruginosa* strains carry metallo-beta-lactamase gene *bla*(VIM) in a level I Iranian burn hospital. *Burns*. 2010;36(6):826–830.
104. Khosravi AD, Mihani F. Detection of metallo-beta-lactamase-producing *Pseudomonas aeruginosa* strains isolated from burn patients in Ahwaz, Iran. *Diagn Microbiol Infect Dis*. 2008;60(1):125–128.
105. Yazdi HR, Nejad GB, Peerayeh SN, Mostafaei M. Prevalence and detection of metallo- β -lactamase (MBL)-producing *Pseudomonas aeruginosa* strains from clinical isolates in Iran. *Ann Microbiol*. 2007;57(2):293–295.
106. Al Johani SM, Akhter J, Balkhy H, El-Saed A, Younan M, Memish Z. Prevalence of antimicrobial resistance among gram-negative isolates in an adult intensive care unit at a tertiary care center in Saudi Arabia. *Ann Saudi Med*. 2010;30(5):364–369.
107. Kempf M, Rolain JM. Emergence of resistance to carbapenems in *Acinetobacter baumannii* in Europe: clinical impact and therapeutic options. *Int J Antimicrob Agents*. 2012;39(2):105–114.
108. Cantón R, Akóva M, European Network on Carbapenemases, et al. Rapid evolution and spread of carbapenemases among *Enterobacteriaceae* in Europe. *Clin Microbiol Infect*. 2012;18(5):413–431.
109. Castanheira M, Deshpande LM, Costello A, Davies TA, Jones RN. Epidemiology and carbapenem resistance mechanisms of carbapenem-non-susceptible *Pseudomonas aeruginosa* collected during 2009–11 in 14 European and Mediterranean countries. *J Antimicrob Chemother*. 2014;69(7):1804–1814.
110. Clinical and Laboratory Standards Institute. *Performance Standards for Antimicrobial Susceptibility Testing*. Wayne, PA: CLSI; 2018:M100-S17.
111. Kuchibiro T, Komatsu M, Yamasaki K, et al. Evaluation of the modified carbapenem inactivation method for the detection of carbapenemase-producing *Enterobacteriaceae*. *J Infect Chemother*. 2018;24(4):262–266.
112. Pirbonyeh N, Zardosht M, Emami A, Rostampour S, Moattari A, Keshavarzi A. Emergence of storm resistant mechanisms in *Pseudomonas aeruginosa* isolated from burn patients hospitalized in Ghotbeddin Shirazi Burn Hospital. *Nova J Med Biol Sci*. 2016;5(1).
113. Karimi A, Haerizadeh M, Soleymani F, Haerizadeh M, Taheri F. Evaluation of medicine prescription pattern using World Health Organization prescribing indicators in Iran: a cross-sectional study. *J Res Pharm Pract*. 2014;3(2):39–45.
114. Mousavi S, Zargarzadeh AH. Rational drug use in Iran: a call for action. *J Pharm Care*. 2014;2(2):47–48.
115. Sefidani Forough A, Hosseini SR, Jabbari S. Antibiotic utilization evaluation of inpatient and outpatient prescriptions in a rural general hospital in Iran. *Int J Basic Clin Pharmacol*. 2015;4(3):531–536.
116. Leite GC, Neto LVP, Gaudereto JJ, et al. Effect of antibiotic combination and comparison of methods for detection of synergism in multi-resistant gram-negative bacteria. *J Inf Dis Ther*. 2015;3(2):207.
117. Munoz-Price LS, Poirel L, Bonomo RA, et al. Clinical epidemiology of the global expansion of *Klebsiella pneumoniae* carbapenemases. *Lancet Infect Dis*. 2013;13(9):785–796.
118. Viau R, Frank KM, Jacobs MR, et al. Intestinal carriage of carbapenemase-producing organisms: current status of surveillance methods. *Clin Microbiol Rev*. 2016;29(1):1–27.

Infection and Drug Resistance**Dovepress****Publish your work in this journal**

Infection and Drug Resistance is an international, peer-reviewed open-access journal that focuses on the optimal treatment of infection (bacterial, fungal and viral) and the development and institution of preventive strategies to minimize the development and spread of resistance. The journal is specifically concerned with the epidemiology of antibiotic

resistance and the mechanisms of resistance development and diffusion in both hospitals and the community. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/infection-and-drug-resistance-journal>