Economic analysis of acupuncture for migraine prophylaxis

Jitka Pokladnikova^{1,2}Petra Maresova³Josef Dolejs³A-La Park⁴Bo Wang⁵Xin Guan⁵Frantisek Musil¹

¹Czech-Chinese Centre for Traditional Chinese Medicine, University Hospital in Hradec Kralove, Hradec Kralove, Czech Republic; ²Faculty of Pharmacy in Hradec Kralove, Charles University, Hradec Kralove, Czech Republic; ³Faculty of Informatics and Management, University of Hradec Kralove, Hradec Kralove, Czech Republic; ⁴Personal Social Services Research Unit, Department of Health Policy, London School of Economics and Political Science, London, UK; ⁵Shuguang Hospital Affiliated to Shanghai University of Traditional Chinese Medicine, Shanghai, China

Background: Acupuncture has become a viable option for migraine prophylaxis in Europe; however, despite its wide use, more data on the short- and long-term cost-effectiveness are needed when considering the perspectives of a paying third-party, the patient, and of society in general. The aim was to evaluate the cost and effectiveness of adjuvant acupuncture to pharmacologic treatment vs pharmacologic treatment alone in migraine patients after a 3-month acupuncture course and a 6-month follow-up from all perspectives.

Methods: The study involved an open-label randomized clinical trial of patients receiving acupuncture (n=42), and a waiting list control group (n=44). The number of migraine days during the last 28 days, as well as direct and indirect costs were considered. The trial was registered under DRKS00009803.

Results: The total cost per patient reached €696 vs €285 after 3 months of acupuncture and €66 vs €132 in the acupuncture and control groups after a 6-month follow-up, respectively ($P=0.071$). The trends observed in effectiveness and costs from all perspectives are discussed.

Conclusion: The inclusion of acupuncture in health care results beneficial mainly for its observed trend in reduced losses of productivity and income, with the latter often exceeding the costs of acupuncture treatment. As such, acupuncture may be recommended as an adjuvant treatment in migraine prophylaxis to standard pharmacotherapy.

Keywords: acupuncture, cost, effectiveness, migraine, prophylaxis, Traditional Chinese Medicine

Introduction

Migraine is classified within a group of highly prevalent brain disorders with a calculated cost of €662 per migraineur and representing the highest proportion of indirect health care costs, accounting for 87% of overall costs in comparison with other conditions.^{1,2} In Europe, one-third of the population presents at least one brain disorder, including migraine, resulting in an overall cost of 798 billion euro per year (2010).² Migraine-associated disability often results in reduced social activities and work capacity, for example, in the UK some 25 million working/school-days are lost every year because of migraine alone.³ Still, headaches might often be overlooked, by both the lay public and some health care professionals, as a minor complaint and its impacts on physical, emotional, social health, as well as economic development, may not always be fully acknowledged. Further, brain disorders in general are identified as a major health economic challenge in Europe, calling for wider attention on behalf of the health authorities and national governments, as well as more funding for brain research.²

A previous meta-analysis proved that acupuncture could become a viable prophylactic treatment option for frequent or uncontrolled migraine, or for migraineurs

Correspondence: Jitka Pokladnikova
Czech-Chinese Centre for Traditional Chinese Medicine, University Hospital in Hradec Kralove, Sokolska 581, Hradec Kralove 50005, Czech Republic
Tel +420 72 858 5562
Fax +420 49 583 2001
Email jitka.pokladnikova@faf.cuni.cz

experiencing drug side effects.^{4,5} Acupuncture is seemingly superior to pharmacologic treatments in terms of the occurrence of fewer side effects and potential long-term effectiveness after its discontinuation.^{6–8} There is also some evidence suggesting that the introduction of acupuncture to an existing pharmacologic treatment can result in a reduced economic impact due to decreased use of medications, number of sickness absence, and medical review appointments.⁹ Despite the evidence for the potential cost-effectiveness of acupuncture in chronic pain management, such as headache, osteoarthritis, and lower back pain, additional data on the long-term cost-effectiveness of acupuncture are needed.^{8–11} Our comparative study is the first economic analysis comparing acupuncture with a standard pharmacologic treatment in the Czech Republic. The aim of the study was to evaluate the effectiveness and cost of acupuncture treatment in patients with migraine after a 3-month course of acupuncture and 6-month follow-up from three perspectives: a third-party payer, the patient, and society.

Methods

Experimental design

The current study is based on an open-label randomized clinical trial (October 2015 to April 2017), taking place at the Czech-Chinese Centre for Traditional Chinese Medicine in the University Hospital Hradec Kralove. The patients were randomly assigned to either acupuncture (AG) or a waiting list control group (CG). During a 12-week therapeutic period, acupuncture treatment was administered 14 times using the semi-standardized therapeutic acupuncture protocol. Patients in both groups were administered with the standard pharmacologic treatment (prophylactic medication and analgesics as needed) following the appropriate guidelines.¹² The design of the trial including flow diagram and intervention has been described in detail elsewhere.¹³

Patients

Patients aged between 18 and 70 years, attending the Neurology Outpatient Clinic at the University Hospital Hradec Kralove, and with a clinical history of migraine of at least 12 months and a minimum of 4 days of migraine in 4 weeks, were enrolled in the study. All patients were diagnosed with migraine by board-certified neurologists according to the criteria set by the International Classification of Headache Disorders.¹⁴ A written informed consent was obtained from all patients. The study was approved by the Ethical Committee of the University Hospital Hradec Kralove, registered under the code DRKS00009803.

Outcomes

The primary outcome was to compare the cost of acupuncture in patients with migraine after a 3-month course of treatment (T12) and a 6-month follow-up (T36) from three perspectives: a paying third party, the patient, and society in general.

Effectiveness

The effectiveness of the treatment was evaluated according to a previous study¹³, and the first parameter evaluated the number of migraine days in the last 4 weeks before randomization (baseline period) (T0), and at 12 (T12) and 36 (T36) weeks after randomization. A second parameter evaluated the reduction in relief medication use, migraine intensity and duration, and the number of patients with reduced migraine frequency ($\geq 50\%$) over a 4-week period. Specific quality of life using the Migraine Disability Assessment (MIDAS) was also recorded.¹⁵

Costs

The present study employs the methodology established by Drummond,¹⁶ dividing the costs into direct and indirect categories. The former includes variable and fixed costs, whereas the latter includes the costs incurred by the patients and their families. In this regard, the psychological costs are also of a similar nature and therefore not negligible; however, these are not the subject of this paper as neither are the costs associated with hospitalization. Loss of productivity was calculated on the basis of “Human Capital Approach”.¹⁷ Direct and indirect costs were not discounted as the intervention period was less than 1 year. Prices of individual items are shown in Table 1.

In the discussion section a further division of costs is used, namely from a third party payer’s, a patient’s, and societal perspectives.^{18,19} The individual items under examination using the categories of direct and indirect costs^{16,20} were set as

Table 1 Unit cost

Type of costs	€ ^a
Treatment	
Acupuncture (cost out-of-pocket)	
Acupuncture package (an initial visit, five acupuncture sessions, and a check-up visit)	166.0
Acupuncture follow-up session	16.6
Medication, package (reimbursement/co-payment) ^b	— ^c
Health care service, unit	0.04
Productivity/income loss, working day	24.52
Travels, km	0.14

Notes: ^a1 EUR = 27.05 CZK. ^bPrescription and over-the-counter drugs were reimbursed based on the Czech National Drug Price List of 2015/2016²³ and priced at the average price calculated from the actual retail prices obtained from a sample of retail and hospital pharmacies in the study location in 2015/2016. ^cThe unit cost depends on the type of medication and so an average price cannot be established.

follows: direct medical costs (medications, outpatient visits, acupuncture visits, emergency visits, adverse drug reactions), nonmedical direct costs (travel costs), and indirect costs (loss of income, productivity lost) (Table 2). Within the framework of the second parameter for costs examination, the individual groups are a third-party payer (outpatient visits, medication reimbursement, acupuncture visits, emergency visits, and adverse drug reactions), the patient's perspective (medication – drugs' surcharge, over-the-counter (OTC) drugs, travel costs, loss of income, acupuncture visits), and the societal perspective (productivity lost).

The health care costs were calculated by multiplying the number of units for intervention by the unit's price. The costs of drugs were calculated on a consumption basis expressed as defined daily doses (DDD) and the price per DDD.²¹ Travel expenses by the patients were calculated as the product of the number of visits to the physician and the price of each visit. Loss of income was recorded as a result of deterioration of symptoms and days off owing to an appointment with the physician.

Resource use

The information concerning demographics data of the patient's clinical condition, comorbidities, medications, physician's time, and costs of health care services (visits to physicians, visits to emergency department [ED], hospitalization) were obtained from electronic patient record system at the Faculty Hospital in Hradec Králové.

The number of units for intervention and their unit cost were based on current legislation of the Czech Republic for

2015/2016.²² Drug reimbursements were determined according to the Numerical Index 2015/2016.²³ However, there is no available source of information concerning the selling price of drugs in the Czech Republic; therefore, the selling price of drugs and OTC drugs were expressed as a mean obtained from a sample of selected hospital and community pharmacies in a given locality (2015/2016). The following data were collected from patients in the form of questionnaires: use of OTC drugs, visits to ED, hospitalization, sick leave, days off from work or school, and time spent at the physician's office, including transit. Travel expenses were calculated based on the distance between the physician's office and the patient's place of residence and the price of fuel per kilometer.²⁴ Productivity lost was expressed by the number of days off per employee, evaluating each day by the proportional mean gross pay for the year 2015/2016 in the Czech Republic and the number of working days in a year according to data available by the Czech Statistical Office in the Czech Republic.²⁵ The costs were expressed in euros using purchasing power parity for the 2015/2016 year.²⁶

Statistical analysis

The statistical analyses of individual costs were performed using the R Project for Statistical Computing v3.3.3, and MS Excel 2016 was used for data manipulation. A P -value ≤ 0.05 was considered significant in all cases and the number of iterations was of 1,000 in the bootstrap method. At first, normality was tested in 42 patients with acupuncture and in 44 patients in the CG for individual item costs. Normality was tested using the Shapiro–Wilk normality test (namely, the function “shapiro.test” was used). Normality was rejected in all cases ($P < 0.05$), and, consequently, to take account of the skewed distributions of the costs, the bootstrap method (namely, the package “boot” version 2017.2 and the function “boot” were used) was used with 95% CIs of the means.²⁷ The null hypothesis, meaning equal in the AG and the CG, was tested against the alternative hypothesis stating that the costs are lower in the AG using the bootstrap method. Consequently, the resulting P -values were one tailed, and the package “boot” version 2017.2 was used in the calculations.

The cost of an acupuncture session was varied from €11 to €35 for each patient, and all other costs were consistent in the analysis.

Results

Sample characteristics

A total of 91 patients were enrolled in the study, and the flow diagram was reported elsewhere.¹³ There were no significant

Table 2 Direct and indirect costs

Direct costs		Indirect costs
Medical	Nonmedical	
Medication	Travel costs	Loss of income Productivity loss
Outpatient visits		
Acupuncture visits		
Emergency visits		
Adverse drug reactions		
Paying third party	Patient's perspective	Society's perspective
Outpatient visits	Medication (drug surcharge, over-the-counter drugs) Travel costs Loss of income Acupuncture visits	Productivity loss
Medication reimbursement		
Acupuncture visits		
Emergency visits		
Adverse drug reactions		

Note: Costs dividing data from Drummond et al¹⁶ and Maresova et al.²⁰

Table 3 Baseline patient characteristics

Characteristics	Acupuncture group N=42	Control group N=44
Age (years)		
Mean (SD)	45.6 (12.8)	46.5 (10.3)
Median	47	47.5
Female, n (%)	37 (88)	39 (89)
Positive family history of migraine, n (%)	24 (57)	29 (66)
Duration of migraine (years)		
Mean (SD)	26.9 (12.9)	23.0 (14.1)
Median	27.5	20.0
Comorbidity, n (%)	23 (54.8)	23 (52.3)
Smokers, n (%)	9 (21)	5 (11)
Migraine days		
Frequency (no. per month)	11.97 (6.6)	12.1 (9.2)
Duration (hours)	12.2 (15.3)	10 (11.4)
Intensity, VAS (mm)	5.2 (1.3)	5.4 (1.8)
Drug consumption (ATC/DDD)		
Rescue medication, mean (SD)	14.8 (14.3)	11.5 (11.8)
Prophylactic medication, mean (SD)	9.6 (11.9)	7.2 (11.9)
Migraine attack (no per month)	6.4 (2.4)	6.0 (2.7)
Migraine disability assessment score	48.9 (38.1)	52.9 (31.9)
Employees (during working hours), n (%)	17 (40.5)	23 (52.3)
Employees (unpaid hours), n (%)	3 (7.1)	5 (11.4)
Place of residency (local patients), n (%)	9 (21.4)	16 (36.4)
Employees (during working hours), n (%)	17 (40.5)	23 (52.3)

Abbreviations: ATC/DDD, anatomical therapeutic classification/defined daily dose; VAS, visual analog scale.

differences between groups when concerning the baseline characteristics (Table 3).

A significant reduction in the number of migraine days per 4 weeks was reached at the end of the intervention ($\Delta -2.0$; 95% CI: -4 to -1) as well as at the 6-month follow up ($\Delta -4.0$; 95% CI: -6 to -2). A significant percentage of responders to treatment was noted in the AG vs CG at the end of the intervention (50% vs 27%; $P < 0.05$) as well as at the 6-month follow-up (81% vs 36%; $P < 0.001$). The AG showed a notable reduction in the intake of symptomatic medication compared with the CG at the end of acupuncture ($\Delta -2.7$; 95% CI: -5.2 to -0.7) and the 6-month follow-up ($\Delta -3.0$; 95% CI: -5.8 to -0.7). Disease-specific quality-of-life score was improved in both groups at both time points, yet a reduction in the MIDAS scores remained significant only in the AG at the 6-month follow-up. Nevertheless, there was a significant difference between the number of patients with mild (grades I and II; mean score 0–10) vs moderate and severe disability (grades III and VI; mean score >10) before and after the intervention, as well as in the follow-up, in favor of the AG.¹³

Use of health care services

The average number of visits to the physician's office remained similar in both groups at both time points (Table 4); on average, patients in the AG visited their acupuncturist 13.5 times (SD 0.7). The number of migraine-related visits to the ED was significantly lower in the AG during acupuncture treatment when compared to the CG, although no between-group differences were found (T0: 0.4 vs T12: 0.1; $P = 0.012$) (Table 4). Also, patients in the AG experienced fewer drug side effects compared to the CG over the course of acupuncture treatment; yet, the reduction did not last till the 6-month follow-up (Table 4). No serious adverse effects were reported following acupuncture treatment.

Costs

The overall mean cost per patient from all the three perspectives is given in Table 5. Neither of the evaluation periods for AG and CG showed a significant difference in the cost components; nevertheless, the absolute amounts of changes in some costs are still interesting. The direct costs were always lower in the AG vs CG; however, at 3 months, AG costs were higher than those of CG (€506.4 vs €64.4) – when the costs associated with acupuncture visits and traveling were included. When these costs were not considered, however, the direct costs were lower in the AG.

At a 6-month follow-up, the direct and indirect total costs were lower in the AG, where direct costs were lower by €20 and the indirect costs by €46.

It is worth mentioning that the costs changed over time for each group. Within the AG, there was a gradual decrease in the loss of income (T0: €6, T12: €14.0, T36: €3.7) and productivity loss (T0: €128.3, T12: €123.3, T36: €10.2). Overall, a decrease in indirect costs could be observed in AG, from €134.3 at the baseline to €13.9 at the 6-month follow-up. Direct costs within the AG were inevitably higher during the intervention period and then dropped (T0: €38.1, T12: €506.4, T36: €51.7). Within the AG direct costs show a gradual increment in all time periods (T0: €39.4, T12: €64.4, T36: €71.9); further, the absolute comparison of AG and CG at the 6-month follow-up period shows that the direct costs are higher by €20.2 in the AG.

Paying third-party perspective

From a third-party payer's perspective, there were no significant changes between both groups. However, there were lower costs for medication reimbursement in AG during the acupuncture treatment period (€25.6 per person) and at the 6-month follow-up (€29.3). Outpatient and emergency

Table 4 Mean use of resources during a 3-month acupuncture treatment and a 6-month follow-up

	Acupuncture group N=42			Control group N=44			P-value
Baseline year	N	Mean	SD	N	Mean	SD	
Rescue/prophylactic medication (ATC/DDD) ^a	42	14.8/9.6	(14.3/11.9)	44	11.5/7.2	(11.8/11.9)	NS
Outpatient visits, n	42	1.0	0.7	44	1.0	0.8	NS
Productivity loss, working days	42	0.1	0.1	44	0.1	0.2	NS
Loss of income, unpaid working days	42	0.0	0.1	44	0.0	0.1	NS
Emergency visits	42	0.4	0.5	42	0.3	0.6	NS
Hospitalization	40	0.0	0.0	41	0.0	0.0	NS
Adverse drug reactions, n (%)	41	9	(22.0)	41	10	(24.4)	NS
Acupuncture treatment (3 months)							
Rescue medication (ATC/DDD) ^a	42	26.1	41.4	42	26.8	27.5	NS
Acupuncture visits, n	42	13.5	0.7				
Productivity loss, working days	42	0.8	1.4				
Loss of income, unpaid working days	42	0.3	0.9				
Outpatient visits, n	42	1.0	1.0	42	1.0	0.9	NS
Productivity loss, working days	42	0.1	0.2	42	0.1	0.1	NS
Loss of income, unpaid working days	42	0.0	0.1	42	0.0	0.1	NS
Emergency visits	42	0.1*	0.4	42	0.2	0.5	NS
Hospitalization	42	0.0	0.0	42	0.0	0.0	NS
Adverse drug reactions, n (%)	42	2	(4.8)	40	6	(15.0)	NS
6-month follow-up							
Rescue medication (ATC/DDD) ^a	40	26.6	50.8	41	28.4	31.9	NS
Outpatient visits, n	40	0.7	0.7	41	1.0	1.1	NS
Productivity loss, working days	40	0.0*	0.1	41	0.1	0.2	NS
Loss of income, unpaid working days	40	0.0	0.0	41	0.0	0.1	NS
Emergency visits	40	0.3	0.6	39	0.4	0.7	NS
Hospitalization	40	0.0	0.0	37	0.0	0.0	NS
Adverse drug reactions, n (%)	39	7	(17.9)	38	8	(21.1)	NS

Notes: ^aRescue medication (ATC: N02, M01, R06). *Level of significance 0.05 (between-group comparison, unpaired t-test, Mann–Whitney test, Fisher's exact test). Level of significance 0.05 (within-group comparison, paired t-test, Wilcoxon signed-rank test).

Abbreviations: ATC/DDD, anatomical therapeutic classification/defined daily dose; NS, nonsignificant.

Table 5 Comparison of mean baseline and overall costs after a 3-month acupuncture treatment and a 6-month follow-up

	Acupuncture group (N=42), cost per patient			Control group (N=44), cost per patient			Bootstrap t P-value (one tailed)	
	N	Mean	95% CI	N	Mean	95% CI	P-value	Mean _{control} – Mean _{acup}
Baseline year								
Direct costs								
Medication reimbursement	42	8.6	(2.8, 14.4)	44	16.4	(0.4, 32.4)	0.89	7.8
Medication co-payment	42	9.8	(3.8, 16)	44	5.5	(2.2, 8.8)	0.89	–4.3
Outpatient visits	42	5.0	(4, 6)	44	4.6	(3.7, 5.6)	0.71	–0.4
Emergency visits	42	6.4	(3.5, 9.2)	44	5.7	(2.6, 8.6)	0.64	–0.7
Adverse drug reactions	42	0.6	(–0.5, 1.8)	44	0.2	(–0.1, 0.6)	0.74	–0.3
Travel costs	42	7.5	(4.3, 10.9)	44	6.8	(3.6, 9.9)	0.62	–0.7
Direct total costs	42	38.1	(–27.8, 65.2)	44	39.4	(19.8, 58.7)	0.45	1.2
Indirect costs								
Productivity loss	42	128.3	(57.2, 199.9)	44	249.2	(68, 430.7)	0.24	120.9
Loss of income	42	6.0	(–2.8, 15.5)	44	4.1	(–2.1, 10.3)	0.65	–1.9

(Continued)

Table 5 (Continued)

	Acupuncture group (N=42), cost per patient			Control group (N=44), cost per patient			Bootstrap t P-value (one tailed)	Mean _{control} – Mean _{acup}
	N	Mean	95% CI	N	Mean	95% CI	P-value	
Indirect total costs	42	134.3	(61.1, 208.7)	44	253.4	(70.9, 439.9)	0.27	119.0
Total costs	42	172.5	(100.4, 282.2)	44	292.8	(110.8, 474.4)	0.26	120.3
Acupuncture treatment (3 months)								
Direct costs								
Medication reimbursement	42	15.8	(5.4, 27)	44	41.5	(-2.0, 84.3)	0.12	25.6
Medication co-payment	42	21.8	(5.9, 37.7)	44	11.1	(4.7, 17.8)	0.89	-10.7
Acupuncture visits, patients	42	308.5	(304.8, 312.1)					
Acupuncture visits, third-party payer	42	25.1	(25.1, 25.1)					
Travel acupuncture costs	42	120.8	(84.9, 155.5)					
Outpatient visits	42	4.0	(2.6, 5.2)	44	4.1	(3.1, 5.1)	0.45	0.1
Travel costs	42	7.8	(4, 11.8)	44	3.7	(2.2, 5.2)	0.97	-4.1
Emergency visits	42	1.8	(0, 3.9)	44	2.9	(1.1, 4.8)	0.20	1.1
Adverse drug reactions	42	0.4	(-0.4, 1.2)	44	0.9	(-0.1, 1.9)	0.24	0.4
Direct total costs (with acupuncture costs)	42	506.4	(463.7, 548.9)	44	64.4	(20.7, 108.2)	0.13	-441.9
Direct total costs (without acupuncture costs)	42	51.9	(29.1, 74)	44	64.4	(24, 105.6)	0.30	12.5
Indirect costs								
Loss of income (acupuncture)	42	12.3	(0.1, 24.9)					
Productivity loss (acupuncture)	42	39.9	(20.3, 60.2)					
Loss of income	42	14.0	(-12.5, 39.9)	44	4.0	(-2.3, 10.4)	0.76	-10.0
Productivity loss	42	123.3	(-19.1, 271.1)	44	216.3	(-1.4, 445.1)	0.24	92.9
Indirect total costs (with acupuncture costs)	42	189.7	(245.1, 564.3)	44	220.3	(-6.6, 452.1)	0.90	30.6
Indirect total costs (without acupuncture costs)	42	137.4	(-2.1, 275.9)	44	220.3	(-1.7, 449.4)	0.26	82.9
Total costs (with acupuncture costs)	42	696.1	(737.4, 1,085.4)	44	284.7	(32.9, 523.7)	0.26	-411.3
Total costs (without acupuncture costs)	42	189.3	(44.1, 334.4)	44	284.7	(34, 533.9)	0.25	95.4
6-month follow-up								
Direct costs								
Medication reimbursement	42	15.5	(6.6, 24.3)	44	44.8	(-5.8, 94.6)	0.12	29.3
Medication co-payment	42	20.7	(2.9, 39)	44	12.0	(4.1, 19.7)	0.81	-8.6
Outpatient visits	42	3.2	(2.2, 4.2)	44	3.7	(2.6, 4.9)	0.25	0.5
Emergency visits	42	4.2	(1.1, 7.4)	44	6.1	(2.3, 9.6)	0.22	1.8
Adverse drug reactions	42	0.1	(-0.1, 0.5)	44	0.3	(-0.3, 1.1)	0.28	0.2
Travel costs	42	7.7	(2.2, 2)	44	4.6	(1.8, 0.0295)	0.86	-3.1
Direct total costs	42	51.7	(28, 76)	44	71.9	(21.8, 124.4)	0.23	20.1
Indirect costs								
Loss of income	42	3.6	(-3, 10.2)	44	8.3	(0.5, 15.9)	0.18	4.6
Productivity loss	42	10.2	(2.7, 18)	44	51.6	(-10.2, 117.8)	0.085	41.4
Indirect total costs	42	13.9	(6.6, 24.3)	44	59.9	(-5.8, 94.6)	0.075	46.0
Total costs	42	65.7	(2.9, 39)	44	131.9	(4.1, 19.7)	0.071	66.2

Note: Mean_{control} – Mean_{acup}, difference between the control group and the group with acupuncture treatment.

Table 6 Sensitivity analysis

Acupuncture group N=42	Control group N=44		Bootstrap t, P-value (one tailed)		
	Mean	95% CI	Mean	95% CI	P-value
Direct costs (min price of acupuncture treatment in the Czech Republic)	398.0	(355.2, 440.6)	64.4	(20.7, 108.2)	0.999
Total cost (min price of acupuncture treatment in the Czech Republic) ^a	805.7	(627, 979.8)	284.7	(32.9, 523.7)	0.970
Direct costs (max price of acupuncture treatment in the Czech Republic)	764.9	(725.8, 806.2)	64.4	(20.7, 108.2)	0.999
Total cost (max price of acupuncture treatment in the Czech Republic) ^a	1,172.6	(1,000.9, 1,346.9)	284.7	(32.9, 523.7)	0.999

Notes: ^aCosts are expressed in 2015/2016 EURO and are varying from €11.0 to €35.0 (cost of acupuncture session) at 0% discount rate.

visits, as well as adverse drug reactions, were nearly constant in both groups over time.

Patient's perspective

From patient's perspective, the essential costs involve the costs for acupuncture therapy (€308.5 on average), the travel expenses related to treatment (€120.8, on average), and the loss of income, which was monitored separately for AG and not in the CG, thus increasing the direct costs in the AG substantially. Other travel costs and medication co-payment, in absolute values, also are slightly higher in the AG, both during the acupuncture treatment period and at the 6-month follow-up. Loss of income is slightly higher in the AG as well but only within the acupuncture treatment period, in the 6-month follow-up these costs are lower.

Societal perspective

From society's perspective, it is clear that the AG shows a lower productivity lost, representing €92.9 during the acupuncture treatment period and €41.4 at the 6-month follow-up. In this regard, it is a positive phenomenon from a societal viewpoint, although the difference in values is not significant.

Sensitivity analysis

There was no significant difference between AG and CG (Table 6). Changes could be observed only in the absolute cost values.

Discussion

The present study compares the costs of acupuncture treatment in addition to standard care in patients with migraine after a 3-month course and a 6-month follow-up, considering the perspectives of a third-party payer, the patient, and the society. No significant differences in both direct and indirect costs were observed between the groups after acupuncture treatment and a 6-month follow-up. Nevertheless, it can be stated that AG presented lower costs for productivity losses

and total indirect costs in comparison with the standard pharmacologic therapy at all time periods. As far as total direct costs are concerned, during the acupuncture treatment period they were higher in the AG, as it was significantly influenced by the inclusion of acupuncture costs per session and the related travel costs. In addition, it was also a double entrance medical examination, both by an internal medicine doctor and an acupuncture specialist, which is a specific feature of the Faculty Hospital Hradec Kralove. The study, nevertheless, points out a potential trend in decreased direct costs in connection with reductions in medication intake, adverse drug reactions, and ER visits by patients who took acupuncture sessions. However, further study in this respect is needed.

Similar trends have been described in other studies as well; a review by Ambrósio et al¹⁰ evaluated seven cost-utility and one cost-effectiveness analyses of acupuncture in the treatment of chronic pain. The conditions treated included low back pain, neck pain, dysmenorrhea, migraine and headache, and osteoarthritis. Only one cost-effectiveness study indicated that there might be both clinical benefits and cost savings associated with acupuncture when concerning migraine;²⁸ however, another six studies indicated that the costs of acupuncture treatment are higher. Notwithstanding, a systematic review by Kim et al⁹ looking at economic evaluations, including cost-effectiveness, cost-utility, and cost-benefit analyses, alongside randomized controlled trials assess the consequences and costs of acupuncture for any medical condition. The cost-effectiveness analysis proved acupuncture to be beneficial at a relatively low cost in six European and Asian studies. However, only one of these studies focused on migraine, demonstrating a positive financial effect on patients in migraine treatment.²⁸ The cost estimates in this study included acupuncture, physician visits, hospital stays, prescription medications, and the patients' incapacity to work. Deng et al¹¹ evaluated different types of economic evaluations for acupuncture treatments for migraine in China. The data were obtained from

a multicenter, randomized controlled trial of acupuncture treatment in patients with migraine. Four-hundred eighty migraineurs were randomly assigned to three branches of treatments with true acupoints. Treatments using specific acupoints in Shaoyang meridians are more cost-effective than that of non-acupoints, representing a dramatic improvement in the quality of life of people with migraine and a significant reduction in costs.

The majority of the above-mentioned studies, including the present one, have concluded that although acupuncture increases health care costs the additional cost can be justified by the associated short- and long-term improvements in clinical outcomes. A study by Wonderling suggested the consideration of acupuncture as a viable intervention for migraine prophylaxis because it was shown to be a cost-effective therapy, relative to a number of interventions recommended by the National Institute for Health and Care Excellence, which are provided by the National Health Service system.²⁹ Such conclusion was also reached by Witt et al in the case of patients with primary headache where acupuncture was shown to be a cost-effective treatment according to international cost-effectiveness threshold values.³⁰

The present study was limited by the absence of a double-blinded method and a small sample size. On the other hand, an advantage of the study is the prospective evaluation of short- and long-term costs alongside a randomized controlled clinical trial. The present study also examined more items involved in direct and indirect costs when compared to the above-mentioned studies. For instance, in comparison with the studies from the review by Ambrósio et al,¹⁰ our study provides additional data on emergency visits, adverse drug reactions, travel costs, productivity loss, and loss of income. In addition, both direct and indirect costs were analyzed from all three viewpoints, that is, those of the patient, a third-party payer, and of society. In conclusion, a 3-month acupuncture course resulted in significant short- and long-term clinical benefits, while it suggested that acupuncture could be a financially attractive treatment option in a longer term, particularly in the case of indirect costs when compared with the absolute values of direct costs. This trend was observed also at a 6-month follow-up, although it was not significant. In the context of what has been hitherto known from the literature about the cost-effectiveness of acupuncture in migraine prophylaxis, the integration of acupuncture into health care is beneficial for society primarily when concerning lower productivity loss and loss of income, which exceed the costs of acupuncture alone. It is, therefore, reasonable

to recommend acupuncture as an adjuvant treatment to the standard pharmacologic therapy in migraine prophylaxis.

Acknowledgments

The authors thank Professor Bohuslav Manek and Doctor Daniel Diaz for translation and proofreading our manuscript and Marketa Talackova for collection of medication pharmacy retail prices. This work was supported by the project (Ministry of Health, Czech Republic) for conceptual development of research organization 00179906 (MH CZ – DRO; grant number UHHK, 00179906).

Disclosure

The authors report no conflicts of interest in this work.

References

- Steiner TJ, Stovner LJ, Birbeck GL. Migraine: the seventh disabler. *Cephalalgia*. 2013;33(5):289–290.
- Diluca M, Olesen J. The cost of brain diseases: a burden or a challenge? *Neuron*. 2014;82(6):1205–1208.
- WHO. Fact sheet: Headache disorders. Available from: <http://www.who.int/mediacentre/factsheets/fs277/en/>. Accessed September 23, 2018.
- Linde K, Allais G, Brinkhaus B, et al. Acupuncture for the prevention of episodic migraine. *Cochrane Database Syst Rev*. 2016;6(6):CD001218.
- Coeytaux RR, Befus D. Role of acupuncture in the treatment or prevention of migraine, tension-type headache, or chronic headache disorders. *Headache*. 2016;56(7):1238–1240.
- Kim MR, Shin JS, Lee J, et al. Safety of acupuncture and pharmacopuncture in 80,523 musculoskeletal disorder patients: a retrospective review of internal safety inspection and electronic medical records. *Medicine*. 2016;95(18):e3635.
- Zhang J, Shang H, Gao X, Ernst E. Acupuncture-related adverse events: a systematic review of the Chinese literature. *Bull World Health Organ*. 2010;88(12):915–921.
- Macpherson H, Vertosick EA, Foster NE, et al. The persistence of the effects of acupuncture after a course of treatment: a meta-analysis of patients with chronic pain. *Pain*. 2017;158(5):784–793.
- Kim SY, Lee H, Chae Y, Park HJ, Lee H. A systematic review of cost-effectiveness analyses alongside randomised controlled trials of acupuncture. *Acupunct Med*. 2012;30(4):273–285.
- Ambrósio EM, Bloor K, Macpherson H. Costs and consequences of acupuncture as a treatment for chronic pain: a systematic review of economic evaluations conducted alongside randomised controlled trials. *Complement Ther Med*. 2012;20(5):364–374.
- Deng ZQ, Zheng H, Zhao L, et al. Health economic evaluation of acupuncture along meridians for treating migraine in China: results from a randomized controlled trial. *BMC Complement Altern Med*. 2012;12:75.
- Evers S, Afra J, Frese A, et al; Members of the task force. EFNS guideline on the drug treatment of migraine – report of an EFNS task force. *Eur J Neurol*. 2006;13(6):560–572.
- Musil F, Pokladnikova J, Pavelek Z, et al. Acupuncture in migraine prophylaxis in Czech patients: an open-label randomized controlled trial. *Neuropsychiatr Dis Treat*. 2018;14:1221–1228.
- Headache Classification Subcommittee of the International Headache Society (IHS). The international classification of headache disorders 3rd edition. *Cephalalgia*. 2013;33:629–808.

15. Stewart WF, Lipton RB, Dowson AJ, Sawyer J. Development and testing of the Migraine Disability Assessment (MIDAS) Questionnaire to assess headache-related disability. *Neurology*. 2001;56(6 Suppl 1): S20–S28.
16. Drummond MF, O'Brien BJ, Stoddart GL. *Method for the Economic Evaluation of Healthcare Programmes*. Oxford, UK: Oxford University Press; 1997.
17. Liljas B. How to calculate indirect costs in economic evaluations. *Pharmacoeconomics*. 1998;13(1 Pt 1):1–7.
18. Wimo A, Reed CC, Dodel R, et al. The GERAS Study: a prospective observational study of costs and resource use in community dwellers with Alzheimer's disease in three European countries – study design and baseline findings. *J Alzheimers Dis*. 2013;36(2):385–399.
19. Pokladnikova J, Krcmova I, Vlcek J. Economic evaluation of sublingual vs subcutaneous allergen immunotherapy. *Ann Allergy Asthma Immunol*. 2008;100(5):482–489.
20. Maresova P, Klimova B, Novotny M, Kuca K. Alzheimer's and Parkinson's diseases: expected economic impact on Europe-a call for a uniform European strategy. *J Alzheimers Dis*. 2016;54(3): 1123–1133.
21. WHO Collaborating Centre for Drug Statistics Methodology. ATC/DDD Index 2017. Available from: https://www.whocc.no/atc_ddd_index/. Accessed September 23, 2018.
22. General Health Insurance Fund of the Czech Republic. Available from: <https://www.vzp.cz/poskytovatele/ciselniky/zdravotni-vykony>. Accessed September 23, 2018.
23. General Health Insurance Fund of the Czech Republic. Available from: <https://www.vzp.cz/poskytovatele/ciselniky/hromadne-vyrabene-lecive-pripravky-a-potravin-y-pro-zvlastni-lekarske-ucely/archiv>. Accessed September 23, 2018.
24. Ministry of Labour and Social Affairs of the Czech Republic. Available from: <http://www.mpsv.cz/cs/78>. Accessed September 23, 2018.
25. CSU. Czech Statistical Office. Available from: <https://www.czso.cz/csu/czso/makroekonomika-fqhqlzbynt>. Accessed September 23, 2018.
26. CSU. Czech Statistical Office. Available from: [http://www.czso.cz/csu/redakce.nsf/i/tab_2_ekonomika_a_infrastruktura_zu/\\$File/2zu_eki.xls](http://www.czso.cz/csu/redakce.nsf/i/tab_2_ekonomika_a_infrastruktura_zu/$File/2zu_eki.xls). Accessed September 23, 2018.
27. Heyse J, Cook J, Carides G. Statistical considerations in analysing healthcare resource utilization and cost data. In: Drummond M, McGuire A, editors. *Economic Evaluation in Healthcare: Merging Theory with Practice*. New York, NY: Oxford University Press; 2001: 215–235.
28. Liguori A, Petti F, Bangrazi A, et al. Comparison of pharmacological treatment versus acupuncture treatment for migraine without aura-analysis of socio-medical parameters. *J Tradit Chin Med*. 2000; 20(3):231–240.
29. Wonderling D, Vickers AJ, Grieve R, McCarney R. Cost effectiveness analysis of a randomised trial of acupuncture for chronic headache in primary care. *BMJ*. 2004;328(7442):747.
30. Witt CM, Reinhold T, Jena S, Brinkhaus B, Willich SN. Cost-effectiveness of acupuncture treatment in patients with headache. *Cephalalgia*. 2008;28(4):334–345.

Neuropsychiatric Disease and Treatment

Publish your work in this journal

Neuropsychiatric Disease and Treatment is an international, peer-reviewed journal of clinical therapeutics and pharmacology focusing on concise rapid reporting of clinical or pre-clinical studies on a range of neuropsychiatric and neurological disorders. This journal is indexed on PubMed Central, the 'PsycINFO' database and CAS,

Submit your manuscript here: <http://www.dovepress.com/neuropsychiatric-disease-and-treatment-journal>

Dovepress

and is the official journal of The International Neuropsychiatric Association (INA). The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.