Molecular epidemiology and clinical significance of *Corynebacterium striatum* isolated from clinical specimens

Jin Woong Suh¹

Yongguk Ju²

Chang Kyu Lee³

Jang Wook Sohn^{1,2}

Min Ja Kim^{1,2}

Young Kyung Yoon^{1,2}

¹Division of Infectious Diseases, Department of Internal Medicine, Korea University College of Medicine, Seoul, Republic of Korea; ²Institute of Emerging Infectious Diseases, Korea University College of Medicine, Seoul, Republic of Korea; ³Department of Laboratory Medicine, Korea University College of Medicine, Seoul, Republic of Korea

Purpose: This study investigated the clinical epidemiology, antimicrobial susceptibility, and molecular epidemiology of *Corynebacterium striatum* isolates.

Patients and methods: An observational study was conducted at a university hospital in the Republic of Korea from August to December 2016. All subjects were patients who tested positive for *C. striatum* clinically. Clinical data were analyzed to evaluate the microbiological and genotypic characteristics of *C. striatum* strains.

Results: Sixty-seven *C. striatum* isolates recovered from non-duplicated patients were characterized. Patients were classified into three groups according to the infection type: nosocomial infection (71.6%), health care-associated infection (8.7%), and community-acquired infection (18.8%). The most common clinical specimens were urine (35.8%) and skin abscesses (32.8%). Fifty-two (77.6%) isolates showed multidrug resistance, defined as resistance to ≥ 3 different antibiotic families. All strains were susceptible to vancomycin and linezolid. Resistance to other antibiotics varied: penicillin (n=65; 97.0%), ampicillin (n=63; 94.0%), cefotaxime (n=64; 95.5%), and levofloxacin (n=61; 91.0%). Phylogenetic analysis identified all 16 S rRNA gene sequences of the 67 isolates as those of *C. striatum*, where 98%–99% were homologous to *C. striatum* ATCC 6940. In multilocus sequence typing for internal transcribed spacer region, *gyrA*, and *rpoB* sequencing, the most predominant sequence types (STs) were ST2, ST3, ST6, and ST5.

Conclusion: *C. striatum* isolates may cause opportunistic infections associated with nosocomial infections through horizontal transmission. The presence of multidrug resistance and intra-hospital dissemination implicate *C. striatum* isolates as a potential target pathogen for infection control and antimicrobial stewardship programs.

Keywords: *Corynebacterium striatum*, multidrug resistant, multilocus sequence typing, opportunistic infections, nosocomial infections

Introduction

Most catalase-positive, gram-positive rods, commonly called coryneform or diphtheroid bacteria, have historically been considered simple contaminants that are unlikely to be pathogenic, when isolated from clinical specimens. Of the >80 species of *Corynebacterium* that have been reported, ~50 species rarely cause infectious diseases in humans.¹ *Corynebacterium striatum* is one of the more commonly isolated coryneform bacteria in clinical microbiology laboratories.

In the last decade, *C. striatum* has been frequently cultured from various surfaces and medical equipment in hospital settings.² In clinical settings, *C. striatum* is increasingly being recognized as a source of opportunistic diseases in immunocompromised

Correspondence: Young Kyung Yoon
Division of Infectious Diseases,
Department of Internal Medicine,
Korea University Anam Hospital, Korea
University College of Medicine, Incheon-ro
73, Seongbuk-gu, Seoul 02841, Republic
of Korea
Tel +82 2 920 5341
Fax +82 2 920 5616
Email young7912@korea.ac.kr

patients suffering from malignancies, human immunodeficiency virus infection, or chronic lung diseases, and in patients wearing prosthetic devices. Furthermore, *C. striatum* is potentially pathogenic in patients with chronic diseases, who are exposed to specific circumstances, such as invasive medical procedures, prolonged use of broad-spectrum antibiotics, or long-term hospitalization.³⁻⁵

Occasionally, *C. striatum* strains are isolated in polymicrobial infections, where their degree of virulence remains largely undetermined. In reality, it is difficult to distinguish between a pathogen causing infection and one causing colonization. Although the clinical significance and prevalence of *C. striatum* remain unclear, this organism has been responsible for a variety of infections, such as bacteremia, arthritis, osteomyelitis, meningitis, endocarditis, breast abscess, peritonitis, wound infections, and prosthetic joint infections in both immunocompetent and immunocompromised patients.³⁻⁶

Infections caused by *C. striatum* are usually considered to originate endogenously. However, recent studies have documented the possibility of patient-to-patient transmission of *C. striatum*. The bacterium may cause serious nosocomial infections in intensive care unit patients and spread from patient to patient via physical contact with attending personnel or via the nosocomial environment itself.⁷⁻⁹

Despite early reports of susceptibility to a wide range of antibiotics, multidrug-resistant phenotypes have been recently reported in most *C. striatum* strains.^{2,10} Moreover, multidrug-resistant *C. striatum* may cause nosocomial transmissions resulting in outbreaks in patients with specific risk factors, leading to increased mortality.^{2,11}

There is limited information on the pathogenicity and clinical implications associated with *C. striatum* strains in a clinical setting. The aim of this study was to investigate the molecular characteristics, clinical significance, and antimicrobial susceptibility of *C. striatum*, with the goal of increased understanding of its pathogenicity.

Patients and methods

Study design

An observational study was conducted at the Korea University Anam Hospital, a 1,048-bed university hospital in Seoul, Republic of Korea (ROK) from August 2016 to December 2016. All subjects were adult patients (>18 years) whose clinical isolates obtained from diagnostic cultures of clinical specimens tested positive for *C. striatum*. If multiple *C. striatum* isolates were recovered from a patient, only the first isolate was included in the study.

This study was performed in compliance with the Declaration of Helsinki. The study protocol was approved by the Institutional Review Board of Korea University Anam Hospital (No. 2018AN0161). As this observational study did not entail any deviations from routine medical practice, the requirement for informed consent was waived.

Clinical data

Clinical data manually extracted from medical records included age, gender, comorbidities, clinical diagnoses, specimen categories, Charlson's comorbidity index, microbiological data, and treatment outcomes. Determination of the clinical significance of *C. striatum* isolates was based on clinical findings, such as fever, white blood cell counts, and C-reactive protein, in addition to whether the patient had or had not received antibiotics for *C. striatum*. Infections were categorized as community acquired, health care associated, or nosocomial, depending on the location that the strains were isolated from. Treatment outcomes were classified on the basis of in-hospital mortality and median length of hospitalization since *C. striatum* isolation. Additionally, demographic, clinical, and microbiological characteristics between the antibiotic treatment group and the observation group were compared.

C. striatum identification

Processing and incubation of all clinical samples was performed according to routine laboratory protocols. *C. striatum* isolates were initially identified using the MicroScan WalkAway-96 Plus system (Beckman Coulter, Inc., Brea, CA, USA) and further confirmed by sequencing of the entire 16S rRNA gene.

The *C. striatum* strains were cultured on blood agar plates for 18 hours at 37°C. The cultured colony was suspended in 200 µL of lysis buffer, incubated overnight at 37°C, and centrifuged for 5 minutes at 3,000 rpm. The pelleted bacteria were suspended in 500 µL of sterile water and boiled for 15 minutes for DNA extraction. *C. striatum* DNA was extracted using a bacterial AccuPrep Genomic DNA Extraction Kit (Bioneer, Daejeon, ROK). To amplify 16S rRNA, universal primers 16SF27 (5'-AGAGTTTGATCMTGGCTCAG) and 16SR1492 (5'-TACGGYTACCTTGTTACGACTT) were used, as previously described.¹² The purified PCR product was sequenced using primer walking with the oligonucleotides using the primers 16SF518 (5'-CCAGCAGCCGCGGTA-ATAC) and 16SR800 (5'-TACCAGGGTATCTAATCC).¹³

The amplified sequence was compared with those available in the GenBank database using the BLAST program.

MicroScan WalkAway-96 Plus system (Beckman Coulter, Inc.), which are considered the gold standard culture media for *Corynebacterium* as recommended by the Clinical and Laboratory Standards Institute (CLSI). The minimum inhibitory concentration (MIC) breakpoint for *Corynebacterium* (CLSI document M45-A2) was applied for the analysis of antimicrobial susceptibility.¹⁵

Multilocus sequence typing (MLST)

For MLST, the internally transcribed spacer 1 (ITS1) region, as well as *gyrA* and *rpoB* were amplified and sequenced for *C. striatum* strains.² The primers used are provided in Table S1. PCR amplification and sequence reaction were performed as previously described.^{16–18}

Notes: Distance estimations were calculated using the Kimura two-parameter model. Bootstrap percentages after 1,000 simulations are shown. The *Actinomyces bovis* (T X8106) sequence was used as an outgroup.

Statistical analyses

Categorical variables are expressed as frequencies and were analyzed using Pearson's chi-squared test or Fisher's exact test. Continuous variables are expressed as medians and IQRs. The two-sample *t*-test or Mann–Whitney *U* test was used as appropriate to compare continuous variables between groups. Significance was set at $P < 0.05$. SPSS Statistics, version 24.0 (IBM Corporation, Armonk, NY, USA) and SAS 9.4 (SAS Institute Inc., Cary, NC, USA) were used for data analyses.

Results

Patients and clinical characteristics

A total of 67 *C. striatum* isolates were identified in different patients. Demographic and baseline characteristics of the 67 patients are summarized in Table 1. Twenty (29.9%) patients had received antibiotics for *C. striatum* isolated

from infection sites. Univariate analysis did not indicate significant differences in age, gender, category of infection, and comorbid diseases between the antibiotic treatment and the observation groups (Table 1).

C. striatum was isolated from various types of specimens (Table 2). Urine isolates (35.8%) were the most common, but only 10% of these came from those who received antibiotic therapy. Antibiotics were most commonly used in skin abscess isolates (55.0%; Table 2). Of the 67 specimens, 25 (38.8%) were polymicrobial isolates (Table 2). Among these, *Escherichia coli* (16.1%), carbapenem-resistant *Acinetobacter baumannii* (16.1%), and *Pseudomonas* species (16.1%) were the most common simultaneously isolated strains.

Antimicrobial susceptibility

All strains were susceptible to erythromycin, vancomycin, linezolid, and daptomycin (Table 3). However, intermediate

Table 1 Demographic and basic characteristics of 67 patients whose clinical specimens tested positive for *Corynebacterium striatum*

Variables	Total (N=67)	Treatment (n=20, 29.8%)	Observation (n=47, 70.1%)	P-value
Male, n (%)	35 (52.2)	14 (70.0)	21 (44.7)	0.058
Age (years), median (IQR)	64 (55–78)	65 (59–75)	69 (55–78)	0.564
Length from admission to <i>C. striatum</i> isolation (days), median (IQR)	15 (3–33)	12 (0–25)	20 (4–37)	0.615
Category of infection, n (%)				0.975
Community onset	13 (18.8)	4 (20.0)	9 (19.1)	
Hospital acquired	6 (8.7)	2 (1.0)	4 (8.5)	
Nosocomial	48 (71.6)	14 (70.0)	34 (72.3)	
Comorbid illness, n (%)				
Cardiovascular	22 (32.8)	6 (30.0)	16 (34.0)	0.785
Central nervous system	29 (43.3)	5 (25.0)	24 (51.0)	0.062
Malignancy	15 (22.4)	5 (25.0)	10 (21.3)	0.756
Renal	4 (6.0)	0	4 (8.5)	0.309
Hepatic	5 (7.5)	2 (10.0)	3 (6.4)	0.631
Respiratory	1 (1.5)	0	1 (2.1)	1.000
Hematology	1 (1.5)	0	1 (2.1)	1.000
Transplantation	1 (1.5)	0	1 (2.1)	1.000
Diabetes mellitus	18 (26.9)	7 (35.0)	11 (23.4)	0.374
Charlson's comorbidity index, median (IQR)	2 (1–3)	2 (0–3)	2 (1–3)	0.503
Risk factors, n (%)				
Recent admission	54 (80.6)	12 (60.0)	42 (89.4)	0.392
Surgery	24 (35.8)	7 (35.0)	17 (36.2)	1.000
Prior use of antibiotics, n (%)	51 (76.1)	14 (70.0)	37 (78.7)	0.534
Steroid	8 (11.9)	3 (15.0)	5 (10.6)	0.687
Foley catheter	33 (49.3)	8 (40.0)	25 (53.2)	0.425
Mechanical ventilator	27 (40.3)	6 (30.0)	21 (44.7)	0.291
ICU care	23 (34.3)	6 (30.0)	17 (36.2)	0.780
Clinical findings				
Fever, n (%)	2 (3.0)	0	2 (4.3)	0.380
WBC ($\times 10^3/\mu\text{L}$), median (IQR)	9.8 (6.1–11.6)	12.1 (6.3–13.2)	8.9 (5.3–11.6)	0.472
CRP (mg/L), median (IQR)	75 (15–130)	109 (31–186)	59 (9–104)	0.437
Procalcitonin (ng/mL), median (IQR)	2.1 (0.1–2.9)	1.7 (0.1–1.6)	1.7 (0.1–1.6)	0.484

Abbreviations: CRP, C-reactive protein; ICU, intensive care unit; WBC, white blood cell.

Table 2 Microbiological characteristics and treatment outcomes of 67 patients whose clinical specimens tested positive for *Corynebacterium striatum*

Parameters	Total (N=67)	Treatment (n=20, 29.8%)	Observation (n=47, 70.1%)	P-value
Category of specimens, n (%)				0.005
Urine	24 (35.8)	2 (10.0)	22 (46.8)	
Skin abscess	22 (32.8)	11 (55.0)	11 (23.4)	
Otitis media	10 (14.9)	2 (10.0)	8 (17.0)	
Blood	6 (9.0)	1 (5.0)	5 (10.6)	
Ascites	4 (6.0)	3 (15.0)	1 (2.1)	
Pleural effusion	1 (1.5)	1 (5.0)	0	
Polymicrobial infections, n (%)	25 (38.8)	13 (65.0)	12 (25.5)	0.003
2	19 (76.0)	9 (69.2)	10 (83.3)	
≥3	6 (24)	4 (30.8)	2 (16.7)	
Distribution of simultaneously isolated strain, n (%)				
<i>Escherichia coli</i>	5 (16.1)	3 (17.6)	2 (14.3)	0.126
Carbapenem-resistant <i>Acinetobacter baumannii</i>	5 (16.1)	4 (23.5)	1 (7.1)	0.011
<i>Pseudomonas</i> species	5 (16.1)	5 (29.4)	0	
Methicillin-resistant <i>Staphylococcus aureus</i>	3 (9.7)	0	3 (21.4)	0.248
<i>Enterococcus</i> species	3 (9.7)	0	3 (21.4)	0.248
Methicillin-susceptible <i>Staphylococcus aureus</i>	2 (6.5)	2 (11.8)	0	0.028
Methicillin-resistant coagulase-negative <i>S. aureus</i>	2 (6.5)	0	2 (14.3)	0.349
<i>Streptococcus</i> species	2 (6.5)	1 (5.9)	1 (7.1)	0.527
<i>Enterobacter</i> species	1 (3.2)	1 (5.9)	0	0.122
<i>Proteus</i> species	1 (3.2)	1 (5.9)	0	0.122
<i>Klebsiella</i> species	1 (3.2)	0	1 (7.1)	0.511
<i>Candida</i> species	1 (3.2)	0	1 (7.1)	0.511
Antibiotic susceptibility, n (%)				
Penicillin	2 (3.0)	0 (0)	2 (4.3)	0.525
Erythromycin	67 (100)	20 (100)	47 (100)	
Levofloxacin	6 (9.0)	3 (15.0)	3 (6.4)	0.845
Vancomycin	67 (100)	20 (100)	47 (100)	
Linezolid	67 (100)	20 (100)	47 (100)	
Multidrug resistance, n (%)	52 (77.6)	14 (70.0)	38 (80.9)	0.106
Treatment				
In-hospital mortality, n (%)	1 (1.5)	1 (5.0)	0	0.290
Length of hospital stay since <i>C. striatum</i> isolation, median days (IQR)	35 (5–77)	31 (1–59)	31 (1–59)	0.695

Table 3 MIC₅₀ and MIC₉₀ values and antimicrobial susceptibility (%) of the clinical isolates of *Corynebacterium striatum*

Antibiotics	MIC (mg/L)			Percentage of susceptible isolates
	MIC ₅₀	MIC ₉₀	Range	
Penicillin	>4	>4	0.12–>4	3.0
Ampicillin	>4	>4	0.25–4	6.0
Cefotaxime	>2	>2	1–>2	4.5
Levofloxacin	>4	>4	≤0.5–>4	9.0
Erythromycin	>0.5	>0.5	≤0.06–>0.5	100
Linezolid	≤1	≤1	≤1	100
Vancomycin	0.5	1	≤0.25–1	100
Daptomycin	<0.25	<0.25	<0.25	100

Abbreviations: MIC, minimum inhibitory concentration; MIC₅₀, MIC for inhibition of 50% of isolates tested; MIC₉₀, MIC for inhibition of 90% of isolates tested.

to high levels of resistance to penicillin (97.0%), ampicillin (94.0%), cefotaxime (95.5%), and levofloxacin (91.0%) were observed (Table 2). Therefore, the 67 strains may be considered multidrug resistant, defined as resistance to ≥3 classes of antibiotics. There was no significant difference in the frequency of multidrug-resistant isolates between the treatment and observation groups (Table 2).

Based on resistance to penicillin, ampicillin, cefotaxime, levofloxacin, and vancomycin, the 67 *C. striatum* strains were classified into five patterns, from I to V (Table 4). Among these, pattern I, which was the multidrug-resistant phenotype susceptible to vancomycin, was the predominant in hospitalized patients (82.8%).

Table 4 Antimicrobial resistance patterns of 67 *Corynebacterium striatum* strains, categorized by susceptibility to penicillin, ampicillin, cefotaxime, levofloxacin, and vancomycin

Resistance pattern (n)	Patients (n)		Antibiotics				
	Inpatients	Outpatient	PEN	AMP	COF	LEV	VAN
I (58)	48	10	R(I)	R(I)	R(I)	R(I)	S
II (5)	I	4	R(I)	R(I)	R(I)	S	S
III (2)	I	I	S	S	S	R(I)	S
IV (1)	I	0	R(I)	S	R(I)	R(I)	S
V (1)	I	0	R(I)	S	R(I)	S	S

Abbreviations: AMP, ampicillin; COF, cefotaxime; I, intermediate; LEV, levofloxacin; PEN, penicillin; R, resistant; S, susceptible; VAN, vancomycin.

Clinical outcomes

Sepsis or septic shock was absent in all cases. Treatment outcomes of the 67 patients are shown in Table 2. Only one (1.5%) patient died during hospitalization. Death was due to a non-infectious cause. Univariate analysis showed no difference between in-hospital mortalities of the treatment and observation groups. The most commonly used therapeutic antibiotics were vancomycin, followed by tigecycline, fluoroquinolones, and piperacillin/tazobactam.

On the other hand, there were no deaths in the 42 patients who had only *C. striatum* isolated from clinical specimens. There was no difference in the length of hospital stay between the treatment and observation groups (9 [IQR, 0–15] vs 20 [IQR, 4–37] days, $P=0.049$).

MLST

Four genes (16S rRNA, ITS1, *gyrA*, and *rpoB*) were analyzed in all strains studied. 16S rRNA was excluded from the reinforcement analysis of MLST due to its high conservation among all strains analyzed. 16S rRNA was used to confirm the identity of *C. striatum* strains. ITS1, *gyrA*, and *rpoB* were used to discriminate among the strains, using a few nucleotide changes within sequences. Distinct allele sequences were assigned arbitrary allele numbers for each locus (Table S2).

In the ITS1 region, allele 2 was the most abundant (52.2%). For *gyrA*, allele 1 was predominant (89.6%). For *rpoB*, allele 2 was the most abundant and was found in 61 strains (91.0%). Twelve sequence types (STs) were identified in these three genes. Of these, the most abundant were ST2 (44.8%), ST3 (22.4%), ST6 (14.9%), and ST5 (4.5%). Among patients with ST2, there were 24 (80.0%) nosocomial infections and 6 (20.0%) community-onset infections. Finally, each ST was grouped and schematized (Figure 2) and analyzed for antibiotic resistance (Table S2). Antibiotic resistance Pattern I occupied an overwhelming number of

predominant STs: 28 (93.3%) in ST2, 13 (86.7%) in ST3, and 10 (100%) in ST6.

Discussion

This study analyzed the molecular epidemiology and clinical significance of *C. striatum* isolates recovered from clinical specimens. The data demonstrate the low virulence of *C. striatum*, reflecting its role as a colonizing opportunistic pathogen. However, the characteristics of multidrug resistance and horizontal transmission of *C. striatum* isolates suggest the possibility that it is an emerging nosocomial pathogen, which should be of interest to medical researchers.

C. striatum is reportedly susceptible to β -lactams.¹⁹ However, presently, most strains (95.5%) display a multidrug-resistant phenotype, although all strains are susceptible to erythromycin, vancomycin, linezolid, and daptomycin. If *C. striatum* isolates are, indeed, causative agents of infections, they may require the use of broad-spectrum antibiotics such as vancomycin.²⁰ Antimicrobial resistance is a major global health issue. In ROK, the prevalence of vancomycin-resistant *Enterococcus faecium* has gradually increased from 22% in 2003 to 31% in 2015.²¹ Since December 2010, infectious diseases caused by six types of multidrug-resistant bacteria, including vancomycin-resistant enterococci (VRE), were legally designated for surveillance as notifiable infectious microorganisms, with the enactment of the Infectious Disease Control and Prevention Act.²² To control antimicrobial resistance, contamination and infection by *C. striatum* isolates should be accurately determined to ensure optimal use of antibiotics.

Currently, there are no guidelines for the treatment of infections caused by *C. striatum*. Optimal antimicrobial therapy is still considered controversial. Our in vitro susceptibility tests demonstrate that vancomycin and linezolid are active against *C. striatum*, indicating their potential

Figure 2 The distribution of STs.

Abbreviation: ST, sequence type.

therapeutic value. However, in the absence of approved breakpoints for *Corynebacterium* species, those antibiotics recommended for *Staphylococcus* species have been used, except for penicillin and ampicillin, for which thresholds for *Listeria* species are used.^{2,23,24}

In our study, *C. striatum* was isolated from various clinical specimens. However, only 20 (29.9%) patients received antibiotic therapy. For several decades, *C. striatum* has been considered as having limited potential for pathogenicity. The present observations confirmed its low virulence. Similar to *C. striatum* strains, VRE, which exhibit low virulence and pathogenicity, were initially considered rare opportunistic pathogens. However, in recent years, the prevalence of VRE has increased and they have joined the list of significant pathogens that frequently cause nosocomial infections with limited treatment options.²⁵ Although several factors contribute to virulence, such as host defense mechanisms and the expression of various microorganism traits, a higher rate of multidrug-resistant bacteria may be associated with a higher rate of infection and mortality. Therefore, efforts to contain the spread of multidrug-resistant bacteria should be given prominence by the development of antimicrobial stewardship programs.

Since the first case of *C. striatum* infection in a patient with chronic lymphocytic leukemia was published in 1980, *C. striatum* strains have been considered as emerging pathogens in clinical settings.^{26,27} In recent years, they have been reported as opportunistic pathogens infecting immunocompromised patients with malignancies, COPD, cardiovascular diseases, and diabetes. Other known risk factors include long-term hospitalization, previous use of broad-spectrum antibiotics, and exposure to invasive devices.^{2,9–11} In our study, 52 patients (77.6%) presented at least one predisposing condition. This suggests that *C. striatum* may have a significant clinical impact on the patients hospitalized for a long term with comorbidity, or with a history of exposure to broad-spectrum antibiotics, or those who are immunosuppressed, as well as critically ill patients with an implanted indwelling device.

In our study, *C. striatum* strains mainly caused nosocomial infections, and the four predominant STs in MLST, which accounted for 86.6% of the strains, were ST2, ST3, ST6, and ST5. The 67 strains represent distinct allele combinations (13 STs, considering only three genes: ITS1, *gyrA*, and *rpoB*). For *gyrA*, allele 1 was predominant (89.6%). For *rpoB*, allele 2 was the most abundant and was found in 61 strains (91.0%). A literature review indicates that these characteristics of *C. striatum* strains may be a major concern for global health

institutions because it is an emergent Gram-positive environmental bacterium which is highly persistent, prevalent, and transmissible person to person and through caregivers, with significant multidrug resistance.² Furthermore, it has often caused nosocomial outbreaks.^{2,7–11}

Identification of *Corynebacterium* species is challenging because there are limitations to distinguishing *Corynebacterium* species based on biochemical profiles using the Api Coryne system. Thus, as in our study, molecular methods, such as gene sequencing, are used as the complementary gold standard for bacterial identification.^{28,29}

Our study had several limitations. Firstly, this study used a single-center design and included a small sample of specimens. Secondly, antibiotic treatment was decided by the treating physician based on his or her interpretation of the microbiological and clinical parameters used in general clinical practice. Finally, the antibiotic susceptibility test did not include gentamicin, as recommended by the CLSI guidelines. Therefore, the possibility that antibiotic resistance of the strains is more extensive cannot be excluded.

Conclusion

C. striatum isolates have been characterized as sources of opportunistic infections associated with nosocomial infections through horizontal transmission. They are only susceptible to a limited number of broad-spectrum antibiotics and can be isolated from clinical specimens due to colonization or contamination. Particularly, characteristics of multidrug resistance and intra-hospital dissemination indicate their potential to be considered as a target pathogen in antimicrobial stewardship programs.

Data sharing statement

Further information and requests for individual deidentified participant data sharing will be fulfilled by contacting the corresponding author YKY (young7912@korea.ac.kr) during the 5 years after publication. Clinical and molecular data and research ethics approval information can be shared.

Acknowledgments

This research was supported by a grant of the Korea Health Technology R&B Project through the Korea Health Industry Development Institute (KHIDI), funded by the Ministry of Health & Welfare, Republic of Korea (grant number: HI16C1048). The funders had no role in the study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Disclosure

The authors report no conflicts of interest in this work.

References

- Bernard K. The genus *Corynebacterium* and other medically relevant coryneform-like bacteria. *J Clin Microbiol*. 2012;50(10):3152–3158.
- Renom F, Gomila M, Garau M, et al. Respiratory infection by *Corynebacterium striatum*: epidemiological and clinical determinants. *New Microbes New Infect*. 2014;2(4):106–114.
- Martínez-Martínez L, Suárez AI, Rodríguez-Baño J, Bernard K, Muniáin MA. Clinical significance of *Corynebacterium striatum* isolated from human samples. *Clin Microbiol Infect*. 1997;3(6):634–639.
- Lee PP, Ferguson DA Jr, Sarubbi FA. *Corynebacterium striatum*: an underappreciated community and nosocomial pathogen. *J Infect*. 2005;50(4):338–343.
- Otsuka Y, Ohkusu K, Kawamura Y, Baba S, Ezaki T, Kimura S. Emergence of multidrug-resistant *Corynebacterium striatum* as a nosocomial pathogen in long-term hospitalized patients with underlying diseases. *Diagn Microbiol Infect Dis*. 2006;54(2):109–114.
- Funke G, von Graevenitz A, Clarridge JE 3rd, Bernard KA. Clinical microbiology of coryneform bacteria. *Clin Microbiol Rev*. 1997;10(1):125–159.
- Baio PV, Mota HF, Freitas AD, et al. Clonal multidrug-resistant *Corynebacterium striatum* within a nosocomial environment, Rio de Janeiro, Brazil. *Mem Inst Oswaldo Cruz*. 2013;108(1):23–29.
- Brandenburg AH, van Belkum A, van Pelt C, Bruining HA, Mouton JW, Verbrugh HA. Patient-to-patient spread of a single strain of *Corynebacterium striatum* causing infections in a surgical intensive care unit. *J Clin Microbiol*. 1996;34(9):2089–2094.
- Leonard RB, Nowowiejski DJ, Warren JJ, Finn DJ, Coyle MB. Molecular evidence of person-to-person transmission of a pigmented strain of *Corynebacterium striatum* in intensive care units. *J Clin Microbiol*. 1994;32(1):164–169.
- Verroken A, Bauraing C, Deplano A, et al. Epidemiological investigation of a nosocomial outbreak of multidrug-resistant *Corynebacterium striatum* at one Belgian university hospital. *Clin Microbiol Infect*. 2014;20(1):44–50.
- Wang J, Wang Y, Du X, et al. Rapid transmission of multidrug-resistant *Corynebacterium striatum* among susceptible patients in a tertiary hospital in China. *J Infect Dev Ctries*. 2016;10(12):1299–1305.
- Lane DJ. 16S/23S rRNA sequencing. In: Stackebrandt E, Goodfellow M, editors. *Nucleic Acid Techniques in Bacterial Systematics*. Chichester: John Wiley and Sons; 1991:115–175.
- Senthilraj R, Prasad GS, Janakiraman K. Sequence-based identification of microbial contaminants in non-parenteral products. *Braz J Pharm Sci*. 2016;52(2):329–336.
- Khamis A, Raoult D, La Scola B. rpoB gene sequencing for identification of *Corynebacterium* species. *J Clin Microbiol*. 2004;42(9):3925–3931.
- Clinical and Laboratory Standards Institute. *Methods for Antimicrobial Dilution and Disk Susceptibility Testing of Infrequently Isolated or Fastidious Bacteria: CLSI Guideline M45*. 3rd ed. Wayne (PA): Clinical and Laboratory Standard Institute; 2016.
- Guasp C, Moore ER, Lalucat J, Bannas A. Utility of internally transcribed 16S-23S rDNA spacer regions for the definition of *Pseudomonas stutzeri* genomovars and other *Pseudomonas* species. *Int J Syst Evol Microbiol*. 2000;50 Pt 4:1629–1639.
- Sierra JM, Martínez-Martínez L, Vázquez F, Giralte E, Vila J. Relationship between mutations in the gyrA gene and quinolone resistance in clinical isolates of *Corynebacterium striatum* and *Corynebacterium amycolatum*. *Antimicrob Agents Chemother*. 2005;49(5):1714–1719.
- Khamis A, Raoult D, La Scola B. rpoB gene sequencing for identification of *Corynebacterium* species. *J Clin Microbiol*. 2004;42(9):3925–3931.
- Soriano F, Zapardiel J, Nieto E. Antimicrobial susceptibilities of *Corynebacterium* species and other non-spore-forming gram-positive bacilli to 18 antimicrobial agents. *Antimicrob Agents Chemother*. 1995;39(1):208–214.
- Hahn WO, Werth BJ, Butler-Wu SM, Rakita RM. Multidrug-resistant *Corynebacterium striatum* associated with increased use of parenteral antimicrobial drugs. *Emerg Infect Dis*. 2016;22(11):1908–1914.
- Kim D, Ahn JY, Lee CH, et al. Increasing resistance to extended-spectrum cephalosporins, fluoroquinolone, and carbapenem in gram-negative bacilli and the emergence of carbapenem non-susceptibility in *Klebsiella pneumoniae*: analysis of Korean Antimicrobial Resistance Monitoring System (KARMS) data From 2013 to 2015. *Ann Lab Med*. 2017;37(3):231–239.
- Centers for Disease Control and Prevention. Case definitions for National Notifiable Infectious Diseases; 2011. Available from: http://www.cdc.gov/cdc/cms/content/58/12558_view.html. Accessed May 17, 2018.
- Weiss K, Laverdière M, Rivest R. Comparison of antimicrobial susceptibilities of *Corynebacterium* species by broth microdilution and disk diffusion methods. *Antimicrob Agents Chemother*. 1996;40(4):930–933.
- Campanile F, Carretto E, Barbarini D, et al. Clonal multidrug-resistant *Corynebacterium striatum* strains, Italy. *Emerg Infect Dis*. 2009;15(1):75–78.
- Chiang HY, Perencevich EN, Nair R, et al. Incidence and outcomes associated with infections caused by vancomycin-resistant *Enterococci* in the United States: systematic literature review and meta-analysis. *Infect Control Hosp Epidemiol*. 2017;38(2):203–215.
- Bowstead TT, Santiago SM. Pleuropulmonary infection due to *Corynebacterium striatum*. *Br J Dis Chest*. 1980;74(2):198–200.
- Alibi S, Ferjani A, Boukadida J, et al. Occurrence of *Corynebacterium striatum* as an emerging antibiotic-resistant nosocomial pathogen in a Tunisian hospital. *Sci Rep*. 2017;7(1):9704.
- Gomila M, Renom F, Gallegos M del C, et al. Identification and diversity of multiresistant *Corynebacterium striatum* clinical isolates by MALDI-TOF mass spectrometry and by a multigene sequencing approach. *BMC Microbiol*. 2012;12:52.
- Alibi S, Ferjani A, Gaillot O, Marzouk M, Courcol R, Boukadida J. Identification of clinically relevant *Corynebacterium* strains by Api Coryne, MALDI-TOF-mass spectrometry and molecular approaches. *Pathol Biol (Paris)*. 2015;63(4–5):153–157.

Supplementary materials

Table S1 Primers used in the molecular analysis of 67 *Corynebacterium striatum* strains

Gene	Primer	Sequence (5'→3')	Function	T ^a (°C)	Length (bp)	Reference
16S rRNA	I6SF27	AGAGTTTGATCMTGGCTCAG	16S rDNA	55	1,400	26
	I6SR1492	TACGGYTACCTTGTACGACTT				
16S rRNA	I6SF518	CCAGCAGCCGCGGTAATAC	16S rDNA	—	—	27
(sequencing)	I6SR800	TACCAGGGTATCTAATCC				
ITS1	I6F945	GGGCCCCGACAAGCGGTGG	Interspace sequence	55	900	9
	23r458	CTTCCCTCACGGTAC	region I			
gyrA	gyrA1	GCGGCTACGTAAAGTCC	Gyrase	55	400	9
	gyrA2	CCGCCGGAGCCGTTTCAT				
rpoB	C2700F	CGWATGAACATYGGBCAGGT	β-subunit of RNA	60	400	9
	C3130R	TCCATYTCRCCRAARCGCTG	polymerase enzyme			

Table S2 STs at three examined loci in the *Corynebacterium striatum* strain and patterns of antimicrobial resistance

Strain	16S rRNA	ITS1	gyrA	rpoB	ST ^a	Resistance pattern
1	1	2	1	2	2	I
2	1	4	1	2	3	I
3	2	2	1	2	2	II
4	1	2	2	2	4	I
5	1	4	1	2	3	I
6	1	2	1	2	2	I
7	1	4	2	2	5	II
8	3	4	1	2	3	I
9	1	2	1	2	2	I
10	4	3	1	2	6	I
11	5	2	1	2	2	I
12	1	4	1	2	3	I
13	1	4	1	2	3	I
14	1	2	1	2	2	I
15	1	4	1	2	3	I
16	1	3	1	2	6	I
17	1	2	1	2	2	I
18	1	2	1	2	2	I
19	1	2	1	2	2	I
20	6	2	3	3	7	II
21	1	2	1	2	2	I
22	5	2	1	2	2	I
23	1	2	1	2	2	I
24	1	2	1	2	2	I
25	5	2	1	2	2	I
26	1	2	1	2	2	I
27	1	2	1	2	2	I
28	1	3	1	2	6	I
29	1	2	2	2	4	I
30	1	2	1	2	2	I
31	1	2	1	2	2	I
32	1	2	1	2	2	I
33	1	4	1	2	3	I
35	1	4	2	2	5	II
36	1	3	2	2	8	I
39	1	4	1	2	3	I
40	1	4	1	2	3	II

(Continued)

Table S2 (Continued)

Strain	16S rRNA	ITS1	gyrA	rpoB	ST ^a	Resistance pattern
41	5	2	1	2	2	I
42	1	4	1	2	3	I
43	1	2	1	2	2	I
44	1	2	1	2	2	I
45	1	4	1	2	3	I
46	1	2	1	2	2	V
47	1	4	2	2	5	III
48	1	4	1	4	9	III
49	1	2	1	2	2	I
50	1	2	1	2	2	I
51	1	2	1	2	2	I
52	1	2	1	2	2	I
53	1	2	1	2	2	I
54	1	4	1	2	3	I
55	1	2	1	4	10	I
56	1	2	1	2	2	I
57	1	3	1	4	11	I
58	7	3	1	2	6	I
59	1	4	1	5	12	I
61	8	3	1	2	6	I
62	1	3	1	2	6	I
63	1	2	1	2	2	I
64	1	4	1	2	3	IV
65	1	3	1	2	6	I
66	1	3	1	2	6	I
67	1	3	1	2	6	I
69	9	3	1	2	6	I
70	1	4	1	2	3	I
71	1	2	1	6	13	I
72	1	4	1	2	3	I
ATCC 6940	1	1	1	1	1	

Note: ^aOnly three of the eight loci analyzed were taken into account in obtaining ST numbers (ITS1 region, *gyrA*, and *rpoB* genes).

Abbreviations: ITS1, internal transcribed spacer 1; ST, sequence type.

Infection and Drug Resistance

Dovepress

Publish your work in this journal

Infection and Drug Resistance is an international, peer-reviewed open-access journal that focuses on the optimal treatment of infection (bacterial, fungal and viral) and the development and institution of preventive strategies to minimize the development and spread of resistance. The journal is specifically concerned with the epidemiology of antibiotic

resistance and the mechanisms of resistance development and diffusion in both hospitals and the community. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/infection-and-drug-resistance-journal>