Melatonin receptor stimulation by agomelatine prevents A β -induced tau phosphorylation and oxidative damage in PC12 cells

This article was published in the following Dove Medical Press journal:
Drug Design, Development and Therapy

Kai Yao
Yong-fei Zhao
Heng-bing Zu

Department of Neurology, Jinshan
Hospital Affiliated to Fudan University,
Shanghai 201508, China

Purpose: As a novel antidepressant drug, agomelatine has good therapeutic effect on the mood disorder and insomnia in Alzheimer's disease (AD). Recent studies have shown the neuroprotective function of agomelatine, including anti-oxidative and anti-apoptosis effect. However, it remains unclear whether agomelatine exerts neuroprotection in AD. Thus, the neuroprotective effect of agomelatine against amyloid beta 25–35 (A β_{25-35})-induced toxicity in PC12 cells was evaluated in this study.

Methods: The concentration of malondialdehyde (MDA), LDH, and ROS was investigated to evaluate oxidative damage. The expression of P-tau, tau, PTEN, P-Akt, Akt, P-GSK3 β , and GSK3 β proteins was assessed by Western blotting. Our results demonstrated that A β_{25-35} significantly increased the content of MDA, LDH, and ROS. Meanwhile, A β_{25-35} upregulated the expression of P-tau and PTEN as well as downregulated P-Akt and P-GSK3 β expression. These effects could be blocked by agomelatine pretreatment. Furthermore, luzindole, the melatonin receptor (MT) antagonist, could reverse the neuroprotective effect of agomelatine.

Conclusion: The results demonstrated that antidepressant agomelatine might prevent the tau protein phosphorylation and oxidative damage induced by A β_{25-35} in PC12 cells by activating MT-PTEN/Akt/GSK3 β signaling. This study provided a novel therapeutic target for AD in the future.

Keywords: agomelatine, Alzheimer's disease, oxidative stress, tau hyperphosphorylation

Introduction

It has been accepted that the extracellular deposition of amyloid beta (A β) plaques and the accumulation of intracellular tau neurofibrillary tangles (NFT) are the most important pathophysiology of AD.¹ Tau, a microtubule-associated protein, is the main component of the intracellular filamentous inclusions, which is essential for the regulation of microtubule structure and axonal transport by binding to the microtubule. In the pathological state, tau protein hyperphosphorylation has been reported to drive tau aggregation and enhance tau-mediated neurotoxicity,² leading to spine collapse and dendritic injury,³ and aggravate neurodegeneration, which is involved in several neurodegenerative diseases, including AD and frontotemporal dementia with parkinsonism-17 (FTDP-17).

Oxidative stress is defined as an imbalance between oxidants and antioxidants, resulting in excessive generation of toxic molecules such as ROS.⁴ When the concentration of reactive species is beyond the control of internal protective mechanisms, oxidative damage occurs to proteins, lipids, and DNA, leading to cytotoxicity.⁵ The level of malondialdehyde (MDA), a marker of lipid peroxidation index, reflects the extent of lipid

Correspondence: Heng-bing Zu
Department of Neurology, Jinshan
Hospital Affiliated to Fudan University,
No 1508 Longhang Road, Jinshan District,
Shanghai 201508, China
Tel +86 21 3418 9990 ext 5490
Fax +86 21 3418 9990
Email hbzyy8@126.com

peroxidation, which is considered as crucial factor in AD.⁶ Besides, glycolytic enzyme LDH increases along with plasma membrane damage, which is often used as an indicator of necrotic cell death caused by a plethora of external stress factors. Oxidative damage to neurons and loss of cholinergic neurons in the forebrain region are observed in AD,⁷ and several investigations have revealed that oxidative stress plays an important role in the pathogenesis of AD.⁸ Furthermore, several anti-oxidative and anti-tau protein hyperphosphorylation therapeutic strategies show great potential in treating AD.^{9,10}

AD often expresses with multiple comorbidities such as depression. AD and depression share some common etiology, including oxidative stress and nitrosative stress;¹¹ therefore, more and more evidence demonstrated that antidepressant exerted neuroprotective effect in the development of AD.^{12,13} As a novel antidepressant drug, agomelatine, widely applied in clinic, is a receptor agonist that affects both MT1 and MT2 melatonin receptors and an antagonist that affects 5-hydroxytryptamine (5HT) 2C receptor. Agomelatine was quite effective not only for insomnia but also for anxiety and depressive symptoms. Recent studies have shown that agomelatine provided neuroprotective effect in multiple disease models, such as ischemic stroke animal model¹⁴ and depression animal model,¹⁵ by anti-oxidative injury,¹⁵ anti-apoptosis, and by promoting neural restoration.^{16,17} However, it remains unclear whether agomelatine exerts neuroprotection in AD.

In the present study, the PC12 cell line was used and aimed to explore 1) effect of agomelatine on tau protein phosphorylation and oxidative damage induced by A β_{25-35} and 2) the neuroprotective mechanism of agomelatine. This study aimed to provide new insights in the therapy of AD.

Materials and methods

Materials

A β_{25-35} (#A4559), agomelatine (#A1362), luzindole (#L2407), the primary antibodies against phospho-tau (Ser396) (#SAB4504557), tau (#SAB4501830), PTEN (#SAB1406331), GAPDH (#SAB2701826), goat antirabbit IgG (#A3687), and antibody antimouse IgG (#M8770) were purchased from Sigma-Aldrich Co., St Louis, MO, USA. The primary antibodies against phospho-Akt (Ser473) (#4060) and Akt (#4691) were purchased from Cell Signaling Technology, Danvers, MA, USA. The primary antibodies against phospho-GSK3 β (Ser9) (Ab131097) and GSK3 β (Ab93926) were purchased from Abcam, Cambridge, UK. Cell counting kit-8 (CCK-8) (#E606335-0500) and ROS assay kit (#50101ES01) were obtained from Sango Biotech

(Shanghai, China). Cell MDA assay kit (#A003-4) and LDH assay kit (#A020-2) were purchased from Nanjing Jiancheng Bioengineering Institute (China).

Cell culture

PC12 cells were purchased from Chinese Academy of Sciences (Shanghai, China) and cultured in DMEM basal culture medium (Thermo Fisher Scientific, Waltham, MA, USA) with 10% FBS (HyClone, Logan, UT, USA) and 1% penicillin–streptomycin at 37°C in 5% CO₂ incubator. In agomelatine pretreatment group, after agomelatine pretreatment at different concentration for 24 hours, PC12 cells were then exposed to A β_{25-35} for 24 hours. In agomelatine posttreatment group, after A β_{25-35} pretreatment for 24 hours, PC12 cells were then exposed to agomelatine for 24 hours. Besides, in luzindole treatment group, PC12 cells were first treated with agomelatine in the presence of luzindole treatment for 24 hours, then exposed to A β_{25-35} for 24 hours.

Cell viability assay

The cell viability was assessed by CCK-8 assay. PC12 cells were first plated into 96-well plates and incubated in DMEM medium at 37°C for 24 hours. After treatment, CCK-8 assay kit was used to determine cell viability, according to the manufacturer's instructions. The assays were performed in triplicate.

Measurement of malondialdehyde (MDA) production

To evaluate lipid peroxidation reactions, intracellular MDA generation was measured using the MDA assay kit. The cells were lysed by lysate buffer, and the lysate was used to measure the level of MDA according to the manufacturer's instructions. The results were expressed as OD values by determining the absorbance at 530 nm. The assays were performed in triplicate.

Lactate dehydrogenase (LDH) measurements

To assess oxidative stress-induced cell injury, LDH activity in the supernatant of PC12 cells was measured by a commercial kit. At the end of the different treatments, the experiment was performed following the manufacturer's instructions. The total absorbance was measured at 490 nm.

Measurement of ROS production

Dichlorodihydrofluorescein diacetate (DCFH-DA), indicator of ROS, was used to analyze intracellular accumulation

of ROS production. After staining with DCFH-DA for 30 minutes at 37°C, the cells were harvested by trypsinization, centrifuged, and suspended in 1 mL PBS buffer, as described in the ROS assay kit. The intracellular accumulation of ROS production was measured by flow cytometry (BD, Franklin Lakes, NJ, USA).

Western blot analysis

After designated treatment, lysates were generated by RIPA lysis buffer, and then Bradford assay was performed to determine total protein concentrations. Subsequently, samples were prepared in sample buffer and heated to 95°C for 5 minutes. Equal amounts of lysates were fractionated using 10% SDS-PAGE and electrotransferred onto nitrocellulose membranes. Gels were run at a constant current (10–15 mA) for 3–4 hours for maximum separation. Wet transfer was performed for 1 hour at constant current (300 mA) using polyvinylidenedifluoride membrane presoaked in methanol. The membrane was blocked in 5% milk in 0.2% Phosphate Buffered Saline Tween-20. The membrane was then washed in Phosphate Buffered Saline Tween-20 $\times 3$ for 15 minutes each. After overnight incubation at 4°C with the primary antibodies (P-tau 1:1,000, tau 1:1,000, PTEN 1:1,000, P-Akt 1:1,000, Akt 1:1,000, P-GSK3 β 1:1,000, GSK3 β 1:1,000, GAPDH 1:5,000), the blots were washed and exposed for 1 hour to corresponding horseradish peroxidase-conjugated secondary antibodies. Chemiluminescent (Bio-Rad, Hercules, CA, USA) detection was then used to detect expression of each protein; GAPDH levels served as internal loading controls.

Statistical analysis

Statistical analysis was performed using SPSS 16.0 statistical software. All data were expressed as mean $\pm$ SEM from at least three independent experiments. *P*-values were determined using one-way ANOVA. Significance was defined as *P* < 0.05.

Results

Agomelatine prevents A β_{25-35} -induced oxidative injury in PC12 cells

To assess A β_{25-35} -induced oxidative damage in PC12 cells, the pretreatment study was conducted first. The concentration of MDA and LDH was determined. Then ROS level was analyzed via flow cytometry analysis. As shown in Figure 1A, the result of CCK-8 assay showed that cell viability was significantly reduced by A β_{25-35} treatment (20 μ M) (*P* < 0.05), and the injury effect was remarkably attenuated by agomelatine with different concentrations (20, 200 μ M). Besides, as

shown in Figures 1B and C, A β_{25-35} (20 μ M) could obviously increase the concentration of MDA, LDH, and ROS in PC12 cells (*P* < 0.01), and agomelatine pretreatment could protect against the oxidative damage (*P* < 0.05). Second, the effect of agomelatine postinsult was investigated. As shown in Figure 1D, after PC cells were already stressed with A β_{25-35} treatment, agomelatine posttreatment could obviously reverse the increase in ROS level induced by A β_{25-35} . The results demonstrated that agomelatine provided protective effect on A β -induced oxidative injury in vitro.

Agomelatine protects against A β_{25-35} -induced tau phosphorylation in PC12 cells

To assess the extent of tau phosphorylation, Western blot analysis was explored. As shown in Figure 2, as compared to control, the protein expression of P-tau and P-tau/tau ratio were significantly upregulated in A β_{25-35} -treated group (20 μ M) and indicated that A β_{25-35} could promote tau phosphorylation in PC12 cells. Furthermore, agomelatine pretreatment could downregulate increased P-tau expression and P-tau/tau ratio induced by A β_{25-35} . Therefore, we concluded that agomelatine could attenuate tau protein hyperphosphorylation induced by A β_{25-35} in AD.

To explore the potential molecular mechanism contributing to the neuroprotective activity of agomelatine, the melatonin receptor antagonist luzindole was used. First, oxidative stress response was evaluated. As shown in Figures 3A and C, A β_{25-35} (20 μ M) obviously promoted the generation of MDA, LDH, and ROS, and agomelatine (20 μ M) pretreatment significantly blocked A β_{25-35} -induced increase in MDA, LDH, and ROS production (*P* < 0.05). Moreover, when cotreated with luzindole (1 μ M), this effect could be reversed (*P* < 0.05). Second, tau protein phosphorylation was determined by Western blot analysis. As shown in Figure 3B, A β_{25-35} (20 μ M) remarkably upregulated the expression of P-tau protein (*P* < 0.01), and A β_{25-35} -induced tau phosphorylation could be blocked by agomelatine (20 μ M). When compared with agomelatine group, P-tau/tau ratio obviously increased by luzindole (1 μ M) cotreatment. According to the results, we found that the anti-oxidative and anti-tau protein phosphorylation effect of agomelatine was prevented by melatonin receptor antagonist luzindole. Therefore, agomelatine may provide neuroprotective function by activating melatonin receptor.

The signaling pathway involved in the neuroprotective effect of agomelatine

PTEN/Akt/GSK3 β signaling pathway plays an important role in the process of oxidative stress and tau protein

hyperphosphorylation. Thus, Western blot analysis was explored to detect the expression of PTEN, P-Akt, Akt, P-GSK3 β , and GSK3 β . As shown in Figure 4, A β_{25-35} (20 μ M) obviously promoted the upregulation of PTEN expression and decreased P-Akt and P-GSK3 β expression ($P < 0.01$). Furthermore, agomelatine pretreatment could partly inhibit the effect of A β_{25-35} . When cotreated with luzindole, agomelatine-induced PTEN/Akt/GSK3 β pathway inhibition was obviously reversed. According to the results, we concluded that agomelatine may exert anti-oxidative effect and prevent tau hyperphosphorylation through activating MT-PTEN/Akt/GSK3 β signaling.

Discussion

As the most prevalent neurodegenerative disease, the etiology of AD is not well understood.^{18,19} The extracellular deposition of A β plaques and the accumulation of intracellular NFT are considered to be the most crucial pathophysiology of AD.¹ Tau protein plays an important role in maintenance of the integrity of genomic DNA,²⁰ regulation of neuronal activity,²¹ and neurogenesis.²² Generally, to maintain physiological functions of tau, an appropriate extent of phosphorylation is necessary, modulated by various protein kinases and phosphatases, including proline-directed protein kinases (PDPKs), non-PDPKs, and tyrosine kinases.²³ In the pathological state,

Figure 1 (Continued)

tau protein hyperphosphorylation promoted microtubule instability,²⁴ axon transport impairment,²⁵ and altered its truncation,²⁶ eventually leading to neurodegeneration. Tau phosphorylation is involved in several neurodegenerative

diseases, including AD and FTDP-17. In the progression of AD, tau protein hyperphosphorylation has been reported to drive tau aggregation, and tau aggregation is essential for tau-induced toxicity.² Several experiments indicated that

Figure 3 Effect of agomelatine and luzindole on oxidative stress and tau phosphorylation induced by Aβ₂₅₋₃₅ in PC12 cells.

Notes: (A, C) Measurement of MDA, LDH, and ROS. PC12 cells untreated or treated with agomelatine (20 μM) for 24 hours in the absence or presence of luzindole (1 μM) for 24 hours were exposed to Aβ₂₅₋₃₅ (20 μM) for 24 hours. (B) Measurement P-tau and tau expression by Western blot. PC12 cells untreated or treated with agomelatine (20 μM) for 24 hours in the absence or presence of luzindole (1 μM) for 24 hours were exposed to Aβ₂₅₋₃₅ (20 μM) for 24 hours. Data represent the mean ± SEM of three independent experiments (n=3). *P<0.05; **P<0.01.

Abbreviations: Aβ₂₅₋₃₅, amyloid beta 25–35; Ago, agomelatine; DCFH-DA, dichlorodihydrofluorescein diacetate; LDH, lactate dehydrogenase; MDA, malondialdehyde.

Aβ exposure could aggravate tau-mediated neurotoxicity by facilitating tau phosphorylation, leading to spine collapse and dendritic degeneration.³ In our study, we found that Aβ could obviously promote tau protein phosphorylation by upregulation of P-tau expression and P-tau/tau ratio in PC12 cells, which suggested that interaction between Aβ and tau might facilitate the development of AD.

A lot of evidence revealed that oxidative stress is involved in the development and progression of AD. It has been widely accepted that oxidative stress damage contributes to initiation and progression of AD,²⁷ through several pathways, including 1) aggravation of Aβ production, secretion, and aggregation;^{28–30} 2) causing mitochondrial dysfunction;³¹ and 3) promoting hyperphosphorylation of tau proteins, tau-mediated toxicity, and intracellular NFT formation.³² Our results showed that Aβ₂₅₋₃₅ treatment could significantly enhance oxidative stress response by increasing the concentration of MDA, LDH, and ROS, which indicate that oxidative stress may play an important role in Aβ-induced toxicity and pathological process of AD.

Furthermore, several anti-oxidative and anti-tau protein hyperphosphorylation therapeutic strategies show great potential in treating AD.^{9,10} In addition, AD often manifests with multiple comorbidities such as depression. More and

more evidence demonstrated that antidepressants exert a neuroprotective effect in the development of AD.¹² As a novel antidepressant drug, agomelatine, widely applied in clinic, could act on the SCN, hippocampus, frontal cortex, and striatum. Agomelatine was quite helpful not only for insomnia through improvement of sleep duration and restoration of the circadian rhythm³³ but also for anxiety and depressive symptoms and has no adverse effects on cognition during the treatment of patients, especially the elderly.³⁴ Agomelatine is effective in the therapy of mood disorder in AD. Furthermore, animal studies have demonstrated that agomelatine treatment could increase generation of brain-derived neurotrophic factor levels and promote hippocampal and prefrontal cortex neurogenesis.¹⁷ On the other hand, agomelatine was found to provide potent antioxidative effect in multiple disease models. It was shown that agomelatine potentially helps reducing testicular damage by decreasing oxidative stress level in STZ-induced type I diabetic rats.³⁵ In ischemic stroke animal model, agomelatine treatment could protect the brain from cerebral I/R injury by anti-apoptosis and antioxidant properties.¹⁴ In rats with chronic mild stress-induced depression, agomelatine exhibited protective effects against brain, kidney, and liver oxidative damage by regulation of the glutathione concentrations and glutathione peroxidase

Figure 4 The signaling involved in the neuroprotective effect of agomelatine. Measurement of PTEN, P-Akt, Akt, P-GSK3β (Ser9), and GSK3β expression by Western blot in PC12 cells.

Notes: PC12 cells untreated or treated with agomelatine (20 μM) for 24 hours in the absence or presence of luzindole (1 μM) for 24 hours were exposed to Aβ₂₅₋₃₅ (20 μM) for 24 hours. Data represent the mean ± SEM of three independent experiments (n=3). *P<0.05, **P<0.01.

Abbreviations: Aβ₂₅₋₃₅, amyloid beta 25–35; Ago, agomelatine.

activity.¹⁵ However, it remains unclear whether agomelatine exerts neuroprotection in AD. In our study, we first reported that agomelatine could significantly prevent $A\beta_{25-35}$ -induced increase in P-tau expression and P-tau/tau ratio. Moreover, as compared to $A\beta_{25-35}$ group, agomelatine pretreatment could alleviate oxidative stress damage caused by $A\beta_{25-35}$ by decreasing the generation of MDA, LDH, and ROS. Our results indicated that agomelatine had great potential for inhibiting tau protein hyperphosphorylation and oxidative damage in AD.

Then the neuroprotective mechanism of agomelatine was investigated. Agomelatine is a receptor agonist that affects both MT1 and MT2 melatonin receptors and an antagonist that affects 5HT_{2C} receptors. Previous studies demonstrated that MT receptors are obviously decreased in multiple brain areas such as the pineal gland, pyramidal and cortical layers, and hippocampus in AD.^{36,37} MT receptors regulate the neurotoxicity and clearance of amyloid- β ,^{38,39} the stabilization of synapses,⁴⁰ the generation of neurofibrillary tangles, and level of oxidative stress in AD.⁴¹ In our study, after luzindole pretreatment, we found the novel results that the decrease in P-tau/tau ratio and oxidative production by agomelatine was obviously upregulated. The study revealed that luzindole, antagonist of melatonin receptor, could reverse the anti-tau phosphorylation and anti-oxidative stress effect provided by agomelatine in $A\beta$ -treated PC12 cells; therefore, we speculated that agomelatine may prevent the pathological injury by activating melatonin receptor in AD.

On the other hand, PTEN/Akt/GSK3 signaling pathway is closely related to the onset and progression of AD. As a serine/threonine protein kinase, GSK3 exists in two isoforms, GSK3 α and GSK3 β , expressed in brain abundantly. GSK3 activity could be modulated by the site-specific phosphorylation of GSK3 β , for example, ser9 of GSK3 β inhibits its activity.⁴² GSK3 signaling is involved in a number of cellular pathological and physiological process, including glucose regulation, inflammation, oxidative stress, and apoptosis.^{43,44} A lot of evidence has shown that both GSK3 isoforms were abnormally activated in the brains of postmortem AD samples and AD animal model,^{45,46} resulting in excessive $A\beta$ formation and aggregation.⁴⁷ Besides, GSK3 activation in AD significantly promotes tau protein hyperphosphorylation, leading to the formation of NFTs and neuronal death.^{48,49} Therefore, GSK3 is widely known as the key kinase responsible for the hyperphosphorylation of tau protein, and GSK3 inhibitor provides potential neuroprotective function in the development of AD.⁴⁸ GSK3 activity is regulated by several upstream kinases, such as PTEN (also named MMAC1/TEP1). PTEN is considered as a dual phosphatase with both protein and lipid

phosphatase activities, modulating cellular growth, survival, and metabolism.⁵⁰ It is now well established that PTEN could inhibit a major cell growth and survival signaling pathway, including PI3K/Akt signaling pathway,⁵¹ and upregulate the expression and activity of GSK3.⁵²⁻⁵⁴ Furthermore, targeted suppression of PTEN activity could protect against oxidative stress damage, neurotoxicity,^{55,56} endoplasmic reticulum stress,⁵⁷ and tau hyperphosphorylation⁵⁶ in the brain of patients with AD. However, little is known about the link between agomelatine, MT receptors, and PTEN/Akt/GSK3 β signaling in AD. According to our study, we found the novel results that $A\beta_{25-35}$ obviously activated PTEN/Akt/GSK3 β signaling by increasing PTEN expression and decreasing P-Akt and P-GSK3 β expression. Moreover, agomelatine could prevent the effect of $A\beta_{25-35}$ through inhibition of PTEN/Akt/GSK3 β axis, which was reversed by melatonin receptor antagonist luzindole. Our study demonstrated that the neuroprotective effect provided by agomelatine was via activating MT-PTEN/Akt/GSK3 β signaling pathway. The results have not been published before and revealed the novel neuroprotective mechanism of agomelatine in AD.

Conclusion

Our data presented that antidepressant agomelatine could prevent the tau protein phosphorylation and oxidative damage induced by $A\beta_{25-35}$ in PC12 cells. Furthermore, agomelatine might exert protective effect by activating MT-PTEN/Akt/GSK3 β signaling pathway. Importantly, the neuroprotective effect of agomelatine must be discussed in vivo in AD animal models in the future.^{58,59} The data above revealed the neuroprotective function of agomelatine in AD-associated pathological injury and provided new insights in the therapy of AD.

Acknowledgment

The authors thank Heng-bing Zu (Department of Neurology, Jinshan Hospital Affiliated to Fudan University, Shanghai, China) for excellent technical assistance in this study. The study was supported by research grants from Research Project of Jinshan District Health and Family Planning Commission (No JSKJ-KTMS-2018-19), Qi Hang project of Jinshan Hospital (No 2018-JSYQ-06).

Disclosure

The authors report no conflicts of interest in this work.

References

1. Querfurth HW, Laferla FM. Alzheimer's disease. *N Engl J Med Overseas Ed*. 2010;362(4):329-344.

2. Ahmed T, Van der Jeugd A, Blum D, et al. Cognition and hippocampal synaptic plasticity in mice with a homozygous tau deletion. *Neurobiol Aging*. 2014;35(11):2474–2478.
3. De Felice FG, Wu D, Lambert MP, et al. Alzheimer's disease-type neuronal tau hyperphosphorylation induced by A β oligomers. *Neurobiol Aging*. 2008;29(9):1334–1347.
4. Sies H, Berndt C, Jones DP. Oxidative stress. *Encyclopedia Stress*. 2007;62(10):45–48.
5. Gagné F. Oxidative stress. In: *Biochemical Ecotoxicology: Principles and Methods*. Amsterdam: Elsevier; 2014:103–115.
6. Gao HM, Zhou H, Hong JS. NADPH oxidases: novel therapeutic targets for neurodegenerative diseases. *Trends Pharmacol Sci*. 2012;33(6):295–303.
7. Limón ID, Díaz A, Mendieta L, et al. Amyloid-beta (25–35) impairs memory and increases NO in the temporal cortex of rats. *Neurosci Res*. 2009;63(2):129–137.
8. Wang X, Wang W, Li L, Perry G, Lee HG, Zhu X. Oxidative stress and mitochondrial dysfunction in Alzheimer's disease. *Biochim Biophys Acta*. 1842;2014(8):1240–1247.
9. Gräff J, Tsai LH. The potential of HDAC inhibitors as cognitive enhancers. *Annu Rev Pharmacol Toxicol*. 2013;53(1):311–330.
10. Kanninen K, Malm TM, Jyrkkänen HK, et al. Nuclear factor erythroid 2-related factor 2 protects against beta amyloid. *Mol Cell Neurosci*. 2008;39(3):302–313.
11. Plotnikoff NP. 2007. Cytokines: stress and immunity. Boca Raton, FL: CRC Press. 405PP.
12. Manji HK, Moore GJ, Chen G. Clinical and preclinical evidence for the neurotrophic effects of mood stabilizers: implications for the pathophysiology and treatment of manic-depressive illness. *Biol Psychiatry*. 2000;48(8):740–754.
13. Noble W, Planel E, Zehr C, et al. Inhibition of glycogen synthase kinase-3 by lithium correlates with reduced tauopathy and degeneration in vivo. *Proc Natl Acad Sci U S A*. 2005;102(19):6990–6995.
14. Chumboatong W, Thummayot S, Govitrapong P, Tocharus C, Jittiwat J, Tocharus J. Neuroprotection of agomelatine against cerebral ischemia/reperfusion injury through an antiapoptotic pathway in rat. *Neurochem Int*. 2017;102:114–122.
15. Demirdaş A, Nazıroğlu M, Ünal GÖ. Agomelatine reduces brain, kidney and liver oxidative stress but increases plasma cytokine production in the rats with chronic mild stress-induced depression. *Metab Brain Dis*. 2016;31(6):1445–1453.
16. Yucel A, Yucel N, Ozkanlar S, et al. Effect of agomelatine on adult hippocampus apoptosis and neurogenesis using the stress model of rats. *Acta Histochem*. 2016;118(3):299–304.
17. Pompili M, Serafini G, Innamorati M, et al. Agomelatine, a novel intriguing antidepressant option enhancing neuroplasticity: a critical review. *World J Biol Psychiatry*. 2013;14(6):412–431.
18. Rosenberg RN, Lambracht-Washington D, Yu G, Xia W. Genomics of Alzheimer disease: a review. *JAMA Neurol*. 2016;73(7):867.
19. Killin LO, Starr JM, Shiue IJ, Russ TC. Environmental risk factors for dementia: a systematic review. *BMC Geriatr*. 2016;16(1):175.
20. Violet M, Delattre L, Tardivel M, et al. A major role for tau in neuronal DNA and RNA protection in vivo under physiological and hyperthermic conditions. *Front Cell Neurosci*. 2014;8(6):84.
21. Gheyara AL, Ponnusamy R, Djukic B, et al. Tau reduction prevents disease in a mouse model of Dravet syndrome. *Ann Neurol*. 2014;76(3):443–456.
22. Hong XP, Peng CX, Wei W, et al. Essential role of tau phosphorylation in adult hippocampal neurogenesis. *Hippocampus*. 2010;20(12):1339–1349.
23. Chung SH. Aberrant phosphorylation in the pathogenesis of Alzheimer's disease. *BMB Rep*. 2009;42(8):467–474.
24. Hanger DP, Anderton BH, Noble W. Tau phosphorylation: the therapeutic challenge for neurodegenerative disease. *Trends Mol Med*. 2009;15(3):112–119.
25. Hoover BR, Reed MN, Su J, et al. Tau mislocalization to dendritic spines mediates synaptic dysfunction independently of neurodegeneration. *Neuron*. 2010;68(6):1067–1081.
26. Dickey CA, Kamal A, Lundgren K, et al. The high-affinity HSP90-CHIP complex recognizes and selectively degrades phosphorylated tau client proteins. *J Clin Invest*. 2007;117(3):648–658.
27. Liu Z, Zhou T, Ziegler AC, Dimitrion P, Zuo L. Oxidative stress in neurodegenerative diseases: from molecular mechanisms to clinical applications. *Oxid Med Cell Longev*. 2017;2017(4):1–11.
28. Sherrington R, Rogaev EI, Liang Y, et al. Cloning of a gene bearing missense mutations in early-onset familial Alzheimer's disease. *Nature*. 1995;375(6534):754–760.
29. Bonda DJ, Wang X, Perry G, et al. Oxidative stress in Alzheimer disease: a possibility for prevention. *Neuropharmacology*. 2010;59(4–5):290–294.
30. Ascertatham GS, Sivasudha T, Jayadevi R, Arul Ananth D. Environmental factors and unhealthy lifestyle influence oxidative stress in humans – an overview. *Environ Sci Pollut Res Int*. 2013;20(7):4356–4369.
31. Abramov AY, Berezhnov AV, Fedotova EI, Zinchenko VP, Dolgacheva LP. Interaction of misfolded proteins and mitochondria in neurodegenerative disorders. *Biochem Soc Trans*. 2017;45(4):1025–1033.
32. Giraldo E, Lloret A, Fuchsberger T, Viña J. A β and tau toxicities in Alzheimer's are linked via oxidative stress-induced p38 activation: protective role of vitamin E. *Redox Biol*. 2014;2:873–877.
33. Laudon M, Frydman-Marom A. Therapeutic effects of melatonin receptor agonists on sleep and comorbid disorders. *Int J Mol Sci*. 2014;15(9):15924–15950.
34. Plesničar BK. Efficacy and tolerability of agomelatine in the treatment of depression. *Patient Prefer Adherence*. 2014;8:603–612.
35. Yigittürk G, Acara AC, Erbas O, et al. The antioxidant role of agomelatine and gallic acid on oxidative stress in STZ induced type I diabetic rat testes. *Biomed Pharmacother*. 2017;87:240–246.
36. Brunner P, Sözer-Topcular N, Jockers R, et al. Pineal and cortical melatonin receptors MT1 and MT2 are decreased in Alzheimer's disease. *Eur J Histochem*. 2006;50(4):311–316.
37. Savaskan E, Ayoub MA, Ravid R, et al. Reduced hippocampal MT2 melatonin receptor expression in Alzheimer's disease. *J Pineal Res*. 2005;38(1):10–16.
38. Chinchalongporn V, Shukla M, Govitrapong P. Melatonin ameliorates A β_{42} -induced alteration of β APP-processing secretases via the melatonin receptor through the Pin1/GSK3 β /NF- κ B pathway in SH-SY5Y cells. *J Pineal Res*. 2018;64(4):e12470.
39. Pappolla MA, Matsubara E, Vidal R, et al. Melatonin treatment enhances A β lymphatic clearance in a transgenic mouse model of amyloidosis. *Curr Alzheimer Res*. 2018;15(7):637–642.
40. Shi Y, Fang YY, Wei YP. Melatonin in synaptic impairments of Alzheimer's disease. *J Alzheimers Dis*. 2018;63(3):1–16.
41. Caballero B, Vega-Naredo I, Sierra V, et al. Favorable effects of a prolonged treatment with melatonin on the level of oxidative damage and neurodegeneration in senescence-accelerated mice. *J Pineal Res*. 2008;45(3):302–311.
42. Dajani R, Fraser E, Roe SM, et al. Crystal structure of glycogen synthase kinase 3 beta: structural basis for phosphate-primed substrate specificity and autoinhibition. *Cell*. 2001;105(6):721–732.
43. Rana AK, Singh D. Targeting glycogen synthase kinase-3 for oxidative stress and neuroinflammation: opportunities, challenges and future directions for cerebral stroke management. *Neuropharmacology*. 2018;139:124–136.
44. Ge XH, Shao L, Zhu GJ. Oxymatrine attenuates brain hypoxic-ischemic injury from apoptosis and oxidative stress: role of p-Akt/GSK3 β /HO-1/Nrf-2 signaling pathway. *Metab Brain Dis*. 2018;33(6):1869–1875.
45. Terwel D, Muyliaert D, Dewachter I, et al. Amyloid activates GSK-3 β to aggravate neuronal tauopathy in bigenic mice. *Am J Pathol*. 2008;172(3):786–798.
46. Leroy K, Boutajangout A, Authélet M, Woodgett JR, Anderton BH, Brion JP. The active form of glycogen synthase kinase-3 β is associated with granulovacuolar degeneration in neurons in Alzheimer's disease. *Acta Neuropathol*. 2002;103(2):91–99.
47. Hooper C, Killick R, Lovestone S. The GSK3 hypothesis of Alzheimer's disease. *J Neurochem*. 2008;104(6):1433–1439.

48. Maqbool M, Mobashir M, Hoda N. Pivotal role of glycogen synthase kinase-3: a therapeutic target for Alzheimer's disease. *Eur J Med Chem*. 2016;107:63–81.
49. Morris G, Berk M, Maes M, Puri BK. Could Alzheimer's disease originate in the periphery and if so how so? *Mol Neurobiol*. Epub 2018 Apr 29.
50. Worby CA, Dixon JE. PTEN. *Annu Rev Biochem*. 2014;83(1):641–669.
51. Robbins HL, Hague A. The PI3K/Akt pathway in tumors of endocrine tissues. *Front Endocrinol*. 2016;6(377–387):188.
52. Ma J, Guo X, Zhang J, et al. PTEN gene induces cell invasion and migration via regulating Akt/GSK-3 β / β -catenin signaling pathway in human gastric cancer. *Dig Dis Sci*. 2017;62(12):3415–3425.
53. Shen H, Li L, Yang S, et al. MicroRNA-29a contributes to drug-resistance of breast cancer cells to adriamycin through PTEN/AKT/GSK3 β signaling pathway. *Gene*. 2016;593(1):84–90.
54. He E, Pan F, Li G, Li J. Fractionated ionizing radiation promotes epithelial-mesenchymal transition in human esophageal cancer cells through PTEN deficiency-mediated Akt activation. *PLoS One*. 2015; 10(5):e0126149.
55. Liu XY, Zhang LJ, Chen Z, Liu LB. The PTEN inhibitor bpV (pic) promotes neuroprotection against amyloid β -peptide (25–35)-induced oxidative stress and neurotoxicity. *Neurol Res*. 2017;39(8):758–765.
56. Pinto-Almazán R, Segura-Urbe JJ, Soriano-Ursúa MA, Farfán-García ED, Gallardo JM, Guerra-Araiza C. Effect of tibolone pretreatment on kinases and phosphatases that regulate the expression and phosphorylation of tau in the hippocampus of rats exposed to ozone. *Neural Regen Res*. 2018;13(3):440.
57. Cui W, Wang S, Wang Z, Wang Z, Sun C, Zhang Y. Inhibition of PTEN attenuates endoplasmic reticulum stress and apoptosis via activation of PI3K/Akt pathway in Alzheimer's disease. *Neurochem Res*. 2017; 42(11):3052–3060.
58. Laferla FM, Green KN. Animal models of Alzheimer disease. *Cold Spring Harb Perspect Med*. 2012;2(11):a006320–a006329.
59. Götz J, Ittner LM. Animal models of Alzheimer's disease and frontotemporal dementia. *Nat Rev Neurosci*. 2008;9(7):532–544.

Drug Design, Development and Therapy

Publish your work in this journal

Drug Design, Development and Therapy is an international, peer-reviewed open-access journal that spans the spectrum of drug design and development through to clinical applications. Clinical outcomes, patient safety, and programs for the development and effective, safe, and sustained use of medicines are the features of the journal, which

Submit your manuscript here: <http://www.dovepress.com/drug-design-development-and-therapy-journal>

Dovepress

has also been accepted for indexing on PubMed Central. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.