Lack of association between *ALOX5AP* genetic polymorphisms and risk of ischemic stroke: evidence from meta-analyses

This article was published in the following Dove Press journal:
Neuropsychiatric Disease and Treatment

Jing-Hui Zheng
Gui-Lan Ning
Wen-Hua Xu
Xin-Cheng Wu
Xiao-Cong Ma

Department of Cardiology, Ruikang
Hospital Affiliated to Guangxi
University of Chinese Medicine,
Nanning 530011, China

Background: In recent years, there has been substantial research evaluating the relationship between arachidonate 5-lipoxygenase-activating protein (*ALOX5AP*) polymorphisms and ischemic stroke (IS). The objective of this study was to systematically review and analyze the existing evidence.

Methods: A comprehensive search of major electronic databases for studies published between 1990 and 2018 was carried out. Data were synthesized as OR and 95% CI using fixed-effects and random-effects models.

Results: A total of 30 studies were available for analysis. The aggregate sample size across all studies was 32,782 (16,294 cases and 16,488 controls). We found no association of the *ALOX5AP* rs10507391 (OR=1.03 for A allele vs T allele; 95% CI: 0.93–1.14; $P=0.557$), rs4769874 (OR=1.13 for A allele vs G allele; 95% CI: 1.00–1.28; $P=0.050$), rs9551963 (OR=1.03 for A allele vs C allele; 95% CI: 0.96–1.11; $P=0.372$), rs17222814 (OR=1.09 for A allele vs G allele; 95% CI: 0.96–1.24; $P=0.195$), rs17222919 (OR=0.89 for G allele vs T allele; 95% CI: 0.75–1.06; $P=0.175$), and rs4073259 (OR=1.20 for A allele vs G allele; 95% CI: 1.00–1.45; $P=0.056$) polymorphisms with IS risk. Haplotype analysis also did not yield significant findings for the HapA (rs17222814G–rs10507391T–rs4769874G–rs9551963A; OR=1.20; 95% CI: 0.91–1.56; $P=0.192$) and HapB (rs17216473A–rs10507391A–rs9315050A–rs17222842G; OR=1.11; 95% CI: 0.90–1.38; $P=0.339$) haplotypes.

Conclusion: Current evidence does not support an association of rs10507391, rs4769874, rs9551963, rs17222814, rs17222919, rs4073259, and HapA and HapB with IS risk.

Keywords: ischemic stroke, *ALOX5AP*, genetic polymorphism, haplotype

Introduction

Stroke is the leading cause of mortality and serious long-term disability in China and worldwide, with ~15 million people throughout the world suffering a stroke every year.¹ Ischemic stroke (IS) accounts for ~85%–90% of all cases and is characterized by the sudden loss of blood circulation to an area of the brain. Strong evidence from twin, family, and animal model studies has consistently suggested a genetic influence on IS risk and prognosis, and in recent years, there has been substantial research evaluating specific genetic risk factors.²

Leukotrienes are short-lived lipid mediators derived from the nuclear membrane of cells that are produced and excreted in response to various immune stimuli. The leukotriene pathway starts with oxidation of arachidonic acid to leukotriene A4 (LTA4) by lipoxygenase-activating protein (*ALOX5AP*).³ The resultant LTA4 can then be converted to leukotriene B4 (LTB4) by leukotriene A4 hydrolase or it can be conjugated

Correspondence: Jing-Hui Zheng
Department of Cardiology,
Ruikang Hospital Affiliated to Guangxi
University of Chinese Medicine,
Nanning 530011, China
Tel +86 771 313 1086
Email drjhzheng@tutanota.com

with reduced glutathione by the leukotriene C4.³ As a crucial regulator of biosynthesis of proinflammatory leukotriene lipid mediators, ALOX5AP plays an important role in the development of atherosclerosis. Pharmacological targeting of ALOX5AP significantly reduced atherosclerosis burden and vascular as well as adipose tissue T-cell inflammation.⁴ In addition, knockout of the *ALOX5AP* gene was associated with ceased leukotriene production and amelioration of stroke damage in a mouse model of middle cerebral artery occlusion.⁵ The findings from clinical trials were consistent with those from animal experiments. Hakonarson et al found that the ALOX5AP inhibitor significantly decreased the concentration of the C-reactive protein, a contributor to the risk of myocardial infarction and IS.⁶ Moreover, ALOX5AP mRNA and protein increased significantly in polymorphonuclear cells from OSA patients vs controls and were associated with carotid luminal diameter and intima-media thickness.⁷

The human *ALOX5AP* is located on chromosome 13q12-13 and consists of five exons and four introns. Genetic variants in *ALOX5AP* gene have been studied in stroke cases, but case-control studies performed in different ethnic populations obtain contradictory results, indicating that the existing evidence regarding the association between *ALOX5AP* polymorphisms and IS risk needs to be systematically reviewed and analyzed. To reconcile inconsistencies across individual studies, we performed a quantitative meta-analysis of the effects of *ALOX5AP* polymorphisms on IS.

Methods

Search strategy

We searched Embase, Medline, PubMed, and China Knowledge Resource Integrated in July 2018 for studies investigating the relationship between *ALOX5AP* polymorphisms and IS risk. Searches were not limited by date restrictions and language. Search terms included: (“*ALOX5AP* polymorphism”, or “*ALOX5AP* variant”, or “*ALOX5AP* genotype”), AND (“cerebral infarction” OR “ischemic stroke”). The “AND” operator was used to combine these terms in varying combinations. Once articles had been collected, bibliographies were then hand-searched for additional references. The review followed the Preferred Reporting Items for Systematic Reviews and Meta-Analyses guidelines.⁸

Inclusion and exclusion criteria

Two investigators independently assessed titles and abstracts for relevance and full reports for inclusion. Discrepancies were resolved by discussion. Studies were included in our meta-analysis if they met the following criteria: 1) reported

on case-control studies in adult humans; 2) published in peer-reviewed journals; and 3) reported genotypic and/or allelic frequencies. The exclusion criteria were as follows: 1) family-based studies; 2) case-only studies; 3) no information on genotypic and/or allelic frequencies; and 4) editorials, narrative reviews or other manuscripts not reporting primary data. If the paper, or author correspondence, suggested overlapping studies, we included only the most comprehensive study for meta-analysis.

Data extraction and quality assessment

Data were tabled in a standardized Excel sheet, and each group comparison was checked by two investigators to confirm accuracy of inclusion. The following information was abstracted from each study: journal, first author, year of publication, participant characteristics, geographical location, dominant ancestry of sample, diagnostic method for IS, numbers of patients and controls, DNA extraction and genotyping methods, allele frequency, and Hardy-Weinberg equilibrium (HWE) in controls. Authors were not contacted to request missing/additional data. For assessing the methodological quality of the primary studies, a quantitative Newcastle-Ottawa Scale (NOS) score was calculated for each study, which was based on the selection of the study groups, the comparability of the groups, and the ascertainment of the exposure.⁹ Each study was graded as either low (scores 0–5) or high quality (scores 6–9).

Statistical analyses

All statistical analyses were performed using Stata 12.0 software (Stata, College Station, TX, USA). Since the majority of studies reported allele frequency instead of genotype data, the relationship between *ALOX5AP* polymorphisms and IS risk was assessed under allele contrast. All associations were presented as OR with the corresponding 95% CI. The significance of the pooled OR was determined using a Z-test. Heterogeneity was assessed by Cochrane's *Q* test of heterogeneity and the *I*² statistic. The *I*² values of 25%, 50%, and 75% corresponded to low, medium, and high between-study heterogeneity. Summary ORs were calculated using random-effects models (DerSimonian and Laird) if there was significant between-study heterogeneity.¹⁰ Random-effects modeling assumes a genuine diversity in the results of various studies and takes into account both within-study and between-study variances. Stratified analyses by sample ancestry were performed in order to assess the potential moderating effect of ancestry. HWE deviation was assessed by using the chi-squared goodness-of-fit test. Sensitivity analysis

was conducted by omitting studies with HWE deviation. Funnel plots, Begg's and Egger's tests were used to assess publication bias. The standard error of log (OR) of each study was plotted against its log (OR) in funnel plots. The funnel plot symmetry was evaluated by Begg's and Egger's tests, which used the correlation between the ranks of effect sizes and the ranks of their variances. A *P*-value of less than 0.05 was considered statistically significant, except for Cochrane's *Q* test, in which a *P*-value less than 0.10 was applied. No corrections for multiple comparisons were performed.

Results

Study characteristics

A total of 302 studies were identified by the search strategy. After duplicates were removed, 176 studies remained. On examination of the full text, 30 papers were eligible for inclusion in the review.^{11–40} Study selection is shown in Figure 1. These 30 studies were reported between 2005 and 2018 and included 32,782 participants. Sample sizes ranged from 200 to 3,727 participants per study and represented Caucasian and Asian populations. Imaging (magnetic resonance imaging/computerized tomography) examination of the brain to support the clinical diagnosis was conducted in all studies. The distribution of genotypes in the controls of all studies was in agreement with HWE except for the studies by Cheng et al, Zhao et al, Xu et al, and Diakite et al.^{21,26,29,37} Table 1 displays the study characteristics for the identified studies.

Data synthesis

Twenty studies provided summaries of data on the *ALOX5AP* rs10507391 polymorphism.^{13–18,20,22,23,25,26,28,29,31,32,34–38}

Random-effects meta-analysis indicated no association of the polymorphism with IS in the overall population (OR=1.03 for A allele vs T allele; 95% CI: 0.93–1.14; *P*=0.557) (Figure 2A; Table 2). There was significant between-study heterogeneity ($I^2=81.6\%$, *P*<0.001). In the subgroup analysis, according to ethnicity, no association between the *ALOX5AP* rs10507391 polymorphism and IS risk was found in Asian (OR=1.10 for A allele vs T allele; 95% CI: 0.98–1.23; *P*=0.109) and Caucasian (OR=0.89 for A allele vs T allele; 95% CI: 0.76–1.02; *P*=0.101) populations (Figure 2A; Table 2).

Thirteen studies contributed to the pooled analysis of the *ALOX5AP* rs4769874 polymorphism (A allele vs G allele).^{13–17,19,23,26,31,32,34–36} Fixed-effect meta-analysis gave an estimated OR of 1.13 (95% CI: 1.00–1.28; *P*=0.050), indicating no significant association with IS (Figure 2B; Table 2). There was no significant between-study heterogeneity ($I^2=10.9\%$, *P*=0.160). Neither Asian (OR=1.10, 95% CI 0.95–1.28, *P*=0.210) nor Caucasian (OR=1.19, 95% CI 0.96–1.48, *P*=0.106) studies found an association with IS (Figure 2B; Table 2).

Twelve studies examined the *ALOX5AP* rs9551963 polymorphism.^{13–15,18,19,23,26,29,31,32,34,36} In the primary pooled analysis (Figure 2C; Table 2), we identified no significant association between the polymorphism and IS risk (OR=1.03 for A allele vs C allele; 95% CI: 0.96–1.11; *P*=0.372). There was significant between-study heterogeneity ($I^2=44.7\%$, *P*=0.047). In the subgroup analysis based on ethnicity, studies with patients of Asian descent or Caucasian descent yielded similar findings (Asian population: OR=1.07, 95% CI 0.96–1.19, *P*=0.230; Caucasian population: OR=0.98, 95%

Figure 1 Flow diagram of study selection.

Abbreviation: CNKI, China Knowledge Resource Integrated.

Table 1 Summary of included studies

Author	Year	Sample size		Diagnostic method	Polymorphisms evaluated	Female (%)		Age		Ethnicity	NOS score
		Cases	Controls			Cases	Controls	Cases	Controls		
Helgadottir et al ¹¹	2004	702	624	CT/MRI confirmed	HapA and HapB	46.9	NM	NM	20–90	Caucasians	8
Helgadottir et al ¹²	2005	450	710	CT/MRI confirmed	HapA and HapB	42.0	49.0	66.8±0.6	67.2±0.4	Caucasians	7
Meschia et al ¹³	2005	377	263	CT/MRI confirmed	rs17222814, rs10507391, rs4769874 and rs9551963	46.4	62.0	64.8±15.0	60.0±14.7	Caucasians	8
Löhmussaar et al ¹⁴	2005	639	736	CT/MRI confirmed	rs17222814, rs10507391, rs4769874, rs9551963 and HapA	36.9	39.3	65.0±18.2	62.0±11.7	Caucasians	7
Zee et al ¹⁵	2006	259	259	CT/MRI confirmed	rs17222814, rs10507391, rs4769874, rs9551963, HapA and HapB	0.0	0.0	60.8±0.3	61.0±0.3	Caucasians	7
Zhang et al ¹⁶	2006	1,285	1,713	CT/MRI confirmed	rs10507391 and rs4769874	36.7	42.7	60.4±9.2	59.6±8.5	Asians	6
Gao and Qi ¹⁷	2008	100	100	CT/MRI confirmed	rs10507391 and rs4769874	35.0	40.0	57.3±6.9	60.0±7.5	Asians	7
Lökvist et al ¹⁸	2008	932	396	CT/MRI confirmed	rs17222814, rs10507391 and rs9551963	44.0	43.2	Age above 65 years: 76.2%	Age above 65 years: 78.8%	Caucasians	7
Quarta et al ¹⁹	2009	294	235	CT/MRI confirmed	rs9551963 and rs4769874	40.1	43.8	68–83	60–79	Caucasians	7
He et al ²⁰	2009	412	368	CT/MRI confirmed	rs10507391	42.7	38.0	71.5±7.8	70.3±6.9	Asians	8
Cheng et al ²¹	2010	280	258	CT/MRI confirmed	rs4073259	37.1	39.5	59.6±12.4	58.5±13.6	Asians	
Domingues-Montanari et al ²²	2010	1,092	781	CT/MRI confirmed	rs10507391	59.3	54.0	Spain cohort: 70.6±11.9 Portugal cohort: 52.4±9.3	Spain cohort: 72.1±6.9 Portugal cohort: 63.0±6.8	Caucasians	8
Gao et al ²³	2010	380	425	CT/MRI confirmed	rs10507391, rs17222814, rs4769874 and rs9551963	37.1	48.2	63.4±23.9	56.8±21.8	Asians	7
Kim et al ²⁴	2011	117	398	CT/MRI confirmed	rs17222919	45.3	51.3	65.5±12.1	53.8±15.2	Asians	6
Lee et al ²⁵	2011	291	278	CT/MRI confirmed	rs10507391 and rs4293222	19.9	41.0	65.4±10.6	64.1±11.1	Asians	7
Zhao et al ²⁶	2012	682	598	CT/MRI confirmed	rs17222814, rs10507391, rs4769874 and rs9551963	40.9	43.8	62.1±9.4	61.8±10.1	Asians	7
Wang et al ²⁷	2012	658	704	CT/MRI confirmed	rs17222919	40.0	44.9	69.4±10.6	69.2±9.0	Asians	6
Sharma et al ²⁸	2013	610	610	CT/MRI confirmed	rs10507391	32.0	32.3	49.3±17.3	49.0±16.8	Asians	8
Xu et al ²⁹	2013	547	794	CT/MRI confirmed	rs9551963 and rs10507391	NM	NM	NM	NM	Asians	7
Zhang et al ³⁰	2014	501	497	CT/MRI confirmed	rs4073259	33.9	39.6	58.7±10.2	53.1±8.6	Asians	8
Yao et al ³¹	2014	420	488	CT/MRI confirmed	rs10507391, rs4769874 and rs9551963	33.3	35.2	64.7±9.1	64.7±9.3	Asians	7
Yang et al ³²	2014	492	490	CT/MRI confirmed	rs10507391, rs4769874, rs9551963 and HapB	41.1	45.3	56.7±8.3	56.2±8.9	Asians	7

Fan et al ³³	2015	1,913	1,814	CT/MRI confirmed	rs17222919	46.6	48.4	Population 1: 56.1±10.6 Population 2: 60.5±7.8	Population 1: 55.3±10.3 Population 2: 60.5±7.8	Asians	8
Shao et al ³⁴	2015	459	462	CT/MRI confirmed	rs9551963, rs4769874, rs4769060 and rs10507391	39.9	40.3	48.6±11.0	63.8±9.2	Asians	7
Wang et al ³⁵	2015	396	300	CT/MRI confirmed	rs4769874 and rs10507391	40.9	35.3	56–71	54–75	Asians	7
Qu et al ³⁶	2015	456	452	CT/MRI confirmed	rs10507391, rs4769874, and rs9551963	34.9	33.4	59.7±11.5	53.7±8.2	Asians	7
Diakite et al ³⁷	2016	175	201	CT/MRI confirmed	rs10507391	43.4	NM	57.1±2.0	54.6±2.0	Caucasians	6
Hu et al ³⁸	2017	479	580	CT/MRI confirmed	rs10507391	11.1	12.4	65.2±5.3	65.1±5.7	Asians	7
Liu et al ³⁹	2017	346	399	CT/MRI confirmed	rs4793222	38.2	43.9	63.1±10.7	61.9±10.7	Asians	6
Shi et al ⁴⁰	2018	550	555	CT/MRI confirmed	rs4073259	49.8	49.9	Population 1: 60.7±12.8 Population 2: 60.5±12.5	Population 1: 60.9±13.3 Population 2: 62.5±13.2	Asians	7

Notes: HapA = SG13S25G–SG13S114T–SG13S89G–SG13S32A. HapB = SG13S377A–SG13S114A–SG13S41A–SG13S35G.

Abbreviations: CT, computed tomography; MRI, magnetic resonance imaging; NM, not mentioned; NOS, Newcastle–Ottawa Scale.

CI 0.90–1.07, $P=0.656$), indicating no association with IS (Figure 2C; Table 2).

Six studies provided data for the *ALOX5AP* rs17222814 polymorphism.^{13–15,18,23,26} Fixed-effect meta-analysis gave an estimated OR of 1.09 (95% CI: 0.96–1.24; $P=0.195$), indicating no association with IS in the overall population (Figure 2D; Table 2). There was no significant between-study heterogeneity ($I^2=25.3\%$, $P=0.244$). In the subgroup analysis by ethnicity, neither Asian nor Caucasian studies yielded significant findings (Table 2).

Four Asian studies in three publications investigated the role of the *ALOX5AP* rs17222919 polymorphism.^{24,27,33} Random-effects meta-analysis indicated no association of the SNP with IS (OR=0.89 for G allele vs T allele; 95% CI: 0.75–1.06; $P=0.175$) (Table 2). Similarly, four Asian studies providing summaries of data on the *ALOX5AP* rs4073259 polymorphism did not find any association with IS risk (OR=1.20 for A allele vs G allele; 95% CI: 1.00–1.45; $P=0.056$) (Table 2).^{21,30,40}

In addition to individual polymorphisms, we performed combined analysis for haplotypes. HapA was defined by rs17222814, rs10507391, rs4769874, and rs9551963 polymorphisms, with alleles G, T, G, and A. No association between HapA and IS risk was found (OR=1.20, 95% CI: 0.91–1.56, $P=0.192$) (Table 2). The pooled effect estimate for HapB (rs17216473A–rs10507391A–rs9315050A–rs17222842G) was similar, indicating no significant association with IS (OR=1.11, 95% CI: 0.90–1.38, $P=0.339$) (Table 2).

Publication bias and sensitivity analysis

Funnel plots did not suggest publication bias (Figure 3). The results of Begg's and Egger's tests for small-study effect biases were not significant, indicating no evidence of publication bias (Table 2). For the *ALOX5AP* rs10507391 and rs9551963 polymorphisms, sensitivity analyses were performed to assess the stability of the results by omission of studies, which were not in HWE. The significance of pooled ORs was not influenced.

Discussion

IS is the leading cause of morbidity and mortality in industrialized countries. Given the limited therapeutic options for IS patients, there remains great interest in identifying novel therapeutic target that may prevent stroke or improve stroke recovery.

The *ALOX5AP* gene encodes a major regulator for 5-lipoxygenase that catalyzes the oxidation of arachidonic acid to LTA4, which is released by inflammatory cells at injured sites and thus plays an important role in atherosclerosis

and other vascular damage.⁴¹ In the literature, there are apparently conflicting associations reported between *ALOX5AP* polymorphisms and IS risk. The conflicting results were possibly because of small effects of *ALOX5AP* polymorphisms on IS risk or the relatively low statistical power of published

studies. To summarize the available evidence, we combined studies involving a total of 32,782 participants. Six single polymorphisms and two haplotypes were analyzed. Our analysis did not support an association of the *ALOX5AP* rs10507391, rs4769874, rs9551963, rs17222814, rs17222919,

Figure 2 (Continued)

Figure 2 Forest plots for the association of *ALOX5AP* polymorphisms with IS risk.

Notes: (A) Meta-analysis of the *ALOX5AP* rs10507391 polymorphism and risk of IS using the random-effects model (A allele vs T allele). (B) Meta-analysis of the *ALOX5AP* rs4769874 polymorphism and risk of IS using the fixed-effects model (A allele vs G allele). (C) Meta-analysis of the *ALOX5AP* rs9551963 polymorphism and risk of IS using the random-effects model (A allele vs C allele). (D) Meta-analysis of the *ALOX5AP* rs17222814 polymorphism and risk of IS using the fixed-effects model (A allele vs G allele). Weights are from random-effects analysis.

Abbreviation: IS, ischemic stroke.

and rs4073259 polymorphisms with IS risk in Asian and Caucasian populations. Haplotype analysis also did not yield significant findings for the HapA (rs17222814G–rs10507391T–rs4769874G–rs9551963A) and HapB (rs17216473A–rs10507391A–rs9315050A–rs17222842G)

haplotypes. To the best of our knowledge, this is the largest and most comprehensive meta-analysis on the association between *ALOX5AP* polymorphisms and IS risk.

Our results are in agreement with a previous meta-analysis published in 2009. Zintzaras et al analyzed the

Table 2 Meta-analysis of association between *ALOX5AP* polymorphisms and IS risk

Polymorphism	Subgroup	Number of studies	Test of heterogeneity		Test of association		Test of publication	
			<i>I</i> ² (%)	P-value	OR (95% CI)	P-value	Begg's test	Egger's test
rs10507391	Total	20	81.6	<0.001	1.03 (0.93–1.14)	0.557	0.974	0.857
	Asians	14	81.2	<0.001	1.10 (0.98–1.23)	0.109	NA	NA
	Caucasians	6	68.7	0.008	0.89 (0.76–1.02)	0.101	NA	NA
	HWE	19	80.8	<0.001	1.05 (0.95–1.16)	0.318	NA	NA
rs4769874	Total	13	28.2	0.160	1.13 (1.00–1.28)	0.050	1.000	0.909
	Asians	9	10.9	0.344	1.10 (0.95–1.28)	0.210	NA	NA
	Caucasians	4	59.2	0.061	1.19 (0.96–1.48)	0.106	NA	NA
rs9551963	Total	12	44.7	0.047	1.03 (0.96–1.11)	0.372	0.451	0.365
	Asians	7	58.6	0.025	1.07 (0.96–1.19)	0.230	NA	NA
	Caucasians	5	0	0.484	0.98 (0.90–1.07)	0.656	NA	NA
	HWE	10	53.3	0.023	1.04 (0.95–1.14)	0.381	NA	NA
rs17222814	Total	6	25.3	0.244	1.09 (0.96–1.24)	0.195	0.256	0.137
	Asians	2	80.4	0.025	1.12 (0.89–1.41)	0.354	NA	NA
	Caucasians	4	0	0.704	1.08 (0.92–1.26)	0.348	NA	NA
rs17222919	Total	4	67.9	0.025	0.89 (0.75–1.06)	0.175	0.308	0.541
	Asians	4	67.9	0.025	0.89 (0.75–1.06)	0.175	0.308	0.541
rs4073259	Total	4	63.7	0.041	1.20 (1.00–1.45)	0.056	0.238	0.156
	Asians	4	63.7	0.041	1.20 (1.00–1.45)	0.056	0.238	0.156
HapA	Total	4	65.6	0.033	1.20 (0.91–1.56)	0.192	0.362	0.621
	Caucasians	4	65.6	0.033	1.20 (0.91–1.56)	0.192	0.362	0.621
HapB	Total	4	0	0.835	1.11 (0.90–1.38)	0.339	0.734	0.251
	Asians	1	NA	NA	1.16 (0.84–1.62)	0.370	NA	NA
	Caucasians	3	0	0.692	1.07 (0.80–1.44)	0.627	NA	NA

Abbreviations: HWE, Hardy–Weinberg equilibrium; IS, ischemic stroke; NA, not applicable.

relationship between *ALOX5AP* polymorphisms and risk of stroke with a total of 5,194 cases and 4,566 controls, finding no association of rs17222814, rs10507391, rs4769874, rs9551963, HapA, and HapB with stroke.⁴² Although the two meta-analyses obtained similar findings, there were still some differences. First, both hemorrhagic and IS patients were included in the Zintzaras et al's study, whereas we only took into account IS patients. Second, Zintzaras et al's meta-analysis overtook some Asian studies^{16,17,20} because they excluded the non-English papers. Third, Zintzaras et al's analysis arose from 12 studies, considerably less in number (5,194 cases and 4,566 controls) than the overall 30 in our study (16,294 cases and 16,488 controls). Besides Zintzaras et al's analysis, the Wang et al's⁴³ meta-analysis involving 10 studies and the Chen et al's⁴⁴ meta-analysis combining seven studies investigated rs10507391.^{43,44} The positive association between rs10507391 and IS risk reported by these two small size meta-analyses was not confirmed by our study applying a large sample.

It should be mentioned that much effort has been performed to conduct appropriately the meta-analysis and avoid any possible source of bias. To avoid the local literature bias, we included both English and non-English articles. We properly assessed deviations from HWE in the control population and performed sensitivity analysis excluding studies where the sample violated HWE to ensure stability of the pooled ORs. In addition, we evaluated small-study effects and publication bias through Begg's and Egger's tests; there was no evidence of systematic missingness of scientific results from the literature. Finally, subgroup analysis by ethnicity confirmed the lack of association between the *ALOX5AP* polymorphisms and IS risk not only in the overall populations but also in each ethnic group.

Previous studies demonstrated the presence of strong linkage disequilibrium in the *ALOX5AP* gene.^{14,15,32} It is believed that haplotype analysis can provide more information than single-locus analysis because it allows us to explore potential interactions among alleles. In this meta-analysis,

Figure 3 Funnel plots to detect publication bias.

Notes: (A) Funnel plot analysis to detect publication bias for the *ALOX5AP* rs10507391 polymorphism and risk of IS. (B) Funnel plot analysis to detect publication bias for the *ALOX5AP* rs4769874 polymorphism and risk of IS. (C) Funnel plot analysis to detect publication bias for the *ALOX5AP* rs9551963 polymorphism and risk of IS. (D) Funnel plot analysis to detect publication bias for the *ALOX5AP* rs17222814 polymorphism and risk of IS.

Abbreviations: IS, ischemic stroke; SE, standard error.

we evaluated the relationship of the HapA (rs17222814G–rs10507391T–rs4769874G–rs9551963A) and HapB (rs17216473A–rs10507391A–rs9315050A–rs17222842G) haplotypes with IS risk. Although no associations of these haplotypes with IS were identified, we could not exclude that other haplotypes may play a role in the development of IS. The study by Zhao et al reported a statistically significant association between the haplotype GCGA constructed by rs17222814, rs10507391, rs4769874, and rs9551963 and IS risk.²⁶ In addition, Yao et al found that the haplotype TGC constructed using rs10507391, rs4769874, and rs9551963 was associated with the increased risk of IS (OR=1.60, 95% CI: 1.28–1.98, $P<0.001$), while the haplotype AGA was

protective (OR=0.66, 95% CI: 0.53–0.81) in a case–control study, involving a total of 420 cases and 488 controls.³¹ Considering the large sample sizes applied by these studies, further studies are warranted to confirm their findings.

Our study has some limitations that should be considered. First, the relationship between the *ALOX5AP* polymorphisms and the levels of leukotriene lipid mediators in IS patients was not evaluated due to lack of published data. Increased levels of products of the 5-lipoxygenase (5-LO)/5-LO-activating protein (FLAP) pathway including LTB₄ had been reported in IS patients.⁴⁵ The evaluation of potential interactions between *ALOX5AP* polymorphisms and leukotriene lipid mediators in IS patients could provide valuable insights

into the development of IS. Second, because there was a paucity of data, we did not assess gender differences in the effects of *ALOX5AP* polymorphisms on IS risk. It is widely known that gender has a complex and interactive effect on IS risk and there are gender differences in the prevalence of IS.⁴⁶ More research is needed to evaluate gender differences in the association of *ALOX5AP* polymorphisms with IS development.

In conclusion, our meta-analysis of 30 studies involving 16,294 IS patients and 16,488 controls indicates no significant association of the *ALOX5AP* polymorphisms rs10507391, rs4769874, rs9551963, rs17222814, rs17222919, and rs4073259 with IS risk. In addition, the HapA and HapB haplotypes also did not show any association with IS.

Acknowledgments

This work was supported, in part, by the National Natural Science Foundation of China (81660776 and 81360535), the Natural Science Foundation of Guangxi Zhuang Autonomous Region (2016JJA140115), and the Science Research and Technology Development Project of Guangxi Zhuang Autonomous Region (1598012-55).

Disclosure

The authors report no conflicts of interest in this work.

References

1. Venketasubramanian N, Yoon BW, Pandian J, Navarro JC. Stroke epidemiology in south, east, and south-east Asia: a review. *J Stroke*. 2017;19(3):286–294.
2. Malik R, Dichgans M. Challenges and opportunities in stroke genetics. *Cardiovasc Res*. 2018;114(9):1226–1240.
3. Mashima R, Okuyama T. The role of lipoxygenases in pathophysiology; new insights and future perspectives. *Redox Biol*. 2015;6:297–310.
4. Bäck M, Sultan A, Ovchinnikova O, Hansson GK. 5-Lipoxygenase-activating protein: a potential link between innate and adaptive immunity in atherosclerosis and adipose tissue inflammation. *Circ Res*. 2007;100(7):946–949.
5. Ström JO, Strid T, Hammarström S. Disruption of the *alox5ap* gene ameliorates focal ischemic stroke: possible consequence of impaired leukotriene biosynthesis. *BMC Neurosci*. 2012;13:146.
6. Hakonarson H, Thorvaldsson S, Helgadóttir A, et al. Effects of a 5-lipoxygenase-activating protein inhibitor on biomarkers associated with risk of myocardial infarction: a randomized trial. *JAMA*. 2005; 293(18):2245–2256.
7. Stanke-Labesque F, Pépin JL, de Jouvenel T, et al. Leukotriene B4 pathway activation and atherosclerosis in obstructive sleep apnea. *J Lipid Res*. 2012;53(9):1944–1951.
8. Moher D, Liberati A, Tetzlaff J, Altman DG, PRISMA Group. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *BMJ*. 2009;339(jul21 1):b2535–336.
9. Stang A. Critical evaluation of the Newcastle-Ottawa scale for the assessment of the quality of nonrandomized studies in meta-analyses. *Eur J Epidemiol*. 2010;25(9):603–605.
10. DerSimonian R, Laird N. Meta-analysis in clinical trials. *Control Clin Trials*. 1986;7(3):177–188.
11. Helgadóttir A, Manolescu A, Thorleifsson G, et al. The gene encoding 5-lipoxygenase activating protein confers risk of myocardial infarction and stroke. *Nat Genet*. 2004;36(3):233–239.
12. Helgadóttir A, Gretarsdóttir S, St Clair D, et al. Association between the gene encoding 5-lipoxygenase-activating protein and stroke replicated in a Scottish population. *Am J Hum Genet*. 2005;76(3):505–509.
13. Meschia JF, Brodt TG, Brown RD, et al. Phosphodiesterase 4D and 5-lipoxygenase activating protein in ischemic stroke. *Ann Neurol*. 2005;58(3):351–361.
14. Löhmußaar E, Gschwendtner A, Mueller JC, et al. *ALOX5AP* gene and the *PDE4D* gene in a central European population of stroke patients. *Stroke*. 2005;36(4):731–736.
15. Zee RY, Cheng S, Hegener HH, Erlich HA, Ridker PM. Genetic variants of arachidonate 5-lipoxygenase-activating protein, and risk of incident myocardial infarction and ischemic stroke: a nested case-control approach. *Stroke*. 2006;37(8):2007–2011.
16. Zhang WL, Yang XM, Shi J, Sun K, Hui RT. Polymorphism of SG13S114T/A in the *ALOX5AP* gene and the risk for stroke in a large Chinese cohort. *Yi Chuan Xue Bao*. 2006;33(8):678–684.
17. Gao ZS, Qc D. To study relationship between polymorphism of SG13S114T/A and SG13S89G/A in the *ALOX5AP* gene and thrombotic stroke. *Ningxia Med J*. 2008;30(5):388–390.
18. Lövkvist H, Smith JG, Luthman H, et al. Ischaemic stroke in hypertensive patients is associated with variations in the *PDE4D* genome region. *Eur J Hum Genet*. 2008;16(9):1117–1125.
19. Quarta G, Stanzione R, Evangelista A, et al. Phosphodiesterase 4D and 5-lipoxygenase activating protein genes and risk of ischemic stroke in Sardinians. *Eur J Hum Genet*. 2009;17(11):1448–1453.
20. He YL, Zhu M, Jin XP, Zhou YL. Relationship of polymorphism of SG13S114A/T in *ALOX5AP* gene with atherosclerotic cerebral infarction. *Zhejiang Da Xue Xue Bao Yi Xue Ban*. 2009;38(6):630–633.
21. Cheng H, Jin QW, Ding XS. Association of *ALOX5AP* and *PDE4D* genes with the risk of lacunar infarction in Nanjing city of Jiangsu province. *Acta Univ Med Nanjing*. 2010;30(1):39–45.
22. Domingues-Montanari S, Fernández-Cadenas I, del Río-Espinola A, et al. Association of a genetic variant in the *ALOX5AP* with higher risk of ischemic stroke: a case-control, meta-analysis and functional study. *Cerebrovasc Dis*. 2010;29(6):528–537.
23. Gao Z, Tian GP. Relationship between *ALOX5AP* gene single nucleotide polymorphism and stroke of Han population in Shenyang City. *J Apop Nerve Dis*. 2010;27(11):1022–1026.
24. Kim DH, Ahn WY, Kim DK, et al. A Promoter polymorphism (rs17222919, -1316T/G) of *ALOX5AP* is associated with intracerebral hemorrhage in Korean population. *Prostaglandins Leukot Essent Fatty Acids*. 2011;85(3–4):115–120.
25. Lee JD, Lee TH, Huang YC, et al. *ALOX5AP* genetic variants and risk of atherothrombotic stroke in the Taiwanese population. *J Clin Neurosci*. 2011;18(12):1634–1638.
26. Zhao J, Wang X, Xu J, et al. Association of inflammatory response gene polymorphism with atherothrombotic stroke in Northern Han Chinese. *Acta Biochim Biophys Sin*. 2012;44(12):1023–1030.
27. Wang Y, Wang GN, Sun H, Chen C, Xiao H, Zhang JS. Association of *ALOX5AP* with ischemic stroke in eastern Chinese. *World J Emerg Med*. 2012;3(2):108–113.
28. Sharma V, Dadheech S, Kaul S, Jyothy A, Munshi A. Association of *ALOX5AP1* SG13S114T/A variant with ischemic stroke, stroke subtypes and aspirin resistance. *J Neurol Sci*. 2013;331(1–2):108–113.
29. Xu C, Qiang L, Limei C, et al. Correlation between cerebral infarction and *ALOX5AP* gene expression. *Cell Biochem Biophys*. 2013; 67(3):899–904.
30. Zhang R, Guo X, Li X, et al. Arachidonate 5-lipoxygenase-activating protein (*ALOX5AP*) gene rs4073259 polymorphism not associated with ischemic stroke in the northeastern Chinese Han population. *Clin Neurol Neurosurg*. 2014;119:64–69.
31. Yao Q, Zhang C, Zhang X, et al. Synergistic effect of *ALOX5AP* polymorphisms and cigarette smoking on the risk of atherosclerotic cerebral infarction in a Northern Han Chinese population. *J Clin Neurosci*. 2014;21(6):975–979.

32. Yang D, He Y, Li M, et al. A novel risk haplotype of ALOX5AP gene is associated with ischemic stroke in Chinese Han population. *J Mol Neurosci*. 2014;53(3):493–499.
33. Fan Y, Chen H, Li A, et al. A promoter polymorphism (rs17222919, -1316T/G) of ALOX5AP gene is associated with decreased risk of ischemic stroke in two independent Chinese populations. *PLoS One*. 2015;10(3):e0122393.
34. Shao M, Yi X, Chi L, Lin J, Zhou Q, Huang R. Ischemic stroke risk in a southeastern Chinese population: insights from 5-lipoxygenase activating protein and phosphodiesterase 4D single-nucleotide polymorphisms. *J Formos Med Assoc*. 2015;114(5):422–429.
35. Wang JN, Yang ZN, Sui Q. The polymorphism of ALOX5AP and PDE4D gene and molecular epidemiology of the susceptibility of cerebral infarction. *J Bengbu Med Coll*. 2015;40(10):1338–1340.
36. Qu Z, Su F, Zhu Y, et al. A tagging ALOX5AP polymorphism and risk of ischemic stroke in a northeastern Chinese Han population. *Int J Clin Exp Med*. 2015;8(11):21343–21350.
37. Diakite B, Hamzi K, Hmimch W, Nadifi S, GMRAVC. Genetic polymorphisms of T-1131C APOA5 and ALOX5AP SG13S114 with the susceptibility of ischaemic stroke in Morocco. *J Genet*. 2016;95(2):303–309.
38. Hu X, Wang J, Li Y, et al. The β -fibrinogen gene 455G/A polymorphism associated with cardioembolic stroke in atrial fibrillation with low CHA₂DS₂-VaSc score. *Sci Rep*. 2017;7(1):17517.
39. Liu D, Liu L, Song Z, Hu Z, Liu J, Hou D. Genetic variations of oxidative stress related genes ALOX5, ALOX5AP and MPO modulate ischemic stroke susceptibility through main effects and epistatic interactions in a Chinese population. *Cell Physiol Biochem*. 2017;43(4):1588–1602.
40. Shi Y, Xu L, Feng Q, et al. Allele-specific methylation contributed by CpG-SNP is associated with regulation of ALOX5AP gene expression in ischemic stroke. *Neurol Sci*. 2018;39(10):1717–1724.
41. Pasterkamp G, van der Laan SW, Haitjema S, et al. Human validation of genes associated with a murine atherosclerotic phenotype. *Arterioscler Thromb Vasc Biol*. 2016;36(6):1240–1246.
42. Zintzaras E, Rodopoulou P, Sakellariadis N. Variants of the arachidonate 5-lipoxygenase-activating protein (ALOX5AP) gene and risk of stroke: a HuGE gene-disease association review and meta-analysis. *Am J Epidemiol*. 2009;169(5):523–532.
43. Wang G, Liu R, Zhang J. The arachidonate 5-lipoxygenase-activating protein (ALOX5AP) gene SG13S114 polymorphism and ischemic stroke in Chinese population: a meta-analysis. *Gene*. 2014;533(2):461–468.
44. Chen Z, Zheng J, Liu W, et al. The SG13S114 polymorphism of the ALOX5AP gene is associated with ischemic stroke in Europeans: a meta-analysis of 8062 subjects. *Neurol Sci*. 2017;38(4):579–587.
45. Wang G, Wang Y, Sun H, et al. Variants of the arachidonate 5-lipoxygenase-activating protein (ALOX5AP) gene and risk of ischemic stroke in Han Chinese of eastern China. *J Biomed Res*. 2011;25(5):319–327.
46. Roy-O'Reilly M, McCullough LD. Age and sex are critical factors in ischemic stroke pathology. *Endocrinology*. 2018;159(8):3120–3131.

Neuropsychiatric Disease and Treatment

Publish your work in this journal

Neuropsychiatric Disease and Treatment is an international, peer-reviewed journal of clinical therapeutics and pharmacology focusing on concise rapid reporting of clinical or pre-clinical studies on a range of neuropsychiatric and neurological disorders. This journal is indexed on PubMed Central, the 'PsycINFO' database and CAS,

Submit your manuscript here: <http://www.dovepress.com/neuropsychiatric-disease-and-treatment-journal>

Dovepress

and is the official journal of The International Neuropsychiatric Association (INA). The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.