Does experimentally induced pain affect attention? A meta-analytical review

Wenxiao Gong^{1,2}

Lu Fan^{1,2}

Fei Luo^{1,2}

¹CAS Key Laboratory of Mental Health, Institute of Psychology, Chinese Academy of Sciences, Beijing 100101, People's Republic of China;

²Department of Psychology, University of Chinese Academy of Sciences, Beijing 100049, People's Republic of China

Background: Recent studies have found that clinical pain is related to cognitive impairment. However, there remains a scarcity of systematic reviews on the influence of acute pain on attention. Laboratory-induced pain is often used to simulate acute pain. The current systematic meta-analysis aimed to evaluate the effect of induced-pain on three components of attention (orienting, alerting, and executive attention) in healthy subjects.

Methods: A systematic search of three databases was performed. Only data from studies that administered laboratory-induced pain and that also included a control group were selected. The effects of experimental pain on orienting attention, alerting attention, and executive attention were analyzed. Two reviewers assessed the studies and extracted relevant data according to the Cochrane Collaboration and Preferred Reporting Items for Systematic Reviews and Meta-Analysis Guidelines.

Results: Eight studies were included in the meta-analysis. Orienting attention was marginally interrupted by pain under the invalid cue and marginally facilitated by pain under the valid cue condition. Performance on alerting attention was decreased by pain. Executive attention was not significantly affected by pain.

Conclusion: There was moderate evidence that experimentally induced pain can produce effects on orienting and alerting attention but not on executive attention. This meta-analysis suggests that experimentally induced pain influences some aspects of attention.

Keywords: experimental pain, attention, meta-analysis

Introduction

Investigations into human cognition have long sought to define attention and to describe the processes involved. Models of attention agree that in addition to cognitive resources, automatic behaviors and thoughts are required. Moreover, there is a consensus that more cognitive resources are allocated to essential information, and multiple tasks are difficult to maintain concurrently. Attention governs our perception of the world and the voluntary control of our thoughts and emotions.¹

Recent neuroimaging studies have provided new perspective on the processes involved in attention.^{2,3} Based on neuroimaging technology, Posner's Attention Networks has been accepted as a framework for the cognitive underpinnings of attention. Posner and Rothbart suggest that there are three distinct networks for attention: orienting, alerting, and executive attention.¹ Orienting involves the selection of information from sensory input, alerting is defined as achieving and maintaining of high sensitivity to approaching stimuli, and executive attention includes a set of mechanisms

Correspondence: Fei Luo
CAS Key Laboratory of Mental Health,
Institute of Psychology, Chinese Academy
of Sciences, 16 Lincui Road, Chaoyang
District, Beijing 100101, People's
Republic of China
Email luof@psych.ac.cn

responsible for monitoring and addressing conflict among feelings, thoughts, and responses.

Pain is a salient signal that demands a high level of cognitive resources. Pain warns an organism of a threat, which may interfere, distract, and demand attention.⁴ Most of the existing reviews on the psychological mechanisms involved in pain have been conducted with chronic pain patients.⁵ It is widely accepted that chronic pain leads to attention deficits, predominantly on tasks that require high attentional demands.⁶ This prior research has contributed greatly to the understanding of pain and has provided avenues for treatment. However, the extent to which and how acute pain interferes with attention remains unclear.

For healthy people, pain is an unpleasant sensory and emotional experience. It is a protective mechanism signaling an immediate response to avoid further injury. Therefore, attention to pain can be positive and adaptive. Despite this, the mechanisms by which acute pain alters attention are not well understood. Prior studies using techniques that mimic pain, such as cold pressure, heat, and electric stimuli, have demonstrated that laboratory-induced pain decreases overall attention performance.⁷ Some studies have revealed that experimentally induced pain draws attention to its location in healthy individuals.⁸

However, the effect of experimentally induced pain on attention differs from chronic pain.^{9,10} Despite frequent reports that laboratory-induced pain leads to poor concentration,¹¹ there have been no systematic reviews of the literature. Our meta-analysis examined the effect of laboratory-induced pain on attention, which is often used to simulate acute pain. Immediate protective responses to acute pain include increased arousal, orientation to the sources of threat, and various safety-seeking behaviors including escape and avoidance. And attention protects the pursuit of current goals, by inhibiting the processing of goal-irrelevant information.¹² These three components are more closely related to acute pain. So the current review focuses on the effects of experimentally induced pain on three components of attention: orienting, alerting, and executive.

Gaining a better understanding of attention performance, as well as the processes that underpin it, within the context of experimentally induced pain may provide additional information that may assist in the process of choosing the most beneficial type of intervention for a given client.

Methods

Data sources

The current meta-analysis was conducted following the Cochrane Collaboration and Preferred Reporting Items for

Systematic Review and Meta-Analyses (“PRISMA”) Statement guidelines. Searches were performed in the following databases: PubMed, Web of Science, and PsychINFO. The following keywords were used as search items: orienting, alerting, executive attention, spatial cue task, continuous performance, Stroop, go/no-go, attention, and intersected with pain. The search results were limited to studies using human participants and studies that are related to the interruptive effect of experimental pain. Only articles written in English were included. Studies related to alerting, orienting, executive attention, and experimentally induced pain were exported to Endnote X7. Full-text articles published in the area of induced pain and attention, which were identified through background reading and systematic searching were retrieved. All the articles were hand-searched for citations containing original data. Figure 1 shows the search and selection process.

Study selection

Studies related to the effect of experimentally induced pain on orienting,^{10,13–16} alerting,^{10,17} and executive attention^{18,19} were selected. Each of the components of attention were measured by well-known paradigms, such as spatial cue test, continuous performance task (CPT), Stroop test,¹⁸ and go/no-go test.¹⁹ Selection criteria were also established, and irrelevant studies were removed.

The spatial cue task, in which a cue signals either a correct (valid cue) or incorrect (invalid cue) location of a target, is used to investigate orienting of attention. Orienting is facilitated when the target appears at the same location as the cue (valid cue). It is inhibited when the target appears at a different location than the cue (invalid cue).²⁰ The difference in reaction times (RTs) between a painful and a nonpainful conditions signifies an effect on orienting. Studies that use pain as a cue were included.^{13–16} In the spatial cue task, a cue was presented on left/right side of the screen, and the cue (or the other location) is replaced by a probe stimulus. Participants are instructed to respond as quickly as possible to the probe. During the orientation, participants respond to the probe as the location of the cue. Pain as a cue would capture participants' attention, which affects the response to the probe. So, pain as a cue is also an immediate effect on orienting. Considering the use of pain as a cue may lead to shift the attention to the location of the painful cues, which would affect the orienting, we singled out these four studies using pain as a cue (see the “Results” section).

Achieving and maintaining a high sensitivity to approach stimuli¹ during an extended task is necessary for alerting

Figure 1 Flowchart of the article search and inclusion process.

attention. Alertness is measured by the CPT where participants are asked to attend to a subject or location during the task and respond to the target. The difference in RT between a painful and a nonpainful conditions signifies an effect on alerting.

Executive attention involves mechanisms for monitoring and resolving conflicts among thoughts, feelings, and responses. Two tests were selected for the analysis of executive attention: the Stroop test and the go/no-go test. The former measures the ability to resolve cognitive conflicts, the latter measures the ability to resolve response conflicts. The difference in RT between painful and nonpainful conditions signifies an effect of pain on executive attention.

All studies included in the meta-analysis compared attention performance under an experimentally induced pain condition with a non-pain condition. Titles and abstracts of articles were screened by two authors (WG and LF). Studies were included if they evaluated orienting, alerting, or executive attention and used spatial cue test, CPT, Stroop test,

or go/no-go test. Irrelevant articles were removed. Studies that used pain-related (eg, pain-related images or words) information instead of the sensory experience of pain were excluded because this topic has been previously studied.⁵ Other exclusion criteria were: 1) child participants; 2) studies that included participants with traumatic brain injury or any disease that would be expected to impair attention. The full texts of potentially eligible studies were retrieved and then formally assessed for eligibility by the two reviewers, using the aforementioned criteria.

Risk of bias assessment

Risk of bias was assessed by two reviewers using a customized risk of bias form. The form was developed a priori and was based on relevant items from a previous meta-analysis that used the Cochrane Collaboration Risk of Bias Tool.²¹ Any differences were resolved by discussion between two reviewers.

Data extraction

Two independent reviewers used a previously piloted, customized form to extract data.²¹ RT results were compared. The rate of correct answers was not analyzed as the tasks were very simple. Extracted data included group-specific data (type of induced pain, sample size in each group, gender, and mean age $\pm$ SD); attention test data (name of attention test), cognitive process evaluated (where it was specified), type of outcome measure of test (RT), interpretation of test (group-specific outcomes on attentional test results; eg, mean differences) (Table 1).

Data analysis

Data were analyzed using Review Manager (Version 5.3). If sufficient data were not reported, the study was not included in the analysis.^{22–29} Studies lacking a control group were excluded from analysis.⁸ Heterogeneity was deemed significant on the basis of $\chi^2 P < 0.01$ and substantial on the basis of $I^2 > 60\%$.

Evaluation of attention was evaluated – tests and test outcomes

From the 1,325 records identified by the search methods, the full texts of 96 studies were retrieved. Of the 96 studies, 17 reported effects of experimentally induced pain on attention but nine of the 17 reported no behavioral data or control groups,^{8,22–29} leading to a total of eight that met all inclusion criteria. In total, 26 different test outcomes resulted from tests of eight different attention constructs. The eight included studies reported outcomes from tests that are commonly used to investigate orienting, alerting, and executive attention: the spatial cue test,^{10,13–16} the CPT,^{10,17} the Stroop task,¹⁸ and the go/no-go task.¹⁹

Orienting

Five studies were included in the meta-analysis on orienting attention.^{10,13–16} All these studies used the spatial cue task to measure RT under valid and invalid conditions. The statistical results of these studies were analyzed separately for each condition. Four of the studies^{13–16} used cues that were modulated by classical conditioning as follows. The conditioned cue was followed by a nociceptive stimulus, thus becoming a signal for pain. These data (pain preceded by a conditioned cue) were included in the pain condition in the meta-analysis. Data from the conditioned cue that was not followed by nociceptive stimulus were included in the no-pain condition in the meta-analysis.^{10,13–16}

Table 1 Characteristics of included studies

Study	Pain stimulation	Matched or tested for baseline in characteristics	Participants				Non-pain			
			Induced pain							
			Age mean ± (SD)	Gender	Number	Age mean ± (SD)	Gender	Number		
Babiloni et al (2004) ¹⁹	Electrical stimulation	Within-group	25 (not described)	Not described	12	25 (not described)	Not described	12		
Cheng et al (2017) ¹⁸	TENS (painful transcutaneous electrical nerve stimuli)	Within-group	20–31	25 M, 26 F	51	20–31	25 M, 26 F	51		
Kurita et al (2015) ¹⁷	Cuff	Within-group	23.5 (2.1)	All M	22	23.5 (2.1)	All M	22		
Koster et al (2005) ¹³	An aversive white noise burst CS	Within-group	19.32 (2.36)	55 F, 11 M	66	19.32 (2.36)	55 F, 11 M	66		
Moore et al (2012), ¹⁰ E1	Thermal pain (ATS)	Within-group	23.5 (3.86)	13 F, 7 M	20	23.5 (3.86)	13 F, 7 M	20		
Moore et al (2012), ¹⁰ E3	Thermal pain (ATS)	Within-group	24.5 (5.59)	16 F, 4 M	20	24.5 (5.59)	16 F, 4 M	20		
Moore et al (2012), ¹⁰ E5	Thermal pain (ATS)	Within-group	27.7 (8.76)	14 F, 6 M	20	27.7 (8.76)	14 F, 6 M	20		
Van Damme et al (2004) ¹⁴	Transcutaneous electrocutaneous stimulus CS	Within-group	19.04 (not described)	44 F, 8 M	52	19.04 (not described)	44 F, 8 M	52		
Van Damme et al (2006) ¹⁵	Electrocutaneous stimulus CS	Within-group	19.3 (1.38)	15 F, 12 M	27	19.3 (1.38)	15 F, 12 M	27		
Van Damme et al (2012) ¹⁶	Electrocutaneous stimulus	Between-group	(Not described)	(Not described)	51 (E1) 50 (E2)	(Not described)	(Not described)	51 (E1) 50 (E2)		

Abbreviations: ATS, advanced thermal stimulator; CS, conditioned stimulus; E, experiment; F, female; M, male.

Alerting

Seven comparisons in two studies were included in the alerting meta-analysis. Two studies used CPT.^{10,17} One of the studies conducted CPT under a pain condition.¹⁰ The other study using CPT was performed under both mild and moderate pain. RT was summarized by percentiles: 10th (fastest RT), 50th (plotted in), and 90th (slowest RT).¹⁷

Executive attention

Three comparisons in two studies were included in the executive attention meta-analysis. One study used the go/no-go test,¹⁹ and the other one used a numerical Stroop test.¹⁸ Using the Stroop test, Cheng et al¹⁸ characterized individuals as A-type (attention dominates) or P-type (pain dominates) based on how pain interfered with task performance. Both groups were included in the meta-analysis.

Results

Risk of bias

All studies included in the meta-analysis were judged to have a high risk of bias, primarily due to ambiguous recruitment methods. None of the included studies reported a priori calculation of the sample size required for adequate power. None of the included study used a double-blind experimental design. Risk of bias data are summarized in Table 2.

Behavioral outcomes

All the included studies reported on RT in one or more tests of attention. The eight studies presented 26 outcomes for RT, across the three components of attention: orienting, alerting, and executive attention. Table 3 summarizes the outcomes, and Table 4 summarizes the comparisons for three components of attention.

Orienting attention

Results are separated into valid and invalid cue conditions.

Valid cues

Valid cues are presented at the same location as the targets. Pooled results from eight comparisons across five studies indicated that induced-pain was associated with facilitated orienting under the valid cue condition. That is, there was a nearly significant effect estimate of -7.02 (95% CI $= -14.36$ to 0.32). Nonsignificant heterogeneity was detected ($\chi^2=3.88$, $P=0.79$, $I^2=0\%$). The results suggest that healthy individuals orient faster when they are in pain vs nonpain under the valid cue condition (Figure 2).

Invalid cue

Invalid cues are presented at a different location than the target. Pooled results from eight comparisons across five studies indicated that induced pain was moderately associated with impaired orienting under the invalid cue condition. That is, there was a marginal significant effect estimate of 7.14 (95% CI $= -0.66$ to 14.93). Nonsignificant heterogeneity was detected for orientation under the invalid cue ($\chi^2=2.52$, $P=0.93$, $I^2=0\%$). The results suggest that the healthy individuals orient slower to the invalid cue when under the pain condition compared with no-pain condition (Figure 3).

Attentional bias to pain

Four of five studies^{13–16} on orienting used a conditioned cue to indicate pain, while the other one⁹ did the experiment under pain condition. The four studies using the painful cue were analyzed in a separate meta-analysis. The results showed that induced pain under the painful valid cue showed a significant effect

Table 2 Risk of bias assessment

Study	Random sequence generation (selection bias)	Sample size was calculated a priori	Blinding of participants and personnel	Cases diagnosed were according to accepted criteria	Confounding variables were controlled	All outcomes and groups were reported
Babiloni et al (2004) ¹⁹	NA	N	N	Y	Y	Y
Cheng et al (2017) ¹⁸	NA	N	N	Y	Y	Y
Kurita et al (2015) ¹⁷	NA	N	N	Y	Y	Y
Koster et al (2005) ¹³	NA	N	N	Y	Y	Y
Moore et al (2012), ¹⁰ E1	NA	N	NA	Y	Y	Y
Moore et al (2012), ¹⁰ E3	NA	N	NA	Y	Y	Y
Moore et al (2012), ¹⁰ E5	NA	N	NA	Y	Y	Y
Van Damme et al (2004) ¹⁴	NA	N	NA	Y	Y	Y
Van Damme et al (2006) ¹⁵	NA	N	N	Y	Y	Y
Van Damme et al (2012) ¹⁶	NA	N	N	Y	Y	Y

Abbreviations: N, no; NA, not applicable; Y, yes.

Table 3 The outcomes for three components of attention

Cognitive process	Test	Outcome	Studies that use this test
Orienting attention	Spatial cue test – one cue (CS+) of the task was a signal for an aversive white noise burst (UCS) and that another cue (CS–) signaled its nonoccurrence	Average time to the target, the test trials consisted of 75% valid trials and 25% invalid trials Acquisition phase: 72 CS+, 72 CS–, 12 catch, and 10 digit trials Extinction phase 2: 36 CS+, 36 CS–, 6 catch, and 5 digit trials	Koster et al. (2005); ¹³ Van Damme et al (2006) ¹⁵
	Spatial cue test – shifting attention toward the direction of an endogenous arrowhead once without pain stimulation, once with a warm sensation, and once under thermal heat pain conditions	Average time to the target, the test trials consisted of valid, invalid and neutral (a horizontal line), or absent Valid: 160 trials; invalid: 40 trials; neutral: 40 trials; absent: 40 trials	Moore et al (2012) ¹⁰
	Spatial cue test – one cue (CS+) of the task was a signal for an aversive white noise burst (UCS) and that another cue (CS–) signaled its nonoccurrence	Average time to the target, the test trials consisted of 96 valid trials, 48 invalid trials, 24 catch trials, and 12 digit trials	Van Damme et al (2004) ¹⁴
	Spatial cue test – visual cues were LEDs presented close to the left or right hand, or centrally between both hands LED cues were followed by a somatosensory stimulus to one of both wrists	Average time to the LED target Experiment 1: 12 valid trials, 12 invalid trials, 12 baseline trials, and 12 catch trials Experiment 2: 36 valid trials, 12 invalid trials, 12 baseline trials, and 12 catch trials	Van Damme et al (2012) ¹⁶
Alerting attention	Continuous performance test	The reaction time to auditory signals at random second intervals under mild and moderate pain. The results were summarized using 10th, 50th, and 90th percentiles, in which 10th represents the fastest and 90th the slowest values	Kurita et al (2015) ¹⁷
	Continuous performance test	The reaction time to respond with a single key press when three consecutive odd or even digits were presented	Moore et al (2012) ¹⁰
Executive attention	Go/no-go task	Average time to go trials under pain or nonpain condition	Babiloni et al (2004) ¹⁹
	Stroop task – numerical interference task counts the number of digits within each box, which the numerical values of the digits did not coincide with the number of digits counted in each box	Average time under no-pain and pain blocks	Cheng et al (2017) ¹⁸

Abbreviations: CS, conditioned stimulus; CS+, conditioned stimulus was presented; CS–, conditioned stimulus disappeared; UCS, unconditioned stimulus.

Table 4 The comparisons for three components of attention

Cognitive process	Paper	Studies	Comparisons
Orienting attention (valid/invalid cue)	Koster et al (2005) ¹³	Acquisition phase	Mean reaction data that used CS+/CS− signals
		Extinction phase 2	Mean reaction data that used CS+/CS− signals
	Moore et al (2012) ¹⁰	Experiment 3	Mean reaction data under pain and baseline conditions
	Van Damme et al (2004) ¹⁴	Experiment phase	Mean reaction data that used CS+/CS− signals
	Van Damme et al (2006) ¹⁵	Experiment 1 acquisition phase	Mean reaction data that used CS+/CS− signals
		Experiment 2 acquisition phase	Mean reaction data that used CS+/CS− signals
	Van Damme et al (2012) ¹⁶	Experiment 1	Mean reaction data under pain and baseline conditions
Experiment 2		Mean reaction data under pain and baseline conditions	
Alerting attention	Kurita et al (2015) ¹⁷	Experiment 1	10th, 50th, and 90th mean reaction data under mild pain and baseline conditions
		Experiment 2	10th, 50th, and 90th mean reaction data under moderate pain and baseline conditions
	Moore et al (2012) ¹⁰	Experiment 1	Mean reaction data under pain and baseline conditions
Executive attention	Babiloni et al (2004) ¹⁹	Behavioral experiment	Mean reaction data under pain and nonpainful conditions
	Cheng et al (2017) ¹⁸	Behavioral experiment (A-type participants)	Mean reaction data under pain and nonpainful conditions
		Behavioral experiment (P-type participants)	Mean reaction data under pain and nonpainful conditions

Abbreviations: CS+, conditioned stimulus was presented; CS–, conditioned stimulus disappeared; A-type, attention dominates; P-type, pain dominates.

Figure 2 Response time (ms) outcomes for valid cues.

Abbreviations: E1, experiment 1; E2, experiment 2.

Figure 3 Response time (ms) outcomes for invalid cues.

Abbreviations: E1, experiment 1; E2, experiment 2.

estimate of -8.21 (95%CI $= -15.72$ to -0.69); while induced pain under the painful invalid cue condition showed no significant estimate (6.85 [95%CI $= -1.16$ to 14.85]) (Figures 4 and 5).

Alerting attention

Pooled results from seven comparisons across two studies suggested that induced pain was significantly associated with impaired alerting. There was a significant effect estimate of 9.04 (95% CI $= 4$ to 14.08). A moderately nonsignificant heterogeneity was detected for alerting attention ($\chi^2=13.09$, $P=0.04$, $I^2=54\%$) (Figure 6).

Executive attention

Pooled results from three comparisons across two studies suggested that induced pain has no significant effect on executive attention (effect estimate $= -4.69$ [95%CI $= -68.91$ to 59.53]). No significant heterogeneity was detected for executive attention ($\chi^2=5.53$, $P=0.06$, $I^2=64\%$) (Figure 7).

Discussion

The present meta-analysis found that induced pain affected orienting and alerting but not executive attention.

Orienting

Valid cue condition

Participants with induced pain responded to valid cues faster ($n=5$ studies; $n=309$ subjects with induced pain, $n=310$ subjects with no induced pain) than those without pain. This finding is consistent with previous research.^{22,30} All the eight included comparisons used a version of exogenous spatial cue test and thus provided the most homogenous data set in the current meta-analysis. Under the condition of induced pain, four of five studies^{13–16} used pain as the cue, while the other one¹⁰ did the experiment under pain condition. Therefore, we may predict that the effect of pain on orienting attention would reveal attentional bias to pain. The result corroborated that participants showed significant bias to pain stimulation.

The valid cue task directs attention toward the correct location of the target and is used to predict the processing of the incoming target. Responses to the targets are therefore faster when subjects know in advance what type of movement they will make – a “motor set.” The “motor set” is a type of top-down attentional control. When individuals are stimulated by pain, a motor set in response to the stimulus site will be generated. Thus, when the target stimulus occurs at the same

Figure 4 Response time (ms) outcomes for bias to pain stimulus.

Abbreviations: E1, experiment 1; E2, experiment 2.

Figure 5 Response time (ms) outcomes for disengagement from pain stimulus.

Abbreviations: E1, experiment 1; E2, experiment 2.

Figure 6 Response time (ms) outcomes for alerting attention.

Figure 7 Response time (ms) outcomes for executive attention.

location with cue, the participants will respond more quickly. Schmidt et al demonstrated that attention was biased toward the location of pain stimulation by eye-tracking.³¹ Similarly, previous research has shown that a significant cue validity effect occurs when pain is used as a pre-cueing stimulus.⁸

Invalid cue condition

Participants experiencing induced pain responded to invalid cues slower than no-pain controls ($n=5$ studies; $n=309$ subjects with induced pain, $n=310$ subjects with no pain). Consistent with a previous study,³² these results provide

evidence that participants with induced pain may experience difficulty in disengaging from pain sensation. The difference between the induced-pain group and the control group was marginally significant.

Under the invalid cue condition, it is necessary to disengage attention from the cued location and reorient attention to the opposite location. As shown in a previous study,³² the biological warning of pain demands attention, leading to disengage from the cue. These results provided evidence for the “motor set” hypothesis: RTs are longer under induced pain compared to no pain when an invalid cue is presented due to a conflict between the location of the target and the “motor set.”

The effects of pain on orienting attention differed under valid and invalid conditions. Pain facilitated processing when the cue was presented in the same location as the target, while pain inhibited processing when the cue was presented in the opposite location from the target. A study by Mangun and Hillyard provided evidence that spatial priming leads to changes in the sensory-perceptual processing of targets.³³ It may be that the experience of pain leads to a disruption in the perceptual processing of reorientation which is necessary under the invalid cue condition. Further studies are needed to investigate this hypothesis.

Alerting attention

The current meta-analysis found that induced pain leads to a significant impairment in alerting attention ($n=2$ studies; $n=158$ subjects with induced pain, $n=158$ subjects with no pain). Seven comparisons used the CPT providing a reliable homogenous data set, suggesting high confidence in group differences.

Alerting attention is defined as the ability to maintain a mental arousal and preparedness for the processing of information with required effort.³⁴ Prior studies have shown that pain interferes with mental processes and disrupts ongoing activities.¹² One possible explanation for the impairment observed under conditions of pain is that while the CPT requires attentional coherence, pain may inhibit or distract the mental processes which results in decreasing capacity to maintain a state of alert arousal. It is not clear whether it is distraction or inhibition that leads to impairment. Future research should focus on separating the two mechanisms. Another possible explanation is that ongoing pain induces fatigue. Fatigue is one of the most common symptoms associated with various pains.³⁵ Fatigue associated with ongoing pain may result in depleted energy necessary for attentional tasks.

Executive attention

Surprisingly, the current meta-analysis found no evidence ($n=2$ studies; $n=57$ subjects with induced pain, $n=57$ subjects with no induced pain) for an effect of induced pain on executive attention. Executive attention is required for the execution of goal-directed behaviors, including the planning of actions, expectations, responding, initiating, and maintaining deliberate behavior, monitoring the results of behavior, and interrupting or modifying behavior.³⁴ Executive attention is necessary for the regulation of cognition and inhibition of responses. Inhibition is important for the successful attentional control of pain. Bjekić et al found that cognitive inhibition measured by the Stroop task is correlated with pain perception.³⁶ In addition, a previous meta-analysis found an impairment in response inhibition in people with chronic pain.²¹ One possible explanation for the inconsistent results between the current study and the previous research in executive attention is the degree of variance between studies. The current meta-analysis included only two studies with three comparisons using different designs.

One study, using the Stroop test, found that pain significantly reduced task speed in P-type individuals but boosted performance in A-type individuals.¹⁸ Another study using the go/no-go test revealed that pain significantly affected performance.¹⁹ The Stroop test involves the ability to suppress cognitive conflict while the go/no-go test involves the ability to suppress behavioral conflict, leading to variation between studies. In addition, previous studies exclusively investigated the effects of chronic pain,⁶ while the current study analyzed the effects of induced pain on response inhibition. Future studies are needed to better understand the relationship between executive attention and laboratory-induced pain.

Strengths and limitations of the current study

The current meta-analysis used a rigorous search procedure to ensure that relevant studies were included, while excluding irrelevant and low-quality studies. Based on Posner's attention theory, we selectively studied the influence of laboratory-induced pain on the three components of attention: orienting, alerting, and executive. Studies using the spatial cue test, CPT, Stroop test, and go-no/go test were included.

Norman and Shallice³⁷ proposed that the attentional system is susceptible to interruption by the presentation of a new superordinate goal in order to protect an organism from danger or harm. Pain is a biological mechanism that can interrupt ongoing attention with the aim of self-protection.³² The current study suggests induced-pain effects orienting attention and

altering attention. However, due to the risk of biases, further studies are needed to confirm the relevance of our findings.

There are limitations that must be considered when interpreting the results of the current meta-analysis. First, attention is a dynamic process that is difficult to measure. Eye tracking techniques are often used to examine the time-course of overt visual attention in humans. Many prior studies have investigated attentional bias to pain-related information,^{38,39} but few have examined attentional bias to actual pain stimulation.³¹ Future research using eye-tracking to investigate the effects of actual pain stimulation on attention would be helpful to further our understanding of the effects of pain on attention performance. Second, we limited the meta-analysis to three components of attention. Other components of attention such as divided attention, selective attention, and working memory were not investigated. Previous research has found that chronic pain decreases performance on tests of these components of attention.^{40,41} Further studies are needed to explore the effects of induced pain on these components of attention. Finally, the behavioral paradigms included in the current meta-analysis may not be sensitive to acute pain and, therefore, the effects reported may be underestimated.

Conclusion

Experimentally induced pain affects some components of attention. Interestingly, the effect of pain can be both facilitative and disruptive. For example, pain disrupts orienting under an invalid painful cue condition while facilitating orienting under a valid painful cue condition. Pain is disruptive to alerting attention, while there is no obvious effect of pain on executive attention. Further studies are necessary to confirm the results of the current meta-analysis.

Disclosure

This work was funded by National Natural Science Foundation of China grant (31471061) to FL. The authors report no other conflicts of interests in this work.

References

1. Posner MI, Rothbart MK. Research on attention networks as a model for the integration of psychological science. *Annu Rev Psychol*. 2007;58(1):1–23.
2. Fan J, McCandliss BD, Fossella J, Flombaum JI, Posner MI. The activation of attentional networks. *Neuroimage*. 2005;26(2):471–479.
3. Crottaz-Herbette S, Menon V. Where and when the anterior cingulate cortex modulates attentional response: combined fMRI and ERP evidence. *J Cogn Neurosci*. 2006;18(5):766–780.
4. Vervoort T, Trost Z, van Ryckeghem DM. Children's selective attention to pain and avoidance behaviour: the role of child and parental catastrophizing about pain. *Pain*. 2013;154(10):1979–1988.
5. Crombez G, van Ryckeghem DM, Eccleston C, Van Damme S. Attentional bias to pain-related information: a meta-analysis. *Pain*. 2013;154(4):497–510.
6. Eccleston C. Chronic pain and attention: a cognitive approach. *Br J Clin Psychol*. 1994;33(4):535–547.
7. Attridge N, Keogh E, Eccleston C. The effect of pain on task switching: pain reduces accuracy and increases reaction times across multiple switching paradigms. *Pain*. 2016;157(10):2179–2193.
8. Van Damme S, Crombez G, Lorenz J. Pain draws visual attention to its location: experimental evidence for a threat-related bias. *J Pain*. 2007;8(12):976–982.
9. Vancleef LM, Peters ML. The interruptive effect of pain on attention. *J Pain*. 2006;7(1):21–22.
10. Moore DJ, Keogh E, Eccleston C. The interruptive effect of pain on attention. *Q J Exp Psychol*. 2012;65(3):565–586.
11. Crombez G, Eccleston C, Baeyens F, Eelen P. The disruptive nature of pain: an experimental investigation. *Behav Res Ther*. 1996;34(11-12):911–918.
12. Vlaeyen JW, Morley S, Crombez G. The experimental analysis of the interruptive, interfering, and identity-distorting effects of chronic pain. *Behav Res Ther*. 2016;86:23–34.
13. Koster E, Crombez G, Van Damme S, Verschuere B, De Houwer J. Signals for threat modulate attentional capture and holding: Fear-conditioning and extinction during the exogenous cueing task. *Cogn Emot*. 2005;19(5):771–780.
14. Van Damme S, Lorenz J, Eccleston C, Koster EHW, De Clercq A, Crombez G. Fear-conditioned cues of impending pain facilitate attentional engagement. *Neurophysiol Clin*. 2004;34(1):33–39.
15. Van Damme S, Crombez G, Eccleston C, Koster EH. Hyper-vigilance to learned pain signals: a componential analysis. *J Pain*. 2006;7(5):346–357.
16. Van Damme S, Legrain V. How efficient is the orienting of spatial attention to pain? An experimental investigation. *Pain*. 2012;153(6):1226–1231.
17. Kurita GP, Malver LP, Andresen T, et al. Does mutual compensation of the cognitive effects induced by pain and opioids exist? An experimental study. *Psychopharmacology*. 2015;232(8):1373–1381.
18. Cheng JC, Bosma RL, Hemington KS, Kucyi A, Lindquist MA, Davis KD. Slow-5 dynamic functional connectivity reflects the capacity to sustain cognitive performance during pain. *Neuroimage*. 2017;157:61–68.
19. Babiloni C, Brancucci A, Arendt-Nielsen L, et al. Cortical sensorimotor interactions during the expectancy of a go/no-go task: effects of painful stimuli. *Behav Neurosci*. 2004;118(5):925–935.
20. Vossel S, Thiel CM, Fink GR. Cue validity modulates the neural correlates of covert endogenous orienting of attention in parietal and frontal cortex. *Neuroimage*. 2006;32(3):1257–1264.
21. Berryman C, Stanton TR, Bowering KJ, Tabor A, McFarlane A, Moseley GL. Do people with chronic pain have impaired executive function? A meta-analytical review. *Clin Psychol Rev*. 2014;34(7):563–579.
22. Buffington AL, Hanlon CA, McKeown MJ. Acute and persistent pain modulation of attention-related anterior cingulate fMRI activations. *Pain*. 2005;113(1):172–184.
23. Filbrich L, Alamia A, Burns S, Legrain V. Erratum to: orienting attention in visual space by nociceptive stimuli: investigation with a temporal order judgment task based on the adaptive psi method. *Exp Brain Res*. 2017;235(9):2901.
24. Fecher M, Schlereth T, Kritzmann S, et al. Stress and thermoregulation: different sympathetic responses and different effects on experimental pain. *Eur J Pain*. 2009;13(9):935–941.
25. Brown CA, Jones AK. A role for midcingulate cortex in the interruptive effects of pain anticipation on attention. *Clin Neurophysiol*. 2008;119(10):2370–2379.
26. Valet M, Sprenger T, Boecker H, et al. Distraction modulates connectivity of the cingulo-frontal cortex and the midbrain during pain-an fMRI analysis. *Pain*. 2004;109(3):399–408.

27. Dowman R, Darcey T, Barkan H, Thadani V, Roberts D. Human intracranially-recorded cortical responses evoked by painful electrical stimulation of the sural nerve. *Neuroimage*. 2007;34(2):743–763.
28. Favril L, Mouraux A, Sambo CF, Legrain V. Shifting attention between the space of the body and external space: electrophysiological correlates of visual-nociceptive crossmodal spatial attention. *Psychophysiology*. 2014;51(5):464–477.
29. Strigo IA, Simmons AN, Matthews SC, Craig AD. The relationship between amygdala activation and passive exposure time to an aversive cue during a continuous performance task. *PLoS One*. 2010;5(11):e15093.
30. Sun ZK, Wang JY, Luo F. Experimental pain induces attentional bias that is modified by enhanced motivation: an eye tracking study. *Eur J Pain*. 2016;20(8):1266–1277.
31. Schmidt K, Gamer M, Forkmann K, Bingel U. Pain affects visual orientation: an Eye-Tracking study. *J Pain*. 2018;19(2):135–145.
32. Eccleston C, Crombez G. Pain demands attention: a cognitive-affective model of the interruptive function of pain. *Psychol Bull*. 1999;125(3):356–366.
33. Mangun GR, Hillyard SA. Modulations of sensory-evoked brain potentials indicate changes in perceptual processing during visual-spatial priming. *J Exp Psychol Hum Percept Perform*. 1991;17(4):1057–1074.
34. Mezzacappa E. Alerting, orienting, and executive attention: developmental properties and sociodemographic correlates in an epidemiological sample of young, urban children. *Child Dev*. 2004;75(5):1373–1386.
35. Theobald DE. Cancer pain, fatigue, distress, and insomnia in cancer patients. *Clin Cornerstone*. 2004;6(1):S15–S21.
36. Bjekić J, Živanović M, Purić D, Oosterman JM, Filipović SR. Pain and executive functions: a unique relationship between Stroop task and experimentally induced pain. *Psychol Res*. 2018;82(3):580–589.
37. Norman DA, Shallice T. *Attention to action: Willed and automatic control of behavior*. Consciousness and self-regulation: Advances in research and theory. New York: Plenum; 1986;(4)1–18.
38. Schoth DE, Ma Y, Lioffi C. Exploring attentional bias for real-world, pain-related information in chronic musculoskeletal pain using a novel change detection paradigm. *Clin J Pain*. 2015;31(8):680–688.
39. Heathcote LC, Vervoort T, Eccleston C, et al. The relationship between adolescents' pain catastrophizing and attention bias to pain faces is moderated by attention control. *Pain*. 2015;156(7):1334–1341.
40. Oosterman JM, Derksen LC, van Wijck AJ, Veldhuijzen DS, Kessels RP. Memory functions in chronic pain: examining contributions of attention and age to test performance. *Clin J Pain*. 2011;27(1):70–75.
41. Grisart JM, Plaghki LH. Impaired selective attention in chronic pain patients. *Eur J Pain*. 1999;3(4):325–333.

Journal of Pain Research

Publish your work in this journal

The Journal of Pain Research is an international, peer reviewed, open access, online journal that welcomes laboratory and clinical findings in the fields of pain research and the prevention and management of pain. Original research, reviews, symposium reports, hypothesis formation and commentaries are all considered for publication.

Submit your manuscript here: <https://www.dovepress.com/journal-of-pain-research-journal>

Dovepress

The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.