Ultrasound-guided fine-needle aspiration biopsy of thyroid nodules <10 mm in the maximum diameter: does size matter?

Yi-jun Lyu^{1,2,*}Fang Shen^{3,*}Yun Yan^{4,5}Ming-zhu Situ^{4,5}Wei-zhu Wu^{1,2}Guo-qiang Jiang³Ya-ya Chen^{4,5}

¹Department of Thyroid and Breast Surgery, Ningbo Medical Center Lihuli Eastern Hospital, Ningbo 315040, Zhejiang, China; ²Department of Thyroid and Breast Surgery, Taipei Medical University Ningbo Medical Center, Ningbo 315040, Zhejiang, China; ³Department of Orthopaedic Surgery's Spine Division, The Affiliated Hospital of Medical School of Ningbo University, Ningbo 315020, Zhejiang, China; ⁴Department of Ultrasound, Ningbo Medical Center Lihuli Eastern Hospital, Ningbo 315040, Zhejiang, China; ⁵Department of Ultrasound, Taipei Medical University Ningbo Medical Center, Ningbo 315040, Zhejiang, China

*These authors contributed equally to this work

Objective: Ultrasound-guided fine-needle aspiration biopsy (US-FNAB) is a safe and effective method of screening malignant thyroid nodules such as papillary thyroid carcinoma. However, not much data are available regarding the diagnostic efficacy of US-FNAB for papillary thyroid microcarcinoma (≤ 10 mm in diameter). We aim to compare the diagnostic efficacy of US-FNAB on thyroid nodules between two groups divided by a diameter of 10 mm by correlating the cytological results of US-FNAB with the histopathologic diagnoses in selected patients.

Patients and methods: Eight hundred twenty-two thyroid nodules (Group A: diameter ≤ 10 mm, $n=620$; Group B: diameter >10 mm, $n=202$) from 797 patients treated between March 2014 and June 2017 were retrospectively evaluated. Only nodules with Thyroid Imaging Reporting and Data System (TIRADS) categories 4–6 were enrolled and sampled by US-FNAB, followed by surgical resection.

Results: According to The Bethesda System for Reporting Thyroid Cytopathology (TBSRTC) diagnostic categories, 94 thyroid nodules were classified as I, III and IV, and were excluded from the analysis. The resultant 728 thyroid nodules from 721 patients were analyzed. The malignant tendency (TBSRTC V and VI) rates on US-FNAB were 88.2% and 84.6% ($P=0.202$) in Group A and Group B, respectively, and the malignant rates were 89.5% and 86.9% ($P=0.330$), respectively, on histopathology. There was a high concordance between cytology and histopathology diagnoses (kappa value $=0.797$), and no statistical difference in terms of US-FNAB accuracy was found between the two groups ($P=0.533$).

Conclusion: For thyroid nodules of TIRADS category 4–6, the diagnostic efficacy of US-FNAB is similar for thyroid nodules either smaller or greater than 10 mm in their maximum diameter.

Keywords: ultrasound-guided fine-needle aspiration biopsy, thyroid nodules, microcarcinoma, histopathology, cytopathology

Introduction

Thyroid nodules are common entities that are detected in 19%–68% of the population by using high-resolution ultrasound (US).¹ Nevertheless, only a relatively small percentage (~5%) of thyroid nodules are malignant, with papillary thyroid carcinoma (PTC) being the most common pathological type.² According to the WHO classification system, papillary thyroid microcarcinomas (PTMCs) are PTCs <10 mm in diameter.³

Early cancer detection and intervention has been suggested to reduce patient's mortality and morbidity. Given its minimal invasiveness and technical simplicity, ultrasound-guided fine-needle aspiration biopsy (US-FNAB) has been widely adopted for characterizing thyroid nodules cytopathologically.⁴ The cytology results will help to determine whether or not subsequent thyroidectomy is necessary. According to the

Correspondence: Ya-ya Chen
Department of Ultrasound, Ningbo Medical Center Lihuli Eastern Hospital,
No. 1111 Jiangnan Road, Ningbo 315040,
China
Tel +86 574 5583 5583
Fax +86 574 5583 5752
Email 67857266@qq.com

American Thyroid Association (ATA) guideline, US-FNAB is recommended for those thyroid nodules with a diameter >10 mm those with intermediate to high suspicion US pattern.⁵ On the other hand, for patients with nodules ≤ 10 mm of suspicious US pattern, active sonographic surveillance is recommended instead.⁶ Despite the fact that the prognostic advantage of PTMC has been an issue of debate in recent studies,^{7–10} early diagnosis and treatment of PTMC might be beneficiary to patient prognosis as small size alone does not guarantee low risk in incidentally found thyroid cancers.^{11,12}

There are several studies that have evaluated the efficacy of US-FNAB for small thyroid nodules.^{13,14} We focused here on the diagnostic efficacy of US-FNAB for ultrasonographically “suspicious”, that is, Thyroid Imaging Reporting and Data System (TIRADS) categories 4–6,¹⁵ and small, that is, measuring ≤ 10 mm as the maximum diameter on US, thyroid nodules. To the best of our knowledge, our research is the first study of this kind with the largest sample size ever reported. This knowledge is important to determine whether US-FNAB should be routinely performed in this subgroup of patients in future clinical practices.

Patients and methods

Patient selection

We retrospectively reviewed 721 patients who had received thyroid US, thyroid nodule US-FNAB and subsequent thyroid surgery at Ningbo Medical Center Lihuili Eastern Hospital and Taipei Medical University Ningbo Medical Center (a tertiary medical center) between March 2014 and June 2017. Thyroid nodules were given cytopathological and histopathologic diagnoses with the criteria listed in the following sections and were grouped according to their maximum diameter as measured on US: Group A (≤ 10 mm) and Group B (>10 mm). Written informed consents were obtained from all patients before performing US-FNABs and surgeries. Written consents were also obtained from the patients regarding the report of their medical data. Ethical approval for conducting this retrospective study was obtained from the Ethics Committee of Ningbo Medical Center Lihuili Eastern Hospital and Taipei Medical University Ningbo Medical Center (ethical approval no. DYLL2016001) in compliance with the Declaration of Helsinki.

Thyroid US evaluation

US examinations were performed at the Department of Ultrasound with Philips Q5 US equipment and a 5–12 MHz linear probe. The reasons for thyroid US scan are as follows: 1) palpable cervical mass and/or enlarged lymph

nodes; 2) patients with a known history of thyroid nodes and referred from other hospitals and 3) on routine health checks that included thyroid US. US patterns of the thyroid lobes and nodules (eg, calcification, echogenicity, volume, shape, dimensions, long axis/short axis ratio, vascularity) were recorded and all nodules were classified according to TIRADS by using five sonographic parameters (ie, composition, echogenicity, shape, margin and echogenic foci). Patients having TIRADS 4–6 thyroid nodules were included for US-FNAB, while the exclusion criteria were: 1) patients who rejected US-FNAB or were not cooperative for the procedure; 2) accompanied with severe cardiovascular or pulmonary conditions and 3) patients with bleeding history of unknown causes or coagulation disorders (prothrombin time >18 seconds, platelet count $<50 \times 10^9/L$, prothrombin activity $<40\%$).

US-FNAB procedure for thyroid nodules

US-FNAB was performed by the same pair of experienced diagnostic sonographers who were licensed to perform the procedure, in order to ensure similarity in techniques employed. Specifically, one was responsible for US guidance and the other for fine-needle aspiration (FNA). Under US guidance, the target nodule was punctured with a 24-G needle connected to a 5 mL syringe without local anesthesia. After confirmation of reaching the target nodule by the needle tip on US, the needle was moved forward and backward ten times under negative pressure to aspirate the sample. Then the needle was withdrawn, negative pressure was released and specimen within the needle was immediately transferred to a liquid-based cytology medium. Generally, only one puncture was needed to sample each individual nodule and a second attempt could be delivered in case of unsatisfactory gross tissue yield by the first attempt as judged by the sonographer performing aspiration as a precaution.

Pathological review of thyroid nodules

Cytological and histopathologic diagnoses were performed at the Ningbo Diagnostic Pathology Center by pathologists specialized in thyroid cancer. The cytological diagnoses were made by two independent specialized cytologists (Dr Jue Zhou and Dr Xian-fa Xu; kappa value $=0.83$), and release of the final report was subjected to the approval of a third senior cytologist (Dr Deng Pan), who was also in charge of making the final decision when there was discrepancy between the two. Cytological results were reported according to The Bethesda System for Reporting Thyroid Cytopathology (TBSRTC) as follows:¹⁶ I, nondiagnostic or unsatisfac-

tory; II, benign; III, atypia of undetermined significance or follicular lesion of undetermined significance; IV, follicular neoplasm or suspicious for a follicular neoplasm; V, suspicious for malignancy and VI, malignant. Surgery was carried out in patients with US TIRADS categories 4–6 at their own choice with signed consents. The histological diagnoses were reported according to the WHO histological classification of thyroid tumors.¹⁷

Statistical analyses

Data were analyzed using the SPSS 22.0 software package (IBM Corporation, Armonk, NY, USA). Continuous variables were presented as mean $\pm$ SD. Categorical variables were presented as percentages. The sensitivity, specificity, positive predictive value (PPV) and negative predictive value (NPV) were given as percentages. The diagnostic performance of US-FNAB was defined by the consistency between cytological and histopathologic results: US-FNAB was considered as accurate when the two results matched and vice versa. Data comparisons between groups were carried out with the Mann–Whitney *U* test. Chi-squared test or Fisher's exact test was used to compare categorical variables. The kappa statistic was used to measure the agreement between the US-FNAB and histopathology diagnoses. The kappa can range from -1 to $+1$: values ≤ 0 indicate no agreement and values 0.01 – 0.20 mean none to slight agreement, 0.21 – 0.40 mean fair agreement, 0.41 – 0.60 mean moderate agreement, 0.61 – 0.80 mean substantial agreement and 0.81 – 1.00 mean almost perfect agreement. A *P*-value < 0.05 was considered statistically significant.

Ethics approval and consent to participate

Ethical approval was obtained from the Ethics Committee of Ningbo Medical Center Lihuili Eastern Hospital and Taipei Medical University Ningbo Medical Center (ethical approval no. DYLL2016001).

Results

Demographics, US and US-FNAB results

A total of 822 thyroid nodules of TIRADS categories 4–6 from 797 patients (772 had 1 nodule and 25 had 2) were initially reviewed during the study period. There were 620 nodules in Group A and 202 nodules in Group B. Cytologically, 94 thyroid nodules were classified as TBSRTC I, III and IV, and were excluded from subsequent statistical analysis. The resultant 728 thyroid nodules from 721 patients (714 had 1 nodule and 7 had 2) were classified as TBSRTC II ($n=92$, 12.6%), V ($n=168$, 23.1%) and VI ($n=468$, 64.3%). Among

those 721 patients, there were 557 (77.3%) females and 164 (22.7%) males. There were 553 nodules in Group A and 175 nodules in Group B. The median nodule size was 6 mm (range: 2–10 mm) for Group A and 15 mm (range: 11–52 mm) for Group B. There was no significant difference in the mean age between the two groups (Group A: 42.3 ± 15.4 vs Group B: 48.9 ± 17.2 , $P=0.018$).

Diagnostic performance of US-FNAB

According to US-FNAB cytology, the percentages of nodes that fell in the diagnostic categories of V (suspicious for malignancy) and VI (malignant), which indicate malignant tendency, were 88.2% in Group A and 84.6% in Group B ($P=0.202$), whereas the histological malignant rates were 89.5% in Group A and 86.9% in Group B ($P=0.330$). There was a high agreement between the US-FNAB cytology and final histopathology (kappa value = 0.797). There was no significant difference in sensitivity, specificity, accuracy, PPV and NPV of US-FNAB between the two groups (Table 1). The diagnostic performance of US-FNAB did not exhibit significant difference between the two groups either ($P=0.533$).

Discussion

Given its minimal invasiveness, technical simplicity and high concordance with histology results, US-FNAB has become a routine method for biopsy of many benign and malignant pathologies, including thyroid nodules.¹⁸ The current recommendations on the size selection criteria for US-FNAB of thyroid nodules have been set at a cutoff value of 10 mm with suspicious sonographic features.^{19,20} ATA guidelines recommend that nodules ≥ 10 mm with high to intermediate suspicious US pattern (or ≥ 15 mm with low suspicious pattern or ≥ 20 mm with very low suspicious pattern) be evaluated by US-FNAB.⁵ Similarly, the Society of Radiologists in Ultrasound recommend performing US-FNAB on thyroid nodules that are > 10 mm and when there

Table 1 Sensitivity, specificity, accuracy, PPV and NPV of US-FNAB for thyroid nodules of TIRADS categories 4–6

Diagnostic parameters	Group A (≤ 10 mm), %	Group B (> 10 mm), %	P-value
Sensitivity	96.8	96.7	1.000
Specificity	84.5	95.7	0.268
Accuracy	95.5	96.6	0.533
PPV	98.2	99.3	0.467
NPV	75.4	81.5	0.596

Abbreviations: NPV, negative predictive value; PPV, positive predictive value; TIRADS, Thyroid Imaging Reporting and Data System; US-FNAB, ultrasound-guided fine-needle aspiration biopsy.

are suspicious US features suggesting cancer.²¹ On the other hand, for nodules <10 mm, ATA recommends further evaluation only for patients with clinical symptoms or associated lymphadenopathy and no routine sonographic follow-up for those with very low suspicious US pattern (weak recommendation, low-quality evidence).²² Therefore, for nodules ≤10 mm that exhibit intermediate to highly suspicious US patterns, routine US follow-ups are justifiable. However, nodule size alone is not predictive of malignancy in patients with Bethesda category III, IV and V thyroid nodules, and due to the existence of thyroid microcarcinomas such as PTMC, early identification by US-FNAB and subsequent surgical intervention might provide clinical benefits for those selected patients.^{23,24} Certain US patterns have been associated with potential malignancy and thus indicate the necessity of US-FNAB. For example, a recent meta-analysis suggested that US features such as microcalcification, a taller than wide shape, irregular margins and absence of elasticity are associated with higher risk of malignancy and have the most satisfactory diagnostic performances.²⁵ Similarly, the Revised Korean Society of Thyroid Radiology Consensus Statement and Recommendations suggest that for nodules >5 mm, FNAB should be performed if the target nodule is solid and hypoechoic, together with any three of the following suspicious US features: microcalcification, nonparallel orientation (taller than wide), spiculated or microlobulated margin.²⁶

According to TBSRTC, the malignancy risk is 60%–75% for of category V (suspicious for malignancy) and 97%–99% for of category VI (malignant).¹⁶ On the other hand, there are only a limited number of reports that have evaluated the diagnostic efficacy of US-FNAB for thyroid nodules <10 mm.^{13,27,28} Unal and Sezer found a concordance rate of 52.4% between US-FNAB cytology and histopathology in 21 nodules <10 mm in diameter. Also, among those inconsistent cases (false negative: cytological category of benign, but histologically malignant), PTMC accounted for the majority of cases.¹⁴ Rosario et al recently specifically addressed the FNA results for a subgroup of “highly suspicious” thyroid nodules (≤10 mm, highly suspicious US features, restricted to the thyroid) in patients who are candidates for active surveillance according to ATA.²⁹ They reviewed a total of 198 nodules and found a very high rate of malignancy on histology for nodules with suspicious/malignant (100%) and indeterminate (81.4%) cytology results. Zhong et al compared the efficacy of US-FNAB for 344 thyroid nodules with different sizes (≤5.0, 5.1–10.0 and >10.0 mm) and found similar diagnostic efficacy regardless of size.³⁰

The main novelty of the current study is the focus on TIRADS 4–6 thyroid nodules and enrolment of the largest sample size to date for comparing the diagnostic efficacy of US-FNAB for nodules that were either ≤10 or >10 mm. Besides, unlike previously studies where the benign nature of the nodule was verified by repetition of cytology or simply by clinical follow-up, all nodes in our study were surgically removed with definite histopathologic diagnoses. Our results showed that the inconsistency rate between US-FNAB and histology was much lower than what has been reported previously: only 21 cytologically “benign” nodules identified by the US-FNAB were later confirmed as PTCs after surgery (false negative) and 10 cytologically “malignant” nodules were later verified as benign nature (false positive). Among the 21 nodes with false-negative US-FNAB results, 5 were >10 mm and 16 were ≤10 mm. This might be caused by sampling of normal thyroid tissue during US-FNAB procedure. Besides, there were false-negative US-FNAB cases in both groups, implicating the need of follow-up of these patients with cytologically “benign” results.

Given the slow-growing nature of PTC, there is still controversy over the clinical value of its early diagnosis by US-FNAB. It was reported that there was a high frequency of occult papillary microcarcinomas in an autopsy study (35.6%).³¹ In our study, the number of nodules ≤10 mm was three times higher than those >10 mm, suggesting the importance of appropriate clinical management of these subcentimeter nodules. The 2015 ATA guidelines, however, do not recommend US-FNAB for subcentimeter nodules unless there is extrathyroidal extension or suspicious lymphadenopathy. Besides, ATA also favors active surveillance in selected low-risk PTMC, instead of immediate surgery. These recommendations are largely based on observational studies done by Japanese researchers that suggested the relatively indolent nature of certain subtypes of PTMC.^{32–34} Nevertheless, perithyroidal lymph node metastasis is a feature seen in certain subtypes of PTMCs which are shown to have poor differentiation.^{12,35,36} However, without US-FNAB, there is no better way to stratify patients with subcentimeter thyroid nodules of intermediate to highly suspicious features into either active surveillance or surgery. Neither there is any reliable method to predict which subsets of PTMC will be more aggressive and thus need to have more timely surgery. A recent study by Gweon et al showed that, among patients with subcentimeter thyroid nodules of highly suspicious US features, those of younger age (<45 years), male gender and with certain US features (microcalcification and a taller-

than-wide shape) might not be good candidates for active surveillance as they are associated with malignancy rate and aggressive biological behavior.³⁷ Besides, prolonged active surveillance itself is not without associated costs in terms of clinical, psychological and economic burdens on the health care system and patients.³⁸ Therefore, ongoing researches are needed to provide evidence for the optimal management of patients with subcentimeter PTMC.

A detailed discussion of the natural history of PTMC is out of scope of the current work,^{24,39–41} and the strength of our results lies in demonstrating that US-FNAB is a convenient and reliable diagnostic procedure for these selected subcentimeter pathologies. However, our study also has several limitations. First, similar to previous work,³⁷ patients with nodules of TIRADS categories 4–6 were offered with US-FNAB and subsequent surgery. Although this provides valuable information on the concordance between cytology and histology results, some physicians would doubt the necessity of surgery for patients with benign US-FNAB cytological findings and would suggest follow-up instead.⁴² This is at least partially because of the associated surgical complications.^{43,44} However, in our practice, we did provide explanations to every patient regarding current ATA guidelines and related researches by Ito et al³³ and offered conservative options including active surveillance.³⁴ After thorough considerations, some patients opted to have US follow-ups, while others still had severe psychological stress and proceeded with surgeries despite our advice. Second, thyroid nodes with TBSRTC I, III and IV cytology (11% of the total cases) were excluded from the analysis because, according to TBSRTC guidelines, a repeated FNA is needed for category I and III nodules, while category IV (follicular or suspicious) is not the focus of the current study. Therefore, our study only focused on thyroid nodules with definite cytology results (either benign or malignant, but excluding follicular neoplasms) yielded by single biopsy attempt. However, this might also lead to selection bias to some extent. Third, we did not exclude patients with coexistence of diffuse thyroid disease, such as thyroiditis or diffuse goiter, and these thyroid conditions might influence the performance of the US-FNABs, especially for small nodules.⁴⁵ Last but not least, our study is retrospective and future prospective study will yield clinical evidence at a higher level to answer the controversy over the benefit of surgery on the prognostic advantage of PTMC.

To conclude, US-FNAB is an effective diagnostic method for thyroid nodules with TIRADS categories >4 regardless of their size. Prospective studies are needed to determine the natural history of thyroid malignancy <10 mm and identify those at particular risk of exhibiting aggressive behavior, in

order to further support the value of its early diagnosis and surgical intervention as well as for providing more optimal and personalized care.

Consent for publication

This manuscript does not contain any individual person's data in any form.

Data sharing statement

The data sets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Acknowledgments

We would like to thank Dr Jue Zhou, Dr Xian-fa Xu and Dr Deng Pan from Ningbo Diagnostic Pathology Center for their help in reviewing pathological diagnosis of the cases.

Disclosure

The authors report no conflicts of interest in this work.

References

- Hay ID, Grant CS, van Heerden JA, Goellner JR, Ebersold JR, Bergstralh EJ. Papillary thyroid microcarcinoma: a study of 535 cases observed in a 50-year period. *Surgery*. 1992;112(6):1139–1146.
- Ito Y, Miyauchi A, Oda H. Low-risk papillary microcarcinoma of the thyroid: a review of active surveillance trials. *Eur J Surg Oncol*. 2018;44(3):307–315.
- Wang Y, Li L, Wang YX, et al. Ultrasound findings of papillary thyroid microcarcinoma: a review of 113 consecutive cases with histopathologic correlation. *Ultrasound Med Biol*. 2012;38(10):1681–1688.
- Singh Ospina N, Brito JP, Maraka S, et al. Diagnostic accuracy of ultrasound-guided fine needle aspiration biopsy for thyroid malignancy: systematic review and meta-analysis. *Endocrine*. 2016;53(3):651–661.
- Haugen BR, Alexander EK, Bible KC, et al. 2015 American thyroid association management guidelines for adult patients with thyroid nodules and differentiated thyroid cancer: the American Thyroid Association guidelines task force on thyroid nodules and differentiated thyroid cancer. *Thyroid*. 2016;26(1):1–133.
- Haugen BR. 2015 American Thyroid Association management guidelines for adult patients with thyroid nodules and differentiated thyroid cancer: what is new and what has changed? *Cancer*. 2017;123(3):372–381.
- Anastasakis AD, Polyzos SA, Makras P, et al. Papillary thyroid microcarcinoma presenting as lymph node metastasis – a diagnostic challenge: case report and systematic review of literature. *Hormones (Athens)*. 2012;11(4):419–427.
- Wang TS, Goffredo P, Sosa JA, Roman SA. Papillary thyroid microcarcinoma: an over-treated malignancy? *World J Surg*. 2014;38(9):2297–2303.
- Gao X, Zhang X, Zhang Y, Hua W, Maimaiti Y, Gao Z. Is papillary thyroid microcarcinoma an indolent tumor?: a retrospective study on 280 cases treated with radioiodine. *Medicine (Baltimore)*. 2016;95(40):e5067.
- Leboulleux S, Tuttle RM, Pacini F, Schlumberger M. Papillary thyroid microcarcinoma: time to shift from surgery to active surveillance? *Lancet Diabetes Endocrinol*. 2016;4(11):933–942.
- Nasir A, Chaudhry AZ, Gillespie J, Kaiser HE. Papillary microcarcinoma of the thyroid: a clinico-pathologic and prognostic review. *In Vivo*. 2000;14(2):367–376.

12. Nam-Goong IS, Kim HY, Gong G, et al. Ultrasonography-guided fine-needle aspiration of thyroid incidentaloma: correlation with pathological findings. *Clin Endocrinol (Oxf)*. 2004;60(1):21–28.
13. Kim DW, Park AW, Lee EJ, et al. Ultrasound-guided fine-needle aspiration biopsy of thyroid nodules smaller than 5 mm in the maximum diameter: assessment of efficacy and pathological findings. *Korean J Radiol*. 2009;10(5):435–440.
14. Unal B, Sezer C. Diagnostic value of ultrasound-guided fine needle aspiration biopsy in malignant thyroid nodules: utility for micronodules. *Asian Pac J Cancer Prev*. 2014;15(20):8613–8616.
15. Horvath E, Majlis S, Rossi R, et al. An ultrasonogram reporting system for thyroid nodules stratifying cancer risk for clinical management. *J Clin Endocrinol Metab*. 2009;94(5):1748–1751.
16. Cibas ES, Ali SZ; NCI Thyroid FNA State of the Science Conference. The Bethesda system for reporting thyroid cytopathology. *Am J Clin Pathol*. 2009;132(5):658–665.
17. Sobin LH. Histological typing of thyroid tumours. *Histopathology*. 1990;16(5):513.
18. Krishnamurthy S, Bedi DG, Caraway NP. Ultrasound-guided fine-needle aspiration biopsy of the thyroid bed. *Cancer*. 2001;93(3):199–205.
19. Ucler R, Usluogullari CA, Tam AA, et al. The diagnostic accuracy of ultrasound-guided fine-needle aspiration biopsy for thyroid nodules three centimeters or larger in size. *Diagn Cytopathol*. 2015;43(8):622–628.
20. Yang GC, Livolsi VA, Baloch ZW. Thyroid microcarcinoma: fine-needle aspiration diagnosis and histologic follow-up. *Int J Surg Pathol*. 2002;10(2):133–139.
21. Shapiro RS. Management of thyroid nodules detected at sonography: society of radiologists in ultrasound consensus conference statement. *Thyroid*. 2006;16(3):209–210.
22. Peli M, Capalbo E, Lovisatti M, et al. Ultrasound guided fine-needle aspiration biopsy of thyroid nodules: guidelines and recommendations vs clinical practice; a 12-month study of 89 patients. *J Ultrasound*. 2012;15(2):102–107.
23. Kiernan CM, Solórzano CC. Bethesda category III, IV, and V thyroid nodules: can nodule size help predict malignancy? *J Am Coll Surg*. 2017;225(1):77–82.
24. Dideban S, Abdollahi A, Meysamie A, Sedghi S, Shahriari M. Thyroid papillary microcarcinoma: etiology, clinical manifestations, diagnosis, follow-up, histopathology and prognosis. *Iran J Pathol*. 2016;11(1):1–19.
25. Remonti LR, Kramer CK, Leitão CB, Pinto LC, Gross JL. Thyroid ultrasound features and risk of carcinoma: a systematic review and meta-analysis of observational studies. *Thyroid*. 2015;25(5):538–550.
26. Shin JH, Baek JH, Chung J, et al. Ultrasonography diagnosis and imaging-based management of thyroid nodules: Revised Korean Society of Thyroid Radiology Consensus Statement and Recommendations. *Korean J Radiol*. 2016;17(3):370–395.
27. Kim DW, Lee EJ, Kim SH, et al. Ultrasound-guided fine-needle aspiration biopsy of thyroid nodules: comparison in efficacy according to nodule size. *Thyroid*. 2009;19(1):27–31.
28. Godazandeh G, Kashi Z, Zargarnataj S, Fazli M, Ebadi R, Kerdabadi EH. Evaluation the relationship between thyroid nodule size with malignancy and accuracy of fine needle aspiration biopsy (FNAB). *Acta Inform Med*. 2016;24(5):347–350.
29. Rosario PW, Silva AL, Calsolari MR. Is fine needle aspiration really not necessary in patients with thyroid nodules ≤ 1 cm with highly suspicious features on ultrasonography and candidates for active surveillance? *Diagn Cytopathol*. 2017;45(4):294–296.
30. Zhong LC, Lu F, Ma F, et al. Ultrasound-guided fine-needle aspiration of thyroid nodules: does the size limit its efficiency? *Int J Clin Exp Pathol*. 2015;8(3):3155–3159.
31. Harach HR, Franssila KO, Wasenius VM. Occult papillary carcinoma of the thyroid. A “normal” finding in Finland. A systematic autopsy study. *Cancer*. 1985;56(3):531–538.
32. Ito Y, Miyauchi A, Inoue H, et al. An observational trial for papillary thyroid microcarcinoma in Japanese patients. *World J Surg*. 2010;34(1):28–35.
33. Ito Y, Miyauchi A, Kihara M, Higashiyama T, Kobayashi K, Miya A. Patient age is significantly related to the progression of papillary microcarcinoma of the thyroid under observation. *Thyroid*. 2014;24(1):27–34.
34. Sugitani I, Toda K, Yamada K, Yamamoto N, Ikenaga M, Fujimoto Y. Three distinctly different kinds of papillary thyroid microcarcinoma should be recognized: our treatment strategies and outcomes. *World J Surg*. 2010;34(6):1222–1231.
35. Zhao C, Jiang W, Gao Y, Niu W, Zhang X, Xin L. Risk factors for lymph node metastasis (LNM) in patients with papillary thyroid microcarcinoma (PTMC): role of preoperative ultrasound. *J Int Med Res*. 2017;45(3):1221–1230.
36. Moon SJ, Kim DW, Kim SJ, Ha TK, Park HK, Jung SJ. Ultrasound assessment of degrees of extrathyroidal extension in papillary thyroid microcarcinoma. *Endocr Pract*. 2014;20(10):1037–1043.
37. Gweon HM, Son EJ, Kim JA, Youk JH. Predictive factors for active surveillance of subcentimeter thyroid nodules with highly suspicious us features. *Ann Surg Oncol*. 2017;24(6):1540–1545.
38. Kandil E, Noureldine SI, Tufano RP. Thyroidectomy vs active surveillance for subcentimeter papillary thyroid cancers – the cost conundrum. *JAMA Otolaryngol Head Neck Surg*. 2016;142(1):9–10.
39. Rodriguez JM, Moreno A, Parrilla P, et al. Papillary thyroid microcarcinoma: clinical study and prognosis. *Eur J Surg*. 1997;163(4):255–259.
40. Arem R, Padayatty SJ, Saliby AH, Sherman SI. Thyroid microcarcinoma: prevalence, prognosis, and management. *Endocr Pract*. 1999;5(3):148–156.
41. Kim JH, Pyo JS, Cho WJ. Clinicopathological significance and prognosis of medullary thyroid microcarcinoma: a meta-analysis. *World J Surg*. 2017;41(10):2551–2558.
42. Ito Y, Urano T, Nakano K, et al. An observation trial without surgical treatment in patients with papillary microcarcinoma of the thyroid. *Thyroid*. 2003;13(4):381–387.
43. Conzo G, Tartaglia E, Avenia N, et al. Role of prophylactic central compartment lymph node dissection in clinically N0 differentiated thyroid cancer patients: analysis of risk factors and review of modern trends. *World J Surg Oncol*. 2016;14:149.
44. Calò PG, Conzo G, Raffaelli M, et al. Total thyroidectomy alone versus ipsilateral versus bilateral prophylactic central neck dissection in clinically node-negative differentiated thyroid carcinoma. A retrospective multicenter study. *Eur J Surg Oncol*. 2017;43(1):126–132.
45. Gao L, Ma B, Zhou L, et al. The impact of presence of Hashimoto's thyroiditis on diagnostic accuracy of ultrasound-guided fine-needle aspiration biopsy in subcentimeter thyroid nodules: a retrospective study from FUSCC. *Cancer Med*. 2017;6(5):1014–1022.

Cancer Management and Research

Publish your work in this journal

Cancer Management and Research is an international, peer-reviewed open access journal focusing on cancer research and the optimal use of preventative and integrated treatment interventions to achieve improved outcomes, enhanced survival and quality of life for the cancer patient. The manuscript management system is completely online and includes

Submit your manuscript here: <https://www.dovepress.com/cancer-management-and-research-journal>

Dovepress

a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.