

Impact of preoperative exercise therapy on surgical outcomes in lung cancer patients with or without COPD: a systematic review and meta-analysis

Xiang Li^{1,*}
 Shaolei Li^{1,*}
 Shi Yan¹
 Yaqi Wang¹
 Xing Wang¹
 Alan DL Sihoe²
 Yue Yang¹
 Nan Wu¹

¹Key Laboratory of Carcinogenesis and Translational Research (Ministry of Education), Department of Thoracic Surgery II, Peking University Cancer Hospital and Institute, Beijing, China; ²Department of Surgery, University of Hong Kong, Hong Kong

*These authors contributed equally to this work

Objectives: This meta-analysis aimed to demonstrate the impact of preoperative exercise therapy on surgical outcomes in patients with lung cancer and COPD. Pulmonary function and muscle capacity were investigated to explore their potential links with outcome improvements after exercise.

Methods: Articles were searched from PubMed, Embase, and the Cochrane Library with criteria of lung cancer patients with or without COPD, undergoing resection, and receiving preoperative exercise training. Key outcomes were analyzed using meta-analysis.

Results: Seven studies containing 404 participants were included. Patients receiving preoperative exercise training had a lower incidence of postoperative pulmonary complications (PPCs; OR 0.44, 95% CI 0.27–0.71) and shorter length of hospital stay (standardized mean difference –4.23 days, 95% CI –6.14 to –2.32 days). Exceptionally, pneumonia incidence remained unchanged. Patients with COPD could not obviously benefit from exercise training to reduce PPCs (OR 0.44, 95% CI 0.18–1.08), but still might achieve faster recovery. No significant difference in pulmonary function was observed between the two groups. However, 6MWD and VO₂ peak were significantly improved after exercise training.

Conclusion: Preoperative exercise training may reduce PPCs for lung cancer patients. However, for patients with COPD undergoing lung cancer resection, the role of exercise is uncertain, due to limited data, which calls for more prospective trials on this topic. Rehabilitation exercise strengthens muscle capacity, but does not improve impaired pulmonary function, which emphasizes the possible mechanism of the protocol design.

Keywords: preoperative exercise, lung cancer, COPD, postoperative pulmonary complications, 6MWD, VO₂ peak

Introduction

Lung cancer is the most fatal cancer worldwide, with an estimated 1.2 million new cases and 1.1 million deaths in 2012.¹ Lung cancer is also the leading cause of cancer death in low-urbanization areas for both males and females. The number of deaths from lung cancer in low-urbanization areas was 35,100, with mortality of 40.71 in 100,000.² Surgical resection remains the key treatment for patients with lung cancer. However, many patients referred for surgery may also have comorbidities, such as COPD or other systematic diseases,³ which increase the risks of postoperative morbidity and mortality. Modern programs for enhanced recovery after surgery often

Correspondence: Nan Wu
 Department of Thoracic Surgery II,
 Peking University School of Oncology,
 Beijing Cancer Hospital & Institute
 52 Fucheng Road, Haidian District,
 Beijing, 100142, China
 Tel +86 10 8819 6569
 Fax +86 10 8812 2437
 Email nanwu@bjmu.edu.cn

include preoperative exercise training to reduce postoperative pulmonary complications (PPCs). Although isolated studies have looked at the impact of such preoperative exercise on patients with lung cancer and COPD, a comprehensive meta-analysis of the available data has hitherto been lacking. Some studies have revealed that intense-exercise therapy delivered in the perioperative period might be effective in helping patients recover well after lung surgery,^{4,5} though the extent of improvement varied among these reports. At the same time, there have been other studies not showing positive effects after rehabilitation training.⁶⁻⁸

PPCs are common after abdominal, cardiac, or thoracic surgery, and are associated with a higher rate of mortality, higher hospital costs, and prolonged hospital length of stay (LOS).⁵ Two systemic reviews have reported that intense exercise before lung surgery might be applied as an effective preoperative therapy to decrease the risk of PCs.^{4,5,9,10} Nevertheless, neither of the reviews conducted a meta-analysis, because of the substantial heterogeneity among the intervention methods. Although statistical analysis was lacking, the outcomes described after exercise were clearly supportive of further meta-analysis. A recently published meta-analysis¹¹ presented optimistic evidence of the impact of preoperative exercise training on surgical outcomes. However, the unselected targeting population and limited sample size indicated that more trials were needed to distinguish the potential benefit.

For the large group of patients with lung cancer and COPD at diagnosis, the role of preoperative exercise in this specific group requires further investigation. Several randomized studies that included patients with COPD for preoperative rehabilitation were published recently, which may add more evidence in this regard. More importantly, detailed exploration of outcome improvement could facilitate protocol design for exercise therapy, such as pulmonary function rehabilitation or muscle-strengthening training. Therefore, we designed this meta-analysis with the aim of determining whether preoperative exercise helps to improve surgical outcomes in patients with lung cancer and COPD. Parameters of pulmonary function and muscle capacity were investigated to explore their potential links with outcome improvements after exercise training.

Methods

This meta-analysis was conducted according to the *Cochrane Handbook for Systematic Reviews of Interventions* and presented based on PRISMA guidance.¹² The meta-analyses of randomized trials adhered to the guidelines outlined in the PRISMA statement. The protocol for this meta-analysis is available from PROSPERO (CRD42018088413).

Eligibility criteria

We only included randomized controlled trials (RCTs) and prospective trials of preoperative exercise training compared with no exercise training for lung cancer patients with or without COPD. We considered studies published in any language. Patients with usual care were set as the control group. After these preoperative interventions and basic therapy, patients received video-assisted thoracoscopic surgery or open thoracotomy, as scheduled. Types of surgery included wedge resection, segmentectomy, lobectomy, and pneumonectomy. Patients received similar clinical monitoring after surgery, and PCs were clearly recorded. A subset of patients in the included trials was diagnosed with mild or moderate COPD, whose data were extracted for further analysis.

Interventions

Protocols of the included studies varied in terms of exercise type, frequency, and intensity, ranging from three times per day for 1 week¹³ to five times per week for 4 weeks.¹⁴ Furthermore, two types of investigation after exercise were involved in the included studies: muscle capacity and pulmonary function analysis. Exercise programs contained aerobic exercise, resistance training, inspiratory muscle training, and education. In some studies, patients were advised to undertake a warm-up before exercise (after a 5-minute warm-up period, 50% of peak work rate was achieved).¹⁵

Outcome measurements

Primary outcomes were PPCs (summarized by type based on included studies: pneumonia, atelectasis, pulmonary embolism, respiratory failure, dyspnea, hemorrhagic drainage, empyema, interstitial pneumonia, bronchial stump dehiscence, bronchospasm, and bronchopleural fistula), pneumonia after surgery (new infiltrate plus either fever [$>38^{\circ}\text{C}$] and white-blood-cell count $>11,000$ or fever and purulent secretions),¹¹ duration of chest tube, and postoperative LOS. Secondary outcomes focused on the patients with lung cancer and COPD, including PPCs and postoperative LOS. Postintervention pulmonary function enhancement was also compared to analyze possible reasons linked with outcome change after preoperative exercise. Three parameters of exercise capacity were also compared after intervention: 6MWD (representing lower-limb muscle strength), VO_2 peak (representing physical performance), and Borg scores (representing severity of dyspnea).

Search methods

PubMed, Medline, and the Cochrane Library (central) were searched from the earliest date of each to June 2017. The

search string used the following keywords and was modified for each database: (“lung cancer” [MeSH] OR non-small-cell lung cancer OR small cell lung cancer) AND (lung surgery OR lobectomy) AND (physical therapy OR physiotherapy OR physical exercise OR physical therapy modalities).

Study selection

Through reading titles and abstracts, relevant articles were selected for full-text reading. Before this, two authors checked the search results and discussed the final list of the included articles. There was low bias in the included articles (Figure 1).





Figure 1 Flow diagram of study selection and funnel plot of studies included. **Abbreviation:** RCT, randomized controlled trial.

Cancer Management and Research downloaded from <https://www.dovepress.com/> by 18.232.188.251 on 01-Apr-2020
For personal use only.

Data extraction

Two authors read the articles and abstracted the data independently, and data were recorded using a predefined evidence table. Outcomes that we concentrated on were PPCs, LOS, and the duration of chest-tube drainage. We summarized information for each study in the list, eg, authors, publication year, type of study, number of participants, specific method of exercise, and index of pulmonary functions before surgery. The use of antibiotics was also clearly recorded.

Risk-of-bias appraisal

Risk of bias was assessed using the Cochrane Risk and Bias Tool, which was developed to evaluate the internal validity of the included RCTs.¹⁶ In RevMan 5.0, the tool contains seven criteria: random-sequence generation (selection bias), allocation concealment (selection bias), blinding of participants and personnel (performance bias), blinding of outcome assessment (detection bias), incomplete outcome data (attrition bias), selective reporting (report bias), and other risks of bias. The seventh criterion includes fraudulent results, other methodological flaws in RCTs, and the potential for industry bias.¹⁷

Statistical analyses

Across all outcomes, a meta-analysis was conducted. Effect estimates are reported as ORs for the dichotomous outcomes. For continuous data, mean difference (MD) was used if outcomes were measured in the same way between trials, and standardized mean difference (SMD) was applied for combining trials measuring the same outcome but using different methods.

For primary outcomes, PPCs were evaluated using pooled RRs with corresponding 95% CIs, and a random-effect model was used to account for potential clinical heterogeneity.¹⁸ Forest plots were constructed with RevMan 5.0, and $P < 0.05$ was regarded as statistically significant. Heterogeneity was evaluated by I^2 : if $I^2 > 50\%$, an article was considered to display substantial heterogeneity, requiring subgroup analysis. To merge the outcomes, the statistic used for analysis was selected for paired analysis. Not all studies supplied complete data. Meanwhile, data that were able to be extracted from existing articles were merged in a forest plot.

Results

Search outcome

Two authors (XL and SLL) performed the literature search, and NW resolved conflicts and discrepancies. The primary literature search produced 590 results, including 20 reviews. After screening of titles and abstracts, 24 studies were found to be possibly relevant and underwent full-text critical appraisal,

resulting in 16 exclusions (Table S1). Reasons for exclusion were outcomes recorded not being PPCs and LOS ($n=12$), no single regular intervention given ($n=1$), therapy was palliative ($n=1$), case reports ($n=2$), and duplicate publication ($n=1$). We included seven studies (Karenovics et al's study was a substudy of Licker et al) in the final analysis, as shown in Table 1. All included studies were assessed by the Grading of Recommendations, Assessment, Development, and Evaluations (GRADE) evidence profile (Table S2). Guidance from GRADE offered a methodology to evaluate the quality of the evidence.

Primary outcomes: PPCs, hospital LOS, chest-drain duration, pneumonia after preoperative exercise

Six studies^{13,14,19–22} had reported PPCs. Preoperative exercise reduced the risk of complications after surgery (OR 0.44, 95% CI 0.27–0.71; Figure 2A). The heterogeneity of this result was acceptable ($I^2=0$, $P < 0.0001$). Five studies^{8,13,14,21,22} reported postoperative pneumonia. Patients who received preoperative exercise before lung cancer resection had no statistically significant differences in the rate of pneumonia after surgery compared with those receiving usual care only (OR 0.47, 95% CI 0.24–0.95; Figure 2B). For LOS, five studies^{13,14,19,21,22} reported that patients receiving preoperative exercise training had shorter postoperative LOS (SMD -4.23 days, 95% CI -6.14 to -2.32 days; Figure 2C). Two studies^{14,19} recorded the duration of chest-tube drainage after surgery. Compared with patients receiving preoperative rehabilitation training, patients receiving usual care had a longer duration of chest-tube drainage (MD -3.28 days, 95% CI -5.21 to -1.36 days; Figure 2D).

Primary outcomes: PPCs and LOS of patients with lung cancer and COPD

Three studies^{19,21,22} included patients diagnosed with COPD and lung cancer simultaneously. Statistically, no significant difference was found for PPC incidence between the exercise-intervention group and usual-care group (OR 0.44, 95% CI 0.18–1.08; Figure 3A). Two studies^{19,22} reported LOS of patients with COPD and lung cancer. Patients receiving preoperative exercise had shorter LOS (MD -6.73 days, 95% CI -9.88 to -3.58 ; Figure 3B).

Secondary outcomes: impact of training exercise on pulmonary function capacity and exercise capacity before surgery

Pulmonary functions of patients in each study are listed in Table 2 and analyzed in Figure 4. There was little difference

Table 1 Overview of included studies

Studies	Participants n (male/female)	Characteristics	I: n (age ± SD) C: n (age ± SD)	Intervention Treatment	Outcome measurements	Comparison	Results
Pehlivan et al ¹³	60 (??)	NSCLC (stage IA–IIIB); open thoracotomy or VATS	I: 30 (54.10±8.53) C: 30 (54.76±8.45)	Preoperative exercise (three times a day), according to patient's tolerance to exercise speed and time; during the walking exercise, warm-up and cooldown included	PPCs LOS Pulmonary function	Usual care	Patients in the intervention group had lower incidence of PPCs (3.3%, 1/30 vs 16.6%, 5/30; P=0.04); hospital stay found to be significantly shortened by intensive physical therapy (P<0.001)
Morano et al ¹⁴	24 (9/15)	NSCLC (stage I–IIIA) and COPD; open thoracotomy or VATS	I: 12 (64.8±8) C: 12 (68.8±7.3)	Strength and endurance training; control group (breathing exercises for lung expansion)	PPCs LOS Pulmonary function	Chest physical therapy; CPT – breathing exercises for lung expansion	Patients in the intervention group had lower coincidence of complications (16.7%, 2/12 vs 77%, 7/12; P=0.01) and had shorter LOS (7.8±4.8 vs 12.2±3.6 days)
Benzo et al ¹⁹	19 (9/10) Two patients (one in each arm) were missing LOS data	NSCLC (stage I–IIIA) and COPD; open thoracotomy or VATS	I: 10 (70.2±8.61) C: 9 (72.0±6.69)	Breathing exercise through the device and sustaining that effort as long as they were able, with a goal of 15–20 minutes of daily use; ten face-to-face sessions of treatment in 1 week (twice a day)	PPCs LOS	Usual care	Patients in the intervention group had lower coincidence of complications (33%, 3/9 vs 63%, 5/8; P=0.23), and had shorter LOS (6.3±3.0 vs 11.0±6.3 days)
Sekine et al ²²	82 (76/6)	NSCLC (stage I–IV) or COPD; open thoracotomy	I: 22 (70.4±4.6) C: 60 (69.0±5.5)	Incentive spirometry, abdominal breathing and breathing exercises (with pursed lips, huffing and coughing after nebulizing for 15 minutes with a bronchodilator five times a day), pulmonary exercise for 30 minutes in the rehabilitation room and walking >5,000 steps every day for 2 weeks preoperatively	PPCs LOS	Usual care	Patients in the intervention group had lower coincidence of complications (18.2%, 4/22 vs 28.3%, 17/60), and had shorter LOS (29.0±9.0 vs 21.0±6.8 days; P=0.0003)
Karenovics et al ²⁰ (substudy of Licker et al)	151 (91/60)	NSCLC (stage I–III); open thoracotomy or VATS	I: 74 (64±?) C: 77 (64±?)	5-minute warm-up at 50% peak work rate achieved during CPET; two 10-minute series of 15-second sprint intervals	PPCs LOS in postanesthesia care unit Death VO ₂ peak	Usual care	Patients in the intervention group had lower coincidence of PPCs (23%, 17/74 vs 44%, 34/77; P=0.018) and a shorter stay in the postanesthesia care unit (median 7 hours, IQR _{25–75} 4–10)
Stefanelli et al ²⁴	40 (23/17)	NSCLC and COPD; lobectomy	I: 20 (<75) C: 20 (<75)	Group R: intensive preoperative PRP; group S, lobectomy only	VO ₂ peak		Peak VO ₂ did not change from T0 to T1 and deteriorated significantly from T1 to T2
Karenovics et al ²⁰	151 (91/60)	NSCLC (stage I–III); open thoracotomy or VATS	I: 74 (64±?) C: 77 (64±?)	5-minute warm-up period at 50% of peak work rate achieved during CPET; two 10-minute series of 15-second sprint intervals	PPCs Pneumonia 6MWD	Usual care	Patients in the intervention group had lower coincidence of PPCs (23%, 27/74 vs 44%, 39/77; P=0.018).

Note: "??": The number of the male and/or female was not clearly recorded in this published article.

Abbreviations: 6MWD, 6-minute walking distance; C, control; CPT, chest physical therapy; CPET, cardiopulmonary exercise testing; I, intervention; LOS, length of stay; NSCLC, non-small-cell lung cancer; PRP, pulmonary rehabilitation programme; PPCs, postoperative pulmonary complications; VATS, video-assisted thoracoscopic surgery.




Figure 2 Forest plots of comparison: intervention group vs control group in lung cancer patients undergoing resection. **Note:** (A) Risk of developing postoperative pulmonary complications; (B) incidence of postoperative pneumonia; (C) postoperative length of hospital stay; (D) duration of chest drainage.

between the two groups. Three studies analyzed 6MWD,^{13,14,21} which was considered an index of lower-limb muscle function.²³ Patients receiving exercise before surgery reported an improved 6MWD compared with those receiving usual care (SMD 71.25, 95% CI 39.68–102.82; Figure 5A). VO₂ peak, reflecting ability of oxygen uptake, was investigated to explore the role of exercise. After training, patients recorded a higher VO₂ peak compared to those without training^{20,24} (SMD 3.26, 95% CI 2.17–4.35; Figure 5B). Patients were less likely to improve in their dyspnea after preoperative exercise therapy^{13,21,24} (MD -0.15, 95% CI -0.66 to 0.36; Figure 5C).

Discussion

Numerous studies with large samples have clearly demonstrated that the presence of COPD is linked with an increase in lung cancer incidence.²⁵ Many patients with lung cancer and COPD will face a significant challenge during the perioperative period. How to guide these patients with COPD before and after cancer surgery with a proper training protocol to improve surgical outcomes remains to be answered. In this meta-analysis, the results imply that preoperative exercise for patients with lung cancer and COPD may be associated with shorter LOS compared with usual care only, but incidence of PPCs remained unchanged




Figure 3 Forest plots of comparison: intervention group vs control group in lung cancer patients with COPD treated with surgery. **Note:** (A) Risk of developing postoperative pulmonary complications; (B) postoperative length of hospital stay.

Table 2 Summary and comparison of pulmonary function in included studies

Pulmonary function test	Pehlivan et al ¹³		Morano et al ¹⁴		Karenovics et al ²⁰		Sekine et al ²²	
	Usual	Rehab	Usual	Rehab	Usual	Rehab	Usual	Rehab
FEV ₁	2.8±0.7	2.3±0.6	1.3±0.3	1.3±0.6	2.45±0.2	2.3±0.2	2.0±0.5	1.8±0.5
FEV ₁ %	–	–	58.8±13	54.8±4.3	87.6±4.3	85.8±5.1	71.6±13.8	63.4±18
D _{LCO}	21.3±6.1	21.1±6.9	–	–	–	–	–	–
FVC	3.2±0.8	3.1±0.6	2.1 (1.5–2.4)	1.7 (1.7–2.8)	3.6±0.2	3.4±0.2	3.1±0.6	3.1±0.7
FVC%	–	–	71 (63–89)	76 (65–79)	104±5	102±4	97.7±16	94.7±18.5

Note: Rehab, patients who had received preoperative exercise before thoracic surgery; usual, patients who had received usual care.

even after exercise training. Accompanied by outcome improvement, several parameters of exercise capacity were found to be strengthening after exercise training, such as 6MWD and VO₂ peak, because lower-limb muscle capacity was improved with higher ability of oxygen uptake. However, pulmonary function was not enhanced after training.

To date, a consistent protocol for preoperative exercise training has been lacking. Although pulmonary function and muscle-strength enhancement were commonly recorded in all the included studies, detailed methods for exercise training differed in terms of exercise duration and content.

Lai et al²¹ designed an exercise protocol containing pharmacotherapy and intense-exercise training (respiratory training and endurance training). Notably, the pharmacotherapy in that study was used to relieve symptoms of dyspnea for patients with lung cancer and COPD, which helped patients to complete the exercise training smoothly. In that study, the small sample and patients in the usual-care group having a certain extent of exercise both contributed to a degree of bias.

Meanwhile, Morano et al¹⁴ concluded that patients receiving preoperative exercise improved more when compared

with those who received chest physical therapy only. The protocol was carefully designed, and contained less extreme endurance training and inspiratory muscle training. The protocol also included flexibility, stretching, and balance exercises in the warm-up and cooldown sections. At the same time, medicine was used to optimize the different preintervention conditions for better comparison. However, patients in the control group received chest physical therapy, which was different from usual care. Another study conducted by Sekine et al reported that COPD patients received breathing-exercise training and walking-exercise training. The reported protocol contained breathing exercises, such as incentive spirometry, abdominal breathing, detailed breathing exercises, and walking exercise, such as walking for 5,000 steps every day.²² Although it was prospectively designed, allocation of the cases in the different groups was not randomized (60 vs 22).

Furthermore, Karenovics et al¹⁵ reported an exercise protocol that included 5 minutes of warm-up and cooldown exercises. Patients were guided by a professional exercise physiotherapist. The workload of exercise was adjusted in each session to target near-maximal heart rate toward the end




Figure 4 Forest plots of comparison of post-intervention pulmonary function: intervention group vs control group in lung cancer patients undergoing resection. **Note:** (A) FEV₁; (B) FEV₁%; (C) FVC%.

of each set of sprints. Notably, all patients were given advice regarding preoperative education and risk management.

Pulmonary rehabilitation has been proven to benefit patients with chronic lung disease, such as COPD.²⁶ Patients with COPD might experience deconditioning and exercise intolerance; therefore, preoperative rehabilitation should target these specific issues to obtain maximum effect. However, the mechanism of effect of exercise on outcome improvement remains unknown. Theoretically, preoperative exercise training is supposed to improve both lung function and muscle strength simultaneously. However, pulmonary function testing in this meta-analysis showed that most of the investigated parameters of pulmonary function were comparable or even slightly lower after exercise, though not significantly. A similar result was reported in other surgical procedures, such as lung transplantation.²⁷ Irreversible lung function after exercise training might represent the severity of the baseline condition of the lung disease and may not be the priority of treatment planning.

Interestingly, skeletal muscle weakness was reported commonly in patients with COPD,²⁸ which raised the question whether targeting muscle capacity may benefit surgical outcome. In this meta-analysis, we found a favorable trend of exercise capacity after training in the Forest plots, which might be related to outcome improvement, represented by improved 6MWD and VO₂ peak. A newly published article reported a strong relationship between peak aerobic capacity, lower-limb muscle function, and lung-diffusion capacity. 6MWD is significantly related to lower-limb muscle function, but not to pulmonary function.²³ Licker et al¹⁵ also reported the importance of VO₂ peak in predicting cardiopulmonary complications. There is a lack of uniform recommendations for a training protocol before surgery for patients with lung cancer and COPD; therefore, these results provide positive evidence that a protocol designed to focus on exercise capacity in the future might have an impact on postoperative complications in this group.




Figure 5 Forest plots of comparison: intervention group vs control group in lung cancer patients undergoing resection.

Note: (A) Postintervention 6MWD (index of lower-limb muscle strength); (B) VO_2 peak (reflecting physical performance); (C) Borg scores (representing dyspnea).

Limitations

This study was subject to several limitations. First, PPCs were set as the main outcome in the analysis. However, different studies had individual definitions of complications, which might have caused bias in counting the PPCs. To reduce this selection bias, we chose those studies defining pneumonia clearly as a standard complication in the analysis, which showed a consistent trend with the overall outcome. Second, the protocol of every study was individually designed and the duration and intensity of exercise varied among the studies. During the process of meta-analysis, the heterogeneity of muscle strength and lung function among studies was still acceptable. Third, the end-point event for preoperative exercise is still obscure among studies, eg, how to decide the outcome of exercise, whether improved or declined, and what the parameters for performance evaluation are, such as lung function and exercise capacity. Exercise-protocol design should be based on information that allows evaluation of the efficacy of the procedure. Fourth, the type of exercise training (aerobic, strength, inspiratory muscle training) and the optimal duration were not consistent in the included studies, which should be balanced against the risk

of delaying cancer resection and consequent extension of the disease. Finally, randomized trials on this topic are still scarce, especially for those patients with lung cancer and COPD, which may limit the power of the meta-analysis. In future, more prospective randomized studies are needed to accumulate further evidence.

Conclusion

The goal of the preoperative exercise is to decrease the risk of complications after surgery and enhance recovery, which can ultimately shorten LOS. This meta-analysis provides some evidence of the effect of preoperative exercise on enhanced recovery after surgery for patients with lung cancer and COPD. Future research should pay attention to details of the exercise protocol, such as intensity, duration, and methods, for those patients preparing for surgery.

Acknowledgments

The authors thank Elixigen Company (Huntington Beach, CA, USA) for proofreading. Beijing Municipal Administration of Hospitals Clinical Medicine Development Fund (grant number ZYLX201509), National Key R&D Program of

China (No. 2018YFC0910700), Beijing Municipal Science & Technology Commission (No. Z161100000516063), Beijing Human Resources and Social Security Bureau (Beijing Millions of Talents Project, 2018A05), Special Fund of Beijing Municipal Administration of Hospitals Clinical Medicine Development (No. XMLX201841), Beijing Municipal Administration of Hospitals' Youth Programme (No. QML20171103), National Natural Science Foundation of China (grant number: 81502578), and Peking University Medicine Seed Fund for Interdisciplinary Research (BMU2018MX008).

Disclosure

The authors report no conflicts of interest in this work.

References

1. Ferlay J, Soerjomataram I, Dikshit R, et al. Cancer incidence and mortality worldwide: sources, methods and major patterns in GLOBOCAN 2012. *Int J Cancer*. 2015;136(5):E359–E386.
2. Chen W, Zheng R, Zhang S, et al. Cancer incidence and mortality in China in 2013: an analysis based on urbanization level. *Chin J Cancer Res*. 2017;29(1):1–10.
3. Licker MJ, Widikker I, Robert J, et al. Operative mortality and respiratory complications after lung resection for cancer: impact of chronic obstructive pulmonary disease and time trends. *Ann Thorac Surg*. 2006;81(5):1830–1837.
4. Rodriguez-Larrad A, Lascurain-Aguirrebena I, Abecia-Inchaurregui LC, Seco J. Perioperative physiotherapy in patients undergoing lung cancer resection. *Interact Cardiovasc Thorac Surg*. 2014;19(2):269–281.
5. Sommer MS, Trier K, Vibe-Petersen J, et al. Perioperative rehabilitation in operation for lung cancer (PROLUCA) – rationale and design. *BMC Cancer*. 2014;14:404.
6. Coats V, Maltais F, Simard S, et al. Feasibility and effectiveness of a home-based exercise training program before lung resection surgery. *Can Respir J*. 2013;20(2):e10–e16.
7. Bobbio A, Chetta A, Ampollini L, et al. Preoperative pulmonary rehabilitation in patients undergoing lung resection for non-small cell lung cancer. *Eur J Cardiothorac Surg*. 2008;33(1):95–98.
8. Benzo R, Wigle D, Novotny P, et al. Preoperative pulmonary rehabilitation before lung cancer resection: results from two randomized studies. *Lung Cancer*. 2011;74(3):441–445.
9. Pouwels S, Fiddelaers J, Teijink JA, Woort JF, Siebenga J, Smeenk FW. Preoperative exercise therapy in lung surgery patients: a systematic review. *Respir Med*. 2015;109(12):1495–1504.
10. Smetana GW. Postoperative pulmonary complications: an update on risk assessment and reduction. *Cleve Clin J Med*. 2009;76(Suppl 4):S60–S65.
11. Cavalheri V, Granger C. Preoperative exercise training for patients with non-small cell lung cancer. *Cochrane Database Syst Rev*. 2017;6:CD012020.
12. Liberati A, Altman DG, Tetzlaff J, et al. The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate health-care interventions: explanation and elaboration. *BMJ*. 2009; 339:b2700.
13. Pehlivan E, Turna A, Gurses A, Gurses HN. The effects of preoperative short-term intense physical therapy in lung cancer patients: a randomized controlled trial. *Ann Thorac Cardiovasc Surg*. 2011;17(5): 461–468.
14. Morano MT, Araújo AS, Nascimento FB, et al. Preoperative pulmonary rehabilitation versus chest physical therapy in patients undergoing lung cancer resection: a pilot randomized controlled trial. *Arch Phys Med Rehabil*. 2013;94(1):53–58.
15. Karenovics W, Licker M, Ellenberger C, et al. Short-term preoperative exercise therapy does not improve long-term outcome after lung cancer surgery: a randomized controlled study. *Eur J Cardiothorac Surg*. 2017;52(1):47–54.
16. Tricco AC, Cogo E, Holroyd-Leduc J, et al. Efficacy of falls prevention interventions: protocol for a systematic review and network meta-analysis. *Syst Rev*. 2013;2(1):38.
17. Tricco AC, Thomas SM, Veroniki AA, et al. Comparisons of interventions for preventing falls in older adults: a systematic review and meta-analysis. *JAMA*. 2017;318(17):1687–1699.
18. DerSimonian R, Laird N. Meta-analysis in clinical trials revisited. *Contemp Clin Trials*. 2015;45(Pt A):139–145.
19. Benzo R, Wigle D, Novotny P, et al. Preoperative pulmonary rehabilitation before lung cancer resection: results from two randomized studies. *Lung Cancer*. 2011;74(3):441–445.
20. Karenovics W, Licker M, Ellenberger C, et al. Short-term preoperative exercise therapy does not improve long-term outcome after lung cancer surgery: a randomized controlled study. *Eur J Cardiothorac Surg*. 2017;52(1):47–54.
21. Lai Y, Su J, Yang M, Zhou K, Che G. [Impact and effect of preoperative short-term pulmonary rehabilitation training on lung cancer patients with mild to moderate chronic obstructive pulmonary disease: a randomized trial]. *Zhongguo Fei Ai Za Zhi*. 2016;19(11):746–753. Chinese.
22. Sekine Y, Chiyo M, Iwata T, et al. Perioperative rehabilitation and physiotherapy for lung cancer patients with chronic obstructive pulmonary disease. *Jpn J Thorac Cardiovasc Surg*. 2005;53(5):237–243.
23. Burtin C, Franssen FME, Vanfleteren LEGW, Groenen MTJ, Wouters EFM, Spruit MA. Lower-limb muscle function is a determinant of exercise tolerance after lung resection surgery in patients with lung cancer. *Respirology*. 2017;22(6):1185–1189.
24. Stefanelli F, Meoli I, Cobuccio R, et al. High-intensity training and cardiopulmonary exercise testing in patients with chronic obstructive pulmonary disease and non-small-cell lung cancer undergoing lobectomy. *Eur J Cardiothorac Surg*. 2013;44(4):e260–e265.
25. Houghton AM. Mechanistic links between COPD and lung cancer. *Nat Rev Cancer*. 2013;13(4):233–245.
26. Ries AL, Bauldoff GS, Carlin BW, et al. Pulmonary rehabilitation: joint ACCP/AACVPR evidence-based clinical practice guidelines. *Chest*. 2007;131(5 Suppl):4S–42S.
27. Pehlivan E, Balci A, Kılıç L, Kadakal F. Preoperative pulmonary rehabilitation for lung transplant: effects on pulmonary function, exercise capacity, and quality of life; first results in turkey. *Exp Clin Transplant*. 2018;16(4):455–460.
28. Abdulai RM, Jensen TJ, Patel NR, et al. Deterioration of limb muscle function during acute exacerbation of chronic obstructive pulmonary disease. *Am J Respir Crit Care Med*. 2018;197(4):433–449.

Supplementary materials

Table S1 Studies excluded in this review

Study	Research for exclusion
Cavalheri et al 2015 ¹	Research for the postoperative exercise
Edvardsen et al 2014 ²	Research for the postoperative exercise
Maeda et al 2015 ³	Duplication; postoperative exercise
Sommer et al 2016 ⁴	Duplication; Included early postoperative exercise training
Esteban et al 2017 ⁵	Not RCT
Cavalheri et al 2017 ⁶	Lacking proper statistics
Nai-WenChang et al 2014 ⁷	Postoperative walking exercise
Missel et al 2015 ⁸	Postoperative exercise
Jones et al 2007 ⁹	Preoperative exercise on cardiopulmonary fitness
Valérie et al 2013 ¹⁰	Lacking proper statistics
Crandal et al 2014 ¹¹	Not RCT
Wiskemann et al 2016 ¹²	Lacking proper statistics
FANG et al 2014 ¹³	Not RCT
Loewen et al 2007 ¹⁴	Not RCT
Weiner et al 1984 ¹⁵	Not RCT and Lacking proper statistics

Abbreviation: RCT, randomized controlled trial.

Table S2 Grading of recommendations, assessment, development, and evaluations

Certainty assessment		Patients, n					Effect		Certainty		
Studies, n	Design	Risk of bias	Inconsistency	Indirectness	Imprecision	Other considerations	Preoperative exercise	Placebo		Relative (95% CI)	Absolute (95% CI)
PPCs											
6 ³⁻⁸	RCT ^a	Not serious	Not serious	Not serious	Serious ^b	All plausible residual confounding would reduce the demonstrated effect	171	211	OR 0.35 (0.21–0.59)	190 fewer per 1,000 (from 108 fewer to 246 fewer)	⊕⊕⊕⊕ High
Pneumonia											
5 ³⁻⁷	RCT ^a	Not serious	Not serious	Serious ^c	Serious	None	97	131	OR 0.45 (0.16–1.25)	67 fewer per 1,000 (from 27 more to 106 fewer)	⊕⊕ Low
Length of hospital stay											
5 ³⁻⁷	RCT ^a	Not serious	Not serious	Not serious	Very serious ^d	None	97	134	–	SMD 1.02 lower (1.31 lower to 0.74 lower)	⊕⊕ Low
Days of chest-tube drainage											
2 ^{3,5}	RCT	Not serious	Not serious	Not serious	Serious ^e	None	21	17	–	MD 3.33 lower (5.35 lower to 1.3 lower)	⊕⊕⊕ Moderate
PPCs of COPD patients											
3 ^{3,6,7}	RCT ^a	Not serious ³	Not serious	Serious ^f	Not serious	None	9/55 (16.4%)	27/92 (29.3%)	–	OR 0.44 (0.18–1.08)	⊕⊕⊕ Moderate
Length of hospital stay in patients with COPD											
2 ^{3,7}	RCT ^a	Serious ⁶	Not serious	Not serious	Very serious ^d	None	31	68	–	MD 6.79 lower (9.69 lower to 3.89 lower)	⊕ Very low
6MWD											
3 ³⁻⁶	RCT	Serious	Not serious	Not serious	Serious ^h	All plausible residual confounding would reduce the demonstrated effect	66	66	–	SMD 0.54 higher (0.19 higher to 0.88 higher)	⊕⊕⊕⊕ High
VO₂ peak (reflecting physical performance)											
2 ^{8,9}	RCT	Serious	Not serious	Not serious	Serious ^h	All plausible residual confounding would reduce the demonstrated effect	94	94	–	SMD 0.96 higher (0.66 higher to 1.27 higher)	⊕⊕ Low
Borg scores (representing dyspnea)											
3 ^{4,6,9}	RCT	Not serious	Serious	Not serious	Serious ^h	All plausible residual confounding would reduce the demonstrated effect	74	74	–	SMD 0.54 higher (0.19 higher to 0.88 higher)	⊕⊕ Low

Note: ¹One of the studies reported by Sekine et al⁶ was prospective; ²PPCs were not defined clearly; ³detected by the clinical laboratory index; ⁴length of stay can be influenced by hospital conditions and other nonclinical problems; ⁵duration of drainage was not consistent in different clinical institutes; ⁶different institutes had different definition of PPCs; ⁷small sample of participants; ⁸distance walked influenced by many factors.

Abbreviations: MD, mean difference; PPCs, postoperative pulmonary complications; RCT, randomized controlled trial; SMD, standardized mean difference.

References

1. Cavalheri V, Jenkins S, Cecins N, et al. Patterns of sedentary behaviour and physical activity in people following curative intent treatment for non-small cell lung cancer. *Chron Respir Dis* 2016;13:82–85.
2. Edvardsen E, Skjonsberg OH, Holme I, et al. High-intensity training following lung cancer surgery: a randomised controlled trial. *Thorax* 2015;70:244–250.
3. Maeda K, Higashimoto Y, Honda N, et al. Effect of a postoperative outpatient pulmonary rehabilitation program on physical activity in patients who underwent pulmonary resection for lung cancer. *Geriatr Gerontol Int* 2016;16:550–555.
4. Sommer MS, Trier K, Vibe-Petersen J, et al. Perioperative Rehabilitation in Operable Lung Cancer Patients (PROLUCA): A Feasibility Study. *Integr Cancer Ther* 2016;15:455–466.
5. Esteban PA, Hernandez N, Novoa NM, et al. Evaluating patients' walking capacity during hospitalization for lung cancer resection. *Interactive cardiovascular and thoracic surgery* 2017;25:268–271.
6. Cavalheri V, Granger CL, Irving LB, et al. Short-term preoperative exercise training: should we expect long-term benefits without postoperative exercise stimulus? *European journal of cardio-thoracic surgery : official journal of the European Association for Cardio-thoracic Surgery* 2017;52:1009.
7. Chang NW, Lin KC, Lee SC, et al. Effects of an early postoperative walking exercise programme on health status in lung cancer patients recovering from lung lobectomy. *J Clin Nurs* 2014;23:3391–3402.
8. Missel M, Pedersen JH, Hendriksen C, et al. Regaining familiarity with own body after treatment for operable lung cancer - a qualitative longitudinal exploration. *Eur J Cancer Care (Engl)* 2016;25:1076–1090.
9. Jones LW, Peddle CJ, Eves ND, et al. Effects of presurgical exercise training on cardiorespiratory fitness among patients undergoing thoracic surgery for malignant lung lesions. *Cancer* 2007;110:590–598.
10. Coats V, Maltais F, Simard S, et al. Home-based rehabilitation program for lung cancer patients. *European Respiratory Journal* 2011.
11. Crandall K, Maguire R, Campbell A, et al. Exercise intervention for patients surgically treated for Non-Small Cell Lung Cancer (NSCLC): a systematic review. *Surgical oncology* 2014;23:17–30.
12. Wiskemann J, Hummler S, Diepold C, et al. POSITIVE study: physical exercise program in non-operable lung cancer patients undergoing palliative treatment. *BMC cancer* 2016;16:499.
13. Fang Y, Ma G, Lou N, et al. Preoperative Maximal Oxygen Uptake and Exercise-induced Changes in Pulse Oximetry Predict Early Postoperative Respiratory Complications in Lung Cancer Patients. *Scandinavian Journal of Surgery* 2014;103:201–208.
14. Loewen GM, Watson D, Kohman L, et al. Preoperative exercise Vo2 measurement for lung resection candidates: results of Cancer and Leukemia Group B Protocol 9238. *Journal of thoracic oncology : official publication of the International Association for the Study of Lung Cancer* 2007;2:619–625.
15. Weiner P, Man A, Weiner M, et al. The effect of incentive spirometry and inspiratory muscle training on pulmonary function after lung resection. *The Journal of thoracic and cardiovascular surgery* 1997;113:552–557.
16. Sekine Y, Chiyo M, Iwata T, et al. Perioperative rehabilitation and physiotherapy for lung cancer patients with chronic obstructive pulmonary disease. The Japanese journal of thoracic and cardiovascular surgery : official publication of the Japanese Association for Thoracic Surgery = Nihon Kyobu Geka Gakkai zasshi 53, 237–243.

Cancer Management and Research

Publish your work in this journal

Cancer Management and Research is an international, peer-reviewed open access journal focusing on cancer research and the optimal use of preventative and integrated treatment interventions to achieve improved outcomes, enhanced survival and quality of life for the cancer patient. The manuscript management system is completely online and includes

Submit your manuscript here: <https://www.dovepress.com/cancer-management-and-research-journal>

a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Dovepress