

# Psychometric properties of the Brief Symptom Inventory in nomophobic subjects: insights from preliminary confirmatory factor, exploratory factor, and clustering analyses in a sample of healthy Italian volunteers

Mohammad Adawi<sup>1</sup>  
 Riccardo Zerbetto<sup>2</sup>  
 Tania Simona Re<sup>2,3</sup>  
 Bishara Bisharat<sup>4,5</sup>  
 Mahmud Mahamid<sup>4</sup>  
 Howard Amital<sup>6,7</sup>  
 Giovanni Del Puente<sup>8</sup>  
 Nicola Luigi Bragazzi<sup>2,3,8,9</sup>

<sup>1</sup>Bar-Ilan Faculty of Medicine, Padeh and Ziv Hospitals, Zafat, Israel; <sup>2</sup>GESTALT Study Center (CSTG), Milan, Italy; <sup>3</sup>UNESCO Chair "Health Anthropology, Biosphere and Healing Systems", University of Genoa, Genoa, Italy; <sup>4</sup>EMMS Nazareth Hospital, Nazareth, Azrieli Faculty of Medicine, Bar-Ilan University, Safed, Israel; <sup>5</sup>The Society for Health Promotion of the Arab Community, The Max Stern Yezreel Valley College, Nazareth, Israel; <sup>6</sup>Sackler Faculty of Medicine, Tel Aviv University, Tel Aviv, Israel; <sup>7</sup>Zabludowicz Center for Autoimmune Diseases, Department of Medicine B, Sheba Medical Center, Tel HaShomer, Israel; <sup>8</sup>DINO GMI, Section of Psychiatry, University of Genoa, Genoa, Italy; <sup>9</sup>Department of Health Sciences, Postgraduate School of Public Health, University of Genoa, Genoa, Italy

Correspondence: Nicola Luigi Bragazzi  
 Department of Health Sciences (DISSAL),  
 University of Genoa, Via Antonio Pastore  
 I, Genoa I6132, Italy  
 Email robertobragazzi@gmail.com

**Background:** The Brief Symptom Inventory (BSI), developed by Derogatis in 1975, represents an important standardized screening instrument that enables one to quantitatively assess psychological distress and psychiatric disorders. The BSI is a 53-item self-report scale, measuring nine dimensions that can be summed up to reflect three global indices, including the General Severity Index (GSI). In the era of new information and communication technologies, nomophobia ("no mobile phobia") is an emerging disorder, characterized by the fear of being out of mobile phone contact. Nothing is known, however, about the factor structure and reliability of the BSI in a population of nomophobic subjects. This study aimed at addressing this gap in knowledge.

**Methods:** A sample of 403 subjects aged  $27.91 \pm 8.63$  years (160 males, 39.7% of the entire sample, and 243 females, 60.3%), recruited via snowball sampling, volunteered to take part in the study. The Italian versions of the Nomophobia questionnaire and the BSI were administered. Exploratory factor analyses, confirmatory factor analyses, and clustering analysis were carried out together with correlation analysis, analysis of variance, and multivariate regression analysis.

**Results:** For each BSI subscale, scores were significantly higher than the norms. The nine subscales exhibited acceptable-to-good Cronbach's alpha coefficients, varying from 0.733 for psychoticism to 0.875 for depression. Overall, the reliability of the entire instrument proved to be excellent (alpha coefficient=0.972). Furthermore, all BSI subscales as well as BSI synthetic indexes correlated with nomophobia in a significant way. Stratifying the population according to the severity of nomophobia (mild, 206 individuals, 51.1% of the sample; moderate, 167 subjects, 41.4%; and severe, 30 individuals, 7.4%), the GSI score could distinguish ( $P < 0.001$ ) between mild and moderate ( $0.99 \pm 0.71$  vs  $1.32 \pm 0.81$ ) and between mild and severe ( $0.99 \pm 0.71$  vs  $1.54 \pm 0.79$ ) nomophobia, although not between moderate and severe nomophobia ( $P > 0.05$ ). Similar patterns could be found for the other subscales of the BSI. Finally, looking at the fit indexes, the second-order 9-factor model best fit the data compared with the Derogatis 1-factor model.

**Conclusion:** The findings of our study show that the BSI is a reliable and valid instrument with acceptable psychometric properties, and can be administered to populations of nomophobic subjects.

**Keywords:** nomophobia, Brief Symptom Inventory, psychometric properties, questionnaire, confirmatory factor analysis

## Introduction

For researchers in the field of psychopathology, the Brief Symptom Inventory (BSI), developed by Derogatis in 1975, represents an important, standardized screening instrument that enables one to quantitatively assess psychological distress and psychiatric disorders.<sup>1-5</sup>

The BSI has been used in a variety of settings, either with adolescents or adults, and, consequently, its psychometric properties have been widely investigated and appraised. The original factor structure has been intentionally designed and developed for adults and adolescents with a range of psychiatric disorders, even though the instrument has also been subsequently tested among cancer patients or individuals suffering from other chronic-degenerative disorders, among others.<sup>6,7</sup>

In the era of new information and communication technologies characterized by widespread and pervasive use of smart phones and mobile devices,<sup>8</sup> nomophobia (“no mobile phobia”) is an emerging psychological concern and disorder.<sup>9-11</sup> Nomophobia can be defined as the irrational fear, stress, or worry of being out of mobile phone contact, that is to say being without one’s own device or being unable to use it due to the absence of a signal or low network coverage, running out of minutes, battery power, or credit, or for some other reasons.

Some studies have investigated the relationship between psychopathological symptoms and technological addictions. For instance, in a sample of 126 university students, Adalier and Balkan<sup>12</sup> found a significant correlation between Internet addiction and psychopathological symptoms like somatization, obsession–compulsion, interpersonal sensitivity, depression, anxiety, hostility, phobic anxiety, paranoid ideation, and psychoticism. In a sample of 334 subjects, Wegmann et al<sup>13</sup> studied the effects of depression and social anxiety on addictive use of social networking sites and found that they were mediated by Internet use expectancies and self-regulation. Similar findings were reported by Stavropoulos et al,<sup>14</sup> who documented a statistically significant association between anxiety levels and Internet addiction among adolescents.

Nothing is known, however, about the factor structure, validity, and reliability of the BSI in a population of nomophobic subjects. This study aimed at addressing this gap in knowledge. On the basis of the previously mentioned studies,<sup>12-14</sup> we hypothesized to find a statistically significant association between nomophobic use of smart phones and psychopathological symptoms as measured by the BSI.

## Materials and methods

### Population

Our sample size well exceeded the minimum number of 300 subjects suggested by Tabachnick and Fidell<sup>15</sup> as a general rule of thumb for properly performing factor analysis. For this cross-sectional study, participants (mainly, undergraduate students and younger subjects) were recruited via an online survey using a snowball approach. Further details concerning the population recruited can be found in our previous publication.<sup>11</sup> Briefly, a sample of 403 subjects aged  $27.91 \pm 8.63$  years (160 males, 39.7% of the entire sample, and 243 females, 60.3%) volunteered to take part in the study. In detail, 45 subjects spent <1 hour on their mobile phone per day (11.2%), 94 spent between 1 and 2 hours (23.3%), 69 spent between 2 and 3 hours (17.1%), 58 spent between 3 and 4 hours (14.4%), 48 spent between 4 and 5 hours (11.9%), 29 spent between 5 and 7 hours (7.2%), 36 spent between 7 and 9 hours (8.9%), and, finally, 24 spent >10 hours (6.0%).

### Instruments

#### Brief Symptom Inventory

The BSI is a 53-item self-report scale designed to evaluate psychopathological and psychological symptoms, measuring nine dimensions (namely, somatization, obsession–compulsion, interpersonal sensitivity, depression, anxiety, hostility, phobic anxiety, paranoid ideation, and psychoticism) that can be summed up to reflect three global indices. These synthetic indices are the General Severity Index (GSI), the Positive Symptom Distress Index, and the Positive Symptom Total. In more detail, the BSI uses a 5-point Likert scale, ranging from 0 (“not at all”) to 4 (“extremely”). The BSI has sound psychometric properties: in the original administration of the questionnaire, internal consistency coefficients ranged from 0.71 to 0.85.

#### Nomophobia questionnaire

Besides the Italian validated version of the BSI, the Italian version of the Nomophobia questionnaire (NMP-Q), translated from the instrument originally developed by Yildirim and Correia,<sup>10</sup> was administered. Exploratory factor analysis (EFA) has previously demonstrated good psychometric properties of the instrument (Cronbach’s alpha coefficient of 0.95, 0.94, 0.89, and 0.88 for the overall questionnaire and for its three factors – factor 1, not being able to access information; factor 2, giving up convenience/losing connectedness; and factor 3, not being able to communicate – respectively). Furthermore, validity of the questionnaire was confirmed

by conducting regression analysis with the number of hours spent on the mobile phone as the regressor.<sup>11</sup> The Italian version was found to have a 3-factor structure, as opposed to the initial version of Yildirim and Correia,<sup>10</sup> and to the translated and validated versions in Spanish<sup>16</sup> and in Persian.<sup>17</sup>

For the purpose of administration, Google Forms, an open-source tool for developing and administering ad hoc online questionnaires/surveys, was utilized. Due to the snowball sampling procedure, we were not able to compute the responder rate. There were no missing items to deal with, and, as such, no imputation analysis was necessary.

Based on the NMP-Q score, the nomophobic level was categorized as “mild nomophobia” (scores in the range 21–59), “moderate nomophobia” (scores in the range 66–99), or “severe nomophobia” (scores  $\geq 100$ ).

### Data analysis strategy

Once the data were collected, before commencing any data handling and processing, they were visually inspected for potential outliers. Normality of data distribution was checked by performing the D’Agostino–Pearson omnibus test. Then, some descriptive analyzes were carried out with the aim to provide information about the general characteristics of the study groups in terms of reported scores. Finally, Cronbach’s alpha coefficients were calculated as estimates of reliability/internal consistency of the instrument. The following rule of thumb was utilized: the coefficient was judged unacceptable if  $< 0.5$ , poor in the range 0.5–0.6, questionable in the range 0.6–0.7, acceptable in the range 0.7–0.8, good in the range 0.8–0.9, and, finally, deemed excellent if  $> 0.9$ .

Correlation analysis was performed between the NMP-Q and BSI scores. The magnitude of the Pearson’s coefficient was interpreted following the rule of thumb developed by Hinkle et al:<sup>18</sup> the strength of the correlation was deemed negligible if the  $r$  coefficient ranged from 0.00 to 0.30, low from 0.30 to 0.50, moderate from 0.50 to 0.70, high from 0.70 to 0.90, and very high from 0.90 to 1.00. Multivariate regression analyses were performed to shed light on the predictors of the overall GSI score and each BSI subscale score. Furthermore, analysis of variance was conducted for the GSI score and each BSI subscale score based on the nomophobic levels.

For all analyses, data with  $P < 0.05$  were considered statistically significant.

The commercial software Statistical Package for Social Sciences (SPSS for Windows, version 24.0, released 2017; IBM Corp., Armonk, NY, USA) was used for carrying out these statistical analyses. Graphs were obtained using the

commercial MedCalc Statistical Software version 17.9.7 (2017; MedCalc Software bvba, Ostend, Belgium; <http://www.medcalc.org>).

### Clustering analysis

Clustering analysis, based on the nomophobic levels, was conducted with the commercial software SPSS. It was carried out in two subsequent steps, hierarchical and k-means clustering techniques, in order to find the optimal number of clusters.

### Exploratory factor analysis

At first, EFA was performed in order to investigate the factor structure of the BSI questionnaire. The Kaiser–Meyer–Olkin measure was computed to assess the sampling adequacy. Ideally, the Kaiser–Meyer–Olkin should be  $> 0.60$ . The likely number of factors was determined by: 1) the number of factors with eigenvalues  $> 1$ ; and 2) a visual inspection of Cattell’s scree plot. After checking the factor loadings, items were deleted in cases of unsatisfactory loading (ie, values  $< 0.45$ ) or loading conflicting with a sound theoretical explanation. Different principal component analyses with varimax rotation runs were, therefore, carried out iteratively until a satisfactory, clearly interpretable solution was finally achieved. Cases of cross-loading were interpreted according to salience and overall explained variance, with theoretical considerations also being taken into account (ie, loadings not conflicting with a sound preestablished theoretical framework).

EFA was conducted utilizing the commercial SPSS software.

### Confirmatory factor analysis

Confirmatory factor analysis (CFA) was carried out using the open-source software Jamovi (version 0.0.03) and the commercial EQS software (version 6.3 for Windows; Multivariate Inc., Temple City, CA, USA). Differently from EFA, CFA enables researchers to quantitatively assess how well an *a priori*, theoretically specified factor model explains the observed pattern of correlations or covariances.

### Goodness of fit indices

As suggested and recommended by many scholars, a wide range of fit indices was calculated and reported, namely discrepancy indices (including the chi-squared test and the Steiger–Lind root mean square error of approximation [RMSEA]), tests comparing the target model with the null model (like the Tucker–Lewis Index [TLI] and Bentler’s Comparative Fit Index [CFI]), and information theory

goodness of fit measures (like the Akaike Information Criterion [AIC] and the Schwarz's Bayesian criterion, known also as Bayesian Information Criterion [BIC]).

Furthermore, the standardized root mean square residual was computed following the recommendation of Jöreskog and Sörbom.<sup>40</sup>

## Cutoff and threshold values

The *P*-value associated with the chi-squared test should exceed 0.05 (ie, it should not be statistically significant). Further, the value of chi-squared divided by the degrees of freedom should ideally be <2.0. As far as the RMSEA is concerned, MacCallum et al in 1996<sup>19</sup> and Steiger in 2000<sup>20</sup> have suggested using 0.01, 0.05, and 0.08 as threshold values to indicate excellent, good, and mediocre fit, respectively. In general, according to Steiger,<sup>20</sup> values higher than 0.10 indicate poor fitting models. Hu and Bentler in 1995<sup>21</sup> recommended a value of RMSEA <0.06.

The TLI should be above 0.95 according to Hu and Bentler.<sup>21</sup> The CFI should exceed 0.95 according to Bentler<sup>22</sup> and to Hu and Bentler,<sup>21</sup> or 0.90 according to other scholars. Acceptable values of the CFI are in the range 0.80–0.90, whereas values <0.80 are unacceptable.

Finally, acceptable values of the AIC and the BIC should ideally be close to 0.

## Ethical clearance

All procedures described in the article and performed in the study were carried out in accordance with the ethical standards of the institutional research committee, and with the 1964 Helsinki Declaration and its subsequent amendments. The study protocol was approved by the ethical committee of the University of Genoa and the UNESCO Chair. Every participant gave written informed consent after being thoroughly advised about the study's aims and procedures.

## Results

Scores as the mean and SD for each subscale of the BSI are reported in Table 1: they ranged from 0.856±0.7986 for interpersonal sensitivity to 1.244±0.7936 for anxiety. In all cases, scores were significantly higher than the norms for adult nonpatients (*P*<0.001), indicating that the BSI is potentially able to distinguish between a non-nomophobic individual and one suffering from nomophobia.

The nine subscales exhibited acceptable-to-good Cronbach's alpha coefficients (as can be seen in Table 2), varying from 0.733 for psychoticism to 0.875 for depression. All coefficients were good except for phobic anxiety and

**Table 1** Scores obtained as mean and SD for each subscale of the Brief Symptom Inventory (BSI)

BSI subscale	Mean	SD	Different from norms for adult nonpatients
Anxiety	1.244	0.7936	<i>P</i> <0.001
Depression	1.225	0.8264	<i>P</i> <0.001
Hostility	0.978	0.8103	<i>P</i> <0.001
Interpersonal sensitivity	0.856	0.7986	<i>P</i> <0.001
Obsession–compulsion	0.884	0.7698	<i>P</i> <0.001
Phobic anxiety	0.931	0.8073	<i>P</i> <0.001
Paranoid ideation	1.101	0.8516	<i>P</i> <0.001
Psychoticism	1.239	0.8854	<i>P</i> <0.001
Somatization	0.945	0.8715	<i>P</i> <0.001

**Note:** Scores have been compared with norms for adult nonpatients.

**Table 2** Reliability statistics for the Brief Symptom Inventory (BSI) among nomophobic subjects

BSI subscale	Cronbach's alpha
Overall	0.972
Somatization	0.846
Obsession–compulsion	0.838
Interpersonal sensitivity	0.847
Depression	0.875
Anxiety	0.863
Hostility	0.810
Phobic anxiety	0.783
Paranoid ideation	0.808
Psychoticism	0.733

psychoticism (0.783 and 0.733, respectively), which were acceptable. Overall, the reliability of the entire instrument was excellent (alpha coefficient=0.972). Correlations among the BSI subscales were statistically significant (Table 3), ranging from *r*=0.568 (*P*<0.0001) for the relationship between hostility and phobic anxiety to *r*=0.810 (*P*<0.0001) between interpersonal sensitivity and paranoid ideation. Furthermore, all BSI subscales as well as the BSI synthetic indexes correlated with the NMP-Q in a statistically significant way (Table 4). Correlation coefficients ranged from *r*=0.115 (*P*=0.0208) to *r*=0.372 (*P*<0.0001) for the relationship between phobic anxiety and factor 1 (not being able to access information) and between the GSI score and factor 2 (giving up convenience/losing connectedness) of the NMP-Q, respectively.

In multivariate regression analysis, the factor 2 score (regression coefficient=0.02877, standard error=0.005494, *r*<sub>partial</sub>=0.2548, *t*=5.236, *P*<0.0001), the number of hours spent on the mobile device (regression coefficient=0.05288, standard error=0.02092, *r*<sub>partial</sub>=0.1262, *t*=2.528, *P*=0.0119),



**Table 3** Correlation analysis among the subscales of the Brief Symptom Inventory

		Anxiety	Depression	Hostility	Interpersonal sensitivity	Obsession-compulsion	Phobic anxiety	Paranoid ideation	Psychoticism
Depression	Correlation coefficient	0.736							
	Significance level <i>P</i>	<0.0001							
Hostility	Correlation coefficient	0.679	0.647						
	Significance level <i>P</i>	<0.0001	<0.0001						
Interpersonal sensitivity	Correlation coefficient	0.740	0.784	0.659					
	Significance level <i>P</i>	<0.0001	<0.0001	<0.0001					
Obsession-compulsion	Correlation coefficient	0.733	0.730	<0.0001	0.717				
	Significance level <i>P</i>	<0.0001	<0.0001	<0.0001	<0.0001				
Phobic anxiety	Correlation coefficient	0.736	0.639	0.568	0.689	0.651			
	Significance level <i>P</i>	<0.0001	<0.0001	<0.0001	<0.0001	<0.0001			
Paranoid ideation	Correlation coefficient	0.731	0.748	0.711	0.810	0.707	0.650		
	Significance level <i>P</i>	<0.0001	<0.0001	<0.0001	<0.0001	<0.0001	<0.0001		
Psychoticism	Correlation coefficient	0.724	0.786	0.646	0.767	0.728	0.723	0.778	
	Significance level <i>P</i>	<0.0001	<0.0001	<0.0001	<0.0001	<0.0001	<0.0001	<0.0001	
Somatization	Correlation coefficient	0.747	0.575	0.611	0.582	0.635	0.675	0.641	0.607
	Significance level <i>P</i>	<0.0001	<0.0001	<0.0001	<0.0001	<0.0001	<0.0001	<0.0001	<0.0001

and the schooling level (regression coefficient=-0.09865, standard error=0.04485,  $r_{\text{partial}}=-0.1100$ ,  $t=-2.200$ ,  $P=0.0284$ ) were statistically significant predictors of the GSI score. For further details, the reader is referred to Table 5. Table 6 reports the predictors for each subscale of the BSI.

Stratifying the population according to the severity of nomophobia (mild, 206 individuals, 51.1% of the sample; moderate, 167 subjects, 41.4%; and severe, 30 individuals, 7.4%), the GSI score could distinguish ( $P<0.001$ ) between mild and moderate ( $0.99\pm0.71$  vs  $1.32\pm0.81$ ) and between mild and severe ( $0.99\pm0.71$  vs  $1.54\pm0.79$ ) nomophobia levels, although it could not differentiate between moderate and severe nomophobia levels ( $P>0.05$ ; Figure 1). Similar patterns could be found for the other subscales of the BSI. These findings were also confirmed by the clustering analysis (Tables 7 and 8), which found two major clusters (162 subjects belonging to the first cluster and 241 individuals to the second). These two clusters approximately coincided with the groups of severe and moderate nomophobia levels (197 subjects) and the group of mild nomophobia level (207 individuals), respectively; 22.029 was the final distance between the two cluster centers.

In the EFA, the 9-factor model explained up to 72.84% of the variance. Factor loadings of the different subscales for different factor models (first-order 1-factor and 9-factor models and second-order 9-factor models) are shown in Table 9.

Finally, CFA confirmed the findings of the EFA. Looking at the fit indexes of the CFA, concerning the first-order factor models, the Derogatis 9-factor model best fit the data compared with the 1-factor model (Table 10). The second-order 9-factor model, however, proved to have a better fit to data. At the subscale level, the following subscales showed the best fit: somatization, obsessive-compulsive, and paranoid ideation.

## Discussion

Nomophobia is an emerging technological addiction or phobia. Personality and psychopathological traits/features could be major determinants of this disorder. The BSI could shed light on this topic, even though its reliability and factor model have not been investigated in nomophobic subjects. Moreover, among scholars it has been debated whether the factor structure of the BSI is unidimensional or multidimensional. Our results show that a second-order 9-factor model fits reasonably well with our data and that the BSI is a psychometrically sound instrument able to distinguish between nonclinical individuals and subjects with behavioral addictions/phobias, such as nomophobia, as shown both by the analysis of variance and clustering analyses. Moreover, the BSI scores correlate with the severity of such a disorder,

**Table 4** Correlation analysis between the Brief Symptom Inventory (BSI) and the Nomophobia questionnaire

BSI subscale		Factor 1 (not being able to access information)	Factor 2 (giving up convenience/losing connectedness)	Factor 3 (not being able to communicate)	Total score
Anxiety	Correlation coefficient	0.237	0.291	0.243	0.314
	Significance level P	<0.0001	<0.0001	<0.0001	<0.0001
Depression	Correlation coefficient	0.222	0.335	0.180	0.281
	Significance level P	<0.0001	<0.0001	0.0003	<0.0001
Hostility	Correlation coefficient	0.184	0.278	0.169	0.241
	Significance level P	0.0002	<0.0001	0.0006	<0.0001
Interpersonal sensitivity	Correlation coefficient	0.221	0.342	0.200	0.291
	Significance level P	<0.0001	<0.0001	0.0001	<0.0001
Obsession–compulsion	Correlation coefficient	0.234	0.344	0.216	0.303
	Significance level P	<0.0001	<0.0001	<0.0001	<0.0001
Phobic anxiety	Correlation coefficient	0.115	0.340	0.138	0.230
	Significance level P	0.0208	<0.0001	0.0057	<0.0001
Paranoid ideation	Correlation coefficient	0.184	0.319	0.184	0.263
	Significance level P	0.0002	<0.0001	0.0002	<0.0001
Psychoticism	Correlation coefficient	0.133	0.323	0.137	0.229
	Significance level P	0.0073	<0.0001	0.0058	<0.0001
Somatization	Correlation coefficient	0.163	0.302	0.194	0.253
	Significance level P	0.0010	<0.0001	0.0001	<0.0001

**Table 5** Multivariate regression analysis for the Global Severity Index of the Brief Symptom Inventory

Independent variable	Coefficient	Standard error	r <sub>partial</sub>	t	P-value
(Constant)	0.7209				
Factor 1 (not being able to access information)	-0.006034	0.006848	-0.04429	-0.881	0.3788
Factor 2 (giving up convenience/losing connectedness)	0.02877	0.005494	0.2548	5.236	<0.0001
Factor 3 (not being able to communicate)	-0.0003147	0.005557	-0.002850	-0.0566	0.9549
Hours	0.05288	0.02092	0.1262	2.528	0.0119
Schooling level	-0.09865	0.04485	-0.1100	-2.200	0.0284
Age	0.001618	0.004620	0.01762	0.350	0.7264
Gender	0.09083	0.07656	0.05959	1.186	0.2362


**Table 6** Predictors for each subscale of the Brief Symptom Inventory

Construct	Predictor(s)
Anxiety	Factor 2 (giving up convenience/losing connectedness), hours, gender
Depression	Factor 2 (giving up convenience/losing connectedness), hours
Hostility	Factor 2 (giving up convenience/losing connectedness)
Interpersonal sensitivity	Factor 2 (giving up convenience/losing connectedness), hours, schooling level, gender
Obsession–compulsion	Factor 2 (giving up convenience/losing connectedness), schooling level
Phobic anxiety	Factor 1 (negative; not being able to access information), factor 2 (giving up convenience/losing connectedness)
Paranoid ideation	Factor 2 (giving up convenience/losing connectedness)
Psychoticism	Factor 2 (giving up convenience/losing connectedness), schooling level
Somatization	Factor 2 (giving up convenience/losing connectedness), hours, gender

further confirming and corroborating the discriminant validity of the instrument.

In the existing scholarly literature, different factor structure models, including structures comprised of five factors (among bereaved patients),<sup>23</sup> six factors (among ethnic groups,

either clinical or nonclinical, or among college and university counseling center clients),<sup>24,25</sup> eight factors (for instance, in subjects with distress),<sup>26,27</sup> and one single factor of general distress (found among patients suffering from epilepsy or adult inpatients with psychiatric disorders),<sup>28–30</sup> have been reported.



**Figure 1** General Severity Index (GSI) broken down according to the severity of nomophobia.

**Table 7** The two clusters with their centers for each Brief Symptom Inventory (BSI) subscale

BSI subscale	Cluster	
	1	2
Somatization	11.57	3.28
Obsession–compulsion	13.06	4.93
Interpersonal sensitivity	9.61	2.66
Depression	13.65	4.54
Anxiety	12.62	4.31
Hostility	9.79	3.77
Phobic anxiety	6.07	0.98
Paranoid ideation	10.93	3.92
Psychoticism	9.06	2.83

However, in most cases, these factor structure models are the results of EFA-based instead of CFA-based investigations. Only few studies, indeed, performed CFA.<sup>31</sup> Furthermore, some methodological differences among studies could explain the different models obtained, including the study design and the populations recruited. Our results are, instead, methodologically more robust, relying on CFA and not on only EFA. Moreover, we also performed a clustering analysis to further corroborate our findings.

The findings of the present investigation are in line with our working hypothesis and the existing scholarly literature. Being out of mobile phone contact, for example being unable to access a mobile device, can lead to an increase in irritability and anxiety. Subjects with technological addiction indeed make unsuccessful attempts to decrease Internet use and to

prevent, or at least mitigate, its negative impact on social, work, and/or academic life, as shown in a sample of 255 university students.<sup>32</sup> Somatization is one of the markers of anxiety: it can be defined as a psychological distress arising from the perception of bodily dysfunction with a strong autonomic component. It is characterized by pain and discomfort, involving different systems, including the cardiovascular, gastrointestinal, respiratory, and muscular systems, among others. A correlation between somatization and Internet use has been found by some scholars.<sup>33,34</sup> Depressive symptoms reflect a dysphoric mood, characterized by loss of interest in daily activities and by a deep feeling of hopelessness and despair. Some studies have found a correlation between depression and Internet use: together with low self-esteem, self-efficacy, and life satisfaction, it is clinically associated with higher levels of technological addiction.<sup>35,36</sup>

Furthermore, other symptoms have been correlated with Internet use, such as obsessive–compulsive symptoms, arising when some thoughts and/or behaviors occur so intrusively as to be perceived as unremitting and irresistible; interpersonal sensitivity, which is represented by feelings of personal inadequacy and inferiority, with scarce social life and interactions; or hostility, characterized by feelings of irritability, urges to break or smash things, and uncontrollable outbursts of temper. Also phobic anxiety, which reflects phobic fears and worries, and psychoticism have been associated with technological addiction.<sup>34–37</sup>

We found that paranoid ideation, which refers to a peculiar mode of thinking dominated by projection, suspiciousness,

**Table 8** Analysis of variance between the two clusters for each Brief Symptom Inventory (BSI) subscale

BSI subscale	Cluster		Error		F	P-value
	Mean square	Df	Mean square	Df		
Somatization	6660.960	1	20.699	401	321.796	<0.001
Obsession–compulsion	6397.213	1	15.617	401	409.639	<0.001
Interpersonal sensitivity	4681.301	1	7.977	401	586.882	<0.001
Depression	8030.578	1	15.742	401	510.121	<0.001
Anxiety	6683.736	1	14.658	401	455.973	<0.001
Hostility	3513.816	1	11.990	401	293.071	<0.001
Phobic anxiety	2508.603	1	8.138	401	308.276	<0.001
Paranoid ideation	4767.526	1	10.655	401	447.451	<0.001
Psychoticism	3767.381	1	9.526	401	395.470	<0.001

**Abbreviation:** Df, degrees of freedom.

**Table 9** Loading factors of the exploratory factor analysis

BSI subscale	Factor loading (second-order 9-factor)	Items	Factor loading (first-order 9-factor)	Factor loading (1-factor)
Anxiety	0.890	1	0.674	0.611
		12	0.819	0.713
		19	0.846	0.735
		38	0.773	0.711
		45	0.824	0.736
Depression	0.868	49	0.671	0.649
		9	0.638	0.579
		16	0.803	0.690
		17	0.834	0.725
		18	0.793	0.661
Hostility	0.802	35	0.807	0.699
		50	0.822	0.751
		6	0.666	0.643
		13	0.790	0.644
		40	0.805	0.577
Interpersonal sensitivity	0.882	41	0.794	0.584
		46	0.709	0.553
		20	0.842	0.731
		21	0.838	0.693
		22	0.834	0.718
Obsession–compulsion	0.855	42	0.799	0.757
		15	0.801	0.709
		26	0.700	0.630
		27	0.814	0.657
		32	0.733	0.683
Paranoid ideation	0.885	36	0.821	0.691
		4	0.686	0.584
		10	0.719	0.607
		24	0.803	0.735
		48	0.773	0.716
		51	0.775	0.641

(Continued)

**Table 9 (Continued)**

BSI subscale	Factor loading (second-order 9-factor)	Items	Factor loading (first-order 9-factor)	Factor loading (1-factor)
Phobic anxiety	0.823	8	0.763	0.518
		28	0.751	0.463
		31	0.794	0.663
		43	0.785	0.714
		47	0.615	0.612
Psychoticism	0.883	3	0.624	0.475
		14	0.736	0.727
		34	0.512	0.404
		44	0.812	0.703
		53	0.801	0.720
Somatization	0.786	2	0.665	0.493
		7	0.674	0.546
		23	0.681	0.540
		29	0.758	0.634
		30	0.755	0.606
		33	0.717	0.525
		37	0.795	0.670

**Abbreviation:** BSI, Brief Symptom Inventory

persecutory and conspiracy beliefs, and fear of loss of control, was associated with nomophobia. Also, this finding is in line with the literature.<sup>38</sup>

Summarizing, according to Taylor et al,<sup>39</sup> who make use of cognitive-behavioral models and social-skills theory, there is a strong relationship between depression and time spent using the Internet, whereas more mixed findings are reported for social anxiety. Loneliness and hostility were also found to correlate with Internet use. We have extended these results to an emerging disorder, nomophobia.

On the other hand, despite its novelty, the present study is not without limitations. The major shortcoming that should be



**Table 10** Fit indices of the confirmatory factor analysis for the overall Brief Symptom Inventory (BSI) questionnaire and for each subscale

Model	CFI	TLI	SRMR	RMSEA	RMSEA 90% CI		AIC	BIC	Exact fit		
					Lower	Upper			$\chi^2$	Df	P
<b>First-order models</b>											
9-factor	0.820	0.806	0.058	0.071	0.068	0.073	53.829	54.561	3.279	1.091	<0.001
1-factor	0.747	0.736	0.063	0.082	0.079	0.085	54.581	54.973	4.186	1.127	<0.001
<b>Second-order models</b>											
9-factor	0.851	0.833	0.047	0.063	0.060	0.065	57.837	58.633	3.188	1.232	<0.001
<b>BSI subscales</b>											
Somatization	0.964	0.946	0.033	0.077	0.054	0.102	8.182	8.266	47.6	14	<0.001
Obsession–compulsion	0.982	0.970	0.025	0.065	0.034	0.097	7.141	7.213	24.2	9	0.004
Interpersonal sensitivity	0.960	0.879	0.031	0.184	0.129	0.246	4.871	4.919	29.3	2	<0.001
Depression	0.928	0.881	0.045	0.152	0.125	0.181	6.880	6.952	92.6	9	<0.001
Anxiety	0.959	0.931	0.036	0.109	0.082	0.139	6.785	6.857	52.3	9	<0.001
Hostility	0.925	0.851	0.048	0.156	0.120	0.195	5.888	5.948	54.0	5	<0.001
Phobic anxiety	0.962	0.925	0.034	0.103	0.067	0.143	5.186	5.246	26.4	5	<0.001
Paranoid ideation	0.980	0.960	0.025	0.077	0.038	0.119	6.078	6.138	16.9	5	0.005
Psychoticism	0.948	0.896	0.044	0.109	0.073	0.149	6.093	6.153	29.0	5	<0.001

**Abbreviations:** AIC, Akaike Information Criterion; BIC, Bayesian Information Criterion; CFI, Comparative Fit Index; Df, degrees of freedom; RMSEA, root mean square error of approximation; SRMR, standardized root mean square residual; TLI, Tucker–Lewis Index.

properly recognized is given by the nonrandomized nature of the recruited sample (snowball sampling procedure). Another drawback is represented by the cross-sectional design of the study. High-quality longitudinal studies should be performed in order to capture the dynamic picture of the relationship between nomophobia and psychopathological symptoms.

## Conclusion

The findings of our study show that the BSI is a reliable instrument with acceptable psychometric properties that can be administered to populations of nomophobic subjects and, as such, can be exploited by researchers in the field of behavioral addictions and technological phobias. However, based on the abovementioned shortcomings, further research in the field is urgently needed.

## Disclosure

The authors report no conflicts of interest in this work.

## References

- Derogatis LR. *Brief Symptom Inventory*. Baltimore, MD: Clinical Psychometric Research; 1975.
- Derogatis LR, Spencer PM. *Administration and Procedures: BSI. Manual I*. Baltimore, MD: Clinical Psychometric Research; 1982.
- Derogatis LR. *The Brief Symptom Inventory (BSI). Administration, Scoring and Procedures Manual II*. Baltimore, MD: Clinical Psychometric Research; 1992.
- Derogatis LR. *Brief Symptom Inventory: Administration, Scoring and Procedures Manual*. 4th ed. Minneapolis, MN: NCS, Pearson, Inc; 1993.
- Derogatis LR, Melisaratos N. The Brief Symptom Inventory: an introductory report. *Psychol Med*. 1983;13(3):595–605.
- Racine NM, Khu M, Reynolds K, Guilcher GMT, Schulte FSM. Quality of life in pediatric cancer survivors: contributions of parental distress and psychosocial family risk. *Curr Oncol*. 2018;25(1):41–48.
- Barbosa F, Mota C, Patrício P, Alcântara C, Ferreira C, Barbosa A. The relationship between alexithymia and psychological factors in systemic lupus erythematosus. *Compr Psychiatry*. 2011;52(6):754–762.
- Bragazzi NL, Dini G, Toletone A, Brigo F, Durando P. Leveraging big data for exploring occupational diseases-related interest at the level of scientific community, media coverage and novel data streams: the example of silicosis as a pilot study. *PLoS One*. 2016;11(11):e0166051.
- Bragazzi N, del Puente G. A proposal for including nomophobia in the new DSM-V. *PRBM*. 2014;7:155–160.
- Yildirim C, Correia A-P. Exploring the dimensions of nomophobia: development and validation of a self-reported questionnaire. *Comput Human Behav*. 2015;49:130–137.
- Adawi M, Bragazzi NL, Argumosa-Villar L, et al. Translation and validation of the Nomophobia questionnaire in the Italian language: exploratory factor analysis. *JMIR Mhealth Uhealth*. 2018;6(1):e24.
- Adalier A, Balkan E. The relationship between Internet addiction and psychological symptoms. *International Journal of Global Education*. 2012;1(2):42–49.
- Wegmann E, Stodt B, Brand M. Addictive use of social networking sites can be explained by the interaction of Internet use expectancies, Internet literacy, and psychopathological symptoms. *J Behav Addict*. 2015;4(3):155–162.
- Stavropoulos V, Gomez R, Steen E, Beard C, Liew L, Griffiths MD. The longitudinal association between anxiety and Internet addiction in adolescence: the moderating effect of classroom extraversion. *J Behav Addict*. 2017;6(2):237–247.
- Tabachnick BG, Fidell LS. *Using Multivariate Statistics*. New York: Harper & Row; 1996.
- González-Cabrera J, León-Mejía A, Pérez-Sancho C, Calvete E. Adaptation of the Nomophobia questionnaire (NMP-Q) to Spanish in a sample of adolescents. *Actas Esp Psiquiatr*. 2017;45(4):137–144.
- Lin CY, Griffiths MD, Pakpour AH. Psychometric evaluation of Persian Nomophobia questionnaire: differential item functioning and measurement invariance across gender. *J Behav Addict*. 2018;7(1):100–108.
- Hinkle DE, Wiersma W, Jurs SG. *Applied Statistics for the Behavioral Sciences*. 5th ed. Boston: Houghton Mifflin; 2003.

19. Maccallum RC, Browne MW, Sugawara HM. Power analysis and determination of sample size for covariance structure modeling. *Psychol Methods*. 1996;1(2):130–149.
20. Steiger JH. Point estimation, hypothesis testing, and interval estimation using the RMSEA: some comments and a reply to Hayduk and Glaser. *Struct Equ Modeling*. 2000;7(2):149–162.
21. Hu LT, Bentler PM. Evaluating model fit. In: Hoyle RH. editor. *Structural Equation Modeling: Concepts, Issues, and Applications*. Thousand Oaks, CA: Sage; 1995:76–99.
22. Bentler PM. Comparative fit indexes in structural models. *Psychol Bull*. 1990;107(2):238–246.
23. Johnson LC, Murphy SA, Dimond M. Reliability, construct validity, and subscale norms of the Brief Symptom Inventory when administered to bereaved parents. *J Nurs Meas*. 1996;4(2):117–127.
24. Daoud FS, Abojedi AA. Equivalent factorial structure of the Brief Symptom Inventory (BSI) in clinical and nonclinical Jordanian populations. *Eur J Psychol Assess*. 2010;26(2):116–121.
25. Hayes JA. What does the Brief Symptom Inventory measure in college and university counseling center clients? *J Couns Psychol*. 1997;44(4):360–367.
26. Ruipérez MA, Ibáñez MI, Lorente E, Moro M, Ortet G. Psychometric properties of the Spanish version of the BSI: contributions to the relationship between personality and psychopathology. *Eur J Psychol Assess*. 2001;3:241–250.
27. Kellett S, Beail N, Newman DW, Frankish P. Utility of the Brief Symptom Inventory in the assessment of psychological distress. *J Appl Res Intellect Disabil*. 2003;16(2):127–134.
28. Endermann M. The Brief Symptom Inventory (BSI) as a screening tool for psychological disorders in patients with epilepsy and mild intellectual disabilities in residential care. *Epilepsy Behav*. 2005;7(1):85–94.
29. Piersma HL, Reaume WM, Boes JL. The Brief Symptom Inventory (BSI) as an outcome measure for adult psychiatric inpatients. *J Clin Psychol*. 1994;50(4):555–563.
30. Piersma HL, Boes JL, Reaume WM. Unidimensionality of the Brief Symptom Inventory (BSI) in adult and adolescent inpatients. *J Pers Assess*. 1994;63(2):338–344.
31. Loutsiou-Ladd A, Panayiotou G, Kokkinos CM. A review of the factorial structure of the Brief Symptom Inventory (BSI): Greek evidence. *Int J Test*. 2008;8(1):90–110.
32. Singh K, Brown RJ. Health-related Internet habits and health anxiety in university students. *Anxiety Stress Coping*. 2014;27(5):542–554.
33. Alavi SS, Maracy MR, Jannatifard F, Eslami M. The effect of psychiatric symptoms on the Internet addiction disorder in Isfahan's university students. *J Res Med Sci*. 2011;16(6):793–800.
34. Alavi SS, Alaghemandan H, Maracy MR, Jannatifard F, Eslami M, Ferdosi M. Impact of addiction to Internet on a number of psychiatric symptoms in students of Isfahan universities, Iran, 2010. *Int J Prev Med*. 2012;3(2):122–127.
35. Xiuqin H, Huimin Z, Mengchen L, Jinan W, Ying Z, Ran T. Mental health, personality, and parental rearing styles of adolescents with Internet addiction disorder. *Cyberpsychol Behav Soc Netw*. 2010;13(4):401–406.
36. Senormancı O, Saraçlı O, Atasoy N, Senormancı G, Koktürk F, Atik L. Relationship of Internet addiction with cognitive style, personality, and depression in university students. *Compr Psychiatry*. 2014;55(6):1385–1390.
37. Lu WH, Lee KH, Ko CH, Hsiao RC, Hu HF, Yen CF. Relationship between borderline personality symptoms and Internet addiction: The mediating effects of mental health problems. *J Behav Addict*. 2017;6(3):434–441.
38. Chen Q, Quan X, Lu H, Fei P, Li M. Comparison of the personality and other psychological factors of students with Internet addiction who do and do not have associated social dysfunction. *Shanghai Arch Psychiatry*. 2015;27(1):36–41.
39. Taylor S, Pattara-Angkoon S, Sirirat S, Woods D. The theoretical underpinnings of Internet addiction and its association with psychopathology in adolescence. *Int J Adolesc Med Health*. Epub 2017 Jul 6.
40. Maiti SS, Mukherjee BN. A note on distributional properties of the Jöreskog–Sörbom fit indices. *Psychometrika*. 1990;55(4):721–726.

## Psychology Research and Behavior Management

### Publish your work in this journal

Psychology Research and Behavior Management is an international, peer-reviewed, open access journal focusing on the science of psychology and its application in behavior management to develop improved outcomes in the clinical, educational, sports and business arenas. Specific topics covered in the journal include: Neuroscience, memory and decision making; Behavior

modification and management; Clinical applications; Business and sports performance management; Social and developmental studies; Animal studies. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/psychology-research-and-behavior-management-journal>

Dovepress