The role of concordant and discordant comorbidities on performance of self-care behaviors in adults with type 2 diabetes: a systematic review

This article was published in the following Dove Press journal:
Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy

Fekadu Aga¹
Sandra B Dunbar²
Tedla Kebede³
Rebecca A Gary²

¹Department of Nursing, School of Nursing & Midwifery, College of Health Sciences, Addis Ababa University, Addis Ababa, Ethiopia; ²Nell Hodgson Woodruff School of Nursing, Emory University, Atlanta, GA 30322, USA;

³Tikur Anbessa Specialized Hospital, Diabetes & Endocrinology Unit, Department of Internal Medicine School of Medicine, College of Health Sciences, Addis Ababa University, Addis Ababa, Ethiopia

Background: Most patients with Type 2 diabetes (T2D) have high number of comorbid chronic conditions that can affect their self-care abilities. Guidelines for diabetes self-care behaviors are disease specific with little attention given to managing T2D with other comorbidities. Identifying comorbidities that either improve or potentially diminish the individual's capacity to perform effective self-care behaviors is essential to enhance clinical outcomes. One such framework conceptualizes comorbidities as concordant or discordant with diabetes pathophysiological pathways and care goals.

Objective: The purpose of this systematic review was to examine the role of diabetes-concordant and discordant chronic conditions on the performance of self-care behaviors in adults with T2D.

Methods: A comprehensive literature search was undertaken to identify published English language articles through the following five electronic databases: PubMed, CINAHL, PsycINFO, ISI Web of Science, and EMBASE. Quantitative studies published from March 2006 to April 2018 were included. Quality of evidence was evaluated using the Joanna Briggs Institutes Critical Appraisal Tools (JBI-CAT) and rated using Quality Assessment Tool for Quantitative Studies (QATQS).

Results: The initial database search identified 1,136 articles but only 33 studies that met the inclusion criteria were included. The most common concordant comorbidity was hypertension while depression was the most common discordant condition. Adherence to medications was the most frequent diabetes self-care behavior reported and tended to be higher among concordant comorbidities. The findings showed mixed results concerning the effect of some concordant comorbidities such as hypertension, hyperlipidemia, retinopathy, and heart failure on diabetes self-care behaviors. But, there is agreement across studies that diabetes-discordant comorbidities have a more detrimental effect on self-care behaviors.

Conclusions: Concordant comorbidities may improve diabetes self-care, but the evidence is inconclusive. Future research using well designed studies are needed to examine the complex relationship between diabetes self-care and comorbidities.

Keywords: comorbidity, concordant comorbidity, diabetes self-care behaviors, discordant comorbidity, multimorbidity, type 2 diabetes

Correspondence: Fekadu Aga
Department of Nursing, School of Nursing & Midwifery, College of Health Sciences, Addis Ababa University, P.O. Box 9083, Addis Ababa
Tel +125 191 103 3684
Email fiqaaduagaa@yahoo.com

Introduction

Diabetes mellitus (DM) is one of the largest global health challenges of the 21st century as the number of cases continues to rise.¹ The epidemic is mainly from type

2 diabetes mellitus (T2D) that is driven by genetic, epigenetic, environmental, and behavioral factors such as sedentary lifestyle, consumption of energy dense diet and obesity.^{1–3} According to the International Diabetes Federation (IDF), about 425 million people between 20 and 79 years of age lived with DM in 2017 worldwide.⁴ The global age-standardized diabetes prevalence is 9.0% in men and 7.9% in women though there is great regional variation.⁵ More than 90% of all DM cases are T2D which is continuously increasing.^{4,6}

Patients with T2D have a higher number of comorbid conditions, also referred to as multimorbidity, which may result in treatment conflict,⁷ reduced glycemic control,^{8,9} increased cost of treatment,^{10,11} and lower quality of life.¹² Multiple studies in T2D have demonstrated multimorbidity prevalence rate as high as 97.5%.^{7,8,10,13–16} Studies show that low socioeconomic status, obesity and job strain,¹⁷ age, race/ethnicity,¹⁸ and low health literacy¹⁹ are associated with increased prevalence of comorbidity in T2D. Furthermore, about 50% of T2D patients face premature mortality from cardiovascular comorbidities such as coronary artery disease, heart failure and stroke, and approximately 10% from renal failure.²⁰

The successful management of T2D requires a patient's engagement in self-care on a daily basis to control its progression and to prevent serious complications, reduce rates of hospitalization and cost of care, and improve quality of life.^{21–24} Diabetes self-care involves sustained engagement in multidimensional behavioral actions including healthy dietary intake, physical activity, blood glucose monitoring, medication adherence, problem solving, and risk reduction such as foot checks and smoking cessation.^{25,26} Antecedent factors such as demographic, psychosocial, clinical, and environmental variables can influence the patient's successful performance of diabetes self-care.^{26–28} The influence of these factors may become even more important when there is another serious comorbid condition or if multimorbidity exists.

There is a need for further research to understand the impact of comorbidities on optimal diabetes self-care in order to maximize self-care for patients with T2D and multimorbidity.²⁹ Guidelines currently are primarily disease-specific and provide little direction for self-care for T2D in the presence of other chronic conditions.³⁰ Interventions that target self-care for T2D in the presence of multimorbidity are also limited. Understanding the relationship between diabetes self-care and other chronic conditions may lead to more effective interventions for this complex population.^{31–33}

A framework developed by Piette and Kerr³⁴ that conceptualizes chronic conditions as concordant or discordant with diabetes care has gained momentum over the last decade to help guide management of T2D and coexisting multimorbidity. Diabetes-concordant comorbidities (such as cardiovascular, metabolic and renal disease) share parts of the same overall pathophysiologic profile and care management as T2D. For example, blood pressure and lipid monitoring are overlapping cardiovascular risk reduction goals for diabetes and cardiovascular disease and likely to lead to better outcomes for both conditions. Magnan and colleagues³⁵ found that having more concordant conditions was linked to better quality of care for diabetes.

Diabetes-discordant comorbidities have no direct relationship with diabetes in their pathophysiologic profile or care management and include chronic conditions such as asthma, cancer and mental illness such as depressive symptoms.^{35–40} Treatment for some of these discordant conditions might interfere with treatment for T2D. For example, using steroids for asthma treatment would increase blood glucose and conflict with the treatment for diabetes while depressive symptoms have been shown to lead to poorer adherence to self-care in chronic conditions, including T2D. Time limitations, conflicting care goals, and other challenges may result in a lower quality of care for diabetes and the discordant condition as compared to having concordance or a single condition. Using a Delphi study method, Magnan and colleagues³⁸ identified 12 concordant and 50 discordant comorbidities in DM. The findings from this study suggested that concordant chronic conditions are more easily managed because care goals are similar and more likely to be identified and addressed by care providers. However, Magnan and colleagues³⁵ noted that the presence of more dominant concordant conditions such as advanced heart failure may reduce diabetes self-care behaviors; others have not shown this effect. Discordant conditions are less likely to be identified and included in care goals which may contribute to poorer diabetes control, additional complications, and higher health care utilization.

The role that the concordance-discordance framework may have in identifying key determinants or challenges in diabetes self-care has received little attention in the literature. Awareness of obstacles faced by patients with T2D who are managing multiple chronic conditions will help inform future strategies for more effective diabetes

self-care. The purpose of this systematic review was to examine the role of diabetes-concordant and discordant chronic conditions on the performance of self-care behaviors in adults with T2D.

Methods

The process of conducting and reporting this systematic review was guided by the Preferred Reporting Items for Systematic Reviews and Meta-Analysis (PRISMA) statements.^{41,42} The PICO (Participant-Intervention/Exposure-Comparison-Outcome) format was used to guide the research question and to establish inclusion and exclusion criteria for articles in this review.

Search strategy

The construction of the search strategy in this review was based on the guideline developed by the Joanna Briggs Institute (JBI).⁴³ A total of 68 search terms were used. Examples include: “Type 2 diabetes,” “adult,” “comorbidity,” “concordant comorbidity,” “discordant comorbidity,” “multi-morbidity,” “myocardial infarction,” “cerebrovascular disease,” “heart failure,” “chronic renal failure,” “asthma,” “depression,” “self-care behavior,” “self-care,” “diet,” “physical activity,” “foot care,” and “medication adherence.” Medical subject heading (MeSH) and free text words were used for searching the databases. Truncation and Boolean operators – AND/OR – were used to combine the different search terms. Finally, the search was refined by publication date (March 2006 to April 2018), English language, and study design (cross-sectional survey, case-control, cohort, non-randomized clinical trial, and randomized clinical trial).

Studies were identified by searching the following electronic databases: PubMed, CINAHL, PsycINFO, ISI Web of Science, and EMBASE. Further search of Google Scholar, Cochrane Library, and websites of pertinent organizations such as World Health Organization (WHO), International Diabetes Federation (IDF), and American Diabetes Association did not yield one additional study. However, scanning of the reference lists of the retrieved articles provided two additional studies. The literature search was grouped into the following comorbidity categories: concordant comorbid conditions only; discordant comorbid conditions only; and both concordant and discordant comorbid conditions.

Inclusion and exclusion criteria

Only quantitative studies that examined the relationship between concordant and discordant comorbid conditions and diabetes self-care behaviors in adults (≥ 18 years) with

T2D were included in this review. Studies that addressed either overall diabetes self-care behavior or specific self-care behaviors such as diet, medication, blood glucose self-monitoring, exercise, foot care, and smoking as an outcome variable were included. Studies not available in English, conference abstracts and books were not included in the review. The time frame for selection of articles was informed by the publication of a conceptual framework for understanding diabetes care in the context of concordant and discordant comorbidities.³⁴

Study selection

All articles retrieved from the search process were saved into the Endnote citation manager. Duplicates were removed, and the remaining records were screened for study inclusion. Eligibility was assessed independently in an unblinded standardized manner by 2 reviewers. Disagreements between reviewers were resolved by consensus. A two-stage screening process was undertaken using a checklist we developed for this purpose. First, the titles and abstracts of the articles were screened for relevance in relation to the PICO question. In the second round, full-text screening was performed. After screening, the studies were assessed for methodological quality in a two-step process. In the first step, the Joanna Briggs Institute (JBI) Critical Appraisal Tools⁴⁴ was used to determine to what extent a study had addressed possible biases and included the PICO of interest in this review. In the second step, the remaining studies were rated as strong, moderate, or weak using the Quality Assessment Tool for Quantitative Studies⁴⁵ in six domains: selection bias, design, confounders, blinding, data collection methods and withdrawals and dropouts. The overall rating for the study was decided by assessing the six domain ratings. Those with no weak ratings and at least four strong ratings were considered strong evidence. Studies with less than four strong ratings and one weak rating were considered moderate while those with two or more weak ratings were considered weak evidence.

Data collection

Data were extracted using a standardized grid format that followed recommendations from the Cochrane Review⁴⁶ and the Joanna Briggs Institute.⁴⁷ Information related to data source, setting, study aim, design, sample characteristics, comorbidities types, outcome measurement, and results were extracted from individual studies.

Data analysis and synthesis

A narrative synthesis method was used in this systematic review because the heterogeneity observed in the study designs and methods of the selected studies did not allow the application of pooled analytical meta-analysis procedures. A narrative synthesis organizes and summarizes studies using words to explain the findings.⁴⁸ In this review, the use of a narrative approach facilitated description of concordant, discordant or the presence of both conditions in T2DM which helped to compare the impact of each on diabetes self-care.

Result

The initial database search identified 1,136 articles. However, only 33 studies met the inclusion criteria and were included in this review. The number of studies excluded at each stage of duplication removal, screening, and eligibility assessment process is shown in Figure 1.

Study characteristics

Table 1 provides a summary of the 33 studies meeting eligibility criteria and included in the systematic review.

Most studies included T2D self-care with discordant comorbidities (n=23) compared to self-care with concordant chronic conditions (n=6). Four studies included T2D self-care with both concordant and discordant chronic conditions. Twenty one studies were cross-sectional,^{49–69} eight were cohort studies,^{70–77} two were case-control studies,^{78,79} one was non-randomized intervention,⁸⁰ and one was randomized controlled trial.⁸¹ Seventeen studies were conducted in the United States (US), four in Canada, two studies each in Malaysia, China and Ireland, and one study each in the Netherlands, Australia, Belgium, Romania, and Jordan. The sample size of the studies ranged from 64 to 740,197 with a median of 615. The studies reported the mean age of the participants ranging from 47.7 to 70.8 years. One study was rated as weak, 27 as moderate, and five as strong.

Twenty studies measured generic or overall diabetes self-care behavior as an outcome. Specific diabetes self-care behaviors were also measured as an outcome and included: medication adherence (N=9), exercise behavior (N=4), dietary behavior (N=2), foot care (N=1), blood glucose monitoring (N=1), and smoking behavior (N=1).

Figure 1 PRISMA flow chart.

Table 1 Summary of the characteristics of studies included in the review

Reference	Setting/ Country	Study aim(s)	Design	Sample characteristics	Outcome of Interest	Result	Concordant/ Discordant
Malik et al ⁶²	6 primary care clinics in USA	To examine the relationship between patient complexity regimen intensity, problems with adherence to medications and control of cardiovascular risk factors among type 2 diabetes patients with and without coronary heart disease (CHD).	Cross-sectional	Age: <ul style="list-style-type: none"> Mean=58.8±11.4 years Sample size: 1,314 Gender: <ul style="list-style-type: none"> Men=40.6% Women=59.4% 	Outcome(s): <ul style="list-style-type: none"> Medication adherence Tool(s): A 13-item measure	<ul style="list-style-type: none"> T2D patients with a history of CHD were 70% less likely to report medication nonadherence (95% CI=0.5,1.0, p=0.04) compared to those without history of CHD. 	Concordant
He et al ⁷³	Tianjin, China	To assess adherence and persistence to insulin therapy and identify its associated factors among Chinese insulin-naïve patients with T2D.	Cohort	Age: <ul style="list-style-type: none"> Mean=58.9±11.6 years Sample size: 24,192 Gender: <ul style="list-style-type: none"> Men=50.5% Women=49.5% 	Outcome(s): <ul style="list-style-type: none"> Medication adherence and persistence Tool(s): Medication Possession Ratio (MPR) Patient had no ≥90 days in insulin therapy	<ul style="list-style-type: none"> About 30.9% (N=7,486) of the total cohort were adherent with a MPR of ≥0.8 in follow-up. Patients with a higher comorbidity index (OR=0.93, 95%CI=0.91, 0.96, p<0.001), or having had comorbid hypertension (OR=0.47, 95% CI=0.44, 0.50, p<0.001) and dyslipidemia (OR=0.74, 95%CI=0.68, 0.80, p<0.001), had a lower likelihood of being adherent. Having comorbid neuropathy (OR=1.33, 95% CI=1.24, 1.43, p<0.001) or nephropathy (OR=1.40, 95%CI=1.30, 1.52, p<0.001) was related to a significantly higher likelihood of adherence. The likelihood of being persistent with medication adherence was higher for patients with comorbid cardiovascular complications, including neuropathy and nephropathy (all HR<1, p<0.001). The likelihood of being persistent was lower for those with comorbid hypertension (adjusted OR=1.80, 95%CI=1.71, 1.88, p<0.001) or dyslipidemia (adjusted HR=1.19, 95% CI=1.14, 1.25, p<0.001). 	Concordant

(Continued)

Table 1 (Continued).

Reference	Setting/ Country	Study aim(s)	Design	Sample characteristics	Outcome of Interest	Result	Concordant/ Discordant
Ahmad et al ⁴⁹	Primary health clinics in Hulu Langat, Selangor, Malaysia	To assess adherence to medications and to identify factors that are associated with non-adherence in type 2 diabetes mellitus patients	Cross-sectional	Age: • Mean=55.95 ±9.13 years • Range=30–82 years Sample size=557 Gender: • Men= 36.8% • Women=63.2%	Outcome(s): • Medication adherence Tool(s): • Medication Compliance Questionnaire (MCQ)	<ul style="list-style-type: none"> • 241 (50.7%) of patients with comorbidities were nonadherent • T2D patients with comorbid hypertension, dyslipidemia, or both were 1.78 (95% CI=1.064, 2.981) times more likely to be nonadherent compared to those without comorbid condition. 	Concordant
Al-Sayah et al ⁵¹	Alberta, Canada	To examine the prevalence and predictors of foot disease, self-care and clinical monitoring in adults with type 2 diabetes in Alberta, Canada	Cross-sectional	Age: • Mean=64.4±10.7 years Sample size=2,040 Gender: • Men=55% • Women=45%	Outcome(s): • Foot care behavior Tool(s): • 4 items from SDSCA	<ul style="list-style-type: none"> • Average foot self-care was 3.9 days per week for the entire sample • 14% of participants performed the recommended foot self-care for ≥6 days per week • The odds of poor foot self-care was 38% (OR=0.62, 95% CI=0.50, 0.80) less in patients with comorbid hyperlipidemia compared to those without hyperlipidemia. 	Concordant
Dixon et al ⁷⁸	Australia	To investigate whether diabetes self-care attitudes, behaviors and perceived burden, particularly related to weight management, diet and physical activity, differ between adults with T2D who are severely obese and matched non-severely obese control subjects.	Case-control	Age: • Mean=58.4±8.9 years Sample size=1,795 Gender: • Men=51.6% • Women=48.4%	Outcome(s): • Overall DSCB Tool(s): • Diabetes Self-Care Inventory-Revised (DSCI-R)	<ul style="list-style-type: none"> • 530 (29.5%) were severely obese (BMI ≥35 kg/m²) and the proportion of class III (BMI ≥40 kg/m²) was greater (13.5%) than the proportion of those in the healthy range (BMI ≤ 25 kg/m², 12.2%). • The group with BMI ≥35 kg/m² was less likely to achieve diet and exercise recommendations, placed less importance on diet and exercise recommendations, and found the burden of diet and exercise recommendations to be greater than the group with BMI <35 kg/m² (p<0.001). • Moderate-to-severe symptoms of depression were independently associated with reduced likelihood of healthy diet and physical activity, and with greater burden associated with diet, physical activity, and weight management (p<0.001). 	Concordant

(Continued)

Table 1 (Continued).

Reference	Setting/ Country	Study aim(s)	Design	Sample characteristics	Outcome of Interest	Result	Concordant/ Discordant
Kroese et al ⁸⁰	the Netherlands	To compare obese and non-obese type 2 diabetes patients at baseline and after participating in an existing self-management intervention (i.e., Beyond Good Intentions) on cognitive, self-care, and behavioral measures to examine whether both groups are equally prepared and able to adopt self-management approaches.	Non-randomized intervention	Age: <ul style="list-style-type: none"> Mean=70.3±7.7 years Range=46–74 years Sample size=64 Gender: <ul style="list-style-type: none"> Men=59.4% Women=40.6% 	Outcome(s): <ul style="list-style-type: none"> Overall DSCB Tool(s): <ul style="list-style-type: none"> SDSCA 	<ul style="list-style-type: none"> At baseline obese T2D patients, compared to nonobese patient, reported lower adherence to self-care recommendations, $F_{(4, 58)}=2.77$, $p=0.036$, effect size (d)=0.16. Intervention showed significant and equal effectiveness both for obese and nonobese T2D patients on adherence to self-care recommendations, $F_{(1, 62)}=24.70$, $p=0.000$, $d=0.29$. 	Concordant
Asuzu et al ⁵²	Primary Clinics in southern United States	To examine the mechanism by which depressive symptoms, diabetes distress and diabetes fatalism together influence diabetes outcomes (self-care and HbA1c) using structured equation modeling.	Cross-sectional	Age: <ul style="list-style-type: none"> Mean=61.3±10.9 years Sample size=615 Gender: <ul style="list-style-type: none"> Men=61.6% Women=38.4% 	Outcome(s): <ul style="list-style-type: none"> Overall DSCB Glycemic control Tool(s): <ul style="list-style-type: none"> SDSCA Monisky Medication Adherence Scale (MMAS) HbA1c 	<ul style="list-style-type: none"> The most frequently reported specific self-care activity was medication adherence ($M=5.9\pm2.0$), followed by general diet ($M=4.7\pm2.0$), blood glucose testing ($M=4.6\pm2.5$), foot care ($M=4.3\pm2.5$), specific diet ($M=4.0\pm1.6$), and exercise ($M=2.6\pm2.2$). Diabetes distress directly significantly related to diabetes self-care ($r=-0.69$, $p<0.001$) and HbA1c ($r=0.69$, $p<0.001$). No significant direct relationship of depressive symptoms with diabetes self-care and HbA1c. Indirect relationship between increased depressive symptoms (mediated by diabetes distress) and decreased self-care ($r=-0.52$, $p<0.001$) and higher HbA1c ($r=0.60$, $p<0.001$). 	Discordant
Schinckus et al ⁶⁶	Belgium	To investigate whether emotional distress or depression moderates the relation between health literacy, self-efficacy and diabetes self-care behaviors.	Cross-sectional	Age: <ul style="list-style-type: none"> Mean=61.87 ±12.27 years Sample size=128 Gender: <ul style="list-style-type: none"> Men=52.4% Women=47.6% 	Outcome(s): <ul style="list-style-type: none"> Overall DSCB Tool(s): <ul style="list-style-type: none"> Diabetes Self-Management Questionnaire (DSMQ) 	<ul style="list-style-type: none"> Depression and diabetes-related distress do not independently predict diabetes self-care behaviors but they moderate the effect of health literacy on these behaviors. The interaction between depression and health literacy ($\beta=-0.322$, $t=-3.260$, $p\leq0.005$) and that between distress and health literacy ($\beta=-0.256$, $t=-2.490$, $p\leq0.05$) predict diabetes self-care behavior. 	Discordant

(Continued)

Table 1 (Continued).

Reference	Setting/ Country	Study aim(s)	Design	Sample characteristics	Outcome of Interest	Result	Concordant/ Discordant
Al-Amer et al ⁵⁰	Endocrinology Clinic, Jordan University Hospital	To assess the relationship between illness perception, depression, social support, religiosity and spiritual coping, self-efficacy, and diabetes self-care activities among T2D patients of an Arabic ethnicity	Cross-sectional	Age: <ul style="list-style-type: none"> Mean=58.2±10.8 years Range=32–82 years Sample size= 220 Gender: <ul style="list-style-type: none"> Men=52.3% Women=47.7% 	Outcome(s): <ul style="list-style-type: none"> Overall diabetes self-care behavior (DSCB) Tool(s): The Summary of Diabetes Self-Care Activities (SDSCA)	<ul style="list-style-type: none"> The mean score of overall self-care activities was 40.5±14.5. The most frequently reported self-care activity was medication adherence (M=6.5±1.4) and the least reported was exercise (M=1.8±2.0). The mean HbA1c value was 8.01±1.8% and 136 (61.8% of the participants had HbA1c value of ≥7%. Depression was indirectly associated to diabetes self-care activities through self-efficacy ($\beta=-0.20$, $p=0.003$). 	Discordant
Chasens et al ⁵³	Pittsburgh, USA	To investigate the association of impaired sleep quality and daytime sleepiness on self-reported diabetes control and psychological and social factors that affect diabetes self-management.	Cross-sectional	Age: <ul style="list-style-type: none"> Mean=52.23 ±9.28 years Range=31–83 years Sample size= 107 Gender: <ul style="list-style-type: none"> Men=42% Women=58% 	Outcome(s): <ul style="list-style-type: none"> Overall DSCB BMI Tool(s): Diabetes Care Profile (DCP) Kg/m ²	<ul style="list-style-type: none"> Participants had higher BMI (M=35.03±6.55 Kg/m²) and higher HbA1c (M=7.33±1.52%). Higher BMI associated with low scores on self-care adherence ($r=-0.42$, $p<0.004$) and self-care ability ($r=-0.24$, $p<0.014$). Impaired sleep quality and daytime sleepiness significantly associated with lower self-care adherence ($r=-0.21$, $p<0.05$). 	Discordant
Chen et al ⁵⁴	Pittsburgh, USA	To evaluate the effects of depression on the number of self-identified behavior change goals and the number of barriers to diabetes self-care among patients with type 2 diabetes.	Cross-sectional	Age: <ul style="list-style-type: none"> Mean=58 years Sample size= 778 Gender: <ul style="list-style-type: none"> Men=39.1% Women=60.9% 	Outcome(s): <ul style="list-style-type: none"> Overall DSCB Tool(s): Diabetes Self-Management Assessment Report Tool (D-SMART)	<ul style="list-style-type: none"> T2D patients with comorbid depression had greater number of barriers to self-care compared to nondepressed compatriot (7 vs 6, $p<0.001$). They also had higher BMI (Median=34.8 (q1-q2=30.9–40.1) vs 33.2 (28.8–39.1), $p=0.03$, but no difference in HbA1c values (Median=7.7 (q1-q2=6.8–9.3) vs 7.7 (6.7–9.3), $p=80$). 	Discordant

(Continued)

Table 1 (Continued).

Reference	Setting/ Country	Study aim(s)	Design	Sample characteristics	Outcome of Interest	Result	Concordant/ Discordant
Chen et al ⁵⁵	Taipei, China	To compare differences in self-efficacy scores and self-care behaviors between outpatients with comorbid schizophrenia and T2D and outpatients with T2D only.	Cross-sectional	Age: <ul style="list-style-type: none"> Mean=47.7±8.7 years Range=28–65 years Sample size=211 Gender: <ul style="list-style-type: none"> Men=39.8% Women=60.2% 	Outcome(s): <ul style="list-style-type: none"> Overall DSCB Glycemic control BMI Tool(s): <ul style="list-style-type: none"> Chinese version of SDSCA (C-SDSCA) HbA1c Kg/m² 	<ul style="list-style-type: none"> T2D patients with comorbid schizophrenia had significantly lower: total self-care scores compared to those with T2D only ($M=35.22\pm12.70$ vs 39.18 ± 12.64, $p=0.02$). They also had significantly lower score in diet ($M=16.60\pm6.83$ vs 18.16 ± 4.07, $p=0.04$), physical activity ($M=6.65\pm3.64$ vs 8.33 ± 3.40, $p<0.001$), and blood glucose self-monitoring ($M=1.40\pm3.05$ vs 3.07 ± 3.29, $p<0.0001$) compared to those with T2D only. Similarly they had significantly higher HbA1c ($M=7.9\pm1.7$ vs 7.4 ± 0.8, $p=0.02$) and higher BMI ($M=27.4\pm4.7$ vs 24.2 ± 2.3, $p<0.0001$) compared to those with T2D only. 	Discordant
Devarajoo and Chinna ⁵⁶	Hulu Selangor, Malaysia	To explore the relationship between depression, diabetes distress, and self-efficacy with diabetes self-care practice.	Cross-sectional	Age: <ul style="list-style-type: none"> Mean=55.33 ± 10.09 years Sample size=371 Gender: <ul style="list-style-type: none"> Men=38.0% Women=62.0% 	Outcome(s): <ul style="list-style-type: none"> Overall DSCB Tool(s): <ul style="list-style-type: none"> Malaya version of SDSCA 	<ul style="list-style-type: none"> The mean self-care (SDSCA) score=3.87±0.82, with 170 (45.8%) categorized as good diabetes self-care practice, diet score=4.70±1.56, exercise score=2.77±1.78, medication adherence score=6.01±1.84, SMBG score=1.38±1.59, and foot care score=5.63±1.98. Depression had no significant direct association with diabetes self-care, but had significant negative effect through self-efficacy ($r=-0.115$, $p<0.01$). The model explained 22% of variance in diabetes self-care. 	Discordant

(Continued)

Table 1 (Continued).

Reference	Setting/ Country	Study aim(s)	Design	Sample char- acteristics	Outcome of Interest	Result	Concordant/ Discordant
Feil et al ⁵⁷	USA, National study	To examine the relationship between cognitive impairment and diabetes self-management in a population-based com- munity sample of older adults with Type 2 diabetes.	Cross- sectional	Age: • Mean=70±7.4 years Sample size=1,398 Gender: • Men=47.2% • Women=52.8%	Outcome(s): • Overall DSCB • Glycemic control Tool(s): • DCP • HbA1c	<ul style="list-style-type: none"> • T2D patients with highest comorbidity scores had higher HbA1c values compared to those with lowest comorbidity score ($M=7.3 \pm 1.5\%$ vs $7.0 \pm 1.1\%$, $p=0.02$). • T2D patients with worsened comorbidities 4% less likely to exercise ($p<0.01$) and 2% less likely to check feet for wounds or sores ($p=0.01$). • Moderate cognitively impaired were 9% and severe cognitively impaired were 38% less likely to follow recommended eating plan (both $p=0.01$). • Moderate cognitively impaired were 27% and severe cognitively impaired were 28% less likely to exercise regularly (both $p=0.05$). 	Discordant
Gonzalez et al ⁵⁸	Boston, USA	To examine the relationships between depression and the full range of diabetes self-care behaviors .	Cross- sectional	Age: • Mean=66.2±12.4 years Sample size=909 Gender: • Men=52% • Women=48%	Outcome(s): • Overall DSCB • BMI Tool(s): • SDSCA • Kg/m ²	<ul style="list-style-type: none"> • T2D patients with probable major depression had significantly highest BMI compared to those not meeting the criteria for major depression ($M=34.19 \pm 6.18$ vs 30.70 ± 6.18, $p<0.001$). • They reported significantly fewer days of adherence to diet, exercise, and glucose self-monitoring ($p<0.01$). • They had 2.3-fold increased odds of missing medication doses in the previous week (95% CI=1.5, 3.6, $p<0.001$). 	Discordant

(Continued)

Table 1 (Continued).

Reference	Setting/ Country	Study aim(s)	Design	Sample characteristics	Outcome of Interest	Result	Concordant/ Discordant
Hernandez et al ⁵⁹	Chicago, USA	To examine the relationships between depressive symptoms and diabetes self-care in African American and Hispanic/Latino patients with Type 2 diabetes and whether the associations, if any, is mediated by diabetes-related self-efficacy.	Cross-sectional	Age: <ul style="list-style-type: none"> Mean=53.2±12.3 years Range=25–86 years Sample size=250 Gender: <ul style="list-style-type: none"> Men=31.6% Women=68.4% 	Outcome(s): <ul style="list-style-type: none"> Overall DSCB Tool(s): <ul style="list-style-type: none"> SDSCA 	<ul style="list-style-type: none"> The mean score of diabetes self-care behavior ranged from a low of 2.5 per week (exercise) to a high of 6.2 days per week (oral medication adherence). Depressive symptoms negatively significantly associated with self-care domains of general diet ($\beta=-0.032$, $p=0.01$), specific diet ($\beta=-0.020$, $p=0.02$), physical activity ($\beta=-0.046$, $p<0.01$), and blood glucose self-monitoring ($\beta=-0.020$, $p=0.01$) in the African American group. Depressive symptoms negatively significantly associated only with specific diet ($\beta=-0.030$, $p<0.01$) in the Hispanic/Latino group. Self-efficacy mediated the association between depression and self-care only for African American group ($\beta=-0.024$, $p=0.01$). 	Discordant
Johnson et al ⁶⁰	Alberta, Canada	To examine the association of diabetes-related distress (DD) and depressive symptoms (DS) with physical activity (PA) and adherence to recommended dietary behaviors in adults with T2D.	Cross-sectional	Age: <ul style="list-style-type: none"> Mean=64.4±10.6 years Sample size=2,040 Gender: <ul style="list-style-type: none"> Men=54.8% Women=45.2% 	Outcome(s): <ul style="list-style-type: none"> Exercise behavior Dietary behavior Tool(s): <ul style="list-style-type: none"> SDSCA 	<ul style="list-style-type: none"> 42% of participants had good adherence of dietary behaviors whereas 79% did not achieve the guideline of 150 minutes of weekly moderate-to-vigorous physical activity (MVPA). T2D patients with DD alone were 1.8 times (95% CI=1.1, 2.9) and those with DD and DS combined were 2.0 times (95% CI=1.1, 3.7) more likely not to meet PA guideline compared to those without DD or DS. DS did not independently significantly associated with meeting PA guideline (OR=1.4, 95% CI=0.7, 3.0). T2D patients with DD alone were 1.5 times (95% CI=1.4, 3.4), DS alone were 5.2 times (95% CI=2.7, 9.7), and DD and DS combined were 2.5 times (95% CI=1.6, 3.8) more likely to have poor adherence to recommended dietary behavior compared to those without DD or DS. 	Discordant

(Continued)

Table 1 (Continued).

Reference	Setting/ Country	Study aim(s)	Design	Sample characteristics	Outcome of Interest	Result	Concordant/ Discordant
Lipscombe et al ⁶¹	Quebec, Canada	To examine the association between physical inactivity and anxiety symptoms in a community-based sample of men and women of with type 2 diabetes mellitus.	Cross-sectional	Age: • Mean=61.7±8.3 years Sample size =1,953 Gender: • Men=50.1% • Women=49.9%	Outcome(s): • Exercise behavior Tool(s): • Number of days over the previous month spent on exercise engagement for at least 15 minutes.	<ul style="list-style-type: none"> • A 27.3% of participants reported being physically inactive. • T2D patients with mild anxiety symptoms and moderate to severe anxiety symptoms were 1.4 times (95%CI=1.05, 1.96) and 1.7 times (95%CI=1.16, 2.50) more likely to report being inactive compared to those without anxiety symptoms. • Women who had mild anxiety symptoms were 1.5 times (95%CI=1.02, 2.31) more likely to report being inactive compared to women without anxiety symptoms. • Men who had moderate to severe anxiety symptoms were 2.5 times (95%CI=1.31, 4.75) more likely to be inactive compared to men without anxiety symptoms. • T2D patients who had >2 comorbidities were 1.5 times (95%CI=1.08, 2.13), obesity were 1.5 times (95%CI=1.08, 2.15), and depression were 1.8 times (95%CI=1.04, 3.13) more likely to be physically inactive. 	Discordant
Mut-Vitcu et al ⁶³	Romania	To evaluate the prevalence of depression and its impact on the quality of diabetes-related self-care activities in elderly patients with type 2 diabetes	Cross-sectional	Age: • Median=64 years • Range=52–86 years Sample size =184 Gender: • Men=43.5% • Women=56.5%	Outcome(s): • Overall DSCB Tool(s): • SDSCA	<ul style="list-style-type: none"> • Participants had a mean HbA1c of 8.14±1.7% and BMI of 30.7±4.3 Kg/m². • Severe depression – compared with moderate and no depression – in T2D patients is associated with a significant decrease in quality of diabetes self-care activities global score (24 vs 33 vs 37 points; $p<0.001$), in the diet score (12 Vs 16 Vs 18 points; $p<0.001$); physical exercise score (5 vs 6 vs 8 points; $p<0.001$); and feet care score (4 vs 5 vs 7 points; $p=0.017$). • Presence of diabetes neuropathy ($\beta=-0.183$; $p=0.016$) and chronic kidney disease ($\beta=-0.230$; $p=0.003$) are associated with decrease in quality of diabetes self-care activities. 	Discordant

(Continued)

Table 1 (Continued).

Reference	Setting/ Country	Study aim(s)	Design	Sample characteristics	Outcome of Interest	Result	Concordant/ Discordant
Nau et al ⁶⁴	Michigan, USA	To determine the main effects, and interactive effect, of gender and depression on patients' adherence to oral diabetes medication	Cross-sectional	Age: • Mean=56.0±11.1 years Sample size =391 Gender: • Men=50.1% • Women=49.9%	Outcome(s): • Medication adherence Tool(s): • A 4-item scale	<ul style="list-style-type: none"> Depression had a significant main effect on medication adherence ($F=4.82$, $p=0.03$) – persons categorized as depressed were less adherence. No significant association between gender and adherence ($F=3.01$, $p=0.08$) But, gender $\times$ depression interaction effect was significant ($F=5.93$, $p=0.01$) – men with depression reported significantly worse medication adherence than men without depression, while women were relatively adherent regardless of depressive symptom severity. 	Discordant
Shin et al ⁶⁷	John Hopkins Hospital, USA	To examine whether problem-solving and diabetes self-management behaviors differ by depression diagnosis in adults with T2D.	Cross-sectional	Sample size =103 Gender: • Men=40.8% • Women=59.2%	Outcome(s): • Overall DSCB Tool(s): • SDSCA	<ul style="list-style-type: none"> T2D patients with major depression (MDD) had lower total self-care ($MD=-10.02$, $p\leq 0.01$), diet ($MD=-5.32$, $p\leq 0.01$), and blood glucose self-monitoring ($MD=-2.69$, $p\leq 0.05$) scores compared to those without depression after adjusting for age, sex, and diabetes duration. 	Discordant
Smith et al ³⁵	Quebec, Canada	To ascertain the association of elevated co-occurring anxiety and depression symptoms, elevated anxiety symptoms or elevated depression symptoms alone with self-care behaviors in people with T2D.	Cross-sectional	Age: • Mean=60.51±8.4 years Sample size =1,990 Gender: • Men=49.8% • Women=50.2%	Outcome(s): • Blood glucose checking • Dietary behavior • Exercise behavior • Smoking behavior Tool(s): • Canadian recommendation for self-care	<ul style="list-style-type: none"> Groups with elevated co-occurring anxiety were 3.39 times (95%CI=2.23, 5.14) and depression symptoms were 3.02 times (95%CI=1.94, 4.71), elevated anxiety symptoms alone were 2.71 times (95%CI=1.36, 5.40) more likely to report poor eating habits compared to those without these problems. Groups with elevated co-occurring depression and anxiety symptoms were 1.85 times (95%CI=1.19, 2.87) and elevated depression symptoms alone were 1.98 times (95%CI=1.23, 3.17) also less likely to meet physical activity recommendations compared to those without depressive symptoms. T2D people with elevated co-occurring depression and anxiety were 1.90 time (95%CI=1.23, 2.95) more likely to be current smokers compared to those with low depression and anxiety scores. 	Discordant

(Continued)

Table 1 (Continued).

Reference	Setting/ Country	Study aim(s)	Design	Sample characteristics	Outcome of Interest	Result	Concordant/ Discordant
Zuberi et al ⁶⁹	Pakistan	To assess the associations of depression with glycemic control and compliance to self-care activities in adult patients with T2D.	Cross-sectional	Age: • Mean=52 years • Range=31–60 years Sample size=286 Gender: • Men=44.8% • Women=55.2%	Outcome(s): • Overall DSCB • Glycemic control Tool(s): • Self-report items • HbA1c	<ul style="list-style-type: none"> • Over half of the study participants practiced self-care activities except exercise. • Depressed compared to nondepressed persons had significantly higher mean HbA1c levels (8.5% Vs 7.7%, $p<0.001$). • Depression was associated with low compliance to self-care activities such as taking dose as advised ($OR=0.32$; 95%CI=0.14–0.73), dietary restrictions ($OR=0.45$; 95%CI=0.26–0.79) and foot care ($OR=0.38$; 95%CI=0.18–0.83). • Depressed patients were 5.6 times (95% CI=2.88–10.76) more likely to have poor glycemic control ($HbA1c \geq 7\%$). 	Discordant
Aikens et al ⁷⁰	Midwestern, USA	To clarify the longitudinal associations between depressive symptoms (DS) and diabetes-related distress (DRD) and key outcomes (self-management behaviors and glycemic control) measured 6 months later.	Cohort	Age: • Mean=57.3±8.3 years • Range=27–88 years Sample size=253 Gender: • Men=50% • Women=50%	Outcome(s): • Overall DSCB • Glycemic control Tool(s): • SDSCA • HbA1c	<ul style="list-style-type: none"> • The mean (SD) HbA1c was $7.6 \pm 1.6\%$. • 52% of the participants were high adherent, 42% medium adherent, and 6% low adherent. • DS significantly predicted future diet behavior ($\beta=-0.15$, $p=0.049$), physical activity ($\beta=-0.25$, $p=0.001$), and glucose testing ($\beta=-0.17$, $p=0.018$). • DRD predicted future medication adherence ($\beta=0.20$, $p=0.011$). • DRD predicted future glycemic control ($\beta=0.30$, $p<0.001$). 	Discordant
Axon et al ⁷¹	USA	To examine whether depression impacts medication non-adherence (MNA) over time and determine if race has a differential impact on MNA in patients with T2D and comorbid depression.	Cohort	Sample size=740,197 Gender: • Men=97.7% • Women=2.3%	Outcome(s): • Medication adherence Tool(s): • Medication Possession Ratio (MPR)	<ul style="list-style-type: none"> • The odds of MNA was 1.12 times (95%CI=1.11, 1.13) higher for those with depression compared to those without depression. • After excluding depression, those with 1 comorbid condition ($OR=0.86$, 95% CI=0.85, 0.87) and 2 comorbidities ($OR=0.91$, 95%CI=0.90, 0.92) had lower MNA compared to those with T2D alone. • Participants with ≥ 3 comorbidities (after excluding depression) had higher MNA ($OR=1.09$, 95% CI=1.08, 1.10) compared to those with diabetes alone. 	Discordant

(Continued)

Table 1 (Continued).

Reference	Setting/ Country	Study aim(s)	Design	Sample characteristics	Outcome of Interest	Result	Concordant/ Discordant
Gonzalez et al ⁷²	Massachusetts and Boston, USA	To examine prospectively the association of depression symptoms with subsequent self-care and medication adherence in patients with T2D.	Cohort	Age: • Mean=65.5±11.6 years Sample size=208 Gender: • Men=51% • Women=49%	Outcome(s): • Overall DSCB Tool(s): • SDSCA	<ul style="list-style-type: none"> T2D patients with higher levels of depression at baseline reported significantly lower adherence to general diet recommendations and specific recommendations for consumption of fruits and vegetables and spacing of carbohydrates, less exercise, and poorer foot care at follow-up (β ranging from -0.12 to -0.23, $p<0.05$). Also, each 1-point increase in baseline depression score was associated with a 1.08-fold increase in the odds of non-adherence to prescribed medication at follow-up (95% CI, 1.001, 1.158, $P=0.047$). 	Discordant
Kalsekar et al ⁷⁴	West Virginia	To examine the impact of depression on adherence to oral hypoglycemic agents (OHAs) in patients newly diagnosed with T2D.	Cohort	Sample size =1,326 Gender: • Men=36.7% • Women=63.3%	Outcome(s): • Medication adherence Tool(s): • MPR	<ul style="list-style-type: none"> Depressed patients were 3–6% less adherent to OHAs than nondepressed patients ($p<0.05$), after adjusting for confounding variables. 	Discordant
Kreyenbuhl et al ⁷⁵	USA	To compare adherence to oral hypoglycemic medications (OHA) for diabetes patients with Vs without comorbid schizophrenia.	Cohort	Sample size =22,014 Gender: • Men=96.5% • Women=3.5%	Outcome(s): • Medication adherence Tool(s): • MPR	<ul style="list-style-type: none"> Having schizophrenia was associated with a 25% lower likelihood of poor adherence compared to with not having schizophrenia (OR=0.75, 95% CI=0.70, 0.80). Those with depression were 1.25 times (95% CI=1.12, 1.38, $p<0.0001$), with CCI≥2 were 1.09 times (95% CI=1.03, 1.16, $p=0.005$) and with alcohol or substance use disorder were 1.45 times (95% CI=1.28, 1.64, $p<0.0001$) likely to be poor adherent to OHA. 	Discordant

(Continued)

Table 1 (Continued).

Reference	Setting/ Country	Study aim(s)	Design	Sample characteristics	Outcome of Interest	Result	Concordant/ Discordant
Wagner et al ⁷⁹	USA	To examine the association between lifetime history of major depressive disorder (L-MDD) and diabetes self-management in women with Type 2 diabetes who were not depressed at the time of assessment.	Case-control	Age: • Mean=60.1±9.7 years Sample size=153	Outcome(s): • Overall DSCB Tool(s): • SDSCA	<ul style="list-style-type: none"> • Total self-care (10 SDSCA core items) mean (SD) score was 37.3±12.2, blood glucose self-monitoring was 4.7±2.4, general diet was 4.3±1.9, specific diet was 4.2±1.6, exercise was 2.7±2.0, and foot care was 4.8±1.4. • The participants had a mean (SD) HbA1c of 6.7 ± 1.2%. • Never depressed T2D women with L-MDD had lower total SDSCA score compared to control (F (1, 110)=8.01, $p<0.05$) after controlling for treatment regimen, depressive symptoms, and conscientiousness. • Self-efficacy was a significant mediator of the influence of L-MDD on SDSCA scores ($z=-2.14$, standard error=1.01, $p<0.05$). 	Discordant
O'Shea et al ⁷⁶	Ireland	To examine whether the type of comorbid condition affects medication persistence and adherence in patients initiating oral anti-hyperglycemic (OAH) therapy.	Cohort	Sample size =21,280 Gender: • Men=54% • Women=46%	Outcome(s): • Medication adherence Tool(s): • MPR	<ul style="list-style-type: none"> • Persistence was 74.0% and 62.6% and adherence was 70.0% and 66.7% for all OAHs at 6 and 12 months, respectively. • Patients with only concordant comorbidity were significantly more likely to be persistent at 6 month (OR=1.45, 95%CI=1.28, 1.65) and 12 month (OR=1.22, 95%CI=1.09, 1.38). • Patients with only discordant comorbidity were significantly less likely to be persistent at 6 month (OR=0.40, 95%CI=0.35, 0.46) and 12 month (OR=0.43, 95%CI=0.38, 0.50). • Patient with only concordant comorbidity were significantly more likely to be adherent at 6 month (OR=1.43, 95%CI=1.28, 1.61) and 12 month (OR=1.42, 95%CI=1.26, 1.62). • Patients with only discordant comorbidity were significantly less likely to be adherent at 6 month (OR=0.47, 95%CI=0.41, 0.54) and 12 month (OR=0.46, 95%CI=0.40, 0.54). 	Both

(Continued)

Table 1 (Continued).

Reference	Setting/ Country	Study aim(s)	Design	Sample characteristics	Outcome of Interest	Result	Concordant/ Discordant
Palakodeti et al ⁷⁷	North California, USA	To identify the predictors and clinical effects among inactive patients with diabetes who become physically active, in the setting of a large integrated health system.	Cohort	Age: <ul style="list-style-type: none"> Mean=60.2 years Sample size =6,853 Gender: <ul style="list-style-type: none"> Men=47.1% Women=52.9% 	Outcome(s): <ul style="list-style-type: none"> Physical activity Tool(s): <ul style="list-style-type: none"> Self-reported items 	<ul style="list-style-type: none"> 62.5% reported being physically inactive, while 16.0% reported achieving the recommended physical activity (≥ 150 minute/week) by the last visit of the study period. Patients with comorbid chronic kidney disease were 22% (OR=0.78, CI=0.65, 0.94), depression were 23% (OR=0.77, CI=0.62, 0.96), obesity of BMI 30–34.9 kg/m² were 24% (OR=0.76, CI=0.60, 0.97), and obesity of BMI ≥ 35 kg/m² were 45% (OR=0.55, CI=0.42, 0.70) less likely to achieve the recommended threshold of 150 min/week of MVPA during the follow-up period. 	Both
Trief et al ⁸¹	Syracuse, New York, USA	To examine the change in adherence to the recommended self-care in response to the diabetes home telemedicine intervention.	Randomized Controlled Trial (RCT)	Age: <ul style="list-style-type: none"> Mean=70.8$\pm$6.6 years Sample size =1,665 Gender: <ul style="list-style-type: none"> Men=37.2% Women=62.8% 	Outcome(s): <ul style="list-style-type: none"> Overall DSCB Glycemic control Tool(s): <ul style="list-style-type: none"> SDSCA HbA1c 	<ul style="list-style-type: none"> The total sample had a mean (SD) BMI of 31.9 $\pm$ 6.8 kg/m² and HbA1c of 7.4$\pm$1.5%. Participants with greater comorbidity over time had poorer adherence to self-care activities (β=−0.0204, p=0.01). Adherence to self-care was a significant mediator of HbA1c (p<0.001). 	Both
O'Shea et al ⁶⁵	Dublin, Ireland	To examine the impact of comorbidity on HRQoL and self-care in T2DM patients attending an outpatient setting	Cross-sectional	Age: <ul style="list-style-type: none"> Range=25–80 years Sample size =159 Gender: <ul style="list-style-type: none"> Men=59.7% Women=40.3% 	Outcome(s): <ul style="list-style-type: none"> Overall DSCB Tool(s): SDSCA	<ul style="list-style-type: none"> T2D patients with both concordant and discordant comorbidities had the lowest level of physical activity (β=−0.32, p<0.0001) compared to those without any comorbidity. But, no difference in other dimensions of DSCB. 	Both

Abbreviations: T2D, type 2 diabetes; USA, United States of America; SDSCA, Summary of Diabetes Self-Care Activities; DSCB, Diabetes Self-Care Behaviors; HbA1c, glycated hemoglobin.

Concordant and discordant comorbidities

The most frequently measured T2D concordant and discordant comorbidities in relation to self-care behaviors are shown in Table 2. Hypertension, obesity and chronic kidney disease were the most frequent concordant conditions with heart failure being the least frequently measured. Depression was the most frequently measured discordant comorbidity.

Frequency of diabetes self-care behaviors in concordant and discordant comorbidities concordant

Medication adherence was the most frequently reported self-care behavior for concordant conditions. Two of the

Table 2 Comorbid conditions investigated by the studies included in this review

Comorbid condition	Number of studies
Diabetes-concordant comorbidities:	
Hypertension	11
Chronic kidney disease	9
Obesity	8
Coronary artery disease	8
Retinopathy	7
Hyperlipidemia	6
Neuropathy	4
Peripheral vascular disease	4
Heart failure	3
Diabetes-discordant comorbidities:	
Depression	26
Osteoarthritis	5
Lung disease	4
Cancer	3
Liver disease	3
Schizophrenia	2
Cognitive impairment	2
Anxiety	2
Asthma	2
Stomach/intestinal ulcer	2
Back pain	2
Anemia	2
Thyroid disease	2
Substance use disorder	2
Sleep disturbance	1
Migraines	1
Addison's disease	1
Cushing's syndrome	1
Fluid/electrolyte disorder	1

four studies that focused on medication adherence in concordant comorbid conditions described that more than 50% of their participants were adherent.^{62,73} The other two studies, however, reported that compared to those without comorbid conditions, T2D patients with concordant comorbidities including hypertension, dyslipidemia and obesity were 1.78 (95% CI=1.064, 2.981) times less likely to be adherent to the medication regimen⁴⁹ and less likely to be physically active ($\beta=-0.32$, $p<0.0001$).⁶⁵ Another study evaluating foot self-care in adults with T2D reported that those with hyperlipidemia were 38% less likely to perform the recommended foot self-care for ≥ 6 days per week.⁵¹ A randomized controlled trial tested the effects of a telemedicine intervention (Informatics for Diabetes Education and Telemedicine Project) on adherence over time in older Hispanic and African Americans reported that greater comorbidity over time had worse adherence to diabetes self-care activities ($\beta=-0.0204$, $p=0.01$) compared to those without comorbidities.⁸¹

Discordant

The most common self-care behavior reported for discordant conditions was also medication adherence followed by dietary and physical activity. Only three studies reported the mean score of the overall diabetes self-care activities.^{50,56,79} Among T2D patients with discordant comorbid depression, the overall mean diabetes self-care score ranged from 3.87 to 40.5. This shows the mean of total diabetes self-care score on the Summary of Diabetes Self-Care Activities (SDSCA) for diet, exercise, medication, blood glucose monitoring, foot care, and smoking cessation which indicates less diabetes self-care performance. Two additional studies also reported that only 50% of their study participants practiced diabetes self-care behaviors in the presence of depressive symptoms.^{69,70}

Regarding specific diabetes self-care behaviors, four studies examined depressive symptoms and found that taking prescribed medications was the most frequently performed while exercise was the least often performed.^{50,52,59,79} In contrast, another study investigating depressive symptoms reported that taking medications as prescribed was the most frequently performed self-care behavior ($M=6.01\pm 1.84$) during the week, followed by exercise ($M=2.77\pm 1.78$) and blood glucose self-monitoring ($M=1.38\pm 1.59$).⁵⁶ Johnson and colleagues⁶⁰ examined dietary and physical activity behaviors among adults with diabetes-related distress and depressive symptoms. In this study, only 42% had good adherence to the recommended diabetes dietary recommendations and

almost 80% did not meet physical activity guidelines. Participants with depressive symptoms were 5.2 less likely to adhere to dietary guidelines compared to those without depressive symptoms, but this was not independently associated with physical activity level.⁶⁰ Persons with mild anxiety symptoms and moderate to severe anxiety symptoms were 1.4 times and 1.7 times more likely to report being physically inactive than persons without anxiety symptoms, respectively. Feil and colleagues⁵⁷ reported that older T2D patients with greater cognitive impairment were less likely to adhere to exercise ($p<0.05$), diet ($p<0.01$) or check feet for wounds ($p=0.01$) compared to those without cognitive impairment. In addition, as cognitive impairment worsened, adherence to each diabetes self-care behavior declined. Axon and colleagues⁷¹ examined whether depression affects medication adherence over time. Findings revealed the odds of being medication nonadherent was 1.12 times higher among those with comorbid depression compared to those with T2D alone. The same study revealed that having three or more comorbid conditions, after excluding depression, was associated with higher medication nonadherence (OR 1.09, [95% CI: 1.08, 1.10]) while having two or less comorbidities was associated with lower medication nonadherence compared to those with T2D alone.

Both concordant and discordant

Two studies that measured exercise and medication adherence behaviors in those with both concordant and discordant comorbid conditions such as depression, cognitive impairment, chronic kidney disease, and obesity found that over 62.5% of their participants did not achieve the recommended weekly moderate-to-vigorous physical activity level⁷⁷ and 30% were medication nonadherent.⁷⁶ In the anxiety study described above, if there were greater than 2 comorbidities including obesity, the likelihood of being physically inactive was almost two-fold⁶¹ compared to those without comorbidities.

Summary of concordant and discordant comorbidity and diabetes self-care behaviors

Two cohort studies^{73,76} and two cross-sectional studies^{51,62} reported that there were significantly higher odds of medication adherence and foot self-care in T2D patients with specific concordant comorbidities such as hyperlipidemia and coronary heart disease compared to those without these comorbidities. In contrast, two cohort,^{73,77} two

cross-sectional,^{53,63} one case-control,⁷⁸ and one interventional⁸⁰ study reported that T2D patients with concordant comorbid hypertension, dyslipidemia, obesity, and chronic kidney disease had significantly lower odds of achieving the recommended diabetes self-care behaviors than those without these comorbid conditions.

Studies have shown the negative association between diabetes-discordant comorbidities and diabetes self-care behaviors. A cohort study⁷⁶ found statistically significant lower odds of medication adherence in T2D patients with discordant comorbidities overtime. Similarly, 14 of the 21 cross-sectional studies reviewed reported that T2D patients with discordant comorbid depression,^{54,58,59,63,64,67} diabetes-related distress,^{52,60} anxiety,^{35,61} schizophrenia,⁵⁵ impaired sleep quality and daytime sleepiness,⁵³ and cognitive impairment⁵⁷ had statistically significant lower odds of performing the recommended diabetes self-care behaviors compared to those without these comorbidities. The two case-control studies also reported that T2D patients with comorbid depression had significantly lower scores for diabetes self-care behaviors compared to those without depression.^{78,79} In addition, five of the eight cohort studies reported that T2D patients with discordant comorbid schizophrenia and alcohol or substance use disorder,⁷⁵ diabetes-related distress,⁷⁰ and depression^{70–72,74,75} had statistically significant lower odds of achieving the recommended self-care behaviors.

In several studies, depression was mediated by self-efficacy or other factors. Among seven studies depression had no direct association with diabetes self-care behavior but was mediated by self-efficacy,^{50,56,79} diabetes-related distress,^{52,60} gender,⁶⁴ and health literacy.⁶⁶

Discussion

Findings from this systematic review indicate that concordant and discordant comorbidities may result in poorer diabetes self-care. To our knowledge, this is the first systematic review detailing differences between concordant and discordant comorbid conditions on diabetes self-care behaviors. The findings are consistent with previous research that multimorbidity has a negative effect on the T2D patients' achievement of the recommended diabetes self-care behaviors. In general, there is support in the literature that concordant conditions may result in higher frequency and adherence of self-care behaviors than discordant. For example, O'Shea and colleagues⁷⁶ directly compared concordant and discordant comorbidities on oral-hypoglycemic medication adherence and determined those with concordant conditions were

more adherent. Other studies, however, do not support these findings. He and colleagues,⁷³ for example, showed a positive association between peripheral neuropathy and adherence to insulin therapy, while another study showed poorer quality of diabetes self-care activities with concomitant peripheral neuropathy.⁶³ The effect of diabetes-concordant comorbidities on diabetes self-care behaviors may vary depending on the type of comorbidity and the self-care behavior being measured. For example, in the two studies reporting medication adherence with existing peripheral neuropathy,^{63,73} there were considerable differences in participant comorbidity type and the self-care behaviors measured. In the He and colleagues⁷³ study, only concordant comorbidities and one specific behavior, medication adherence, was included. The Mut-Vitcu and colleagues⁶³ included both concordant and discordant chronic conditions and overall diabetes self-care behaviors were measured which may explain in part, the disparate findings.

Too much of an emphasis on the features of single comorbid illnesses may result in failure to see larger patterns in the ways that treatments for T2D and comorbidities interact.³⁸ An increased number of comorbidities was associated with a decline in diabetes self-care behaviors and is consistent with previous research.^{61,71} Importantly, not all comorbid conditions are the same nor have the same consequences on diabetes self-care performance. Periodic evaluation of comorbidities, particularly as they worsen, may change patient self-care priorities from diabetes to a more dominant comorbid condition resulting in poor disease control and higher risk for complications.³⁶ The use of the concordance and discordance framework is beneficial toward this end and may help to identify which chronic conditions are most likely to result in poorer outcomes based on care goals of one of more dominant comorbidities.³⁸

Results from this systematic review support previous research concerning the negative impact of diabetes-discordant comorbidities on diabetes self-care behaviors. Most of the diabetes-discordant comorbidities involved mental health and neuropsychiatric disorders such as depression, anxiety, diabetes-related distress, cognitive decline and schizophrenia. Analyses identified that these comorbidities had a more negative impact on diabetes self-care behaviors compared to patients with T2D alone. Most of the discordant conditions were examined independently with diabetes self-care behaviors rather than with other multiple conditions which may have influenced findings. Screening T2D patients for neuropsychological multimorbidity such as depression

and cognitive decline is important because they have been shown to reduce the capability of the individual to perform effective diabetes self-care. Simple screening tests such as the Patient Health Questionnaire-9 (PHQ-9)⁸² and the Montreal Cognitive Assessment (MoCA)⁸³ can be administered in clinic settings and have been previously validated in the diabetes population.

The frequency of diabetes self-care performance was lower in T2D patients with comorbid conditions compared to those without comorbidities. This is true both for the overall or global diabetes self-care behaviors and specific components including diet, exercise, blood glucose self-monitoring, medication adherence, foot care, and smoking behaviors. There is agreement among studies^{52,56,59,79} that medication adherence is the most frequently performed specific diabetes self-care behaviors whereas exercise is the least performed. Thus, barriers to engagement in exercise self-care among T2D patients with comorbid conditions need to be explored in future research.

Medication adherence was the most frequently performed diabetes self-care behavior in both concordant and discordant chronic conditions, but little is known about how adults with T2D rank the importance of their specific self-care behaviors, management, and outcomes relative to those of other comorbid conditions such as hypertension or depression. A better understanding of the individual's self-care priorities and the process used for decision-making when managing other complex chronic conditions may provide additional insight on adherence to self-care. Future research is warranted to examine which strategies best organize care for T2D patients with comorbid conditions to optimize clinical outcomes and quality of life, as well as identifying self-care priorities from the patient's perspective. The number of adults experiencing T2D with multimorbidity is anticipated to increase making integration of care an essential component of diabetes management. Guidelines and interventions that encompass concordant conditions if they are found to be associated with improved diabetes self-care would be beneficial and help to personalize care goals. Worsened outcomes associated with discordant chronic conditions could be used to highlight areas where patients may need additional care to prevent or delay complications.^{35,36}

Strengths and limitations

Comparing the effect of concordant and discordant comorbid conditions on diabetes self-care behaviors was strength because no studies have previously described diabetes self-care using this approach. Using the diabetes-concordant and

discordant framework can provide clinicians additional insight into which comorbid conditions may lead to better or poorer adherence to diabetes self-care behaviors. The majority of studies were cross-sectional and nonrandomized which limited the quality of evidence and interpretation of findings. Whether all confounders known to influence T2D self-care behaviors were controlled in these studies is unknown. In addition, self-reported measures were used for most studies which are less accurate and reliable. The heterogeneity of the samples, designs and measures of self-care across studies made it impossible to perform pooled analytical procedures or meta-analysis. The majority (N=26) of studies included in this review were conducted in high-income countries which limits the generalizability of the findings to middle and low-income countries where the rates of comorbid conditions may be higher in T2D.^{84,85} For instance, there was no study reported from Africa concerning the effect of comorbidities on diabetes self-care behaviors.

Conclusion

Diabetes-concordant comorbidities are associated with higher adherence to diabetes self-care behaviors than discordant, but future research is needed using more rigorous designs to better understand these complex relationships with diabetes self-care behaviors. Given the increase in multimorbidity observed among patients with T2D, guidelines are urgently needed that integrate diabetes self-care with other chronic conditions, especially discordant conditions in order to optimize clinical outcomes and quality of life.

Disclosure

The authors report no conflicts of interest in this work.

References

1. Zimmet PZ. Diabetes and its drivers: the largest epidemic in human history? *Clin Diabetes Endocrinol*. 2017;3(1). doi:10.1186/s40842-40016-40039-40843
2. Jaacks LM, Siegel KR, Gujral UP, Narayan KMV. Type 2 diabetes: a 21st century epidemic. *Best Pract Res Clin Endocrinol Metab*. 2016;30:331–343. doi:10.1016/j.beem.2016.05.003
3. Unnikrishnan R, Pradeepa R, Joshi SR, Mohan V. Type 2 diabetes: demystifying the global epidemic. *Diabetes*. 2017;66:1432–1442. doi:10.2337/db16-0766
4. IDF. *IDF Diabetes Atlas*. International Diabetes Federation: 2017.
5. NCDRiskFactorCollaboration. Worldwide trends in diabetes since 1980: a pooled analysis of 751 population-based studies with 4.4 million participants. *Lancet*. 2016;387:1513–1530. doi:10.1016/S0140-6736(16)00618-8
6. WHO. *Global Reports on Diabetes*. Geneva: WHO; 2016.
7. Caughey GE, Roughead EE, Vitry AI, McDermott RA, Shakib S, Gilbert AL. Comorbidity in the elderly with diabetes: identification of areas of potential treatment conflicts. *Diabetes Res Clin Pract*. 2010;87:385–393. doi:10.1016/j.diabres.2009.10.019
8. Lang VB, Markovic BB. Prevalence of comorbidity in primary care patients with type 2 diabetes and its association with elevated HbA1c: a cross-sectional study in Croatia. *Scand J Prim Health Care*. 2016;34(1):66–72. doi:10.3109/02813432.2015.1132886
9. Luijckx H, Biermans M, Bor H, et al. The effect of comorbidity on glycemic control and systolic blood pressure in type 2 diabetes: a cohort study with 5 year follow-up in primary care. *PLoS One*. 2015;10(10):e0138662. doi:10.1371/journal.pone.0138662
10. O'Shea M, Teeling M, NBennet K. The prevalence and ingredient cost of chronic comorbidity in the Irish elderly population with medication treated type 2 diabetes: a retrospective cross-sectional study using a national pharmacy claims database. *BMC Health Services Res*. 2013;13(23). Available from: <http://www.biomedcentral.com/1472-6963/1413/1423>
11. Li R, Bilik D, Brown MB, et al. Medical costs associated with type 2 diabetes complications and comorbidities. *Am J Manag Care*. 2013;19(5):421–430.
12. Adriaanse MC, Drewes HW, van der Heide I, Struijs JN, Baan CA. The impact of comorbid chronic conditions on quality of life in type 2 diabetes patients. *Qual Life Res*. 2016;25(1):175–182. doi:10.1007/s11136-015-1061-0
13. Iglay K, Hannachi H, Howie PJ, et al. Prevalence and co-prevalence of comorbidities among patients with type 2 diabetes mellitus. *Curr Med Res Opin*. 2016;32(7):1243–1252. doi:10.1185/03007995.2016.1168291
14. Alonso-Moran E, Orueta JF, Esteban JIF, et al. Multimorbidity in people with type 2 diabetes in the Basque Country (Spain): prevalence, comorbidity clusters and comparison with other chronic patients. *Eur J Intern Med*. 2015;26:197–202. doi:10.1016/j.ejim.2015.02.005
15. Chen H, Zhang Y, Wu D, et al. Comorbidity in adult patients hospitalized with type 2 diabetes in Northeast China: an analysis of hospital discharge data from 2002 to 2013. *Biomed Res Int*. 2016;2016:1671965. doi:10.1155/2016/1671965
16. Luijckx H, Schremer T, Bor H, et al. Prevalence and incidence density rates of chronic comorbidity in type 2 diabetes patients: an exploratory cohort study. *BMC Med*. 2012;10(128). Available from: <http://www.biomedcentral.com/1741-7015/1710/1128>
17. Ervasti J, Kivimäki M, Dray-Spira R, et al. Comorbidity and work disability among employees with diabetes: associations with risk factors in a pooled analysis of three cohort studies. *Scand J Public Health*. 2016;44(1):84–90. doi:10.1177/1403494815605245
18. Lopez JMS, Bailey RA, Rupnow MFT. Demographic disparities among medicare beneficiaries with type 2 diabetes mellitus in 2011: diabetes prevalence, comorbidities, and hypoglycemia events. *Popul Health Manag*. 2015;18(4):283–289. doi:10.1089/pop.2014.0115
19. Naskar S, Victor R, Nath K. Depression in diabetes mellitus—a comprehensive systematic review of literature from an Indian perspective. *Asian J Psychiatr*. 2017;27:85–100. doi:10.1016/j.ajp.2017.02.018
20. Dieren S, Bealens JWJ, Schouw Y, Grobbee DE, Neal B. The global burden of diabetes and its complications: an emerging pandemic. *Eur J Cardiovasc Prev Rehabil*. 2010;17(Suppl 1):S3–S8. doi:10.1097/01.hjr.0000368191.86614.5a
21. Adepoju OE, Bolin JN, Phillips CD, et al. Effects of diabetes self-management programs on time-to-hospitalization among patients with type 2 diabetes: a survival analysis model. *Patient Educ Couns*. 2014;95:111–117. doi:10.1016/j.pec.2014.01.001
22. Osborn CY, Bains SS, Egede LE. Health literacy, diabetes self-care, and glycemic control in adults with type 2 diabetes. *Diabetes Technol Ther*. 2010;12(11):913–919. doi:10.1089/dia.2010.0058
23. Song M. Diabetes mellitus and the importance of self-care. *J Cardiovasc Nurs*. 2010;25(2):93–98. doi:10.1097/JCN.0b013e3181c5a364

24. Tshiananga JKT, Kocher S, Weber C, Emy-Albrecht K, Berndt K, Neeser K. The effect of nurse-led diabetes self-management education on glycosylated hemoglobin and cardiovascular risk factors: a meta-analysis. *Diabetes Educ.* 2012;38(1):108–123. doi:10.1177/0145721711423978
25. Haas L, Maryniuk M, Beck J, et al. National standards for diabetes self-management education and support. *Diabetes Educ.* 2012;38(5):619–629. doi:10.1177/0145721712455997
26. Shrivastava SR, Shrivastava PS, Ramasamy J. Role of self-care in management of diabetes mellitus. *J Diabetes Metab Disord.* 2013;12(14). Available from: <http://www.jmdonline.com/content/12/11/14>
27. Ahola AJ, Groop P-H. Barriers to self-management of diabetes. *Diabetic Med.* 2013;30:413–420. doi:10.1111/dme.12105
28. Onwudiwe NC, Mullins CD, Winston RA, et al. Barriers to self-management of diabetes: a qualitative study among low-income minority diabetics. *Ethn Dis.* 2011;21:27–32.
29. Zulman DM, Asch SM, Martins SB, Kerr EA, Hoffman BB, Goldstein MK. Quality of care for patients with multiple chronic conditions: the role of comorbidity interrelatedness. *J Gen Intern Med.* 2014;29(3):529–537. doi:10.1007/s11606-013-2616-9
30. Roland M, Paddison C. Better management of patients with multimorbidity. *Br Med J.* 2013;346:f2510. doi:10.1136/bmj.f1164
31. Dunbar SB, Butts B, Reilly CM, et al. A pilot test of an integrated self-care intervention for persons with heart failure and concomitant diabetes. *Nurs Outlook.* 2014;62:97–111. doi:10.1016/j.outlook.2013.09.003
32. Dunbar SB, Reilly CM, Gary RA, et al. Randomized clinical trial of an integrated self-care intervention for persons with heart failure and diabetes: quality of life and physical functioning outcomes. *J Card Fail.* 2015;21(9):719–729. doi:10.1016/j.cardfail.2015.05.012
33. Smith SM, Soubhi H, Fortin M, Hudon C, O'Dowd T. Managing patients with multimorbidity: systematic review of interventions in primary care and community settings. *Br Med J.* 2012;345:e5205. doi:10.1136/bmj.e5205
34. Piette JD, Kerr EA. The impact of comorbid chronic conditions on diabetes care. *Diabetes Care.* 2006;29(3):725–731.
35. Magnan EM, Palta M, Johnson HM, Bartels CM, Schumacher JR, Smith MA. The impact of a patient's concordant and discordant chronic conditions on diabetes care quality measures. *J Diabetes Complications.* 2015;29(2):288–294. doi:10.1016/j.jdiacomp.2014.10.003
36. Aung E, Donald M, Coll J, Dower J, Williams GM, Doi SA. The impact of concordant and discordant comorbidities on patient-assessed quality of diabetes care. *Health Expect.* 2015;18(5):1621–1632. doi:10.1111/hex.12151
37. Lin P-J, Kent DM, Winn AN, Cohen JT, Neuman PJ. Multiple chronic conditions in type 2 diabetes mellitus: prevalence and consequences. *Am J Manag Care.* 2015;21(1):e23–e34.
38. Magnan EM, Gittelsohn R, Bartels CM, et al. Establishing chronic condition concordance and discordance with diabetes: a Delphi study. *BMC Fam Pract.* 2015;16:42. doi:10.1186/s12875-12015-10253-12876
39. Ricci-Cabello I, Stevens S, Kontopantelis E, et al. Impact of the prevalence of concordant and discordant conditions on the quality of diabetes care in family practices in England. *Ann Fam Med.* 2015;13:514–522. doi:10.1370/afm.1848
40. Teljeur C, Smith SM, Paul G, Kelly A, O'Dowd T. Multimorbidity in a cohort of patients with type 2 diabetes. *Eur J Gen Pract.* 2013;19(1):17–22. doi:10.3109/13814788.2012.714768
41. Moher D, Liberati A, Tetzlaff J, Altman DG; The PRISMA Group. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med.* 2009;6(7):e1000097. doi:10.1371/journal.pmed.1000097
42. Liberati A, Altman DG, Tetzlaff J, et al. The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate health care interventions: explanation and elaboration. *J Clin Epidemiol.* 2009;62:e1–e34. doi:10.1016/j.jclinepi.2009.06.006
43. Aromataris E, Ritano D. Constructing a search strategy and searching for evidence: a guide to the literature search for a systematic review. *Am J Nurs.* 2014;114(5):49–56. doi:10.1097/01.NAJ.0000446779.99522.f6
44. Joanna Briggs JBI. *Institute Reviewer's Manual 2014 Edition.* Australia: JIB; 2014.
45. Thomas BH, Ciliska D, Dobbins M, Micucci S. A process for systematically reviewing the literature: providing the research evidence for public health nursing interventions. *Worldviews Evid Based Nurs.* 2004;1(3):176–184. doi:10.1111/j.1524-475X.2004.04006.x
46. Higgins J, Deeks J. Selecting studies and collecting data. In: Higgins J, Green S, editors. *Cochrane Handbook for Systematic Reviews of Interventions. Version 5.1.0 [Updated March 2011].* The Cochrane Collaboration; 2011. Available from: www.cochrane-handbook.org
47. Munn Z, Tufanaru C, Aromataris E. Data extraction and synthesis the steps following study selection in a systematic review. *Am J Nurs.* 2014;114(7):49–54. doi:10.1097/01.NAJ.0000451683.66447.89
48. Popay J, Roberts H, Sowden A, et al. *Guidance on the Conduct of Narrative Synthesis in Systematic Reviews: A Product from the ESRC Methods Programme.* 2006.
49. Ahmad NS, Ramil A, Islahudin F, Paraidathathu T. Medication adherence in patients with type 2 diabetes mellitus treated at primary health clinics in Malaysia. *Patient Prefer Adherence.* 2013;7:525–530. doi:10.2147/PPA.S44698
50. Al-Amer R, Ramjan L, Glew P, Randall S, Salamonson Y. Self-efficacy, depression, and self-care activities in adult Jordanians with type 2 diabetes: the role of illness perception. *Issues Ment Health Nurs.* 2016;37(10):744–755. doi:10.1080/01612840.2016.1208692
51. Al-Sayah F, Soprovich A, Qiu W, Edwards AL, Johnson JA; Diabetic Foot Disease. Self-care and clinical monitoring in adults with type 2 diabetes: the Alberta's Caring for Diabetes (ABCD) cohort study. *Can J Diabetes.* 2015;39:S120–S126. doi:10.1016/j.cjcd.2015.05.006
52. Asuzu CC, Walker RJ, Williams JS, Egede LE. Pathways for the relationship between diabetes distress, depression, fatalism and glycaemic control in adults with type 2 diabetes. *J Diabetes Complications.* 2017;31:169–174. doi:10.1016/j.jdiacomp.2016.09.013
53. Chasens ER, Korytkowski M, Sereika SM, Burke LE. Effect of poor sleep quality and excessive daytime sleepiness on factors associated with diabetes self-management. *Diabetes Educ.* 2013;39(1):74–82. doi:10.1177/0145721712467683
54. Chen H-Y, Ruppert K, Charron-Prochownik D, Noullet WV, Zgibor JC. Effects of depression and antidepressant use on goal setting and barrier identification among patients with type 2 diabetes. *Diabetes Educ.* 2011;37(3):370–380. doi:10.1177/0145721711400662
55. Chen S-R, Chien Y-P, Kang C-M, Jeng C, Chang W-Y. Comparing self-efficacy and self-care behaviours between outpatients with comorbid schizophrenia and type 2 diabetes and outpatients with only type 2 diabetes. *J Psychiatr Ment Health Nurs.* 2014;21:414–422. doi:10.1111/jpm.12101
56. Devarajoon C, Chinna K, Khamseh ME. Depression, distress and self-efficacy: the impact on diabetes self-care practices. *PLoS One.* 2017;12(3):e0175096. doi:10.1371/journal.pone.0175096
57. Feil DG, Zhu CW, Sultzer DL. The relationship between cognitive impairment and diabetes self-management in a population-based community sample of older adults with type 2 diabetes. *J Behav Med.* 2012;35:190–199. doi:10.1007/s10865-011-9344-6
58. Gonzalez JS, Safren SA, Cagliero E, et al. Depression, self-care, and medication adherence in type 2 diabetes: relationships across the full range of symptom severity. *Diabetes Care.* 2007;30:2222–2227. doi:10.2337/dc07-0158
59. Hernandez R, Ruggiero L, Prohaska TR, et al. A cross-sectional study of depressive symptoms and diabetes self-care in African Americans and hispanics/latinos with diabetes: the role of self-efficacy. *Diabetes Educ.* 2016;42(4):452–461. doi:10.1177/0145721716654008

60. Johnson ST, Al-Sayah F, Mathe N, Johnson JA. The relationship of diabetes-related distress and depressive symptoms with physical activity and dietary behaviors in adults with type 2 diabetes: a cross-sectional study. *J Diabetes Complications*. 2016;30:967–970. doi:10.1016/j.jdiacomp.2016.02.019
61. Lipscombe C, Smith KJ, Garipey G, Schmitz N. Gender differences in the relationship between anxiety symptoms and physical inactivity in a community-based sample of adults with type 2 diabetes. *Can J Diabetes*. 2014;38:444–450. doi:10.1016/j.cjcd.2013.12.002
62. Malik S, Billimek J, Greenfield S, Sorkin DH, Ngo-Metzger Q, Kaplan SH. Patient complexity and risk factor control among multimorbid patients with type 2 diabetes: results from the R2D2C2 study. *Med Care*. 2013;51:180–185. doi:10.1097/MLR.0b013e318273119b
63. Mut-Vitcu G, Timar B, Timar R, Oancea C, Citu IC. Depression influences the quality of diabetes-related self-management activities in elderly patients with type 2 diabetes: a cross-sectional study. *Clin Interv Aging*. 2016;11:471–479. doi:10.2147/CIA.S104083
64. Nau DP, Aikens JE, Pacholski AM. Effects of gender and depression on oral medication adherence in persons with type 2 diabetes mellitus. *Gen Med*. 2007;4(3):205–213.
65. O'Shea MP, Teeling M, Bennett K. Comorbidity, health-related quality of life and self-care in type 2 diabetes: a cross-sectional study in an outpatient population. *Iran J Med Sci*. 2015;184:623–630. doi:10.1007/s11845-014-1190-4
66. Schinckus L, Dangoisse F, Broucke S, Mikolajczak M. When knowing is not enough: emotional distress and depression reduce the positive effects of health literacy on diabetes self-management. *Patient Educ Couns*. 2018;101:324–330. doi:10.1016/j.pec.2017.08.006
67. Shin N, Hill-Briggs F, Langan S, Payne JL, Lyketsos C, Golden SH. The association of minor and major depression with health problem-solving and diabetes self-care activities in a clinic-based population of adults with type 2 diabetes mellitus. *J Diabetes Complications*. 2017;31:880–885. doi:10.1016/j.jdiacomp.2017.01.026
68. Smith KJ, Pedneault M, Schmitz N. Investigation of anxiety and depression symptom co-morbidity in a community sample with type 2 diabetes: associations with indicators of self-care. *Can J Public Health*. 2015;106(8):e496–e501. doi:10.17269/CJPH.106.5170
69. Zuberi SI, Syed EU, Bhatti JA. Association of depression with treatment outcomes in type 2 diabetes mellitus: a cross-sectional study from Karachi, Pakistan. *BMC Psychiatry*. 2011;11(27). Available from: <http://www.biomedcentral.com/1471-1244X/1411/1427>
70. Aikens JE. Prospective associations between emotional distress and poor outcomes in type 2 diabetes. *Diabetes Care*. 2012;35:2472–2478. doi:10.2337/dc12-0181
71. Axon RN, Gebregziabher M, Hunt KJ, et al. Comorbid depression is differentially associated with longitudinal medication nonadherence by race/ethnicity in patients with type 2 diabetes. *Medicine*. 2016;95(25):1–7. doi:10.1097/MD.00000000000003983
72. Gonzalez JS, Safren SA, Delahanty LM, et al. Symptoms of depression prospectively predict poorer self-care in patients with type 2 diabetes. *Diabetic Med*. 2008;25:1102–1107. doi:10.1111/j.1464-5491.2008.02535.x
73. He X, Chen L, Wnag K, Wu H, Wu J. Insulin adherence and persistence among chinese patients with type 2 diabetes: a retrospective database analysis. *Patient Prefer Adherence*. 2017;11:237–245. doi:10.2147/PPA.S123389
74. Kalsekar ID, Madhavan SS, Amonkar MM, et al. Depression in patients with type 2 diabetes: impact on adherence to oral hypoglycemic agents. *Ann Pharmacother*. 2006;40:605–611. doi:10.1345/aph.1G606
75. Kreyenbuhl J, Dixon LB, McCarthy JF, Suliman S, Ignacio RV, Valenstein M. Does adherence to medications for type 2 diabetes differ between individuals with vs without schizophrenia? *Schizophr Bull*. 2010;36(2):428–435. doi:10.1093/schbul/sbn106
76. O'Shea MP, Teeling M, Bennett K. An observational study examining the effect of comorbidity on the rates of persistence and adherence to newly initiated oral anti-hyperglycaemic agents. *Pharmacoepidemiol Drug Saf*. 2013;22:1336–1344. doi:10.1002/pds.3535
77. Palakodeti S, Uratsu CS, Schmittiel JA, Grant RW. Changes in physical activity among adults with diabetes: a longitudinal cohort study of inactive patients with type 2 diabetes who become physically active. *Diabetic Med*. 2015;32:1051–1057. doi:10.1111/dme.12748
78. Dixon JB, Browne JL, Mosely KG, et al. Severe obesity and diabetes self-care attitudes, behaviours and burden: implications for weight management from a matched case-controlled study. Results from diabetes MILES—Australia. *Diabetic Med*. 2014;31:232–240. doi:10.1111/dme.12306
79. Wagner JA, Tennen H, Osborn CY. Lifetime depression and diabetes self-management in women with type 2 diabetes: a case-control study. *Diabetic Med*. 2010;27:713–717. doi:10.1111/j.1464-5491.2010.02996.x
80. Kroese FM, Adriaanse MA, Ridder TDD. Are self-management interventions suitable for all? Comparing obese versus nonobese type 2 diabetes patients. *Health Educ Behav*. 2012;40(5):552–558. doi:10.1177/1090198112454285
81. Trief PM, Izquierdo R, Eimicke JP, et al. Adherence to diabetes self care for white, African-American and Hispanic American telemedicine participants: 5 year results from the IDEATel project. *Ethn Dis*. 2013;18(1):83–96. doi:10.1080/13557858.2012.700915
82. Bruce DG, Davis WA, Starkstein SE, Davis TME. Clinical risk factors for depressive syndrome in type 2 diabetes: the Fremantle Diabetes Study. *Diabetic Med*. 2018;35:903–910. doi:10.1111/dme.13631
83. Alagiakrishnan K, Zhao N, Mereu L, Senior P, Senthilselvan A. Montreal cognitive assessment is superior to standardized mini-mental status exam in detecting mild cognitive impairment in the middle-aged and elderly patients with type 2 diabetes mellitus. *Biomed Res Int*. 2013;2013:1–5. Article ID 186106. doi:10.1155/2013/186106
84. Mendenhall E, Norris SA, Shidhaye R, Prabhakaran D. Depression and type 2 diabetes in low and middle income countries: a systematic review. *Diabetes Res Clin Pract*. 2014;103(2):276–285. doi:10.1016/j.diabres.2014.01.001
85. Mendenhall E, Kohrt BA, Norris SA, Ndetel D, Prabhakaran D. Non-communicable disease syndemics: poverty, depression, and diabetes among low-income populations. *Lancet*. 2017;389:951–963. doi:10.1016/S0140-6736(17)30402-6

Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy**Dovepress****Publish your work in this journal**

Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy is an international, peer-reviewed open-access journal committed to the rapid publication of the latest laboratory and clinical findings in the fields of diabetes, metabolic syndrome and obesity research. Original research, review, case reports, hypothesis formation, expert opinion

and commentaries are all considered for publication. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/diabetes-metabolic-syndrome-and-obesity-targets-and-therapy-journal>