Influence of thoracic drainage fluid on proliferation, migration, apoptosis, and drug resistance in lung cancer cell lines

Yuqiang Mao¹Ying Yu²Yun Han¹

¹Department of Thoracic Surgery,
Shengjing Hospital of China Medical
University, Shenyang 110004, China;

²Liaoning Medical Device Test
Institute, Shenyang 110179, China

Background: This study aimed to clarify the effect of thoracic drainage fluid (DF) on lung cancer cells in vitro.

Methods: We assessed the influence of DF on the proliferation and migration of lung cancer cells (LTEP-a-2 and A549) using the MTT cell proliferation assay and scratch wound assay. Cell apoptosis was determined by flow cytometric analysis. We also investigated the effect of DF on drug chemosensitivity, assessing viability of LTEP-a-2 and A549 cells.

Results: The proliferative rates of cancer cells in the DF-treated group were significantly higher than those of the control group. Similar results were obtained for cell migration of lung cancer cells. Cells in the DF-treated groups showed a lower percentage of apoptosis than those of the control groups. Chemosensitivity of lung cancer cells to doxycycline and cisplatin (DDP) was lowered by DF.

Conclusion: These findings suggest that DF affects lung cancer cells by promoting proliferation and migration, inhibiting apoptosis, and increasing drug resistance.

Keywords: lung cancer, drainage fluid, proliferation and migration, apoptosis, drug resistance

Introduction

Lung cancer has the highest incidence and mortality rate among all cancers in China and worldwide.^{1,2} Currently, surgery is the most important and only method for curing lung cancer. With the improvement of diagnostic and treatment procedures, such as endoscopy, radical surgery, radiochemotherapy, targeted therapy, and other integrated treatments,^{3,4} more accurate staging and early diagnosis have significantly increased the survival of lung cancer patients.⁵⁻⁷ Also, advances in surgical techniques, such as thoracoscopy, have decreased surgical trauma and incidence of complications in patients, improving their quality of life.⁸⁻¹¹

In recent years, many studies on lung cancer surgery have been put forward. With the number of lung cancer patients in China increasing every year, treatment efficiency in large hospitals has become a medical and even a social problem, awaiting resolution. Against this background, the concept of rapid rehabilitation surgery was born. In recent years, this concept has been explored and applied to gastrointestinal, thoracic, and other surgeries, in accordance with clinical diagnosis and treatment models.¹²⁻¹⁴ Among these is the thoracic surgery domain of lung cancer surgery, where many thoracic surgeons have proposed that the chest tube can be removed earlier after radical lung cancer surgery, especially after thoracoscopic lobectomy, to achieve early postoperative discharge of patients, increase bed turnover, and reduce medical costs. There are many studies showing that early removal of chest tubes does not increase the incidence of complications in patients during the perioperative period, and a small number of

Correspondence: Yun Han

Department of Thoracic Surgery,
Shengjing Hospital of China Medical
University, No. 39 Huaxiang Road, Tiexi
District, Shenyang, Liaoning 110022,
China
Tel +86 24 133 5249 0777
Email hany_sjhosp@126.com

studies reported that eliminating the chest tube drainage of 500 mL/day also does not increase complications.¹⁵ However, these practices entail reabsorption of excess pleural effusion into the systemic circulation. As malignant tumors are systemic diseases, the presence of chemical substances in plural effusion, and even residual tumor cells, has potential effects on local or systemic recurrence, and development of drug resistance. These effects are still unknown.

Previous studies have found that the drainage fluid (DF) from breast cancer patients promoted proliferation and migration of breast cancer cells, and this effect showed some degree of dose-dependence. One study showed that surgery may have negative effects on malignant tumors.¹⁶ Based on this, we designed an experiment to examine the effects of thoracic DF obtained after lung cancer surgery on the biological behavior of lung cancer cells, including chemotherapeutic drug resistance. This provides experimental evidence of the biological characteristics of postoperative DF from various malignancies and may help guide the timing of postoperative tube removal and chemotherapy.

Results

Influence of DF on cell death of LTEP-a-2 and A549 cells

We evaluated the cell death of LTEP-a-2 and A549 cells by MTT assay, taking the untreated cells as reference. The results showed remarkably decreased cell death in the DF-treated group compared to the untreated cells (Figure 1). The increased survival rates of A549 cells in 0.1%, 0.5%, and 1.0% DF-treated groups were 111.00% ($\pm 10.09\%$), 115.08% ($\pm 11.15\%$), and 114.78% ($\pm 11.26\%$), respectively. These were higher than the untreated group and the differences were

statistically significant ($P=0.0036$, $P=0.010$, and $P=0.011$, respectively; Figure 1A and B). Similarly, the survival rates of LTEP-a-2 cells in 0.1%, 0.5%, and 1% DF-treated groups were 160.13% ($\pm 34.91\%$), 158.40% ($\pm 19.90\%$), and 161.31% ($\pm 17.33\%$), respectively, and these were also significantly higher than the untreated group (0.1%, 0.5%, and 1.0% DF, all $P<0.001$; Figure 2B).

Effect of DF on chemosensitivity

We tested the effect of DF on drug chemosensitivity, assessing the viabilities of LTEP-a-2 and A549 cells (Figure 2). In the doxycycline chemosensitivity group, the viability of DF-treated (0.1% and 1%) LTEP-a-2 cells was higher than that in the untreated groups for both drug concentration levels (5 and 10 mg/L) and the differences were statistically significant ($P=0.047$ and $P=0.007$, respectively, for 5 mg/L doxycycline; $P<0.001$ and $P=0.002$, respectively, for 50 mg/L doxycycline; Figure 2A). Our results with DDP treatment were similar to those obtained for the doxycycline group. Cell viability in DF-treated groups was significantly higher than that in the control group, with statistically significant differences ($P=0.039$ and $P=0.034$, respectively, for 5 mg/L DDP; $P=0.010$ and $P=0.041$, respectively, for 10 mg/L DDP; Figure 2B). We observed a similar effect of DF on the drug chemosensitivity of A549 cells (Figure 2C and D). In the doxycycline group, viability of A549 cells with 0.1% and 1% DF was significantly higher than that in the untreated groups ($P=0.021$ and $P=0.013$, respectively, for 5 mg/L doxycycline; $P=0.048$ and $P=0.002$, respectively, for 10 mg/L doxycycline, Figure 2C). We obtained similar results with DDP. Cell viability in DF-treated groups was significantly higher than in control groups ($P=0.046$ and $P=0.038$, respectively,

Figure 1 MTT proliferation assay: DF increases proliferation of LTEP-a-2 and A549 cells.

Notes: (A) Increases proliferation by DF in LTEP-a-2. (B) Increases proliferation by DF in A549. Proliferation of LTEP-a-2 and A549 cells was assessed by MTT proliferation assay, taking the untreated cells as reference. Proliferation was significantly increased in 0.1%, 0.5%, and 1% DF-treated groups compared to the control groups.

Abbreviation: DF, drainage fluid.

Figure 2 Chemosensitivity test: DF weakened the doxycycline and DDP chemosensitivity of lung cancer cells, taking the control groups as reference.

Notes: In the doxycycline chemosensitivity group, viability of LTEP-a-2 cell groups treated with DF (0.1% and 1%) were higher than those of the control groups at both drug concentration levels (5 and 50 mg/L), and the differences were statistically significant (**A**). Treatment with DDP produced results similar to those obtained with doxycycline (**B**). DF similarly weakened chemosensitivity of A549 cells (**C** and **D**).

Abbreviations: DDP, cisplatin; DF, drainage fluid; N, negative control.

for 5 mg/L DDP; $P=0.033$ and $P=0.047$, respectively, for 10 mg/L DDP; Figure 2D).

Flow cytometry analysis

We performed a flow cytometric analysis to determine the percentage of apoptotic cells in each individual group. Cells in the groups treated with 0.1% and 1% DF showed lower apoptosis percentages than those of the control groups (Figure 3).

Influence of DF on motility of lung cancer cells

The potential motilities of LTEP-a-2 and A549 cells were evaluated by the scratch wound assay. The results showed that DF tends to stimulate migration of lung cancer cells in both 0.1% and 1% DF groups compared with the control group (Figure 4).

Discussion

In this study, we showed that the effects of DF on lung cancer cells include promoted proliferation and migration, inhibited apoptosis, and increased drug resistance. These results were

consistent with the findings of previous studies,^{16–18} which have suggested that DF acts as a form of tumor microenvironment and that its constituent cytokines and other components act to promote malignant biological behavior of tumor cells. In the local primary tumor, these negative effects provide an environment conducive to recurrence of residual tumor cells, while excessive DF absorption and circulation increase the probability of systemic recurrence of the tumor.

In a situation where China's economy and national income are still not ideal, chemotherapy still remains an important adjuvant treatment for lung cancer as compared with targeted therapy,^{7,19} which has an indispensable function in the prevention of local and systemic recurrence. Our experimental results showed that DF not only promotes proliferation and inhibits apoptosis of lung cancer cells but also increases drug resistance. This suggests that the presence of local DF after surgery may increase recurrence and decrease the therapeutic efficacy of chemotherapy. Furthermore, DF was found to inhibit both early and late apoptosis in lung cancer cells, providing an environment conducive to the survival and recurrence of residual cancer cells. Therefore, there is still no definite conclusion as to whether chest tubes should be removed soon after surgery.

Figure 3 Flow cytometry analysis: LTEP-a-2 and A549 cells in the groups treated with 0.1% and 1% DF showed lower percentages of apoptosis than the control groups.

Notes: (A) Representational figures for apoptosis of LTEP-a-2 and A549 cells. (B) Statistical results for apoptosis of LTEP-a-2 cells. (C) Statistical results for apoptosis of A549 cells.

Abbreviations: DF, drainage fluid; NC, negative control; PI, propidium iodide.

Previous studies have reported the major components of DF,¹⁶ which include cytokines, chemokines, and matrix metalloproteinases. These are influenced by chemotherapy, TNM staging, pathological type, and molecular subtype. Among cytokines, the concentration of IL-6 is the highest, and previous studies have suggested that IL-6 can mediate tumor recurrence and metastasis through STAT3, MMP3, and other relevant pathways.^{20–23} Therefore, it is conceivable that several complex components of DF can mediate the malignant biological behavior of tumors through a variety of signaling pathways.

More in-depth studies to elucidate the specific underlying mechanisms of the biological effects of DF are warranted. The ability of cells to promote MTT reduction was only evaluated at one time point (48 hours). The MTT reduction is dependent not only on the number of cells but also on the metabolic performer of cells connected with their mitochondrial activity. The results can only improve the mitochondrial functionality or decreased number of cells that died after seeding without affecting the duplication time. Thus, additional assays are required to evaluate the effect of DF on the proliferation rate of cancer cells. The assessment of cell death

Figure 4 Scratch wound tests: DF promoted migration of LTEP-a-2 and A549 cells.

Notes: Scratch wounds for LTEP-a-2 and A549 cells at 0 and 48 hours after introducing the wound, and treatment with DF. Scratch fold changes in 0.1% and 1% DF-treated groups were significantly higher than in the control group. The scratch closure change area of cells treated with 1% DF tended to be larger than with 0.1% DF.

Abbreviation: DF, drainage fluid.

by apoptosis and the assessment of cell resistance to apoptotic drugs allow limited conclusions. Apoptosis involves the activation of caspases, which is a group of cysteine proteases. The inclusion of the effects on the activity of the initiator and effector caspases should also be considered.

In addition, primary tumor cells from diverse lung cancer patients need to be tested in order to further investigate the drug susceptibility and biological effects of DF. Samples from different types of lung cancers are necessary to understand the differences in chemotherapeutic susceptibility between different clinicopathological types. This research helps improve chemotherapeutic regimens for different lung cancer types and also provides guidance on the best time for chest tube removal. At the same time, the design of rigorous animal experiments and continuous follow-up on the tube removal status and survival of patients are necessary.

Materials and methods

Patients and collection of DF

We obtained 16 DF samples from 16 lung cancer patients who underwent lobectomy. All patients had no known underlying

diseases except lung neoplasm. Sixteen DF samples were used for the cell MTT proliferation test, and nine samples were used for the chemosensitivity test in vitro.

Supernatant from centrifuged DF samples was collected under sterile conditions and preserved at -80°C without additives. Written informed consent was obtained from the patients, and the study protocol was conducted in accordance with the Declaration of Helsinki and approved by the Ethics Committee of China Medical University.

Cell incubation

The DF specimens were sterilized and pretreated at 0°C before co-culture assays in vitro. Sixteen DF specimens were used to evaluate the influence of DF on lung cancer cell lines. LTEP-a-2 and A549 lung cancer cell lines were acquired from American Type Culture Collection (ATCC, Manassas, VA, USA). LTEP-a-2 and A549 cells were cultured in RPMI 1640 medium (ATCC) and in F-12K medium (ATCC) supplemented with 10% FBS (Sigma-Aldrich Co., St. Louis, MO, USA). Each experiment was repeated three times.

Cell death test and in vitro chemosensitivity test

We tested the survival potential and drug chemosensitivity of LTEP-a-2 and A549 cells by MTT assay. LTEP-a-2 and A549 cells (5×10^4 /mL for the proliferation test and 10×10^4 /mL for the in vitro chemosensitivity test) were seeded onto 96-well plates in 100 μ L complete medium. Lung cells were incubated for 24 hours at 37°C, after which we replaced the medium with new complete medium for both groups. At this point, DF (0.1%, 0.5%, and 1%) samples were added to the treated groups' medium. Some cells were used for in vitro chemosensitivity tests, which included two concentration levels (5 and 10 mg/L) of doxycycline or DDP. The remaining cells were used for proliferation tests. After 48 hours of incubation at 37°C, 10 μ L MTT dye was added to each well and co-cultured with cells for 4 hours. The MTT solution was then removed. Formazan crystals were dissolved by adding 200 μ L dimethyl sulfoxide to each well. Absorbance was measured at 570 nm with a microplate reader (Titertek Multiskan PLUS, MK II; Labsystems, McLean, VA, USA). Each experiment was repeated three times.

Flow cytometry analysis

LTEP-a-2 and A549 cells (1×10^5 cells/well) were seed into 24-well plates and incubated in complete medium overnight. Culture medium was then replaced with new complete medium for both groups, while DF (0.1% and 1%) samples were added to the treated groups' medium at 37°C for 24 hours. The cells were then washed with PBS and sloughing cells were removed. The recombinant annexin V conjugated to green-fluorescent FITC dye was used to detect the externalization of phosphatidylserine, as a sign of apoptotic cells. The remaining cells were stained with anti-FITC and anti-PI, and the percentages of apoptotic cells in each individual group were determined using flow cytometry analysis. Each experiment was repeated three times.

Scratch wound test

LTEP-a-2 and A549 cells with 90% confluence were plated uniformly in complete medium in 24-well plates. Scratched cells were located at the bottom of the wells after 24 hours of incubation. Cell suspensions were freshly prepared for visualization in PBS. Initial scratch wounds (0 hour) were imaged at once. The medium was replaced, while DF (0.1% and 1%) was added to the treated groups' medium. Scratch wounds after 48 hours incubation with DF were visualized under a

microscope (Leica DM4000B; Leica Microsystems, Wetzlar, Germany). Each experiment was repeated three times.

Statistical analysis

We used IBM SPSS statistical version 22.0 software (IBM Corporation, Armonk, NY, USA). We made statistical figures using PRISM (version 5; GraphPad Software, Inc., La Jolla, CA, USA). The differences among groups were compared by analysis of variance, followed by a Student Newman Keuls post hoc test. *P*-values <0.05 were considered statistically significant.

Acknowledgment

This study was funded by the Liaoning Natural Science Foundation (No. 20180530020 and 20170520024).

Disclosure

The authors report no conflicts of interest in this work.

References

1. Chen W, Zheng R, Baade PD, et al. Cancer statistics in China, 2015. *CA Cancer J Clin*. 2016;66(2):115–132.
2. Torre LA, Bray F, Siegel RL, Ferlay J, Lortet-Tieulent J, Jemal A. Global Cancer statistics, 2012. *CA Cancer J Clin*. 2015;65(2):87–108.
3. Cetinkaya E, Usluer O, Yilmaz A, et al. Is endobronchial ultrasound-guided transbronchial needle aspiration an effective diagnostic procedure in restaging of non-small cell lung cancer patients? *Endosc Ultrasound*. 2017;6(3):162–167.
4. Chinese Thoracic Society, Chinese Alliance against Lung Cancer. Chinese expert consensus statement on issues related to small specimen sampling of lung cancer. *Endosc Ultrasound*. 2017;6(4):219–230.
5. Chaudhary T, Chahar A, Sharma JK, Kaur K, Dang A. Phytomedicine in the treatment of cancer: a health technology assessment. *J Clin Diagn Res*. 2015;9(12):XC04–XC09.
6. Castro-Pocas FM, Araújo TP, Ferreira ml, Saraiva MM. The role of endoscopic ultrasound in a case of lung cancer with jaundice. *Endosc Ultrasound*. 2016;7(4):279–281.
7. Burdett S, Rydzewska LHM, Tierney JF; NSCLC Meta-analysis Collaborative Group. Preoperative chemotherapy for non-small-cell lung cancer: a systematic review and meta-analysis of individual participant data. *Lancet*. 2014;383(9928):1561–1571.
8. Gonzalez-Rivas D. Uniportal thoracoscopic surgery: from medical thoracoscopy to non-intubated uniportal video-assisted major pulmonary resections. *Ann Cardiothorac Surg*. 2016;5(2):85–91.
9. Harris K, Modi K, Kumar A, Dhillon SS. Endobronchial ultrasound-guided transbronchial needle aspiration of pulmonary artery tumors: a systematic review (with video). *Endosc Ultrasound*. 2015;4(3):191–197.
10. Aswanetmanee P, Limswat C, Kabach M, Alraiyes AH, Kheir F. The role of sedation in endobronchial ultrasound-guided transbronchial needle aspiration: systematic review. *Endosc Ultrasound*. 2016;5(5):300–306.
11. Dziedzic D, Peryt A, Szolkowska M, Langfort R, Orlowski T. Evaluation of the diagnostic utility of endobronchial ultrasound-guided transbronchial needle aspiration for metastatic mediastinal tumors. *Endosc Ultrasound*. 2016;5(3):173–177.
12. Kehlet H. Fast-track colorectal surgery. *Lancet*. 2008;371(9615):791–793.

13. Muller S, Zalunardo MP, Hubner M, Clavien PA, Demartines N; Zurich Fast Track Study Group. A fast-track program reduces complications and length of hospital stay after open colonic surgery. *Gastroenterology*. 2009;136(3):842–847.
14. Dong Q, Zhang K, Cao S, Cui J. Fast-track surgery versus conventional perioperative management of lung cancer-associated pneumonectomy: a randomized controlled clinical trial. *World J Surg Oncol*. 2017; 15(1):20.
15. Cerfolio RJ, Bryant AS. Results of a prospective algorithm to remove chest tubes after pulmonary resection with high output. *J Thorac Cardiovasc Surg*. 2008;135(2):269–273.
16. Wang D, Hu K, Gao N, et al. High throughput screening of cytokines, chemokines and matrix metalloproteinases in wound fluid induced by mammary surgery. *Oncotarget*. 2015;6(30):29296–29310.
17. Valeta-Magara A, Hatami R, Axelrod D, et al. Pro-oncogenic cytokines and growth factors are differentially expressed in the post-surgical wound fluid from malignant compared to benign breast lesions. *Springerplus*. 2015;4:483.
18. Zhang Y, Yan D, Zhang H, et al. Influence of wound fluid on chemotherapy sensitivity in primary breast cancer cells. *Oncotarget*. 2016;7(40):65034–65041.
19. Barton MK. Patients of all ages with advanced non-small cell lung cancer are not receiving chemotherapy. *CA Cancer J Clin*. 2015;65(5):337–338.
20. Lesina M, Kurkowski MU, Ludes K, et al. STAT3/SOCS3 activation by IL-6 transsignaling promotes progression of pancreatic intraepithelial neoplasia and development of pancreatic cancer. *Cancer Cell*. 2011;19(4):456–469.
21. Grivennikov S, Karin E, Terzic J, et al. IL-6 and STAT3 are required for survival of intestinal epithelial cells and development of colitis-associated cancer. *Cancer Cell*. 2009;15(2):103–113.
22. Cutler SJ, Doecke JD, Ghazawi I, et al. Novel STAT binding elements mediate IL-6 regulation of MMP-1 and MMP-3. *Sci Rep*. 2017;7(1):8526.
23. He G, Dhar D, Nakagawa H, et al. Identification of liver cancer progenitors whose malignant progression depends on autocrine IL-6 signaling. *Cell*. 2013;155(2):384–396.

Cancer Management and Research

Publish your work in this journal

Cancer Management and Research is an international, peer-reviewed open access journal focusing on cancer research and the optimal use of preventative and integrated treatment interventions to achieve improved outcomes, enhanced survival and quality of life for the cancer patient. The manuscript management system is completely online and includes

Submit your manuscript here: <https://www.dovepress.com/cancer-management-and-research-journal>

Dovepress

a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.