Botulinum toxin (Botox) A for reducing the appearance of facial wrinkles: a literature review of clinical use and pharmacological aspect

This article was published in the following Dove Press journal:
Clinical, Cosmetic and Investigational Dermatology

Bagus Komang Satriyasa

Department of Pharmacology, Faculty of
Medicine, Udayana University, Bali,
Indonesia

Background: Botulinum toxin (Botox) consists of 7 types of neurotoxins; however, only toxins A and B are used clinically. Botox A is used for several disorders in the field of medicine, particularly in dermatology, for cosmetic purposes. It is produced by the bacterium *Clostridium botulinum* and can be used as a treatment to reduce the appearance of wrinkles in the upper areas of the face, elevate the eyebrows and treat problems such as hyperhidrosis, lichen simplex, pompholyx (dyshidrotic eczema) and acne vulgaris.

Objectives: This article provides a literature review regarding the general issue of Botox as a treatment for reducing facial wrinkle.

Discussion: Botox works by blocking the release of acetylcholine, resulting in paralysis of the local muscles, which usually occurs 24 hrs to two weeks following Botox injection. This effect will last three to six months. The optimal dose of cosmetic Botox in dermatology is 20 units. Botox is relatively safe and does not result in any adverse side effects. However, in certain circumstances, the effect of Botox will gradually resolve, resulting in reduced muscle paralysis over time.

Conclusion: Botox is good and safe medicine to reduce the appearance of facial wrinkles.

Keywords: Botox, botulinum toxin, drug, facial wrinkles

Introduction

Botulinum toxin (Botox) is a drug made from a toxin produced by the bacterium *Clostridium botulinum*. In large amounts, this toxin can cause botulism, an illness that affects the nerves. Botox has been used since the 1970s in the field of ophthalmology, and in the last 20 years, its use has expanded to various health scopes, especially dermatology.^{1,2}

Botox consists of 7 types of neurotoxins; however, only toxins A and B are used clinically. Botox A is used for several disorders in the field of medicine, particularly in dermatology, for cosmetic purposes.³ The first type of Botox introduced to the market was onabotulinum toxin A. In 2002, it was recommended to be used as a cosmetic treatment for glabellar frown lines by the Food and Drug Administration (FDA).^{2,4,5} The second formulation of onabotulinum toxin A, which was produced in France, obtained its license to be used for esthetic purposes from the European Union in 2006 and was approved by the FDA in 2009.^{5,6} Botox type A has become a term used by the society to describe all ingredients used in cosmetic treatments.⁷

A study in 1994 reported the effectiveness of Botox A for reducing the appearance of facial wrinkles; since then, it has been used as a cosmetic treatment.⁸ Botox injections can be used to treat glabellar frown lines, wrinkles

Correspondence: Bagus Komang Satriyasa
Department of Pharmacology, Faculty of
Medicine, Udayana University, Bali,
Indonesia
Email bksatriyasa@unud.ac.id, 80223

around the lips (smoker's lines) and marionette lines, platysmal bands in the neck, strabismus, blepharospasm, cervical dystonia, hyperhidrosis as well as synkinesis following facial surgery.^{1,2,9}

The objective of this article is providing a good literature review regarding Botox as a treatment for reducing facial wrinkles and also comparing it with other modalities such as fillers, skin peptide and facial muscle exercise. The main focuses are the mechanism of action, indication, contraindication, dosage, clinical effect, safety, side effect, and complication.

Mechanism of action

The mechanism of action of Botox includes the following four main steps:

1. The first step is binding of the toxin to specific receptors on the surface of the presynaptic cells, mediated by the C-terminal of the heavy chain. This step occurs over approximately 30 mins.¹⁰
2. The second step is internalization, which is an energy-dependent receptor-mediated endocytic process. In this step, the plasma membrane of nerve cells invaginates around the toxin-receptor complex, forming a vesicle containing a toxin in nerve terminal.¹⁰
3. The third step is translocation. Following internalization, the disulfide bond is cleaved and the 50-kDa light chain of the toxin is released across the endosomal membrane of the endocytic vesicle into the cytoplasm of the nerve terminal.¹⁰
4. The last step is blocking. The light chain of serotypes A and E inhibit the release of acetylcholine by cleaving the cytoplasmic protein (SNAP-25) needed for the docking of acetylcholine vesicles on the inner side of the nerve membrane of the nerve terminal.¹⁰

Following the injection, the toxin diffuses into the tissue until it binds selectively and reversibly in the presynaptic terminal of the neuromuscular junction and then attaches to the specific protein-membrane responsible for acetylcholine excretion.^{5,6} The toxin immediately inhibits the release of acetylcholine in the neuromuscular junction causing relaxation of local muscles that is reversible, resulting in reduced facial wrinkles/lines, of which some are due to constant facial muscle contractions.^{11,12}

Indication

In the field of dermatology, Botox is generally injected into the muscles of facial expression.² Most of these muscles are attached to soft tissues rather than bones, and by contracting, they pull across the skin to give facial expressions.^{2,13} In esthetics, Botox is used for reducing glabellar frown lines, crow's feet at the side of the eyes, horizontal forehead creases, wrinkles around the mouth, nasolabial folds and smoothing out neck and chest/cleavage wrinkles.^{2,14} It also can be used to elevate the eyebrows and treat problems such as hyperhidrosis, lichen simplex, pompholyx (dyshidrotic eczema) and acne vulgaris.^{7,15} Botox cannot be used to prevent other signs of aging such as dry skin, pigmentation disorders and vascular abnormalities.¹⁶

Contraindication

Contraindications with the use of Botox include patients with myasthenia gravis, amyotrophic lateral sclerosis, multiple sclerosis, Eaton Lambert syndrome, women who are pregnant and breastfeeding, neonate and children, patients with focal and systemic infections, patients who are hypersensitive or allergic to Botox and patients who had previously undergone lower eyelid surgery.^{17–20}

Dosage

The standard dose for Botox usage is 20 Units. The minimum dose for treating glabellar frown lines is 20 Units as it was found that dosing 20–40 Units of Botox was more effective than 10 Units alone for reducing glabellar frown lines.¹⁶ For male patients, Botox A dosage is effective when started from 40 Units.²¹ Men require higher doses of Botox as they have a greater muscle mass than women.^{22,23}

Clinical effects

The clinical effects of Botox are seen on the first to the fourth days after injection, followed by 1–4 weeks of maximum effect, which will resolve after 3–4 months. In order to prolong the effects of Botox from six months to one year, the treatment should be repeated for one year or more.²⁴ The duration of Botox effect varies among individuals due to differences in muscle arrangements, meaning that different individuals may require different doses of Botox. The effect will last up to 120 days.^{25,26}

Safety

Botox is a drug with a broad margin of safety (lethal dose 50% (LD50) in humans can reach up to 40 U/kg BW).

Therefore, its use in cosmetics is relatively safe. Botox is relatively safe and effective for treating facial wrinkles.^{27,28} Botox A does not cause persistent changes at the nerve terminals and targeted muscles. In general, it does not cause any long-term adverse or side effects in the field of dermatology.^{29,30}

Side effects

The possible side effects of Botox include bleeding, swelling, erythema and pain at the injection sites.³¹ These side effects can be avoided by using thinner needles and diluting Botox with saline. Headaches may also occur following Botox injections but will resolve after 2–4 weeks. This side effect can be treated using systemic analgesics.^{27,28} Other side effects that have also been reported to occur include malaise, nausea, influenza-like symptoms and ptosis.³¹ Ptosis usually occurs in patients who use Botox to treat the glabellar area and is caused by local diffusion of Botox, which can remain for several weeks but can be treated with alpha-adrenergic agonist ophthalmic drops. Ectropion may also develop due to the local diffusion process of Botox when injected into the lower eyelids. Also, patients who undergo Botox injections to treat crow's feet or bunny lines (periorbital) may experience strabismus caused by inadvertent injection of Botox and local diffusion of Botox.^{31,32} Nevertheless, all of these side effects will gradually resolve after the paralytic effect of the toxin dissipates.^{33,34}

Complications

Complications with cosmetic Botox injections rarely occur. The most common complications are ecchymosis and purpura, which could be minimized by compressing ice on the injection sites before and after Botox injection.^{27,28} Botox should be injected in minimal concentrations, with the appropriate dose and injected at least 1 cm from the superior, inferior or lateral margin of the orbital bone. Following treatment, patients should not manipulate the injected sites for 2–3 hrs and should remain in an upright seating or standing position for 3–4 hrs.³⁵

Botox derivatives-acetyl hexapeptide-8

Acetyl hexapeptide-8 or acetyl hexapeptide-3 which is more known by the name Argireline® is the most peptide used to treat wrinkles on the skin as an alternative to botox therapy.³⁶ Acetyl hexapeptide-3 is very similar to the N-terminal end of SNAP-25 protein, competing with the

N-ethylmaleimide-sensitive factor attachment protein receptor complex which is an important protein in muscle contraction and the occurrence of wrinkles on the face.³⁷ When compared with Botox, acetyl hexapeptide-8 has a toxic potential 4,000 times lower and can be applied topically in the form of skin creams or via injections.^{35,36}

An animal study conducted by Wang et al³⁸ regarding the efficacy of Argireline for forty 42-month-old aging mice induced by skin injection using D-galactose showed that rats given topical Argireline 10% in oil-and-water emulsions twice a day in six weeks had lower type I collagen compared with the control group (69.82 ± 4.38 Vs 73.53 ± 8.21 ; $p < 0.01$) and higher amount of collagen type III compared to the control group (19.71 ± 7.52 Vs 18.43 ± 5.66 ; $p < 0.05$).

Another study conducted by Raikou et al³⁹ on the use of 10% Argireline topical twice daily on 24 facial skins of women aged 30–60 years for 60 days, found a significant decrease in transepidermal water loss (TEWL) in the Argireline group compared to the placebo group. Water content from the skin is very dependent on TEWL and the hydration level of the epidermis, which contributes to the health of skin and young skin appearance. The decrease in TEWL that occurred in the study was thought to be due to the use of topical Argireline which caused an increase in moisturization of the stratum corneum. There were no side effects observed in this study such as flushing, stinging, burning, redness, or desquamation of the facial skin.

The explanation provides a new solution that acetyl hexapeptide-8 can be an alternative therapy of botox for person who does not want to use the injection method because it can be used as a topical agent. Also which has a better level of safety compared to botox.

Comparison of Botox with dermal fillers

The most widely used types of dermal fillers are those made from hyaluronic acid, which is a glycosaminoglycan unit, which is composed of replicated glucuronic acid and N-acetyl-glucosamine units.⁴⁰ Hyaluronic acid is naturally a polysaccharide found in body tissues such as the skin and cartilage. It is very hydrophilic (attracts water), which causes pressure swelling which can withstand compressive forces that make hyaluronic acid the ideal substance as dermal fillers.⁴¹

The contraindications to use are quite similar to botox such as hypersensitivity to the components of fillers, bleeding disorders, history of anaphylaxis because hyaluronic acid

products are made from bacterial fermentation, the history of allergies in gram-positive bacteria is contraindicated.⁴²

Complications that are very likely to occur in the use of fillers because they are done by injection are bruising and hematoma. The use of filler is very dependent on operator expertise so that there are several side effects associated with injecting techniques such as unrealistic patient expectation, undercorrection, overcorrection and lumping.⁴³

Lumping is something that is very often caused by the placement of non-homogeneous filler material and lack of massage in the injection area. The filler when injected will fill a narrow space in the skin by doing massage on the injection area, the filler will be compacted and then mixed to form a closer and smoother texture. Failure in this process can cause irregularities of the skin that can be seen and touched (lumping).⁴⁴

A serious complication that can occur in the use of fillers is focal necrosis, although it is very rare in 0.001% of cases.^{41,42} The glabella area is the most common place for focal necrosis in filler use. Early signs of focal tissue necrosis usually appear very quickly within 24–48 hrs.^{43,44} This is caused by the occlusion caused by the direct effect of intravascular injection. This can be addressed if the clinician is aware of impending necrosis sign, and the first antidote treatment is injection of hyaluronidase 10–30 U per 2-cm² skin area.⁴⁴

Both treatment, botox and fillers are very operator-dependent and have very similar side effects. Both of these therapies have antidotes in dealing with the conditions of side effects that occur.

Comparison of Botox with facial muscle exercise for facial rejuvenation

Botox has an anti-wrinkle effect through its mechanism of action on the neuromuscular junction which causes relaxation of facial muscles thereby reducing facial wrinkles. But facial exercise therapy provides a different perspective by exercising, strengthening, moving or manipulating the facial muscles.^{45,46} This approach is a trend because it is not invasive, cheap and the services are mostly carried out by non-medical professions. The effects of anti-wrinkle obtained by the facial exercise method are still a debate, but it is believed that the effects obtained are a result of increased muscle tone, the increase in muscle volume obtained from facial exercises will tighten the skin which causes a reduction

of facial wrinkles. Moreover, the skin will experience the benefits of facial exercise due to improvements in waste product drainage and improvement of tissue regeneration as a result of improved blood flow and facial lymphatics circulation.^{47,48}

A study conducted by De Vos et al⁴⁹ involving 18 Caucasian women with an average age of 40 years. The samples were divided into two groups (interventions and controls). The intervention group was given facial exercise for seven weeks and then evaluated the condition of the face (judged by taking photographic of the faces) on conditions before and after the intervention. The assessment was carried out using a visual analog scale (VAS), which is a 100 mm line score with the “young” and “old” panels on the right and left ends. The facial exercise intervention has given focuses on 5 points, (1) exercise to reduce horizontal wrinkles on the forehead by exercising frontalis muscle, (2) exercise to reduce vertical wrinkles above the lips by training the superior part of the orbicularis oris muscle, (3) exercise to reduce nasolabial folds by training the orbicularis oris muscle and minor zygomaticum muscle, (4) exercise to reduce double chin by training masseter, sternocleidomastoid, and mylohyoid muscle. Assessment is done by comparing the five areas targeted in the intervention. It was found that the VAS value was significantly younger in the forehead and nasolabial folds in the post-intervention condition, but there was no improvement in the upper lip area, jaw line and under the chin area, and this improvement only occurred in the intervention group and not in the control group.

Facial exercise may provide a new alternative in reducing facial wrinkles, does not require invasive procedures and can be done without medical profession expertise. However, because there are still very limited clinical trials that compare the effects of facial exercise and the mechanism of action in reducing wrinkles that are still vague, this therapy has not been widely used compared to botox and dermal fillers.

Conclusion

Based on this review, it can be concluded that Botox is a good safe medication in reducing facial wrinkles. There are several issues regarding side effect and complication following the injection. However, there are several techniques to reduce side effect and complication rate after the injection.

Acknowledgments

The current review did not receive any specific grant from the government or any private sector.

Disclosure

The author reports no conflicts of interest in this work.

References

- Trindade de Almeida AR, Secco LC, Carruthers A. Handling botulinum toxins: an updated literature review. *Dermatol Surg*. 2011;37(11):1553–1565.
- Awan KH. The therapeutic usage of botulinum toxin (Botox) in non-cosmetic head and neck conditions – an evidence based review. *Saudi Pharm J*. 2017;25(1):18–24. doi:10.1016/j.jsps.2016.04.024
- Rohrer TE, Beer K. Background to botulinum toxin. In: Carruthers A, Carruthers J, editors. *Botulinum Toxin*. USA: Elsevier Inc.; 2005:9–18.
- Carruthers A, Carruthers J. Botulinum toxin. In: Bologna JL, Jorizzo JL, Rapini RP, editors. *Dermatology*. 2nd ed. New York: Mosby Elsevier; 2008:2381–2390.
- Glogau RG. Botulinum toxin. In: Wolf K, Goldsmith LA, Kartz SI, Gilchrist BA, Paller AS, Leffell DJ, editors. *Fitzpatrick's Dermatology in General Medicine*. 7th ed. New York: McGraw Hill; 2008:2389–2395.
- Flynn TC. Advances in the use of botulinum neurotoxins in facial esthetics. *J Cosmet Dermatol*. 2012;11(1):42–50. doi:10.1111/j.1473-2165.2011.00593.x
- Klein AW. Complications with the use of botulinum toxin. *Dermatol Clin*. 2004;22(2):197–205.
- Markey AC. Botulinum A exotoxin in cosmetic dermatology. *Clin Exp Dermatol*. 2000;25(3):173–175.
- Rzany B, Zielke H. Overview of botulinum toxin. In: de Maio M, Rzany B, editors. *Botulinum Toxin in Aesthetic Medicine*. New York: Springer-Verlag Berlin Heidelberg; 2007:1–10.
- Wollina U, Konrad H, Petersen S. Botulinum toxin in dermatology—beyond wrinkles and sweat. *J Cosmet Dermatol*. 2005;4(4):223–227. doi:10.1111/j.1473-2165.2005.00195.x
- Khawaja HA, Hernandez-Perez E. Botox in dermatology. *Int J Dermatol*. 2001;40(5):311–317.
- Lowe NJ. Minimally invasive treatments and procedures for aging skin. In: Burns T, Breathnach S, Cox N, Griffiths C, editors. *Rook's Textbook of Dermatology*. 8th ed. London: Blackwell Publishing; 2010:80.1–80.14.
- Joel LC, Freeman SR. Botulinum toxins. In: Draelos ZD, editor. *Cosmetic Dermatology Products and Procedures*. Oxford: Blackwell Publishing; 2010:342–351.
- Rasaldo H, Niforos F-R, Gassia V, et al.; Consensus Group. Lower-face and neck antiaging treatment and prevention using onabotulinumtoxin A: the 2010 multidisciplinary French consensus—part 2. *J Cosmet Dermatol*. 2011;10(2):131–149. doi:10.1111/j.1473-2165.2011.00560.x
- Michael AC, Kane MD. Botox injections for lower facial rejuvenation. *Oral Maxillofac Surg Clin North Am*. 2005;17(1):41–49. doi:10.1016/j.coms.2004.09.003
- Cohen JL, Dayan SH, Cox SE, Yalamanchili R, Tardie G. Onabotulinumtoxin A dose-ranging study for hyperdynamic perioral lines. *Dermatol Surg*. 2012;38(9):1497–1505. doi:10.1111/j.1524-4725.2012.02456.x
- Rzany B. Requirements and rules. In: Philipp M, editor. *Botulinum Toxin in Aesthetic Medicine*. Berlin: Springer; 2007:21–24.
- Gart MS, Gutowski KA. Overview of botulinum toxins for aesthetic uses. *Clin Plast Surg*. 2016;43(3):459–471. doi:10.1016/j.cps.2016.03.003
- Cula GO, Bargo PR, Nkengne A, Kollias N. Assessing facial wrinkles: automatic detection and quantification. *Skin Res Technol*. 2013;19(1):e243–e251. doi:10.1111/j.1600-0846.2012.00635.x
- Simon C, Resto V, Quinn FB Jr, Quinn MS. *Botulinum Toxin for Cosmetic Use*. USA: UTMB Health ; 2010.
- Bauman L. Botulinum toxin. In: Baumann L, Elsaie ML, Grunebaum L, editors. *Cosmetic Dermatology*. 2nd ed. New York: McGraw Hill; 2009:169–190.
- de Maio M, Rzany B. Injection technique. In: Rzany B, editor. *Botulinum Toxin in Aesthetic Medicine*. New York: Springer-Verlag Berlin Heidelberg; 2007:25–26.
- Carruthers JA, Lowe NJ, Menter MA, et al. A multicenter, double-blind, randomized, placebo-controlled study of the efficacy and safety of botulinum toxin type A in the treatment of glabellar lines. *J Am Acad Dermatol*. 2002;46(6):840–849.
- Krishtul A, Waldorf HA, Blitzer A. Complications of cosmetic botulinum toxin therapy. In: Carruthers A, Carruthers J, editors. *Botulinum toxin*. USA: Elsevier Inc.; 2005:12132.
- Klein AW. Contraindications and complications with the use of botulinum toxin. *Clin Dermatol*. 2004;22(1):66–75. doi:10.1016/j.clindermatol.2003.12.026
- Carruthers A, Carruthers J. Upper face treatment. In: Carruthers A, Carruthers J, editors. *Botulinum Toxin*. 1st ed. Philadelphia: Elsevier Saunders; 2005:31–43.
- Ascher B, Talarico S, Cassuto D, et al. International consensus recommendations on the aesthetic usage of botulinum toxin type A (Speywood Unit)—part II: wrinkles on the middle and lower face, neck and chest. *J Eur Acad Dermatol Vene reol*. 2010;24(11):1285–1295. doi:10.1111/j.1468-3083.2010.03728.x
- Maio M, Rzany B. Patient selection. In: Acquilla R, Patel T, Gonzales M, editors. *Botulinum Toxin in Aesthetic Medicine*. New York: Springer; 2007:11–19.
- Carruthers A, Carruthers J, Said S. Dose-ranging study of botulinum toxin type A in the treatment of glabellar rhytids in females. *Dermatol Surg*. 2005;31(4):414–422.
- Hexsel C, Hexsel D, Porto MD, Schilling J, Siega C. Botulinum toxin type A for aging face and aesthetic uses. *Dermatol Ther*. 2011;24(1):54–61. doi:10.1111/j.1529-8019.2010.01378.x
- Cohen JL, Freeman SR. Botulinum toxins. In: Draelos ZD, editor. *Cosmetic Dermatology Products & Procedures*. United Kingdom: Wiley-Blackwell Publishing Ltd; 2010.
- Cox SE, Adigun CG. Complications of injectable fillers and neurotoxins. *Dermatol Ther*. 2011;24(6):524–536. doi:10.1111/j.1529-8019.2012.01455.x
- Klein AW. Complications, adverse reactions, and insights with the use of botulinum toxin. *Dermatol Surg*. 2003;29(5):549–556.
- Ogden S, Griffiths TW. A review of minimally invasive cosmetic procedures. *Br J Dermatol*. 2008;159(5):1036–1050. doi:10.1111/j.1365-2133.2008.08845.x
- Rzany B, Zielke H. Safety of botulinum toxin in aesthetic medicine. In: de Maio M, Rzany B, editors. *Botulinum Toxin in Aesthetic Medicine*. New York: Springer-Verlag Berlin Heidelberg; 2007:119–125.
- Wang Y, Wang M, Xiao S, Pan P, Li P, Huo J. The anti-wrinkle efficacy of argireline, a synthetic hexapeptide, in Chinese subjects: a randomized, placebo-controlled study. *Am J Clin Dermatol*. 2013;14:147–153. doi:10.1007/s40257-013-0009-9
- Lim SH, Sun Y, Thiruvallur TM, Rosa V, Kang L. Enhanced skin permeation of anti-wrinkle peptides via molecular modification. *Sci Rep*. 2018;8:1596. doi:10.1038/s41598-017-18454-z
- Wang Y, Wang M, Xiao XS, Huo J, Zhang WD. The anti-wrinkle efficacy of Argireline. *J Cosmet Laser Ther*. 2013;15:237–241. doi:10.3109/14764172.2013.769273
- Raikou V, Varvaresou A, Panderi I, Papageorgiou E. The efficacy study of the combination of tripeptide-10-citrulline and acetyl hexapeptide-3. A prospective, randomized controlled study. *J Clin Dermatol*. 2017;15:271–278.
- Dayen SH. Complication from toxins and fillers in the dermatology clinic, recognition, prevention, and treatment. *Facial Plast Surg Clin N Am*. 2013;21:663–673. doi:10.1016/j.fsc.2013.07.008

41. Alam M, Tung R. Injection technique in neurotoxins and fillers: indications, products, and outcomes. *J Am Acad Dermatol*. 2018;79(3):423–435. doi:10.1016/j.jaad.2018.01.037
42. Hassouneh B, Newman JP. Laser, fillers, and neurotoxins avoiding complication in the cosmetic facial practice. *Facial Plast Surg Clin N Am*. 2013;21:585–598. doi:10.1016/j.fsc.2013.07.002
43. Bellomo R, Harrington LK. Neurotoxins and dermal fillers: choosing the right product. *Physician Assist Clin*. 2016;1:333–345. doi:10.1016/j.cpha.2015.12.008
44. Niamtu J. Complications in fillers and Botox. *Oral Maxillofacial Surg Clin N Am*. 2009;21:13–21. doi:10.1016/j.coms.2008.11.001
45. Kim K, Jeon S, Kim J-K, Hwang JS. Effects of Kyunghee Facial Resistance Program (KFRP) on mechanical and elastic properties of skin. *J Dermatol Treat*. 2016;27(2):191–196. doi:10.3109/09546634.2015.1056078
46. Ezure T, Hosoi J, Amano S, Tsuchiya T. Sagging of the cheek is related to skin elasticity, fat mass and mimetic muscle function. *Skin Res Technol*. 2009;15:229–305. doi:10.1111/j.1600-0846.2009.00364.x
47. Castillo-Garzon MJ, Ruiz JR, Ortega FB, Gutierrez A. Anti-aging therapy through fitness enhancement. *Clin Interv Aging*. 2006;1:213–220.
48. Clark HM, O'Brien K, Callej A, Corrie SN. Effects of directional exercise on lingual strength. *J Speech Lang Hear Res*. 2009;52:1034–1047. doi:10.1044/1092-4388(2009/08-0062)
49. De Vos M-C, Den Brande HV, Boone B, Borsel JV. Facial exercises for facial rejuvenation: a control group study. *Folia Phoniatr Logop*. 2013;65:117–122. doi:10.1159/000354083

Clinical, Cosmetic and Investigational Dermatology

Dovepress

Publish your work in this journal

Clinical, Cosmetic and Investigational Dermatology is an international, peer-reviewed, open access, online journal that focuses on the latest clinical and experimental research in all aspects of skin disease and cosmetic interventions. This journal is indexed on CAS.

The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/clinical-cosmetic-and-investigational-dermatology-journal>